

✓
June 10, 1949.

Searched Rio Tuas 1 mile above Petaca. Found
Arroyo No Agua east of Tres Piedros dry.
Went back to below to look for spawning
but there was none. Returned then to upper
Vallecitos creek where we angled for a short while
and caught 1 rainbow ~~each~~.

WJK49-06-10-02
Upper Vallecitos

June 10, 1949

WJK49-06-10-01

Locality N. M., RIO ARRIBA CO.

Tusas Cr. 1 mi. above Petaca.

Elev. 7100 Date June 10, 1949 Collector WJK, K. R. Coburn

Size 10-30 ft wide 4 (-6?) ft deep mostly 15 ft wide x 4 in.

Temp. Air 60 Water 53 Time 4:00 Weather cloudy

Water W S T Current m-str Bottom r in rf, sd in P.

Vegetation none

Shore willow, cult.

Watershed PJ

Capture Method 6 & 10' S., angling 1/2 h. Time 8:15-11:15

Dist. from shore S-S

Depth -3 ft. Orig. Pres. 10% F

Pantosteus plebeius

Yg. scattered, ads. mostly in deeper P. some ripe; very small a dwarf population, color described: back and head mottled olive brown; a distinct narrow golden stripe from upper corner of opercle to base of C., beneath this is a slightly wider orange red streak which becomes wider and more diffuse at the base of the C. below the red is a wide blackish band. The lower third of the head and body is whitish. Fins plain lowermost suffused with orange. Upper part of iris olive brown; lower silvery, pupil margined with orange.

Gila nigrescens

scarce in P.

Rhinichthys cataractae transmontanus

scattered; ads. were only fish in fastest water. both sexes ripe.

Salmo (trutta fario?)

1 ♀ seen above diversion dam, hooked and lost on a dry fly.

Nov. 31, 1951

V.M., Rio Arriba Co., Tusas Creek 1 mile
above Petaca.

June 10, 1949

W.V.K., K.R. Colman.

Pantosteus plebeius S & G.

199 yg-ad. 24-109 m SL.

A dwarf population. Sev. small
tuberculate individuals. Heavily parasitized
with Nereis.

Rhinichthys cataractae transmontanus Cope

244 yg-ad. 24-68 m SL.

Heavily parasitized with Nereis
Gila nigrescens G.

3 juv 52-58.

Rana pipiens DeRoeber

1 imago 44 m SL.

Bufo [coquatus or compactilis]

1 leg
