

June 12, 1950

WJK50-06-12-01

Went to Jersey about 3 miles above
mouth of Guadalupe creek.

Arrived about 1400

Rhinichthys spawning slightly below
sp. area of last off Friday. Watched until 1630.
when interest in sp. seemed to wane. After 1600
only a few guarding pdds remained
General pattern the same. Most

individuals were concentrated in a small area
and seemed to be without territories. Several
do maintained territories. There were ~~scattered~~
~~near the margin~~ as far as a meter away
from the main mass. All terr. were
~~off~~ in front of or between pieces of rubble
rather than on the smaller gravel.

Two Gila were present; one most
of the time till about 1610, the other was
present less than half of the time. They were
more easily frightened than Rhin. Mostly
they stayed at the lower edge of the mass
of Rhin which placed them in the "rough" water
just below the glossy "V" of the ripple.

June 12, 1950 #2.

X 90 W 80 T 1445

X 86 W 79.5 T 1500 a few clouds obscuring the sun.

X 89 W 79 T 1600

X 88 W 78 T 1630

Went out at ~~1600~~ 1630 - 1700 to catch Cula on small bait. The stream has seemingly been stocked recently with small (5-6") rainbows which took the bait more readily than the Cula. These were preserved. Switched to a fly and released most of the trout. One Cula female was taken on a fly - was ripe. They are hard to hook on even a number 14 fly. Even with bait, one must wait longer than with a trout.

Three brown trout above legal size were taken, the two largest on a dry. The heads & series & some scales were saved. Lengths: 1. 160 2. 198. 3 276.