

2016

MRI for Pediatric Appendicitis Reduces Pediatric Radiation Exposure

Edward Lam

Juliana Starcevich

Allyson Richards

Jessica Williams

Bryan Upham

See next page for additional authors

Follow this and additional works at: https://digitalrepository.unm.edu/rad_pubs

Recommended Citation

Lam, Edward; Juliana Starcevich; Allyson Richards; Jessica Williams; Bryan Upham; and Susan Williamson. "MRI for Pediatric Appendicitis Reduces Pediatric Radiation Exposure." (2016). https://digitalrepository.unm.edu/rad_pubs/2

This Poster is brought to you for free and open access by the Radiology at UNM Digital Repository. It has been accepted for inclusion in Radiology Research and Scholarship by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.

Authors

Edward Lam, Juliana Starcevich, Allyson Richards, Jessica Williams, Bryan Upham, and Susan Williamson

MRI for Pediatric Appendicitis Reduces Pediatric Radiation Exposure

Author Contact Information
 • Edward Lam
 • University of New Mexico Department of Radiology MSC10 5530
 • University of New Mexico
 Albuquerque, NM 87131-0001
 • E-mail: edlam@salud.unm.edu

Edward Lam, MD¹; Juliana Starcevich, MD¹; Allyson Richards, MD¹; Jessica Williams, MD¹; Bryan Upham, MD²; Susan Williamson, MD¹
¹University of New Mexico Hospital Department of Radiology, ²University of New Mexico Hospital Department of Emergency Medicine

Current standard of care

- ACR appropriateness criteria supports initial ultrasound followed by CT if needed.
- Role of MRI as yet undefined, with the value to be determined as MRI becomes more clinically available.

Pros and Cons of evaluating for appendicitis with MRI

Pros:

- No ionizing radiation
- Better contrast resolution

Cons:

- MRI less available
- Slower than CT
- Technically more difficult to interpret
- Unclear results may prompt CT

Objectives

- Determine if MRI can sufficiently assess for appendicitis and preclude the radiation exposure of CT.

Methods

- Retrospective review of the first 100 MRIs for appendicitis ordered by the UNMH Pediatric ED (January-September 2013).
- Interpretations categorized as:
 - Positive
 - Equivocal, favor positive
 - Equivocal
 - Equivocal, favor negative
 - Negative

Results

Final Interpretations:

- Positive: 22
- Equivocal, but favor positive: 5
- Equivocal: 20
- Equivocal, but favor negative: 11
- Negative: 42

The 22 positive interpretations


- 21 patients taken to surgery within 48 hours

The 36 Equivocal cases


- Equivocal, favor positive : Of the 5 studies, 3 were taken to the OR and found to have appendicitis.
- Equivocal: Of the 20 patients, 15 were discharged without further imaging. 5 admitted for observation before discharge.
- Equivocal, favor negative: All 11 patients were discharged from the ED.

The 42 Negative Cases


- All patients were either discharged or admitted for non-appendiceal pathology.




Positive exam: Enlarged, fluid filled appendix with adjacent free fluid.



Equivocal exam: Appendix not seen. Bilateral lower quadrant free fluid.



Negative exam: Normal appendix. No free fluid.



Follow up CT

- Ordered in 10% of patients, most of which had equivocal MRI interpretations
- None of these patients went to surgery for appendicitis

Conclusion:

- MRI is effective in the positive identification of appendicitis. No case dictated as equivocal, equivocal favor negative, or negative required surgery for appendicitis.
- The use of MRIs in the evaluation of right lower quadrant pain/suspected appendicitis provides valuable diagnostic information without ionizing radiation.

References

1. ACR Appropriateness Criteria® right lower quadrant pain—suspected appendicitis. Rosen MP, Ding A, Blake MA, Baker ME, Cash BD, Fidler JL, Grant TH, Greene FL, Jones B, Katz DS, Lalani T, Miller FH, Small WC, Spottwood S, Sudakoff GS, Tulchinsky M, Warshauer DM, Yee J, Coley BD. J Am Coll Radiol. 2011 Nov;8(11):749-55.
2. MRI for clinically suspected pediatric appendicitis: an implemented program. Moore MM, Gustas CN, Choudhary AK, Methratta ST, Hulse MA, Geeting G, Egli KD, Boal DK. Pediatr Radiol. 2012 Sep;42(9):1056-63.
3. MR imaging of the acute abdomen and pelvis: acute appendicitis and beyond. Singh A, Danrad R, Hahn PF, Blake MA, Mueller PR, Novelline RA. Radiographics. 2007 Sep-Oct;27(5):1419-31.