

University of New Mexico

UNM Digital Repository

The Kidney Institute of New Mexico

HSC Data Community

2021

Overlapping roles of NADPH Oxidase 4 (Nox4) for diabetic and gadolinium-based contrast agent-induced systemic fibrosis-Animal Equivalent Dosing of Gadolinium-based contrast agents Supplementary Figure 2

Francesca Bruno

Eboli Hospital "Maria SS Addolorata," UOC of Nephrology and Dialysis, Eboli, Italy

Joshua DeAguero

Kidney Institute of New Mexico, University of New Mexico Health Science Center, New Mexico Veterans Administration Health Care System

Catherine Do

Kidney Institute of New Mexico, University of New Mexico Health Science Center, New Mexico Veterans Administration Health Care System

Doug Yoon Lee

University of Texas Health System

Chunyan Tan

University of Texas Health System

Follow this and additional works at: <https://digitalrepository.unm.edu/kinm>

See next page for additional authors


Recommended Citation

Bruno, Francesca; Joshua DeAguero; Catherine Do; Doug Yoon Lee; Chunyan Tan; G Patricia Escobar; and Brent Wagner. "Overlapping roles of NADPH Oxidase 4 (Nox4) for diabetic and gadolinium-based contrast agent-induced systemic fibrosis-Animal Equivalent Dosing of Gadolinium-based contrast agents Supplementary Figure 2." (2021). <https://digitalrepository.unm.edu/kinm/3>

This Article is brought to you for free and open access by the HSC Data Community at UNM Digital Repository. It has been accepted for inclusion in The Kidney Institute of New Mexico by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.

Creators

Francesca Bruno, Joshua DeAgüero, Catherine Do, Doug Yoon Lee, Chunyan Tan, G Patricia Escobar, and Brent Wagner



Supplementary Figure 2. Renal changes induced by gadolinium and experimental diabetes. Periodic acid acid-Schiff stained photomicrographs were analyzed by a blinded veterinary pathologist (**A-C**). Regions were sectioned at identical thicknesses, stained similarly, and photographed at the same magnification. For each mouse there were $n = 3$ figures of different glomeruli. Background darkness due to inadequate illumination was removed using the “levels” command and each figure was brought to the same size and resolution using the “image size” command. Glomerular area (**A**), number of nuclei in each glomerulus (**B**), and area of strongly staining mesangial matrix (**C**) were determined for each. **D**. DHE fluorescence from renal cortex. Reactive oxygen species generation in the renal cortex from gadolinium treatment and experimental diabetes. Fluorescence was quantitated using ImageJ 1.52d (National Institutes of Health, USA (22)), $n = 4$ per group. *, $P < 0.05$, *** $P < 0.001$ from lane 1, # $P < 0.05$, ## $P < 0.01$, and ### $P < 0.001$ from lane 2 by analysis-of-variance and Tukey honestly significant post-hoc testing.