

Summer 7-11-2018

# The Effect of Breathy and Strained Vocal Quality on Vowel Perception

Sarah Nieman  
*University of New Mexico*

Follow this and additional works at: [https://digitalrepository.unm.edu/shs\\_etds](https://digitalrepository.unm.edu/shs_etds)



Part of the [Speech and Hearing Science Commons](#)

---

## Recommended Citation

Nieman, Sarah. "The Effect of Breathy and Strained Vocal Quality on Vowel Perception." (2018). [https://digitalrepository.unm.edu/shs\\_etds/22](https://digitalrepository.unm.edu/shs_etds/22)

This Thesis is brought to you for free and open access by the Electronic Theses and Dissertations at UNM Digital Repository. It has been accepted for inclusion in Speech and Hearing Sciences ETDs by an authorized administrator of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).

Sarah Rose Nieman, B.A.

*Candidate*

Department of Speech and Hearing Sciences

*Department*

This thesis is approved, and it is acceptable in quality and form for publication:

*Approved by the Thesis Committee:*

Amy T. Neel, Ph.D., CCC-SLP, Chairperson

Phyllis Palmer, Ph.D., CCC-SLP

Rick Arenas, Ph.D.

**THE EFFECT OF BREATHY AND STRAINED VOCAL  
QUALITIES ON VOWEL PERCEPTION**

**by**

**SARAH R. NIEMAN**

**B.A. SPEECH AND HEARING SCIENCES, SPANISH**

**THESIS**

Submitted in Partial Fulfillment of the  
Requirements for the Degree of

**MASTER OF SCIENCE  
SPEECH-LANGUAGE PATHOLOGY**

The University of New Mexico  
Albuquerque, New Mexico

**JULY, 2018**

**ACKNOWLEDGEMENTS**

I am extremely grateful to Dr. Neel for her time, expertise, and support throughout this process. I'd like to thank her for welcoming me into her lab as an undergraduate, for allowing me to be her graduate assistant, and for asking me to do a thesis. I never would have thought I could do it. Our ability to hold full conversations without content words is one of my favorite memories of graduate school.

Thank you to the listeners in this study for their time. Thank you to members of my thesis committee, Dr. Arenas and Dr. Palmer, for their technological know-how and research savvy. Thank you to Sofiya Krasilshchikova for her tenacious data-crunching.

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

The Effect of Breathy and Strained Vocal Quality on Vowel Perception

Sarah R. Nieman

B.A., Speech and Hearing Sciences, Spanish, University of New Mexico,

2016

M.S., Speech-Language Pathology, University of New Mexico, 2018

## ABSTRACT

**INTRODUCTION:** Research into speech intelligibility in dysarthria historically focuses on articulation deficits. However, voice quality deficits associated with motor speech disorders may also impact speech perception. This study investigates how breathy and strained vocal quality affects vowel identification and ratings of vowel goodness.

**METHODS:** A healthy speaker recorded vowels with normal, simulated breathy and simulated strained voice quality. Acoustic, physiologic, and perceptual measures confirmed the presence of the desired voice deficits. 16 volunteer listeners participated in three perceptual tasks: vowel identification, vowel goodness ratings, and voice quality ratings.

**RESULTS:** In the voice quality rating task, listeners detected voice quality deficits with ease. Breathy and strained stimuli were rated as significantly poorer in voice quality than normal stimuli. The voice quality deficits did not appear to impact vowel identification: identification accuracy for all three sets was high (95% and above) and scores did not differ significantly across the three sets of vowels. Listener judgments of vowel goodness, however, were affected by voice quality. Breathy and strained vowels were rated as significantly poorer than normal vowels. In addition, listeners needed more

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

time to rate the articulatory goodness of the disordered stimuli and replayed them more often while making their goodness judgments.

**CONCLUSION:** Simulation of voice quality deficits appears to be a valid way of assessing the impact of speech factors beyond articulation on the perception of disordered speech. Stimuli with simulated breathiness and strain were rated as poorer in voice quality than normally voiced vowels, indicating that voice quality is salient to listeners. Although identification accuracy was not affected by voice quality deficits, breathy and strained vowels were judged as poorer in articulatory goodness than normally voiced vowels. Abnormal voice quality appeared to interfere with listener judgments of the articulatory goodness of vowels. Voice quality deficits associated with dysarthria may affect speech perception by causing increased listener effort even if speech intelligibility is not directly affected. Further study of the effect of voice quality in more realistic listening conditions (e.g., in noise) with more complex speech stimuli (e.g., sentences or conversation) will help determine the need for phonatory treatment of dysarthric speech.

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

## TABLE OF CONTENTS

<b>ACKNOWLEDGEMENTS .....</b>	<b>iii</b>
<b>ABSTRACT.....</b>	<b>iv</b>
<b>LIST OF FIGURES .....</b>	<b>viii</b>
<b>LIST OF TABLES .....</b>	<b>ix</b>
<b>INTRODUCTION/LITERATURE REVIEW .....</b>	<b>1</b>
Speech Production .....	1
Speech Intelligibility .....	2
Dysarthria in Neurogenic Disorders .....	4
Vowels in Dysarthria .....	6
Vowels in Parkinson Disease .....	7
LSVT and Loud Speech.....	8
Aims .....	9
<b>METHOD .....</b>	<b>12</b>
Participants.....	12
Stimuli.....	12
Objective Measures.....	13
Subjective Measures .....	14
Procedure .....	15
Training.....	15
Tasks .....	15
Vowel Identification .....	15
Vowel Goodness .....	16
Quality Goodness.....	17

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Time/Replays .....	17
Analysis.....	17
<b>RESULTS .....</b>	<b>19</b>
Vowel Quality Acoustic Measures .....	19
Voice Quality Measures.....	23
Objective Measures.....	23
Subjective Measures .....	25
Vowel Identification .....	26
Vowel Goodness Ratings .....	27
Voice Quality Ratings.....	29
Reliability Ratings .....	30
Intra-rater Reliability .....	30
Inter-rater Reliability .....	31
<b>DISCUSSION .....</b>	<b>34</b>
Did Vowel Sets Differ in Quality?.....	34
Were Vowels Similar in Other Aspects of Production? .....	36
Did Listeners Notice Quality Differences?.....	37
Did Voice Quality Affect Vowel Goodness Judgements?.....	37
Did Voice Quality Affect Vowel Identification? .....	38
Limitations .....	39
Clinical Implications .....	40
Directions of Future Research .....	41
<b>LIST OF APPENDICES .....</b>	<b>43</b>



**REFERENCES.....48**

**LIST OF FIGURES**

Figure 1 Alvin 3 screenshot of identification task .....16

Figure 2 Alvin 3 screenshot of vowel goodness task.....17

Figure 3 Means and SD for F1.....19

Figure 4 Means and SD for F2.....20

Figure 5 Dynamic vowel chart.....21

Figure 6 Means and SD for F2.....22

Figure 7 Means and SD for Duration.....23

Figure 8 Cape-V Perceptual Scale .....26

Figure 9 Means and SD for vowel goodness ratings .....28

Figure 10 Means and SD for voice quality ratings .....29

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

## LIST OF TABLES

Table 1 Acoustic measures .....	21
Table 2 Repeated Measures ANOVA for acoustic measures .....	24
Table 3 CAPE-V Perceptual Scale .....	26
Table 4 Repeated Measures ANOVA for experimental tasks .....	27
Table 5 ICC Comparisons: Intra-rater .....	30
Table 6 ICC Comparisons: Inter-rater .....	32

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

## Literature Review

**Hypothetical Case Study.** Mr. Montoya is a 65-year-old man living with Parkinson disease (PD). His chief complaint is that his wife and grandchildren can't understand him. Like 90% of individuals with PD, Mr. Montoya has a concomitant motor speech disorder called hypokinetic dysarthria (Tylakova et al., 2017; Duffy, 2013). Due to the dysarthria, Mr. Montoya has deficits in all four subsystems required for functional speech: respiration, phonation, resonance and articulation. At the moment, there is insufficient scientific evidence to determine which aspect of speech production should be targeted to attain optimal intelligibility gains in the shortest amount of time. The present study examines phonatory effects on intelligibility to support future clinical decisions for patients like Mr. Montoya, beginning with an overview of current literature on speech intelligibility and brief analysis of the gaps in the research.

**Speech Production.** The production of intelligible speech requires the interaction of two components, the glottal source and the vocal tract filter (Fant, 1960). For vowel sounds, the glottal source consists of the complex tone produced by the vibrating vocal folds. The filter consists of the vocal tract above the level of the vocal folds and the articulators (e.g., lips, tongue, jaw). Acting as an acoustic resonator, the filter allows specific bands of frequencies, called resonances or formants, to pass into the air with higher energy than other frequencies. Speakers change formant frequencies by modifying the shape of the vocal tract filter. Moving the tongue, jaw, and lips changes the frequencies that are best resonated by the vocal tract (Behrman, 2018). Listeners use formant frequency patterns to perceive vowel sounds. The first formant frequency, or F1, is associated with tongue height. Vowels with a high tongue position, such as /i/ and /u/, have low F1 frequencies,

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

whereas low vowels, such as /ɑ/, have high F1 frequencies. F2, or the second formant frequency, is associated with tongue advancement. Front vowels, such as /i/ and /ɪ/, have high F2 frequencies, and back vowels, such as /u/ and /ʊ/ have low F2 values (Stemple et al., 2014).

**Speech intelligibility.** Most studies of speech intelligibility focus on the articulation of speech sounds associated with the vocal tract filter (Skodda, Visser & Schlegel, 2011; Kim, Hasegawa-Johnson, & Perlman, 2011; Platt, Andrews, Young, & Quinn, 1980).

Although both consonants and vowels are important for speech intelligibility, the focus of this paper is on vowels. In neurogenic disorders such as PD, weak or uncoordinated articulators are unable to shape the vocal tract correctly, resulting in speech sound distortions, substitutions of one phoneme for another, or omissions of phonemes.

Impaired shaping of the vocal tract can affect production of vowels, leading to reduced speech intelligibility (Monsen, 1983; Whitehead & Wirz, 1979).

Characteristics of the glottal source may also influence the ability of listeners to understand speech (Dyle, Danhauer & Reed, 1988; Eadie et al., 2013; DeBodt, 2002; Ramig, 1992). Vocal source characteristics include fundamental frequency, vocal intensity, and vocal quality. Fundamental frequency ( $f_0$ ), is the source characteristic perceived as vocal pitch. The rate of vibration of the vocal folds is directly related to the  $f_0$  (Stemple et al., 2014). For example, when the vocal folds vibrate at a rate of 200 times per second, the fundamental frequency is 200 Hz. Pitch changes are used to impart suprasegmental information to listeners. Per Kent (1988), prosody is informed by voice quality, intensity variation, pitch level and pitch variation. Pitch contours are necessary for listeners to interpret meaning, and monopitch, or lack of pitch contours, negatively

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

affects intelligibility (Haycock, 1933; Greene, 1956; Hood, 1966; Monsen, 1979; DeBodt, 2002). Restricted fundamental frequency ranges have been well-documented in PD and other neurogenic disorders (Canter, 1963; 1965; Kent & Rosenbek, 1982). Reduced fundamental frequency range has been shown to decrease intelligibility in both healthy speakers and in dysarthric speakers (Laures & Weismer, 1999). Though Bunton (2006) found fundamental frequency to typically be a redundant cue in healthy speakers, listeners in the study used it to identify vowels when listening to dysarthric speech.

Vocal intensity, perceived as loudness by listeners, is a measure of the sound pressure level of the voice. Vocal intensity is a function of subglottal pressure and the degree of laryngeal adduction. The duration of vocal fold closure, degree of closure and closure speed are specific factors that affect intensity. An increased closed duration will increase subglottal pressure, resulting in greater intensity. Similarly, a tight laryngeal adduction will increase subglottal pressure, while an incomplete closure reduces pressure build-up. The faster the closure speed of the vocal folds, the more energy passes into the air at the mouth (Behrman, 2018). Intensity is important for speech intelligibility for two reasons: for the audibility of the speech signal to listeners and for its prosodic functions. Intensity is affected in neurogenic disorders such as PD due to disordered laryngeal and respiratory function. Vocal fold bowing or other glottal incompetence prevents complete glottal closure, which inhibits build-up of subglottal pressure (Ramig, 1992). Reduced vocal intensity is a well-known characteristic of PD (Canter, 1963; Kent & Rosenbek, 1982; Ludlow & Bassich, 1983), as is vocal fold bowing (Hansen et al., 1984; Smith et al., 1995). Several recent therapeutic techniques (LSVT, SpeakOut, Clear Speech) focus

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

on increasing loudness in speakers with PD by facilitating improved vocal fold adduction (Watts, 2016; Cleveland, et al., 2015; Ramig, Fox & Sapis, 2011; Lam & Tjaden, 2016)

Another parameter associated with glottal tone is voice quality. Voice quality refers to the auditory perception of multi-dimensional factors including laryngeal adduction, respiration, muscle tension, fundamental frequency, and others. These factors come together in varying ratios, leading to different signal perception and descriptive interpretation (Kreiman, 2008). This makes research on vocal quality difficult because listeners will perceive vocal quality differently. For the purposes of this paper, Laver's (1980) method of distinguishing physiological differences at the laryngeal and supralaryngeal levels will be used to define voice qualities. Glottic leakage due to hypoadduction of the vocal folds, a space-occupying lesion, or bowed vocal folds leads to perception of breathy vocal quality (Stemple et al., 2014; Barsties von Latoszek et al., 2017), due to the increased obstruction. Strained vocal quality is perceived when the vocal folds are hyperadducted or "pressed" tightly (Stemple et al., 2014; Barsties von Latoszek et al., 2017). Voice quality has been shown to affect intelligibility in alaryngeal speakers (Doyle, Danhauer, & Reed, 1988) and in deaf and hard of hearing speakers (Whitehead & Wirz, 1979; Monsen, 1983). To date, there is limited documented research on the effect of voice quality and intelligibility in individuals with motor speech disorders, though it has been noted clinically (Ramig, 1992).

**Dysarthria in Neurogenic Disorders.** Motor speech disorders caused by damage at some point along the motor pathway are classified as dysarthrias (Duffy, 2013).

Compromised laryngeal and/or supralaryngeal neural integrity leads to the development of characteristic voice quality, articulatory, and respiratory patterns signaling the location

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

of the damage (Stemple et al., 2014). Dysarthria may result in both source and filter deficits depending on the specific etiology and affected motor pathways (ASHA, 2017). Individuals with hypokinetic dysarthria and flaccid dysarthria have low-intensity, excessively breathy phonation and insufficient prosody due to source deficits. Hypokinetic dysarthria, typically associated with PD, may be further characterized by imprecise consonants and a variable rate due to filter deficits (Gazewood, Richards, & Clebak 2013; Duffy, 2013). Flaccid dysarthria, typically associated with amyotrophic lateral sclerosis (ALS) and brainstem strokes, may also be characterized by short phrases, reduced speaking rate and mild consonant imprecision (Duffy, 2013). Hyperkinetic dysarthria and spastic dysarthria lead to strained-strangled voice quality, inappropriate loudness, and monopitch. Hyperkinetic dysarthria, typical of Huntington's chorea and spasmodic dysphonia, is further characterized by voice stoppages, distorted vowels and imprecise consonants (Duffy, 2013).

The present study is primarily interested in research concerning PD since approximately 90% of individuals with the disease will develop dysarthria, which may affect respiration, phonation, resonance and/or articulation (Tylakova et al., 2017; Duffy, 2013). With multiple subsystems degenerating simultaneously and a limited timeline, it is important to identify the most important intervention target. The most common form of dysarthria in PD is hypokinetic dysarthria, though hyperkinetic dysarthria and mixed dysarthria are possible (Tjaden, 2008; Duffy, 2013).

Research has examined both source and filter deficits in PD. In a study of 31 hypokinetic dysarthric subjects Zwirner & Barnes (1992) found a higher ratio of source deficits to filter deficits. Similarly, in a study of 200 subjects Logeman & Fisher (1981)

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

found all subjects with articulation deficits had voice quality deficits and 45% of the 200 subjects had voice quality deficits only. Overall, 89% of subjects in Logeman & Fisher's study experienced voice quality deficits. Ho et al. (1999) supported Logeman & Fisher's findings with a descriptive study of 200 individuals with PD. They found voice deficits were the prominent deficits experienced by their subjects, at 65.5%. Extensive research supports the presence monopitch, reduced fundamental frequency, and reduced vocal intensity in PD (Hanson, Gerratt & Ward, 1984; Logeman & Fisher, 1978; Darley, 1996; Boshes, 1996), though intensity may be influenced by both source and filter.

Filter deficits have been associated with lower rates of speech intelligibility in PD (Skodda, Visser & Schlegel, 2011; Kim, Hasegawa-Johnson, & Perlman, 2011; Platt, Andrews, Young, & Quinn, 1980). Logeman & Fisher (1978) found imprecise articulation patterns led to stops and affricates being produced as fricatives, repetitions of syllables (i.e., fluency deficits), and inappropriate rushes of speech referred to as festinated speech (Duffy, 2013). Kim, Hasegawa-Johnson, & Perlman (2011) contrasted with Logeman & Fisher, finding voicing and place errors to be more frequent than manner errors. They found non-uniform error patterns overall. Lower intelligibility was associated with voicing and place errors over manner errors. Reduced jaw movement, velopharyngeal movement and voice onset time (VOT) have also been documented in the research (Weismer, 1984; Canter, 1965; Logeman & Fisher, 1978; Caligiuri, 1987; Conner et al., 1989)

**Vowels in Dysarthria** Vowels are frequently the focus of dysarthria research. Vowels are targeted because they are relatively long time periods of voiced speech with limited filter effects from vocal tract shaping, allowing for controlled perceptual assessment of



## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

source characteristics (Kreiman, 2008). Vowels are distinguished by producing them with varying tongue heights and tongue advancement. These variations correspond to well-documented, distinct vowel formant frequency patterns, F1 and F2 (Peterson & Barney, 1952). Vowel perception is important to overall speech intelligibility. As distinctiveness of a given vowel compared to other vowels is reduced, intelligibility is also reduced (Kim et al., 2011; Savageau et al., 2015; Monsen, 1983).

**Vowels in Parkinson Disease** There has been extensive research on vowels in PD.

Vowel Space Area (VSA) is a common research metric to determine the distinctiveness of an individual's vowel productions. VSA measures the distance between vowels in a vowel quadrilateral plot. Reduced vowel space area reflects reduced tongue movement and correspondingly-altered measures of F1 and F2. Results have been inconsistent, with some studies finding VSA could differentiate between normal and dysarthric speakers and others finding it could not (Sapir et al., 2011). Skodda and colleagues (2011, 2012) found vowel space scores deteriorated with progression of PD. Other studies support vowel movement to the center of the vowel quadrilateral (vowel centralization) and reduced vowel space in PD, even early on in the disease (Bang et al., 2013; Rusz, et al., 2013). VSA correlates with filter deficits.

Teasing apart source and filter effects in speakers with PD is important in understanding what causes deficits in speech intelligibility. When a clinician is faced with abnormal phonation, articulation, resonance, and prosody simultaneously, it is difficult to know what to treat first to obtain the best functional gains, especially in a limited time frame. Comprehending the effect of voice quality deficits on vowel perception begins the process of learning how vocal quality impacts speech intelligibility.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

**LSVT and Loud Speech** One well-supported treatment for individuals with speech deficits due to PD is the Lee-Silverman Voice treatment (LSVT LOUD), described in one study as the “most effective treatment for PD in reducing the impact of hypokinetic dysarthria on functional communication” (Constantinescu, Theodoros, et al., 2011). LSVT LOUD targets the development of high-intensity, effortful speech production in an effort to “recalibrate” self-perception of intensity and motor recruitment (Ramig, Fox & Sapir, 2011). Per Ramig and colleagues, increased intensity improves laryngeal deficits, respiratory deficits and orofacial movement, leading to improved vocal quality and improved vowel and consonant articulation (Ramig, 1992; Schulman, 1985; Dromey & Ramig, 1998; Dromey, Ramig & Johnson, 1995; Sapir et al., 2007).

Research shows LSVT LOUD and other loud speech techniques, like Speak Out, affect both the source and filter including vocal intensity, vocal quality and articulatory accuracy (Watts, 2016; Cleveland, et al., 2015; Ramig, Fox & Sapir, 2011) . LSVT LOUD is the most studied loud speech technique and is correlated with both source and filter changes. Stroboscopy indicates tighter, more symmetrical vocal fold adduction following LSVT LOUD (Smith et al., 1995). Increased vowel space in many speakers (Bunton, 2006; Neel & Beveridge, 2006) also supports source change. Greater tongue strength (Ward et al., 2000;) and greater articulator movement (Schulman, 1989; Dromey & Ramig, 1998; Sapir & Ramig et al., 2002) support filter changes.

The study that inspired this thesis was conducted by Shimon Sapir and colleagues in 2007. The randomized control trial included a treatment group of individuals with PD, a control group with PD, and a control group of age-matched, neurotypical peers. Subjects in the treatment group attended hour-long, individual therapy sessions four times

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

a week for four weeks (16 sessions). Subjects also performed a daily home exercise regimen consisting of loud phonation for as long as possible, high-amplitude speech exercises, and phonation at both maximum high- and low-pitch. Sapir and colleagues concluded changes in vowel articulation, a by-product of high-intensity speech as noted above, were responsible for better vowel goodness ratings in the treatment group compared to the no-treatment group. Sapir's study did not control for the improved vocal quality that is an expected byproduct of forceful vocal fold adduction, however. It is possible listeners were not able to distinguish between voice quality improvement and vowel articulation improvement when rating vowel "goodness," leading to confounded results.

We do not currently have a full understanding of how source characteristics and articulatory dimensions impact speech intelligibility (Kent et al., 2003). Ramig (1992) points to research on speech of deaf individuals and alaryngeal speakers to support the theory that vocal quality affects speech intelligibility ratings. Voice quality deficits and articulation deficits both contribute to reduced speech intelligibility in deaf children (Monsen, 1983). Similarly, in laryngectomees voice quality deficits have been coupled with lower speech intelligibility (Dyle, Danhauer & Reed, 1988; Eadie et al., 2013). With dysarthric speech, DeBodt et al., (2002) found functional intelligibility is improved linearly by combining speech dimensions. While prosody and articulation were found to be the most influential speech dimensions correlated with intelligibility estimations, voice quality and prosody were also implicated in improved intelligibility.

**Aims.** This current study aims to identify the impact of disordered vocal quality on vowel perception in listeners. In order to ascertain the effects of breathy and strained voice

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

quality on vowel perception, researchers must use vowel stimuli that differ from one another only in voice quality. The vowel stimuli must not differ in the other source characteristics, pitch and intensity, nor should they differ in filter characteristics such as F1 (tongue height), F2 (tongue advancement) and duration. It would be useful to employ computer-created synthetic or resynthesized vowels to carefully control all elements of vowel production. However, it is difficult to produce a set of several realistic synthetic or resynthesized breathy and strained vowels. Therefore, this study employs simulated breathy and strained vowel stimuli as well as vowels produced with normal voice quality spoken by a healthy speaker who attempted to hold other source and filter characteristics constant while producing the three voice qualities. In contrast to Sapir et al., (2007), this study will clearly demonstrate if vowel goodness changes are due to voice quality apart from articulation. Results will guide further research as we determine if voice quality can affect speech intelligibility. Clinicians will then be better able to assess client need and select appropriate intervention targets in order to support function in a timely, efficient manner. To this end, the specific aims of this project are to:

**Specific Aim #1:** To determine the effect of each simulated deficit on vowel identifiability, identification scores, time needed to identify tokens and the number of replays needed.

**Specific Aim #2:** To determine the effect of each simulated deficit on vowel “goodness” rating, average ratings, time needed to rate “goodness” and the number of replays needed.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Specific Aim # 3: To determine the effect of each simulated deficit on voice quality rating, average ratings, time needed to rate voice quality and the number of replays needed.

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

## Methods

This research project was approved by the Institutional Review Board (IRB) of the University of New Mexico.

### Participants

Sixteen volunteer undergraduate and graduate students in the Speech and Hearing Sciences Department at the University of New Mexico were recruited via email list. Participants were self-reported native English speakers with no history of speech, language, or hearing problems. Each participant passed a pure-tone audiometry test ( 500 Hz, 1,000 Hz, 2,000 Hz, and 4, 000 Hz at 20 decibels) prior to beginning the tasks.

### Stimuli

One healthy female speaker, this study's principal investigator, produced three sets of 10 vowels (/i ɪ ε æ ʊ u ʌ oʊ ɑ eɪ/) within the carrier phrase, "say hood again", for a total of 30 phrases. Breathy vocal quality was produced by reducing vocal fold contact, therefore increasing turbulent airflow during phonation. Strained vocal quality was produced by hyperadducting the vocal folds. Stimuli were recorded via Audacity (Mazonni, 1999) using an EG2-PCX model electroglottograph (EGG) with 35-mm dual channel electrodes and accompanying Glottal Enterprises M80 omnidirectional headset microphone in a sound-treated booth. A harmless electrical current passed through the speaker's vocal folds to record vocal fold movement to find the Contact Quotient (i.e., CQ: contact time of the vocal folds divided by cycle length) measures. Cycle length is determined based on a preset threshold (e.g., contact begins at 25% of maximum amplitude and ends at 25% amplitude). CQ50% is a physiological measure with a threshold at 50% of the amplitude.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

CQ50% has been shown to distinguish breathy, strained, and normal phonation (Liu et al., 2017).

*Objective Measures.* PRAAT software (Boersima & Weenik, 2015) was used to annotate stimuli in order to isolate vowels. Average intensity for each vowel in each condition was extracted via PRAAT software. A script customized by Dr. Richard Arenas extracted  $f_0$ , F1, F2, duration, and Cepstral Peak Prominence-Smoothed (CPPs) measures for each marked vowel. Standard PRAAT settings track 5 formants up to 55k Hz, but settings needed to be adjusted for some stimuli (i.e., up to 5k Hz). Formant measures were double-checked by hand and changed as needed to ensure accurate formant tracking ( $<10$ ). Several acoustic measures were used to determine that vowel productions were similar in articulation and differed only in voice quality. F1 and F2 measures informed tongue height and advancement across conditions. Fundamental frequency measures confirmed similar pitch across all conditions. To prepare the stimuli for the perceptual tests, a second PRAAT script extracted each vowel as a separate wav file. Vowels were equated for loudness (mean RMS intensity) using Adobe Audition (Audacity.sourceforge.net, 2015) software to ensure that intensity differences did not influence perceptual judgments.

Two measures were used to determine that the three types of stimuli did differ in voice quality as desired: CPPs and CQ50%. A Fourier transformation converts waveform frequency into a time domain leading to a “spectral representation of the spectrum” (Heman-Ackah et al., 2003), called a cepstrum. Smoothed cepstral peak prominence (CPPs) is the highest amplitude in a given cepstrum and is the acoustic measure most strongly associated with breathy voice quality (Latozek et al., 2016). Breathly voices have

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

a flatter cepstrum overall, meaning CPPs measurements are smaller than for non-breathy voice quality (Hillenbrand & Houde, 1996). Small CPPs have been correlated with strained vocal quality (Lowell et al., 2012). CPPs measures were obtained using Praat. The physiologic measure CQ50% was used to determine differences in vocal fold closure patterns across the three voice quality types. Vocal fold closure patterns differ across voice quality. In typical phonation, the vocal folds oscillate fluidly horizontally, vertically, and longitudinally. Electroglottograph measures pass an electrical current through the vocal folds to record when vocal folds are open vs. closed. Software translates these readings into a contact quotient (CQ) by dividing cycle length by total time the vocal folds were closed. For this study, cycle is defined by 50% peak to 50% peak in the EGG signal. CQ50% for typical phonation shows about a 2:3 ratio of contact time to open time (Liu et al., 2012). Strained phonation occurs due to hyperadduction of the vocal folds, limiting the fluid motion of the vocal fold edge (Stemple et al., 2014). CQ50% for strained phonation is expected to be the largest of the three vocal qualities, with a higher proportion of contact time to open time (Liu et al., 2012). In breathy phonation the vocal folds do not fully approximate, meaning EGG measures for breathy phonation are limited. Although anterior vocal fold contact may be sufficient for contact readings, a posterior vocal fold gap is likely in this condition and may not be sensed by the EGG.

*Subjective Measures.* To further confirm that the desired voice quality deficits were achieved, two experienced clinical speech-language pathologists assessed four vowels and four sentence-level experimental stimuli using the CAPE-V perceptual scale,



## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

a reliable subjective measure of voice quality (Zraick et al., 2010). CAPE-V scores confirmed moderate deficits in breathiness and strain present in respective stimuli

### **Procedure**

**Training.** Participants received a short training session from the investigator to ensure understanding of goodness concepts, voice quality concepts and the international phonetic alphabet (IPA) (Appendix B). During the training, participants were presented with examples of normal, breathy, and strained stimuli in /hVd/ contexts. They practiced Visual Analog Scale (VAS) rating tasks and vowel identification tasks using speech samples from non-experimental stimuli. Research has found no significant difference in perceptual judgements of dysarthric speech between expert and naïve listeners (Sussman & Tjaden, 2012), so undergraduate and graduate students were selected as the research participants.

### **Tasks**

**Vowel Identification.** Participants performed 10-alternative forced choice tasks for vowel identification. For each trial, one of ten vowels was presented via headphones at a comfortable listening level. Participants identified which of the ten possible vowels was presented via a mouse click on the perceived vowel (Figure 1). Vowel choices were presented orthographically (e.g., had) and phonetically (e.g., /hæd/). There was a total of 30 stimuli (10 vowels X 3 conditions X 1 speaker) presented in random order 3 times for interrater reliability. Each participant therefore completed 90 vowel identification trials. Delivery and response collection was managed by Alvin 3 experiment software (J.M. Hillenbrand & Gayvert, 2005). Percent correct vowel identification was calculated for each voice type.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION



Figure 1. Screenshot of Alvin 3 experiment software display for identification task

**Vowel Goodness.** Participants completed vowel “goodness” ratings of the same stimuli sets in /hVd/ contexts using a visual analog scale (VAS) ranging from “poor example” to “good example” of each vowel under each condition (Figure 2). Each stimulus was presented 3 times in random order to determine interrater reliability. There were a total of 90 vowel goodness trials per participant. The Alvin 3 program translated mouse clicks on the analog scale into a number between 0 (good example) and 100 (poor example). Vowel goodness measurements will provide more fine-tuned data on how close a given vowel is to the listener’s concept of an excellent exemplar. Measuring both intelligibility and vowel goodness may result in observable trends not visible with only one measure (Franklin & Stoel-Gammon, 2014).

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION



Figure 2. Screenshot of Alvin 3 experiment software display for vowel goodness task

**Quality Goodness.** Participants completed voice quality “goodness” ratings of the same stimuli sets in /hVd/ contexts using a visual analog scale (VAS) ranging from “poor example” to “good example” of each vowel under each condition. The intended vowel was displayed orthographically so participants were able to compare. Each stimulus was presented in 3 times randomly to determine interrater reliability. There were a total of 90 quality goodness trials per participant. The Alvin 3 program translated mouse clicks on the analog scale into a number between 0 (good example) and 100 (poor example).

**Time/Replays.** The Alvin 3 program recorded time needed to respond in milliseconds and recorded number of replays for each stimuli. Time and number of replays were compared for breathy, normal and strained tokens in each task (identification, goodness rating and quality rating).

### Analysis

1. To determine the effect of each simulated deficit on vowel identifiability, identification scores, time needed to identify tokens and the number of replays were averaged across the listeners, transformed into RAU, and submitted to a mixed ANOVA

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

using IBM SPSS Statistics 25 (SPSS Inc., Chicago, IL) with within-subjects factor of condition, which includes the normal voice quality and the two simulated deficits.

2. To determine the effect of each simulated deficit on vowel “goodness” rating, average ratings, time needed to rate “goodness” and the number of replays were submitted to repeated-measures ANOVA with the within-subjects factor of condition (breathy, normal, or strained). Ratings were also subjected to Bonferroni-adjusted pairwise comparison.

3. To determine the effect of each simulated deficit on voice quality rating, average ratings, time needed to rate voice quality and the number of replays were submitted to repeated-measures ANOVA with the within-subjects factor of condition (breathy, normal, or strained). Ratings were also subjected to Bonferroni-adjusted pairwise comparison.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

### Results

**Vowel quality acoustic measures.** To confirm that vowels were similar for parameters other than vocal quality, measures were taken of F1, F2, and fundamental frequency at 20%, 50% and 80% of the vowel. F1, F2, fundamental frequency, and vowel duration were submitted to repeated-measures analysis of variance (ANOVA) using IBM SPSS Statistics 25 (SPSS, Inc., Chicago, IL) with a within-subject factor of voice type (breathy, normal, or strained). There was a significant effect of voice type for F1,  $F(1,2)=6.57$ ,  $p=.005$ ,  $\eta^2=.422$  (Table 2, Figure 3). Pairwise Bonferroni adjusted comparisons indicate a significant difference of 99 Hz between the breathy and strained conditions,  $p=0.016$ . There was no significant difference among the three voice quality types for F2,  $F(1,2)=2.681$ ,  $p=.096$  (Table 2, Figure 4).




Figure 3. Means and SDs for F1.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION




Figure 4. Means and SDs for F2.

Results of the dynamic vowel chart (Figure 5) shows F1 and F2 values for each vowel at 20%, 50%, and 80% of vowel duration. This F1 X F2 vowel scatterplot reveals differences in production for a few vowels across the three voice quality types. The strained voice quality version of /eɪ/ was produced with a lower tongue position and /ʊ/ was produced with a higher tongue position than for the breathy and normal versions. For normal voice quality, /u/ was produced with a more backed tongue position than the /u/ in breathy or strained voice quality. Table 1 shows formant frequencies by vowel and voice type.

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION




Figure 5. F1 (y-axis) and F2 (x-axis) formant frequencies at 20%, 50% and 80% of each vowel for every vowel in each condition. Note: 20% of the vowel is indicated by the largest marker in a series. 80% of the vowel is indicated by the smallest marker in the series.

**Table 1**

Measures of F1 (Hz), F2 (Hz),  $f_0$  (Hz), and duration (sec) by vowel and voice type

Vowel	F1			F2			$f_0$			Duration		
	B	N	S	B	N	S	B	N	S	B	N	S
i	306	268	423	2766	2827	2860	209	243	258	0.13	0.11	0.2
ɪ	399	509	494	2286	2137	2217	233	240	242	0.1	0.08	0.13
eɪ	439	455	706	2578	2631	2404	229	220	208	0.14	0.13	0.16
ε	726	802	729	2001	2052	1994	206	207	221	0.1	0.08	0.19
æ	941	993	909	1946	1680	1864	198	199	169	0.11	0.12	0.24
ʌ	664	783	828	1716	1595	1508	196	203	190	0.08	0.07	0.18
ɑ	943	948	974	1564	1404	1487	190	204	183	0.13	0.12	0.20
oʊ	416	424	548	971	1007	969	177	204	216	0.12	0.12	0.23
ʊ	427	515	511	1434	1446	1254	179	227	228	0.1	0.08	0.19
u	326	255	452	1276	925	1103	202	230	224	0.12	0.12	0.24

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

For fundamental frequency, Greenhouse-Geisser-corrected measures of  $f_0$  were not significantly different,  $F=3.090$ ,  $p=.100$  (Table 1, Table 2, Figure 6). This indicates fundamental frequency (vocal pitch) was held relatively constant across the three voice quality types.




Figure 6. Means and SDs for  $f_0$

Greenhouse-Geisser-corrected measures of vowel duration were significantly different across the three voice quality types,  $F=79.024$ ,  $p=.000$ . There was a large effect size,  $\eta^2 = .898$ . Bonferroni-adjusted pairwise comparison indicated strained condition



## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

was longer than both breathy and normal,  $p=.000$ . On average, strained vowels were 0.08 ms longer than the normal breathy vowels. Although breathy vowels were not significantly different than normal vowels ( $p=.052$ ) they were, on average, 0.01ms longer than normal vowels (Tab. 1; Fig. 7). Vowel duration was not held constant across the three conditions, but differences were relatively small (less than 82 ms).




Figure 7. Means and SDs for duration.

### Voice quality measures

**Objective measures.** To determine that the speaker did in fact vary voice quality across the three conditions, one acoustic measure and one physiologic measure were compared. Cepstral peak prominence-smoothed (CPPs) measures were subjected to repeated-measures analysis of variance (ANOVA) with the within-subjects factor of

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

voice type (breathy, normal, or strained). As expected, CPPs differed significantly with voice condition,  $F=100.857$ ,  $p=.000$ ,  $\eta^2 =.918$ . Bonferroni-adjusted pairwise comparison indicated CPPs for the strained voice vowels was larger than for normally voiced vowels, which was larger than the breathy voiced vowels ( $S>N>B$ ), with  $p=.000$  for each comparison (Table 2). This indicates that breathy and strained vocal quality conditions were successfully simulated.

**Table 2**

*Repeated Measures ANOVA with within-subjects factor of voice type (breathy, normal, or strained) statistics for acoustic and physiological measures.*

Measure	F (1,2)	p	$\eta^2_p$
Duration	79	.000*	.9
F1	7	.005*	.4
F2	3	0.961	N/A
F0	3	.100	N/A
CQ50%	17	.000*	.6
CPPs	100	0.000*	0.9

Electroglottogram measures (EGG) of contact quotient with 50% criterion (CQ50%) were subjected to repeated-measures analysis of variance (ANOVA) with within-subject factor of voice type (breathy, normal, or strained). Statistics showed a significant effect for voice type at the midpoint of the vowel,  $F(1,2)=16.777$ ,  $p=.000$  and  $2=.617$  (Table 2). Bonferroni-adjusted pairwise comparison indicated CQ50% at vowel midpoint was significantly different between breathy and strained conditions ( $p=.002$ )

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

and between normal and strained conditions ( $p=.001$ ), but not between breathy and normal conditions (Table 2). Measuring CQ50% over the whole vowel, a significant effect was again present for voice type  $F(1,2)= 5.278$ ,  $p=.016$ ,  $\eta^2 =.370$ . Bonferroni-adjusted pairwise comparison indicated CQ50% across the whole vowel was significantly different between normal and strained conditions ( $p=.006$ ).

Pearson Product Moment correlations for CPPs and CQ50% of the whole vowel (CQ\_all) and CQ50% at the midpoint of the vowel (CQ\_mid) were included to confirm that acoustic and physiological measures were in agreement. The correlation between CPPs and CQ\_all was significant (.525), as was the correlation between CPPs and CQ\_mid (.760).

***Subjective measure.*** To further confirm that the desired voice quality differences were achieved, two licensed speech-language pathologists experienced in voice disorders performed the Consensus Auditory-Perceptual Evaluation of Voice (CAPE-V) on four vowels and four sentences under each condition. The speech-pathologists were blinded to the condition, to each others' ratings, and to the purpose of the present study. The average scores for overall severity, breathiness and strain (Figure 8, Table 3) indicate a moderate severity for the breathy condition, a moderate-to-severe rating for the strained condition and no abnormal quality for the normal condition. For degree of breathiness, the breathy condition was rated as moderately breathy, the strained condition as mildly breathy and the normal condition was rated to be without breathiness. For degree of strain, the breathy condition was rated mildly strained, the normal was rated without strain, and the strained condition was rated moderate-to-severely strained. Therefore, to the trained ear, the appropriate perceptual qualities were present in the stimuli.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION




Figure 8. Average CAPE-V ratings by experienced clinicians. Note Normal condition (orange) does not show in the figure due to zero scores for all measures.

**Table 3**

*Expert clinician judgements on the CAPE-V perceptual test out of 100 points each*

	Breathy	Normal	Strained
Overall Severity	31.5	0	60
Breathiness	55	0	14.5
Strain	7.5	0	66

**Voice quality and vowel identification.** There was no significant effect of voice type,  $F(1,2) = 1.797$ ,  $p = .183$ . The time taken to identify vowels was not significantly different

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

across the three voice quality types,  $F(1,2)=0.51$   $p=.52$ , nor were number of replays  $F(1,2)=3.17$ ,  $p=.081$ .

Comprehensive confusion matrices (Appendix B) show both vowel identification accuracy for each vowel and the nature of confusions. In all conditions, /eɪ/ stimuli had the largest number of errors (9 for breathy, 4 for normal, 6 for strained). This vowel was confused for /ɪ/ and, more often, /ɛ/.

**Table 4**

*Repeated Measures ANOVA with within-subjects factor of voice type (breathy, normal, or strained) for experimental tasks.*

Task	F(1,2)	p	$\eta^2$
Vowel ID	1.797	.183	.107
Vowel ID Time	0.51	0.52	N/A
Vowel ID Replays	3.17	0.081	N/A
Vowel Goodness	11.72	$p<.001^*$	.439
Vowel Goodness Time	15.57	.000*	.509
Vowel Goodness Replays	6.13	.006*	.290
Voice Quality	61.645	.000*	0.804
Voice Quality Time	1.77	.198	N/A
Voice Quality Replays	3.33	.073	N/A

\* Significant at the .05 level

**Vowel goodness ratings.** Vowel goodness ratings were collected to understand how vocal quality affected listeners' perception of phoneme production. Sphericity-assumed results indicate a significant difference,  $F(1,2)=11.72$ ,  $p < .001$ . Bonferroni-adjusted pairwise comparison indicated goodness ratings for the breathy and strained conditions

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

were not significantly different from each other, but were significantly different from the goodness ratings for the normal condition (Table 4, Figure 9). Time needed to rate “goodness” was significantly different  $F(1,2)=15.57$ ,  $p=.000$ ,  $\eta^2 =.509$  (Table 4). Pairwise comparisons indicate that listeners took more time to respond to breathy stimuli than to normal stimuli, with a mean difference of 1411 ms ( $p < .001$ ). Listeners also took significantly more time for strained stimuli compared to normal stimuli, with a mean difference of 683 milliseconds ( $p=.018$ ). Number of replays were significantly different across voice type,  $F(1,2)=6.13$ ,  $p=.006$ ,  $\eta^2 =.290$  (Table 6). Pairwise comparison indicate that listeners used significantly more replays for breathy than for the normal stimuli, with a mean difference of .229 replays ( $p=.008$ ) There was no significant difference between strained and normal vowel stimuli replays (Table 4).




Figure 9. Means and SDs for vowel goodness ratings (0=good, 100=poor).

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

**Voice quality ratings.** Sphericity-assumed results indicate a significant difference across voice type,  $F(1,2)=61.645$ ,  $p=.000$  (Table 4). Bonferroni-adjusted pairwise comparison indicated goodness ratings for the breathy and strained conditions were not significantly different from each other, but both breathy and strained vowels were rated significantly more poorly in voice quality than the normal vowels (Figure 10).

Time needed to rate voice quality did not differ across the three voice quality types,  $F(1,2)=1.77$ ,  $p=.198$ . The number of replays was not significant,  $F(1,2)=3.33$ ,  $p=.073$  (Table 6).




Figure 10. Means and SDs for voice quality ratings (0=good, 100=poor).

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

### Reliability analysis

***Intra-rater agreement.*** Each of the 16 listeners heard each of the 30 vowel tokens three times in each experimental task. Therefore, voice quality and vowel goodness ratings were compared across the three repetitions to assess intra-judge agreement for each listener. Pearson correlations for the first and second set of stimuli, the first and third set of stimuli, and the second and third set of stimuli for each listener (Table 5) Mean intra-rater agreement for voice quality ratings was 85% and for vowel goodness ratings was 56%. Rater 2 was excluded from calculations for vowel goodness because all but one vowel token was rated as 0. For voice quality ratings, one hundred percent of intra-rater correlations for voice quality ratings were statistically significant at the .05 level, and 66% percent for vowel quality ratings were significant (Table 5).

**Table 5**

*Intraclass correlation coefficients by rater and two-way comparisons for vowel goodness task.*

Speaker	Breathy-Strained	Strained-Normal	Breathy-Normal	Mean
1	0.35	0.620329*	0.669*	0.546443
2	N/A	N/A	N/A	N/A
3	-0.06083	-0.05577	0.888608*	0.257338
4	0.064834	0.439371**	0.570384*	0.358197
5	0.421959**	0.267259	0.393916**	0.361044
6	0.608371*	0.508227*	0.776765*	0.631121
7	0.764553*	0.800996*	0.726545*	0.764031
8	0.584298*	0.256765	0.22831	0.356458
9	0.756957*	0.64932*	0.910892*	0.77239



## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

10	0.796281*	0.948324*	0.79225*	0.845618
11	0.358954	0.269817	0.288486	0.305752
12	0.612807*	0.738722*	0.867579*	0.739703
13	0.283751	0.564258*	0.603566*	0.483858
14	0.675691*	0.841362*	0.777172*	0.764741
15	0.457366**	0.730945*	0.525493*	0.571268
16	0.734064*	0.666522*	0.892367*	0.764318
Mean	0.493937	0.549763	0.660756	0.568152
Range	-0.06083	-0.05577	0.22831	0.257338
Minimum				
Range	0.796281	0.948324	0.910892	0.845618
Maximum				

\*indicates significance at .05

\*\* indicates significance at .01

In addition, the percentage of tokens receiving scores of close agreement was calculated. Close agreement was defined as a difference of less than 10 points on the 100 point rating scale for the two ratings. For vowel quality, close agreement for normal vowels was 73%, but for breathy and strained vowels it was 16% and 14% respectively. For vowel goodness, close agreement for normal vowels was 81%, breathy vowels was 51% and strained vowels were 64%.

***Inter-rater reliability.*** To assess agreement across the 16 listeners, interclass correlation coefficients (ICC) (two-way mixed model) were calculated for voice quality and vowel goodness ratings. ICCs are the ratio of rating variance and variance sum and error sum. This measure allows environmental variables and other error sources to be considered leading to a highly generalizable measure of inter-rater reliability (Sheard,

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Adams & Davis, 1991). Absolute agreement ICC for voice quality was .957 ( $p < .001$ ), indicating excellent reliability (Koo & Li, 2016). For vowel goodness ratings, absolute agreement ICC was .726 ( $p < .001$ ), indicating moderate agreement.

**Table 6**

*Interclass correlation coefficients by rater and two-way comparisons for voice quality ratings.*

Speaker	Breathy-Strained	Strained-Normal	Breathy-Normal	Mean
1	0.906*	0.921*	0.875*	0.900667
2	0.717*	0.467*	0.47*	0.551333
3	0.608*	0.635*	0.57*	0.604333
4	0.845*	0.896*	0.805*	0.848667
5	0.782*	0.826*	0.794*	0.800667
6	0.858*	0.885*	0.861*	0.868
7	0.977*	0.992*	0.975*	0.981333
8	0.712*	0.733*	0.72*	0.721667
9	0.923*	0.889*	0.879*	0.897
10	0.972*	0.988*	0.968*	0.976
11	0.973*	0.974*	0.98*	0.975667
12	0.758*	0.824*	0.883*	0.821667
13	0.926*	0.926*	1*	0.950667
14	0.985*	0.993*	0.987*	0.988333
15	0.757*	0.83*	0.751*	0.779333
16	0.931*	0.977*	0.967*	0.958333
Mean	0.851875	0.85975	0.842813	0.851479
Range	0.608	0.467	0.47	0.551333
Minimum				

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Range	0.985	0.993	1	0.988333
Maximum				

\*indicates significance at .05

\*\* indicates significance at .01

### Discussion

#### Summary

The aim of this study was to identify the impact of disordered vocal quality on vowel perception in listeners. In addition to the better-studied articulatory deficits, breathy and strained vocal quality are also common in individuals with dysarthria. As dysarthria is such a common symptom of degenerative neurogenic disorders, it is important to understand exactly what factors affect intelligibility so clinicians can support functional communication with timely, efficacious interventions. Vowel stimuli were used in this pilot study to establish the feasibility of simulated voice quality deficits for future research at word and sentence levels. By using voice quality deficits simulated by a healthy speaker, natural speech tokens could be used in perception tasks.

#### **Did the vowel sets differ in voice quality?**

In order to carry out the study, we had to ensure that the three sets of vowels - breathy normal, and strained - did differ in voice quality. Using the CAPE-V, a standard assessment of voice quality, two experienced clinicians judged that the three sets of vowel stimuli differed from one another. Breathly stimuli were judged as moderately breathy and strained vowel stimuli were judged moderate-to-severely strained. Normal stimuli were judged to be free of perceptual deficits.

Currently, no single measure, acoustic or physiologic, accounts for both breathy and strained vocal quality. Therefore, both acoustic and physiological measures were performed to confirm clinician judgements of perceptual deficits. The acoustic measure CPPs confirmed the breathy and normal stimuli were representative of the vocal quality conditions. As expected, breathy vowels had the lowest CPPs values, indicating lower acoustic energy and periodicity (Heman-Ackah, et al., 2002; Barsties von Latoszek et al.,

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

2016). Strained vowels had the highest CPPs values in our study, indicating higher amounts of acoustic energy, or periodicity, present in the stimuli. This contrasts with Lowell and colleagues (2012), who found strained-pressed phonation had lower CPPs peaks than normal phonation in a sentence-level task and in a sustained vowel task. We are unsure why our measures differed. The efficacy of CPPs measurements for strained vocal quality is limited due to scarcity of research correlating strained phonation and CPPs.

Physiologically, electroglottograph measures were taken by passing an electrical current through the speaker's vocal folds to record when vocal folds are open vs. closed. The included software translated these readings into a contact quotient (CQ) by dividing cycle length by total time the vocal folds were closed. In keeping with prior research, we set the cycle length threshold at 50% (e.g., cycle defined by 50% peak to 50% peak in the EGG signal). CQ50% has been shown to distinguish breathy, strained, and normal phonation (Liu et al., 2017). In women for sustained vowel tasks, vocal folds were found to be closed 35% of the cycle for breathy phonation, 40% of the cycle for normal phonation and 53% of the time for strained phonation (Liu et al., 2017). In our study, vocal folds were found to be closed 40% of the cycle for breathy phonation, 42% of the cycle for normal phonation and 60% of the time for strained phonation. EGG measures showed significant effects of voice type on CQ50%. Strained vowels were significantly different from both normal and breathy vowels at the midpoint of the vowel. Breathly and normal CQ50% were not significantly different from one another, though breathly phonation had a slightly lower CQ50% when looking at the middle three pitch pulses of the EGG signal. It was difficult to measure CQ50% reliably using vowels produced in

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

words with normal duration compared to sustained vowels, like in Liu et al., which may be why our results differ somewhat. Similar to CPPs for strained phonation, CQ50% is not well-suited to measure breathy vocal quality due to the possibility of recording contact due to anterior vocal fold adduction even with a posterior vocal fold gap.

With the exception of the acoustic measure (CPPs) for the strained voice vowels and CQ50% for breathy phonation, the three sets of stimuli differed as expected in expert perceptual judgements, acoustic measures and physiologic measures of voice quality.

### **Were vowels similar in other aspects of production?**

In order to control for confounding variables, it was necessary to hold all factors other than voice quality constant if possible. Several acoustic measures were used to confirm consistency of vowel production. F1 and F2 measures were used to compare tongue height and tongue advancement over time across conditions. F2 at 50% of the vowel duration did not differ significantly, indicating similar tongue advancement for three groups of stimuli. However, there was a significant difference for F1: strained stimuli were about 100 Hz lower than breathy stimuli on average, indicating the tongue and jaw were positioned higher for strained stimuli. Higher positioning may be due to overall increased muscle tension required to produce strained voice, limiting the speaker's ability to lower the jaw. The dynamic vowel formant plot in which F1 and F2 values are shown at 20%, 50%, and 80% of vowel duration for each vowel revealed lowering of /eI/ and raising of /ʊ/ in the strained condition compared to the other two conditions. In addition, /u/ was produced with greater tongue backing in the normal condition compared to the fronted /u/ vowels for the breathy and strained conditions. This varied vowel production may have affected accurate identification of /eI/, which was the

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

most commonly misidentified vowel in the strained condition, though it was also misidentified relatively frequently in breathy and normal conditions. The vowels /ɔ/ and /u/ did not have a high incidence of misidentification in the normal condition, so the formant differences were not large enough to affect their identifiability.

Measures of fundamental frequency ( $f_0$ ) were used to determine whether vocal pitch was held constant across conditions. There was no significant difference for  $f_0$  across quality types, indicating vocal pitch was similar across all three groups of stimuli. Vowel duration was measured to determine vowel length was relatively constant. Strained vowels were an average of 0.08ms longer than normal and breathy vowels. While duration differences were relatively small (about 90 ms longer at most), the large effect size suggested a consistent effect occurring on all ten vowels.

### **Did listeners notice voice quality differences?**

Voice quality was highly salient to listeners. Ratings for strained and breathy conditions were significantly higher (poorer) than for the normally voiced vowels. On average, breathy vowels were rated as poorer than normal vowels by 58 points on a 100 point scale. Strained vowels were rated as poorer than normal stimuli by 68 points and poorer than breathy vowels by 10 points. Listeners had excellent inter-rater and intra-rater reliability for this task. The presence of many "0" ratings for normal stimuli, however, may have inflated the ICC score, however. Time and number of replays needed were not significantly different across the three types of vowels, supporting the idea that listeners made voice quality determinations with ease.

### **Did voice quality affect vowel goodness judgements?**

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Voice quality affected vowel goodness judgements. Both breathy and strained vowels were rated significantly higher (poorer) than vowels in the normal voice quality condition. On average, breathy vowels were rated poorer than normal vowels by 18 points on a 100 point scale and poorer than strained by 2 points. Strained vowels were rated poorer than normal stimuli by 16 points. Inter-rater and intra-rater reliability for this task was worse than for the voice quality rating task. Time to make decision and number of replays needed were significantly different across the three types of vowels. Vowels with breathy vocal quality provoked more replays and listeners took longer to enter ratings than for the normal stimuli. Vowels with strained vocal quality also took longer for listeners to rate, but the number of replays was not significantly greater than for the normal stimuli. Listeners also reported vowel goodness rating task was more difficult than the voice quality rating task. These results suggest that voice quality deficits interfered with judgments of vowel goodness.

### **Did voice quality affect vowel identification?**

Simulated voice quality deficits did not appear to affect vowel identification accuracy, which was above 95% for all conditions. Neither time needed to identify vowels nor number of stimuli replays were significantly different across the three voice quality types. However, the high rates of accuracy (above 95% for all three types of vowel stimuli) suggest that a ceiling effect may have occurred - the task may simply have been too easy to reveal subtle effects of voice quality on vowel identification. Playing the vowel stimuli in noise may reduce accuracy and eliminate the ceiling effect, providing a more accurate understanding of the potential impact of vocal quality deficits on vowel identification (Nabelek, 1988).



## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

### **Limitations**

One of the biggest issues with this study is the use of a single speaker to provide stimuli. Having vowels produced by more speakers would support better generalization of results. Several speakers attempted to perform the stimuli recording, but they were unable to hold vowel production relatively constant while simulating breathy and strained voice quality. Female voices have higher intelligibility overall (Kwon, 2010) and intelligibility in dysarthria may be vulnerable to sex effects, so it will be important to expand to male speakers (Kent et al., 1994). Secondly, the inconsistencies in vowel production with regards to F1 and duration (Table 2) reduces the level of variable control. It is possible that differences in tongue position, duration, or fundamental frequency in addition to the desired voice quality differences contributed to poorer goodness ratings for breathy or strained vowels. Thirdly, the vowel identification task may have been vulnerable to the ceiling effect. Because identification accuracy was so high for all three conditions, a ceiling effect may have masked difficulties in perceiving breathy and strained vowels. Fourthly, our CPP values for strained voice did not match those for Lowell and colleagues (2012) for an unknown reason that needs to be further explored. Current literature on acoustic correlations of strained voice is limited, but using spectral tilt measures like long-term average spectrum (LTAS) may be useful in guiding research. Finally, the focus of this study on vowel perception at the phoneme level limits its applicability to speech intelligibility in clinical practice. The results of this study should be regarded as an early step in understanding the effect of voice quality deficits on real-world communication skills.

### **Clinical Implications**

The results of this study indicate that voice quality deficits are both readily apparent to listeners and difficult to separate from judgments of vowel goodness. We know that vowel perception is important to overall speech intelligibility (Kim et al., 2011; Savageau et al., 2015; Monsen, 1983) and we know listeners need multiple acoustic cues to judge vocal effort (Tasko, 2008), but the effects of disordered vocal quality on vowel perception at the word or sentence level are still largely unknown.

Listeners in the present study did require more time and replays during the relatively simple vowel goodness task for breathy and strained stimuli, suggesting they may have exerted more effort on tokens with disordered voice quality. Sapir and colleagues' (2007) study on vowel goodness comparisons of loud speech and habitual speech in PD may have been confounded by vocal quality changes. The loud speech stimuli likely had better vocal quality than the habitual speech stimuli because the strong vocal fold adduction required to produce high-intensity phonation also reduces breathiness. The findings of the present study support that improved vocal quality may have affected vowel goodness ratings in that study and, therefore, the authors' conclusions reporting improved intelligibility due to louder speech and articulation changes may be incomplete.

If vowel goodness rating is affected by voice quality deficits, it is logical to assume that speech intelligibility will also be impeded by quality deficits due to distraction or possibly increased effort required to understand speech with disordered vocal quality. Research on tracheosophageal speech (Nagle & Eadie, 2012), and dysarthria (Landa, Pennington et al., 2014; Cote-Reschny & Hodge, 2010) have found increased

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

listener effort associated with vocal quality deficits. Listener burden is a particular concern with older populations. As part of the typical, healthy aging process older adults have a higher chance of attention deficits, including both selective attention and sustained attention (Zanto & Gazzaley, 2014). These subtle deficits have been shown to reduce the ability to perform effortful listening tasks, particularly in “suboptimal” conditions (Philips, 2016), which presumably may extend to disordered vocal quality. Considering individual with some dysarthria etiologies are typically older adults and therefore, their communication partners are more likely to be older adults, voice quality could have a larger effect on these populations than expected. These findings support continued research into the effects of voice quality deficits on intelligibility.

### **Directions of Future Research**

Future research should increase task difficulty to eliminate the potential ceiling effects for the vowel identification task. For example, vowel stimuli could be played in noise that more closely replicates real-life conversational settings, improving the external validity of results. Future research may expand tasks to sentence and conversation level stimuli. Utilizing more “real-world” tasks like transcription of sentences produced with simulated breathy and strained voice quality and intelligibility or effort judgments of connected speech with voice quality deficits rather than vowel tasks will further elucidate the impact of voice quality on functional communication. Increasing the number of speakers simulating voice deficits and recruiting a larger sample size of listeners would also improve the generalizability of the results.

In conclusion, this study found some data supporting the hypothesis that voice quality deficits affect vowel perception in listeners. While data did not find increased

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

difficulty with vowel identification in the presence of disordered vocal quality, voice quality was salient to listeners and goodness ratings were more difficult to make quickly when stimuli had quality deficits. Functionally, these findings suggest listeners expend more effort when listening to disordered vocal quality. This may affect overall intelligibility, but study limitations require cautious interpretation until research is expanded.

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

## List of Appendices

Appendix A Training Script.....46

Appendix B Communication Matrices .....47

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

### Appendix A. Study Script for training

#### **Vowel Goodness**

1. Double-click on Alvin3 icon to open program
2. File>Open Experiment
3. Click on appropriate .luax file
4. Enter subject code
5. Click on “start” button
6. Move slider
7. Click on “okay”
8. At end, click “Main menu”

#### Instructions

##### VIDtest-Practice for vowel identification task

- You are about to hear some single-syllable words spoken by a person using different speech techniques. When you hear a word, you will also see ten words on the screen written in standard English and in the International Phonetic Alphabet (IPA). Click on the word that you heard.

##### VGtest-Practice for goodness judgements on single vowel

- You are going to hear some vowels spoken by a person using different speech techniques. You are going to judge the goodness of each vowel you hear. The intended word will appear on the screen. If the vowel you hear is a good example of that vowel, click on the left side of the scale towards “good example.” If the vowel you hear is not a good example of the vowel, click on the right-hand side of the line towards “poor example.” Only pay attention to vowel quality, ignore any elements of nasality, resonance, or vocal quality.

##### QualTest-Practice for goodness judgements on vocal quality

- You are going to hear some words spoken by a person using different speech techniques. You are going to judge the goodness of the vocal quality you hear. If the voice sample you hear is a good example of healthy voice quality, click on the left side of the scale towards “good example.” If the vowel you hear is not a good example of healthy vocal quality, click on the right-hand side of the line towards “poor example.” Only pay attention to vocal quality, ignore any elements of nasality, resonance, or vowel quality.

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Appendix B. *Confusion matrix for breathy vowels, normal vowels, strained vowels*

	i	ɪ	eɪ	ɛ	æ	ʌ	ɑ	oʊ	ʊ	u
i	46	0	2	0	0	0	0	0	0	0
ɪ	0	48	0	0	0	0	0	0	0	0
eɪ	0	4	39	5	0	0	0	0	0	0
ɛ	0	0	0	45	3	0	0	0	0	0
æ	0	0	0	0	48	0	0	0	0	0
ʌ	0	0	0	0	0	47	0	0	1	0
ɑ	0	0	0	0	1	1	46	0	0	0
oʊ	1	0	0	0	0	0	0	47	0	0
ʊ	0	0	0	0	0	1	0	0	47	0
u	0	0	0	0	0	1	0	2	1	44

# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Normal

	i	ɪ	eɪ	ɛ	æ	ʌ	ɑ	oʊ	ʊ	u
i	45	2	1	0	0	0	0	0	0	0
ɪ	0	48	0	0	0	0	0	0	0	0
eɪ	0	1	44	3	0	0	0	0	0	0
ɛ	0	0	0	45	3	0	0	0	0	0
æ	0	0	0	0	48	0	0	0	0	0
ʌ	0	0	0	0	0	48	0	0	0	0
ɑ	0	0	0	0	1	0	47	0	0	0
oʊ	0	0	0	0	0	0	0	47	1	0
ʊ	0	0	0	0	0	1	0	0	47	0
u	0	0	0	0	0	0	0	1	1	46



# VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

Strained

i            ɪ            eɪ            ε            æ            ʌ            ɑ            oʊ            ʊ            u

i	47	0	1	0	0	0	0	0	0	0
ɪ	0	48	0	0	0	0	0	0	0	0
eɪ	0	2	42	4	0	0	0	0	0	0
ε	0	0	0	48	0	0	0	0	0	0
æ	0	0	0	0	48	0	0	0	0	0
ʌ	0	0	0	0	0	48	0	0	0	0
ɑ	0	0	0	0	2	0	46	0	0	0
oʊ	0	0	0	0	0	0	0	47	1	0
ʊ	0	0	0	0	0	1	0	0	47	0
u	0	0	0	0	0	0	0	1	2	45

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

### References

- American Speech and Hearing Association. *Dysarthria*. Retrieved from <http://www.asha.org/public/speech/disorders/dysarthria/>
- Bang Y., Min K., Sohn Y., Cho S. (2013). Acoustic characteristics of vowel sounds in patients with Parkinson disease. *NeuroRehabilitation*, 32, 649–654.
- Barsties v. Latoszek, B., Maryn, Y, Gerrits, E., & De Bodt, M. (2017). The acoustic breathiness index (ABI): A multivariate acoustic model for breathiness. *Journal of Voice*. doi: 10.1016/j.jvoice.2016.11.017.
- Behrman, A. (2018). *Speech and voice science*. 3. San Diego: Plural Publishing.
- Boersma, P., & Weenink, D. (2004). Praat: Doing phonetics by computer (Version 4.3) [Computer software]. Retrieved from <http://www.praat.org>.
- Boshes, B. (1996). Voice changes in Parkinsonism. *Journal of Neurosurgery*, 24, 286-288.
- Bunton, K. (2006). Fundamental frequency as a perceptual cue for vowel identification in speakers with Parkinson's disease. *Folia Phoniatica et Logopaedica*, 58(5), 323-339. DOI: [10.1159/000094567](https://doi.org/10.1159/000094567)
- Caligiuri, M. (1989). The influence of speaking rate on articulatory hypokinesia in parkinsonian dysarthria. *Brain and Language*, 36(3).
- Canter, G. J. (1963). Speech characteristics of patients with parkinson's disease: I Intensity, pitch, and duration. *J Speech Hear Disord*, 28(3), 221-229. doi: 10.1044/jshd.2803.221.
- Canter, G. J. (1965). Speech characteristics of patients with parkinson's disease: III. Articulation, diadochokinesis, and over-all speech adequacy. *J Speech Hear Disord*, 30(3), 217-224. doi: 10.1044/jshd.3003.217.
- Cleveland, L., Pimentel, J., Talarico, M. & Solomon, J. (2015). Vocal intensity and speech intelligibility pre- and post-SPEAK OUT!® treatment. Poster presented at the annual American Speech-Language-Hearing Association Convention, Denver, CO.
- Connor, N.P., Abbs, J., Cole, K., Gracco, V. (1989). Parkinsonian deficits in serial multiarticulate movements for speech. *Brain*, 112, 997-1009.
- Constantinescu, G., Theodoros, D., Russell, T., Ward, E., Wilson, S., & Wootton, R. (2011). Treating disordered speech and voice in parkinson's disease online: A

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

- randomized controlled non-inferiority trial. *International Journal of Language & Communication Disorders*, 46(1), 1-16. doi:10.3109/13682822.2010.484848
- Cote-Reschny, K. J., & Hodge, M. M. (2010). Listener effort and response time when transcribing words spoken by children with dysarthria. *Journal of Medical Speech-Language Pathology*, 18(4), 24-35.
- Darley, F. L., Aronson, A. E., & Brown, J. R. (1969). Clusters of deviant speech dimensions in the dysarthrias. *J Speech Hear Res*, 12(3), 462-496. doi: 10.1044/jshr.1203.462.
- De Bodt, M., Hernández-Díaz, M., Van De Heyning, P. (2002). Intelligibility as a linear combination of dimensions in dysarthric speech. *Journal of Communication Disorders*, 35(3). 283-292.
- Doyle, P. C., Danhauer, J. L., & Reed, C. G. (1988). Listeners' perceptions of consonants produced by esophageal and tracheoesophageal talkers. *J Speech Hear Disord*, 53(4), 400-407. doi: 10.1044/jshd.5304.400.
- Duffy, J. (2013). *Motor speech disorders : Substrates, differential diagnosis, and management*. St. Louis, Mo.: Elsevier Mosby.
- Dromey, C., & Ramig, L. (1998). Intentional changes in sound pressure level and rate: Their impact on measures of respiration, phonation, and articulation. *Journal Speech and Hearing Research*, 41, 1003–1018.
- Dromey, C., Ramig, L. O., & Johnson, A. B. (1995). Phonatory and articulatory changes associated with increased vocal intensity in Parkinson disease: A case study. *Journal of Speech, Language, and Hearing Research*, 38, 751–764. <https://doi.org/10.1044/jshr.3804.751>
- Doyle, P., Danhauer, J. L., & Reed, C. G. (1988). Listeners' perceptions of consonants produced by esophageal and tracheoesophageal talkers. *Journal of Speech & Hearing Disorders*, 53(4), 400-407. <http://dx.doi.org/10.1044/jshd.5304.400>
- Eadie, T., Day, A., Swain D., Lamvik, K., & Doyle, P. (2013). Auditory-perceptual speech outcomes and quality of life after total laryngectomy. *Otolaryngology – Head and Neck Surgery*.
- Evitts, P., Starmer, H., Teets, K., Montgomery, C., Calhoun, L., Schulze, A., MacKenzie, A., & Adams, L. (2016). The impact of dysphonic voices on healthy listeners: Listener reaction times, speech intelligibility and listener comprehension. *American Journal of Speech-Language Pathology*, 25, 561-575.
- Fant, G (1960). *Acoustic theory of speech production*. The Hague: Mouton.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

- Franklin, A. D. & Stoel-Gammon, C. (2014). Using multiple measures to document change in English vowels produced by Japanese, Korean, and Spanish speakers: The case for goodness and intelligibility. *Am J Speech Lang Pathology*, 23(4), 625-640. doi: 10.1044/2014\_AJSLP-13-0144.
- Ferrand, C. T. (2018). *Speech science: an integrated approach to theory and clinical practice*. NY, NY: Pearson.
- Gayvert, R., & Hillenbrand, J. (2003). Open source software for speech perception research. *The Journal of the Acoustical Society of America*, 113, 2260.
- Gazewood, J., Richards, R., & Clebak, K. (2013). Parkinson disease: An update. *American Family Physician*, 87(4), 267-73.
- Greene, D.S. (1956). Fundamental frequency of the speech of the profoundly deaf individuals: Unpublished doctoral dissertation. Purdue University, West Lafayette, IN
- Hansen, D., Gerrat B., & Ward. P. (1984). Cinegraphic observations of laryngeal function in Parkinson's disease. *Laryngoscope* 94, 348-353.
- Haycock, G. (1933). *The teaching of speech*. Stoke-on-Trent: Hill & Ainsworth.
- Heman-Ackah, Y., Michael, D., & Goding, G. (2002). The relationship between cepstral peak prominence and selected parameters of dysphonia. *Journal of Voice*, 16(1), 20-27. doi:10.1016/S0892-1997(02)00067-X
- Henton, C. G., & Bladon, R. A. W. (1985). Breathiness in normal female speech: Insufficiency versus desirability. *Language and Communication: An Interdisciplinary Journal* 5(3), 221-227.
- Hillenbrand, J., Cleveland, R. A., & Erickson, R. L. (1994). Acoustic correlates of breathy vocal quality. *Journal of Speech Language and Hearing Research*, 37(4), 769.
- Hillenbrand, J. M., & Gayvert, R. T. (2005). Open source software for experiment design and control. *Journal of Speech, Language and Hearing Research*, 48(1), 45–60.
- Hillenbrand, J. & Houde, R. (1997). Acoustic correlates of breathy vocal quality: dysphonic voices and continuous speech. *Journal of Speech Language and*

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

*Hearing Research*, 39, 311– 321.

Ho, A., Bradshaw, J., Ianseck, R., & Alfredson, R. (1999). Speech volume regulation in Parkinson's disease: Effects of implicit cues and explicit instructions.

*Neuropsychologia*, 37(13).

Hood, R.B. (1966). Some physical concomitants of the perception of speech rhythm of the deaf: Unpublished doctoral dissertation. Stanford University: Stanford, CA.

Hustad, K. (2008). The relationship between listener comprehension and intelligibility scores for speakers with dysarthria. *Journal of Speech, Language, and Hearing Research*, 51, 562–573.

IBM Corp. (2013). IBM SPSS Statistics for Windows (Version 22.0) [Computer software]. Armonk, NY: IBM Corp.

Jacewicz, E., Fox, R., & Salmons, J. (2006). Prosodic prominence effects on vowels in chain shifts. *Language Variation and Change*, 18(3), 285-316.

Kent, R. D., & Read, C. (1992). *The acoustic analysis of speech*. San Diego, CA: Singular Publishing Group.

Kent, R., & Rosenbek, J. (1982). Prosodic disturbance and neurologic lesion. *Brain and Language*, 15, 259-291. 10.1016/0093-934X(82)90060-8.

Kim, H., Hasegawa-Johnson, M., & Perlman, A. (2011). Vowel contrast and speech intelligibility in dysarthria. *Folia Phoniatrica et Logopaedica*, 63, 187–194.

Kitzing, P. (1986). LTAS criteria pertinent to the measurement of voice quality. *Journal of Phonetics*, 14, 477-482.

Koo, T. K., & Li, M. Y. (2016). A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *Journal of Chiropractic Medicine*, 15(2), 155–163. <http://doi.org/10.1016/j.jcm.2016.02.012>

Kreiman, J. , Vanlancker-Sidtis, D. & Gerratt, B. R. (2008). Perception of voice quality. *The Handbook of Speech Perception*. doi:10.1002/9780470757024.

Lam J., Tjaden K. (2016). Clear speech variants: An acoustic study in parkinson's disease. *Journal of Speech, Language, and Hearing Research*, 59(4), 631-646. doi:10.1044/2015\_JSLHR-S-15-0216.

Lambacher, S. G., Martens, W. L., Kakehi, K., Margasinghe, C. A., & Molholt, G. (2005). The effects of identification training on the identification and production

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

of American English vowels by native speakers of Japanese. *Applied Psycholinguistics*, 26, 227–247.

- Landa, S., Pennington, L., Miller, N., Robson, S., Thompson, V., & Steen, N. (2014). Association between objective measurement of the speech intelligibility of young people with dysarthria and listener ratings of ease of understanding. *International Journal of Speech-Language Pathology*, 16(4), 408-416.
- Laures, J. S. & Weismer, G. (1999). The effects of a flattened fundamental frequency on intelligibility at the sentence level. *J Speech Lang Hear Research*, 42(5), 1148-1156. doi: 10.1044/jslhr.4205.1148.
- Laver, J. (1980). The phonetic description of voice quality. *Cambridge studies in linguistics*, 31. Cambridge England: Cambridge University Press.
- Liu, D., Kankare, E., Laukkanen, A., & Alku, P. (2017). Comparison of parametrization methods of electroglottographic and inverse filtered acoustic speech pressure signals in distinguishing between phonation types. *Biomedical Signal Processing and Control*, 36, 183-193. doi:10.1016/j.bspc.2017.04.001
- Logemann, J., & Fisher, H. (1981). Vocal tract control in parkinson's disease. *Journal of Speech and Hearing Disorders*, 46(4), 348-348. doi:10.1044/jshd.4604.348
- Lowell, S. Y., Kelley, R. T., Awan, S. N., Colton, R. H., & Chan, N. H. (2012). Spectral- and cepstral-based acoustic features of dysphonic, strained voice quality. *Annals of Otology, Rhinology and Laryngology*, 121(8), 539-548.
- Ludlow, C.L., & Bassich, C.J. (1983). The results of acoustic and perceptual assessment of two types of dysarthria. *Clinical dysarthria*, 121-153. San Diego: College-Hill Press.
- Mazonni, D. (1999). *Audacity: Free audio editor and recorder*. [Computer software]. Retrieved from <http://audacity.sourceforge.net/>.
- Martel-Sauvageau, V., Roy, J. P., Langlois, M., & Macoir, J. (2015). Impact of the LSVT on vowel articulation and coarticulation in Parkinson's disease. *Clinical Linguistics & Phonetics*, 29(6), 424–440
- Michaelis, D., Gramss, T., & Strube, H. (1997) Glottal-to-noise excitation ratio: A new measure for describing pathological voices. *Acustica/Acta Acustica*, 83, 700–706.
- Miller, N., Noble, E., Jones, D., & Burn, D. (2006). Life with communication changes in Parkinson's disease. *Age and Ageing*, 35(3), 235–239. <https://doi.org/10.1093/ageing/afj053>
- Monsen, R. (1979). Acoustic qualities of phonation in young hearing-impaired children.

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

*Journal of Speech and Hearing Research* 48. 270-288.

- Monsen, R. (1983). Voice quality and speech intelligibility among deaf children. *American Annals of the Deaf*, 12-19.
- Nabelek, A.K., & Letowski, T.R. (1988). Similarities of vowels in nonreverberant and reverberant fields. *The Journal of the Acoustical Society of America*, 83 5, 1891-9.
- Nagle, K. F., & Eadie, T. L. (2012). Listener effort for highly intelligible tracheoesophageal speech. *Journal of Communication Disorders*, 45(3), 235-245.
- Neel, A.T. (2009). Intelligibility of loud, amplified, and habitual speech in Parkinson Disease, *Journal of Speech, Language, and Hearing Research*, 52, 1021-1033.
- Neel, A.T. & Beveridge, A.C. (2006, November). Identification of loud- and habitual-level vowels in Parkinson disease. Poster presented at the American Speech-Language-Hearing Convention, Miami, FL
- Noll, A. (1967). Cepstrum pitch determination. *The Journal of the Acoustical Society of America*, 41(2), 293-309. doi:10.1121/1.1910339
- Peterson, G.E. & Barney, H. (1952). Control methods used in a study of the vowels. *Journal of the Acoustical Society of America* 24. 175–184.
- Phillips, N. (2016). The implications of cognitive aging for listening and the FUEL model. *Ear Hear* 37, 44–51.
- Platt, L. J., Andrews, G., & Howie, P. M. (1980). Dysarthria of adult cerebral palsy: Phonemic analysis of articulation errors. *Journal of Speech and Hearing Research*, 23, 41–55.
- Ramig, L. (1992) The role of phonation in speech intelligibility: A review and preliminary data from patients with Parkinson's disease. *Intell. Speech Disorders Theory, Measurement, and Management*. 119-155.
- Rogers, C. L., & Dalby, J. (2005). Forced-choice analysis of segmental production by Chinese-accented English speakers. *Journal of Speech, Language, and Hearing Research*, 48, 306–322.
- Rusz J., Cmejla R., Tykalova T., Ruzickova H., Klempir J., Majerova V., et al. (2013). Imprecise vowel articulation as a potential early marker of Parkinson's disease: effect of speaking task. *J. Acoust. Soc. Am.*, 134(3). 2171–81.
- Salkind, N. (2010). *Encyclopedia of research design*. Thousand Oaks, Calif.: Sage.
- Samlan, R., Story, B., & Bunton, K. (2013) Relation of perceived breathiness to laryngeal

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

- kinematics and acoustic measures based on computational modeling. *Journal of Speech, Language, & Hearing Research*, 56, 1209–1223. [doi.org/10.1044/1092-4388\(2012/12-0194\)](https://doi.org/10.1044/1092-4388(2012/12-0194))
- Sapir S., Ramig L. O., & Fox C. M. (2011). Intensive voice treatment in Parkinson's disease: Lee Silverman Voice Treatment. *Expert Review of Neurotherapeutics*, 11(6), 815–830. <https://doi.org/10.1586/ern.11.43>.
- Sapir, S., Spielman, J., Ramig, L., Story, B., & Fox, C. (2007). Effects of intensive voice treatment (LSVT®) on vowel articulation in dysarthric individuals with idiopathic Parkinson's disease: Acoustic and perceptual findings. *Journal of Speech Language and Hearing Research*, 50, 899-912.
- Schulman, R. (1989). Articulatory dynamics of loud and normal speech. *Journal Acoustic Society of America*, 85, 295–312.
- Sheard, C., Adams, R. D., & Davis, P. J. (1991). Reliability and agreement of ratings of ataxic dysarthric speech samples with varying intelligibility. *Journal of Speech Language and Hearing Research*, 34(2), 285.
- Skodda, S., Flasskamp, A., & Schlegel, U. (2011a). Instability of syllable repetition in parkinson's disease—Influence of levodopa and deep brain stimulation. *Movement Disorders*, 26(4), 728-730. doi:10.1002/mds.23439
- Skodda S, Visser W, Schlegel U. (201b). Vowel articulation in Parkinson's Disease. *J. Voice*, 25, 467–472.
- Skodda S., Gronheit W., Schlegel U. (2012) Impairment of vowel articulation as a possible marker of disease progression in parkinson's disease. *PLoS ONE*, 7(2).
- Smith, M., & Ramig, L. (1995) Intensive voice treatment in Parkinson's disease: Laryngostroboscopic findings. *J Voice*, 9. 453-459.
- Smith, S., & Shaffer, D. (1995). Speed of speech and persuasion: Evidence for multiple effects. *Personality and Social Psychology Bulletin*, 21, 1051-1060. doi:10.1177/01461672952110006.
- Spector, P., Subtelney, J., Whitehead, R., & Wirz, S. (1979). Description and evaluation of a training program to reduce vocal tension in adult deaf speakers. *The Volta Review*, 81(2), 81-90.
- Stemple, J. C., Roy, N., & Klaben, B. (2014). *Clinical voice pathology: theory and management*. San Diego, CA: Plural Publishing.
- Sussman, J. E., & Tjaden, K. (2012). Perceptual measures of speech from individuals



## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

with Parkinson's Disease and multiple sclerosis: Intelligibility and beyond. *Journal of Speech Language and Hearing Research*, 55(4), 1208.

Tasko, S., Parker, M., & Hillenbrand, J. (2008). Acoustic correlates of vocal effort. *The Journal of the Acoustical Society of America*, 124(4), 2496-2496.

Titze, I. (2000). *Principles of voice production*. Iowa City, Ia: National Center for Voice and Speech.

Tjaden, K. (2008). Speech and swallowing in Parkinson's Disease. *Topics in Geriatric Rehabilitation*, 24(2), 115-126.

<http://doi.org/10.1097/01.TGR.0000318899.87690.44>

Tykalova, T., Rusz, J., Klempir, J., Cmejla, R., & Ruzicka, E. (2017). Distinct patterns of imprecise consonant articulation among parkinson's disease, progressive supranuclear palsy and multiple system atrophy. *Brain and Language*, 165, 1-9. doi:10.1016/j.bandl.2016.11.005

Ward, E. C., Theodoros, D. G., & Murdoch, B. (2000). Changes in articulatory pressures following the Lee Silverman Voice Treatment Program. Paper presented at the Conference on Motor Speech, San Antonio, TX.

Watts, C. R. (2016). A retrospective study of long-term treatment outcomes for reduced vocal intensity in hypokinetic dysarthria. *BMC Ear, Nose, and Throat Disorders*, 16, 2. <http://doi.org/10.1186/s12901-016-0022-8>.

Watts, C.R. & Awan, S.N. (2011). Use of spectral/cepstral analyses for differentiating normal from hypofunctional voices in sustained vowel and continuous speech contexts. *Journal of Speech, Language, and Hearing Research*, 54(6), 1525-37. doi:10.1044/1092-4388.

Weismer G., Kent R., Hodge M., & Martin R (1988). The acoustic signature for intelligibility test words. *The Journal of the Acoustical Society of America*. 84. 1281-91.

Zanto T. P., Gazzaley A. A. C. Nobre, S. Kastner. (2014). Attention and ageing: The Oxford handbook of attention. Oxford, United Kingdom: *Oxford University Press*. 927-971.

Zraick, R. I., Kempster, G. B., Connor, N. P., Thibeault, S., Klaben, B. K., Bursac, Z., & Glaze, L. E. (2011). Establishing validity of the consensus auditory-perceptual evaluation of voice (CAPE-V). *American Journal of Speech-Language Pathology*, 20(1), 14-22. doi: [10.1044/1058-0360\(2010/09-0105\)](https://doi.org/10.1044/1058-0360(2010/09-0105)).

Zwirner, P., & Barnes, G. (1992). Vocal tract steadiness: A measure of phonatory and

## VOCAL QUALITY DEFICITS AND VOWEL PERCEPTION

upper airway motor control during phonation in dysarthria. *Journal of Speech and Hearing Research*, 35(4), 761-