

1-1-2015

Basic Structure of Some Classes of Neutrosophic Crisp Nearly Open Sets & Possible Application to GIS Topology

A. Salama

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Salama, A.. "Basic Structure of Some Classes of Neutrosophic Crisp Nearly Open Sets & Possible Application to GIS Topology." *Neutrosophic Sets and Systems* 7, 1 (2019).
https://digitalrepository.unm.edu/nss_journal/vol7/iss1/4

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu.



Basic Structure of Some Classes of Neutrosophic Crisp Nearly Open Sets & Possible Application to GIS Topology

A. A. Salama

Department of Mathematics and Computer Science, Faculty of Sciences, Port Said University, Egypt

drsalama44@gmail.com

Abstract. Since the world is full of indeterminacy, the neutrosophics found their place into contemporary research. The fundamental concepts of neutrosophic set, introduced by Smarandache in [30, 31, 32] and Salama et al. in [4-29]. In Geographical information systems (GIS) there is a need to model spatial regions with indeterminate boundary and under indeterminacy. In this paper the structure of some classes of neutrosophic

ic crisp nearly open sets are investigated and some applications are given. Finally we generalize the crisp topological and intuitionistic studies to the notion of neutrosophic crisp set. Possible applications to GIS topological rules are touched upon.

Keywords: Neutrosophic Crisp Set; Neutrosophic Crisp Topology; Neutrosophic Crisp Open Set; Neutrosophic Crisp Nearly Open Set; Neutrosophic GIS Topology.

1 Introduction

The fundamental concepts of neutrosophic set, introduced by Smarandache [30, 31, 32] and Salama et al. in [4-29], provides a natural foundation for treating mathematically the neutrosophic phenomena which exist pervasively in our real world and for building new branches of neutrosophic mathematics. Neutrosophy has laid the foundation for a whole family of new mathematical theories generalizing both their classical and fuzzy counterparts [1, 2, 3, 20, 21, 22, 23, 34] such as a neutrosophic set theory. In this paper the structure of some classes of neutrosophic crisp sets are investigated, and some applications are given. Finally we generalize the crisp topological and intuitionistic studies to the notion of neutrosophic crisp set.

2 Terminologies

We recollect some relevant basic preliminaries, and in particular, the work of Smarandache in [30, 31, 32] and Salama et al. [4-29]. Smarandache introduced the neutrosophic components T, I, F which represent the membership, indeterminacy, and non-membership values respectively, where $]0, 1^+$ is non-standard unit interval. Salama et al. [9, 10, 13, 14, 16, 17] considered some possible definitions for basic concepts of the neutrosophic crisp set and its operations. We now improve some results by the following. Salama extended the concepts of topological space and in-

tuitionistic topological space to the case of neutrosophic crisp sets.

Definition 1. 2 [13]

A neutrosophic crisp topology (NCT for short) on a non-empty set X is a family Γ of neutrosophic crisp subsets in X satisfying the following axioms

- i) $\phi_N, X_N \in \Gamma$.
- ii) $A_1 \cap A_2 \in \Gamma$ for any A_1 and $A_2 \in \Gamma$.
- iii) $\cup A_j \in \Gamma \quad \forall \{A_j : j \in J\} \subseteq \Gamma$.

In this case the pair (X, Γ) is called a neutrosophic crisp topological space (NCTS for short) in X . The elements in Γ are called neutrosophic crisp open sets (NCOSs for short) in X . A neutrosophic crisp set F is closed if and only if its complement F^C is an open neutrosophic crisp set.

Let (X, Γ) be a NCTS (identified with its class of neutrosophic crisp open sets), and $NCint$ and $NCcl$ denote neutrosophic interior crisp set and neutrosophic crisp closure with respect to neutrosophic crisp topology

3 Nearly Neutrosophic Crisp Open Sets

Definition 3.1

Let (X, Γ) be a NCTS and $A = \langle A_1, A_2, A_3 \rangle$ be a NCS in X , then A is called

Neutrosophic crisp α -open set iff $A \subseteq NC \text{int}(NCcl(NC \text{int}(A)))$,

Neutrosophic crisp β -open set iff $A \subseteq NCcl(NC \text{int}(A))$.

Neutrosophic crisp semi-open set iff $A \subseteq NC \text{int}(NCcl(A))$.

We shall denote the class of all neutrosophic crisp α -open sets NCT^α , the calls all neutrosophic crisp β -open sets NCT^β , and the class of all neutrosophic crisp semi-open sets NCT^s .

Remark 3.1

A class consisting of exactly all a neutrosophic crisp α -structure (resp. $NC \beta$ -structure). Evidently $NCT \subseteq NCT^\alpha \subseteq NCT^\beta$.

We notice that every non-empty neutrosophic crisp β -open has NC^α -nonempty interior. If all neutrosophic crisp sets the following $\{B_i\}_{i \in I}$ are $NC \beta$ -open sets, then $\{\cup_{i \in I} B_i\}_{i \in I} \subset NCcl(NC \text{int}(B_i)) \subset NCcl(NC \text{int}(B_i))$, that is A $NC \beta$ -structure is a neutrosophic closed with respect to arbitrary neutrosophic crisp unions.

We shall now characterize NCT^α in terms NCT^β

Theorem 3.1

Let (X, Γ) be a NCTS. NCT^α Consists of exactly those neutrosophic crisp set A for which $A \cap B \in NCT^\beta$ for $B \in NCT^\beta$

Proof

Let $A \in NCT^\alpha$, $B \in NCT^\beta$, $p \in A \cap B$ and U be a neutrosophic crisp neighbourhood (for short $NCnbd$) of p . Clearly $U \cap NC \text{int}(NCcl(NC \text{int}(A)))$, too is a neutrosophic crisp open neighbourhood of p , so $V = (U \cap NC \text{int}(NCcl(NC \text{int}(A)))) \cap NC \text{int}(B)$ is non-empty. Since $V \subset NCcl(NC \text{int}(A))$ this implies $(U \cap NC \text{int}(A) \cap NC \text{int}(B)) = V \cap NC \text{int}(A) = \phi_N$. It follows that $A \cap B \subset NCcl(NC \text{int}(A) \cap NC \text{int}(B)) = NCcl(NC \text{int}(A \cap B))$ i.e. $A \cap B \in NCT^\beta$. Conversely, let

$A \cap B \in NCT^\beta$ for all $B \in NCT^\beta$ then in particular $A \in NCT^\beta$. Assume that $p \in A \cap (NC \text{int}(NCcl(A) \cap NC \text{int}(A)))^c$. Then $p \in NCcl(B)$, where $(NCcl(NC \text{int}(A)))^c$. Clearly $\{p\} \cup B \in NCT^\beta$, and consequently $A \cap \{\{p\} \cup B\} \in NCT^\beta$. But $A \cap \{\{p\} \cup B\} = \{p\}$. Hence $\{p\}$ is a neutrosophic crisp open. As $p \in (NCcl(NC \text{int}(A)))$ this implies $p \in NC \text{int}(NCcl(NC \text{int}(A)))$, contrary to assumption. Thus $p \in A$ implies $p \in NCcl(NC \text{int}(A))$ and $A \in NCT^\alpha$. This completes the proof. Thus we have found that NCT^α is complete determined by NCT^β i.e. all neutrosophic crisp topologies with the same $NC \beta$ -structure also determined the same NC^α -structure, explicitly given Theorem 3.1.

We shall that conversely all neutrosophic crisp topologies with the same NC^α -structure, so that NCT^β is completely determined by NCT^α .

Theorem 3.2

Every neutrosophic crisp NC^α -structure is a neutrosophic crisp topology.

Proof

NCT^β Contains the neutrosophic crisp empty set and is closed with respect to arbitrary unions. A standard result gives the class of those neutrosophic crisp sets A for which $A \cap B \in NCT^\beta$ for all $B \in NCT^\beta$ constitutes a neutrosophic crisp topology, hence the theorem. Hence forth we shall also use the term NC^α -topology for NC^α -structure two neutrosophic crisp topologies deterring the same NC^α -structure shall be called NC^α -equivalent, and the equivalence classes shall be called NC^α -classes.

We may now characterize NCT^β in terms of NCT^α in the following way.

Proposition 3.1

Let (X, Γ) be a NCTS. Then $NCT^\beta = NCT^{\alpha\beta}$ and hence NC^α -equivalent topologies determine the same $NC \beta$ -structure.

Proof

Let $NC^\alpha-cl$ and $NC^\alpha-int$ denote neutrosophic closure and Neutrosophic crisp interior with respect to

$N\Gamma^\alpha$. If $p \in B \in N\Gamma^\beta$ and $p \in B \in N\Gamma^\alpha$, then $(NC\text{int}(NCcl(NC\text{int}(A))) \cap NC\text{int}(B)) \neq \emptyset_N$. Since $NC\text{int}(NCcl(NC\text{int}(A)))$ is a crisp neutrosophic neighbourhood of point p . So certainly $NC\text{int}(B)$ meets $NCcl(NC\text{int}(A))$ and therefore (being neutrosophic open) meets $NC\text{int}(A)$, proving $A \cap NC\text{int}(B) \neq \emptyset_N$ this means $B \subset NC\alpha cl(NC\text{int}(B))$ i.e. $B \in N\Gamma^{\alpha\beta}$ on the other hand let $A \in N\Gamma^{\alpha\beta}$, $p \in A$ and $p \in V \in N\Gamma$. As $V \in N\Gamma^\alpha$ and $p \in NCcl(NC\text{int}(A))$ we have $V \cap NC\text{int}(A) \neq \emptyset_N$ and there exist a neutrosophic crisp set $W \in \Gamma$ such that $W \subset V \cap NC\alpha\text{int}(A) \subset A$.

In other words $V \cap NC\text{int}(A) \neq \emptyset_N$ and $p \in NCcl(NC\text{int}(A))$. Thus we have verified $N\Gamma^{\alpha\beta} \subset N\Gamma^\alpha$, and the proof is complete combining Theorem 1 and Proposition 1. We get $N\Gamma^{\alpha\alpha} = N\Gamma^\alpha$.

Corollary 3.2

A neutrosophic crisp topology $N\Gamma$ a $NC\alpha$ -topology iff $N\Gamma = N\Gamma^\alpha$. Thus an $NC\alpha$ -topology belongs to the $NC\alpha$ -class if all its determining a Neutrosophic crisp topologies, and is the finest topology of finest neutrosophic topology of this class. Evidently $N\Gamma^\beta$ is a neutrosophic crisp topology iff $N\Gamma^\alpha = N\Gamma^\beta$. In this case $N\Gamma^{\beta\beta} = N\Gamma^{\alpha\beta} = N\Gamma^\beta$.

Corollary 3.3

$NC\beta$ -Structure B is a neutrosophic crisp topology, then $B = B^\alpha = B^\beta$.

We proceed to give some results an the neutrosophic structure of neutrosophic crisp $NC\alpha$ -topology

Proposition 3.4

The $NC\alpha$ -open with respect to a given neutrosophic crisp topology are exactly those sets which may be written as a difference between a neutrosophic crisp open set and neutrosophic crisp nowhere dense set

If $A \in N\Gamma^\alpha$ we have $A = (NC\text{int}(NCcl(NC\text{int}(A))) \cap (NC\text{int}(NCcl(NC\text{int}(A))) \cap A^C)^C$, where $(NC\text{int}(NCcl(NC\text{int}(A))) \cap A^C)$ clearly is neutrosophic crisp nowhere dense set, we easily see that $B \subset NCcl(NC\text{int}(A))$ and consequently $A \subset B \subset NC\text{int}(NCcl(NC\text{int}(A)))$ so the proof is complete.

Corollary 3.4

A neutrosophic crisp topology is a $NC\alpha$ -topology iff all neutrosophic crisp nowhere dense sets are neutrosophic crisp closed.

For a neutrosophic crisp $NC\alpha$ -topology may be characterized as neutrosophic crisp topology where the difference between neutrosophic crisp open and neutrosophic crisp nowhere dense set is again a neutrosophic crisp open, and this evidently is equivalent to the condition stated.

Proposition 3.5

Neutrosophic crisp topologies which are $NC\alpha$ -equivalent determine the same class of neutrosophic crisp nowhere dense sets.

Definition 3.2

We recall a neutrosophic crisp topology a neutrosophic crisp extremely disconnected if the neutrosophic crisp closure of every neutrosophic crisp open set is a neutrosophic crisp open.

Proposition 3.6

If $NC\alpha$ -Structure B is a neutrosophic crisp topology, all a neutrosophic crisp topologies Γ for which $\Gamma^\beta = B$ are neutrosophic crisp extremely disconnected.

In particular: Either all or none of the neutrosophic crisp topologies of a $NC\alpha$ -class are extremely disconnected.

Proof

Let $\Gamma^\beta = B$ and suppose there is a $A \in \Gamma$ such that $NCcl(A) \notin \Gamma$. Let $p \in NCcl(A) \cap (NC\text{int}(NCcl(A)))^C$ with $B = \{p\} \cup NC\text{int}(NCcl(A))$, $M = (NC\text{int}(NCcl(A)))^C$ we have $\{p\} \subset M = (NC\text{int}(NCcl(A)))^C = NCcl(NC\text{int}(M))$, $\{p\} \subset NCcl(A) = NCcl(NC\text{int}(NCcl(A))) \subset NCcl(NC\text{int}(B))$. Hence both B and M are in Γ^β . The intersection $B \cap M = \{p\}$ is not neutrosophic crisp open since $p \in NCcl(A) \cap M^C$, hence not $NC\beta$ -open so $\Gamma^\beta = B$ is not a neutrosophic crisp topology. Now suppose B is not a topology, and $\Gamma^\beta = B$. There is a $B \in \Gamma^\beta$ such that $B \notin \Gamma^\alpha$. Assume that $NCcl(NC\text{int}(B)) \in \Gamma$. Then $B \subset NCcl(NC\text{int}(B)) = NC\text{int}(NCcl(NC\text{int}(B)))$ i.e. $B \in \Gamma^\alpha$, contrary to assumption. Thus we have produced a neutrosophic crisp open set whose neutrosophic crisp closure is not neutrosophic crisp open, which completes the proof.

Corollary 3.5

A neutrosophic crisp topology Γ is a neutrosophic crisp extremely disconnected if and only if Γ^β is a neutrosophic crisp topology.

4 Conclusion and future work

Neutrosophic set is well equipped to deal with missing data. By employing NSs in spatial data models, we can express a hesitation concerning the object of interest. This article has gone a step forward in developing methods that can be used to define neutrosophic spatial regions and their relationships. The main contributions of the paper can be described as the following: Possible applications have been listed after the definition of NS. Links to other models have been shown. We are defining some new operators to describe objects, describing a simple neutrosophic region. This paper has demonstrated that spatial object may profitably be addressed in terms of neutrosophic set. Implementation of the named applications is necessary as a proof of concept.

References

- [1] K. Atanassov, intuitionistic fuzzy sets, in V.Sgurev, ed., VII ITKRS Session, Sofia (June 1983 central Sci. and Techn. Library, Bulg. Academy of Sciences (1984).
- [2] K. Atanassov, intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 20, 87-96,(1986).
- [3] K. Atanassov, Review and new result on intuitionistic fuzzy sets, preprint IM-MFAIS-1-88, Sofia, (1988).
- [4] S. A. Alblowi, A.A.Salama and Mohamed Eisa, New Concepts of Neutrosophic Sets, *International Journal of Mathematics and Computer Applications Research (IJMCAR)*, Vol. 4, Issue 1, (2014)pp59-66.
- [5] I.M. Hanafy, A.A. Salama and K. Mahfouz, Correlation of Neutrosophic Data, *International Refereed Journal of Engineering and Science (IRJES)* , Vol.(1), Issue 2 .(2012) PP.39-33
- [6] I.M. Hanafy, A.A. Salama and K.M. Mahfouz, Neutrosophic Classical Events and Its Probability, *International Journal of Mathematics and Computer Applications Research(IJMCAR)* Vol.(3),Issue 1, (2013)pp171-178.
- [7] A. A. Salama and S.A. Alblowi, Generalized Neutrosophic Set and Generalized Neutrosophic Topological Spaces, *Journal Computer Sci. Engineering*, Vol. (2) No. (7), (2012)pp129-132 .
- [8] A. A. Salama and S. A. Alblowi, Neutrosophic Set and Neutrosophic Topological Spaces, *ISOR J. Mathematics*, Vol.(3), Issue(3), (2012) pp31-35.
- [9] A. A. Salama, Neutrosophic Crisp Point & Neutrosophic Crisp Ideals, *Neutrosophic Sets and Systems*, Vol.1, (2013) pp. 50-54.
- [10] A. A. Salama and F. Smarandache, Filters via Neutrosophic Crisp Sets, *Neutrosophic Sets and Systems*, Vol.1, (2013) pp. 34-38.
- [11] A.A. Salama and S.A. Alblowi, Intuitionistic Fuzzy Ideals Topological Spaces, *Advances in Fuzzy Mathematics* , Vol.(7), Number 1, (2012) pp 51- 60.
- [12] A.A. Salama, and H. Elagamy, Neutrosophic Filters, *International Journal of Computer Science Engineering and Information Technology Research (IJCSEITR)*, Vol.3, Issue(1), (2013) pp307-312.
- [13] A. A. Salama, F.Smarandache and Valeri Kroumov, Neutrosophic crisp Sets & Neutrosophic crisp Topological Spaces, *Neutrosophic Sets and Systems*, Vlo.(2),(2014),pp25-30.
- [14] A. A. Salama, Florentin Smarandache and Valeri Kroumov, Neutrosophic Closed Set and Neutrosophic Continuous Functions *Neutrosophic Sets and Systems*, 2014, Vol. (4) pp4-8.
- [15] A. A. Salama, Mohamed Eisa and M. M. Abdelmoghny, Neutrosophic Relations Database, *International Journal of Information Science and Intelligent System*, 3(1) (2014)pp33-46 .
- [16] A. A. Salama , Florentin Smarandache and S. A. Alblowi, New Neutrosophic Crisp Topological Concepts, *Neutrosophic Sets and Systems*, Vol(4), (2014)pp50-54.
- [17] A. A. Salama, Said Broumi and Florentin Smarandache, Neutrosophic Crisp Open Set and Neutrosophic Crisp Continuity via Neutrosophic Crisp Ideals, *I.J. Information Engineering and Electronic Business*, Vol.3, (2014)pp1-8
- [18] A.A. Salama, Florentin Smarandache and S.A. Alblowi. The Characteristic Function of a Neutrosophic Set, *Neutrosophic Sets and Systems*, 2014, Vol. (3), pp14-18.
- [19] A. A. Salama, Neutrosophic Crisp Points & Neutrosophic Crisp Ideals, *Neutrosophic Sets and Systems*, 2013, Vol.(1) pp50-53
- [20] A. A. Salama, Mohamed Abdelfattah and S. A. Alblowi, Some Intuitionistic Topological Notions of Intuitionistic Region, Possible Application to GIS Topological Rules, *International Journal of Enhanced Research in*

- Management & Computer Applications, (2014), Vol.(3), no. (6) pp1-13.
- [21] A. A. Salama, Mohamed Abdelfattah and Mohamed Eisa, A Novel Model for Implementing Security over Mobile Ad-hoc Networks using Intuitionistic Fuzzy Function, International Journal of Emerging Technologies in Computational and Applied Sciences (IJETCAS), 2014, Vol.(6) pp1-7.
- [22] A. A. Salama and Said Broumi, Roughness of Neutrosophic Sets, Elixir Appl. Math. 74 (2014) pp26833-26837.
- [23] A. A. Salama, Mohamed Abdelfattah and Mohamed Eisa, Distances, Hesitancy Degree and Flexible Querying via Neutrosophic Sets, International Journal of Computer Applications, Volume 101– No.10, (2014) pp0975 – 8887
- [24] A.A. Salama, Haithem A. El-Ghareeb, Ayman. M. Maine and Florentin Smarandache. Introduction to Develop Some Software Programs for dealing with Neutrosophic Sets, Neutrosophic Sets and Systems, 2014, Vol(4), pp51-52.
- [25] A. A. Salama, F. Smarandache, and M. Eisa. Introduction to Image Processing via Neutrosophic Technique, Neutrosophic Sets and Systems, 2014, Vol. (5) pp59-63.
- [26] A. A. Salama, Haitham A. El-Ghareeb, Ayman M. Manie and M. M. Lotfy, Utilizing Neutrosophic Set in Social Network Analysis e-Learning Systems, International Journal of Information Science and Intelligent System, 3(2), (2014) pp61-72.
- [27] A. A. Salama, O. M. Khaled, and K. M. Mahfouz. Neutrosophic Correlation and Simple Linear Regression, Neutrosophic Sets and Systems, 2014, Vol. (5) pp3-8.
- [28] A. A. Salama, and F. Smarandache. Neutrosophic Crisp Set Theory, Neutrosophic Sets and Systems, 2014, Vol. (5) pp27-35.
- [29] A. A. Salama, Florentin Smarandache and S. A. Alblawi, New Neutrosophic Crisp Topological Concepts, Neutrosophic Sets and Systems, 2014, Vol(4) pp50-54.
- [30] Florentin Smarandache, Neutrosophy and Neutrosophic Logic, First International Conference on Neutrosophy, Neutrosophic Logic, Set, Probability, and Statistics University of New Mexico, Gallup, NM 87301, USA(2002).
- [31] Florentin Smarandache, A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability. American Research Press, Rehoboth, NM, (1999).
- [32] Florentin Smarandache, Neutrosophic set, a generalization of the intuitionistic fuzzy sets, Inter. J. Pure Appl. Math., 24 (2005), 287 – 297.
- [33] Debasis Sarker, Fuzzy ideal theory, Fuzzy local function and generated fuzzy topology, Fuzzy Sets and Systems 87, 117 – 123. (1997).
- [34] L.A. Zadeh, Fuzzy Sets, Inform and Control 8, 338-353.(1965).

Received: October 3, 2014. Accepted: December 20, 2014.