

1-1-2016

Restricted Interval Valued Neutrosophic Sets and Restricted Interval Valued Neutrosophic Topological Spaces

Anjan Mukherjee

Mithun Datta

Sadhan Sarkar

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Mukherjee, Anjan; Mithun Datta; and Sadhan Sarkar. "Restricted Interval Valued Neutrosophic Sets and Restricted Interval Valued Neutrosophic Topological Spaces." *Neutrosophic Sets and Systems* 12, 1 (). https://digitalrepository.unm.edu/nss_journal/vol12/iss1/6

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu.



Restricted Interval Valued Neutrosophic Sets and Restricted Interval Valued Neutrosophic Topological Spaces

Anjan Mukherjee¹, Mithun Datta², Sadhan Sarkar³

¹ Department of Mathematics, Tripura University, Suryamaninagar, Agartala-799022, Tripura, India, Email: anjan2002_m@yahoo.co.in

² Department of Mathematics, Tripura University, Suryamaninagar, Agartala-799022, Tripura, India, Email: mithunagt007@gmail.com

³ Department of Mathematics, Tripura University, Suryamaninagar, Agartala-799022, Tripura, India, Email: sadhan7_s@rediffmail.com

Abstract: In this paper we introduce the concept of restricted interval valued neutrosophic sets (RIVNS in short). Some basic operations and properties of RIVNS are discussed. The concept of restricted interval valued neutrosophic topology is also introduced together with restricted interval valued neutrosophic finer and

restricted interval valued neutrosophic coarser topology. We also define restricted interval valued neutrosophic interior and closer of a restricted interval valued neutrosophic set. Some theorems and examples are cited. Restricted interval valued neutrosophic subspace topology is also studied.

Keywords: Neutrosophic Set, Interval Valued Neutrosophic Set, Restricted Interval Valued Neutrosophic Set, Restricted Interval Valued Neutrosophic Topological Space.

AMS subject classification: 03E72

1 Introduction

In 1999, Molodtsov [10] introduced the concept of soft set theory which is completely new approach for modeling uncertainty. In this paper [10] Molodtsov established the fundamental results of this new theory and successfully applied the soft set theory into several directions. Maji et al. [8] defined and studied several basic notions of soft set theory in 2003. Pie and Miao [14], Aktas and Cagman [1] and Ali et al. [2] improved the work of Maji et al. [9]. The intuitionistic fuzzy set is introduced by Atanasiu [4] as a generalization of fuzzy set [19] where he added degree of non-membership with degree of membership. Neutrosophic set introduced by F. Smarandache in 1995 [16]. Smarandache [17] introduced the concept of neutrosophic set which is a mathematical tool for handling problems involving imprecise, indeterminacy and inconsistent data. Maji [9] combined neutrosophic set and soft set and established some operations on these sets. Wang et al. [18] introduced interval neutrosophic sets. Deli [7] introduced the concept of interval-valued neutrosophic soft sets.

In this paper we introduce the concept of restricted interval valued neutrosophic sets (RIVNS in short). Some basic operations and properties of RIVNS are discussed. The concept of restricted interval valued neutrosophic topology is also introduced together with restricted interval valued neutrosophic finer and restricted interval valued neutrosophic coarser topology. We also define restricted interval valued neutrosophic interior and closer of a restricted interval valued neutrosophic set. Some theorems and examples are cited. Restricted interval valued neutrosophic subspace topology is also studied. We establish some properties of restricted interval valued neutrosophic soft topological space with supporting proofs and examples.

2 Preliminaries

Definition 2.1[17] A neutrosophic set A on the universe of discourse U is defined as

$$A = \left\{ \langle x, \mu_A(x), \gamma_A(x), \delta_A(x) \rangle : x \in U \right\}, \text{ where}$$

$\mu_A, \gamma_A, \delta_A : U \rightarrow]^{-}0, 1^{+}[$ are functions such that the condition:

$\forall x \in U, \quad -0 \leq \mu_A(x) + \gamma_A(x) + \delta_A(x) \leq 3^+$ is satisfied.

From philosophical point of view, the neutrosophic set takes the value from real standard or non-standard subsets of $]^{-}0, 1^+[$. But in real life application in scientific and engineering problems it is difficult to use neutrosophic set with value from real standard or non-standard subset of $]^{-}0, 1^+[$. Hence we consider the neutrosophic set which takes the value from the subset of $[0, 1]$.

Definition 2.2 [6] An interval valued neutrosophic set A on the universe of discourse U is defined as $A = \{ \langle x, \mu_A(x), \gamma_A(x), \delta_A(x) \rangle : x \in U \}$, where $\mu_A, \gamma_A, \delta_A : U \rightarrow]^{-}0, 1^+[$ are functions such that the condition: $\forall x \in U, \quad -0 \leq \sup \mu_A(x) + \sup \gamma_A(x) + \sup \delta_A(x) \leq 3^+$ is satisfied.

In real life applications it is difficult to use interval valued neutrosophic set with interval-value from real standard or non-standard subset of $Int(]^{-}0, 1^+[)$. Hence we consider the interval-valued neutrosophic set which takes the interval-value from the subset of $Int([0, 1])$ (where $Int([0, 1])$ denotes the set of all closed sub intervals of $[0, 1]$).

Definition 2.3 [15] Let X be a non-empty fixed set. A generalized neutrosophic set (GNS in short) A is an object having the form $A = \{ \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle : x \in X \}$ Where $\mu_A(x), \sigma_A(x)$ and $\gamma_A(x)$ which represent the degree of member ship function (namely $\mu_A(x)$), the degree of indeterminacy (namely $\sigma_A(x)$), and the degree of non-member ship (namely $\gamma_A(x)$) respectively of each element $x \in X$ to the set A where the functions satisfy the condition $\mu_A(x) \wedge \sigma_A(x) \wedge \gamma_A(x) \leq 0.5$.

We call this generalized neutrosophic set [15] as **restricted neutrosophic set**.

Definition 2.4 [15] Let A and B be two RNSs on X defined by

$$A = \{ \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle : x \in X \} \quad \text{and}$$

$$B = \{ \langle x, \mu_B(x), \sigma_B(x), \gamma_B(x) \rangle : x \in X \} .$$

Then union, intersection, subset and complement may be defined as

(i) The union of A and B is denoted by $A \cup B$ and is defined as

$$A \cup B = \{ \langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \gamma_A(x) \wedge \gamma_B(x) \rangle : x \in X \}$$

or

$$A \cup B = \{ \langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \gamma_A(x) \wedge \gamma_B(x) \rangle : x \in X \} .$$

(ii) The intersection of A and B is denoted by $A \cap B$ and is defined as

$$A \cap B = \{ \langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \gamma_A(x) \vee \gamma_B(x) \rangle : x \in X \}$$

or

$$A \cap B = \{ \langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \gamma_A(x) \vee \gamma_B(x) \rangle : x \in X \}$$

or

$$A \cap B = \{ \langle x, \mu_A(x) \cdot \mu_B(x), \sigma_A(x) \cdot \sigma_B(x), \gamma_A(x) \cdot \gamma_B(x) \rangle : x \in X \}$$

(iii) A is called subset of B , denoted by $A \subseteq B$ if and only if

$$\mu_A(x) \leq \mu_B(x), \sigma_A(x) \geq \sigma_B(x),$$

$$\gamma_A(x) \geq \gamma_B(x)$$

or

$$\mu_A(x) \leq \mu_B(x), \sigma_A(x) \leq \sigma_B(x),$$

$$\gamma_A(x) \geq \gamma_B(x).$$

(iv) The complement of A is denoted by A^c and is defined as

$$A = \{ \langle x, \gamma_A(x), 1 - \sigma_A(x), \mu_A(x) \rangle : x \in X \}$$

or

$$A = \{ \langle x, \gamma_A(x), \sigma_A(x), \mu_A(x) \rangle : x \in X \}$$

or

$$A = \{ \langle x, 1 - \mu_A(x), \sigma_A(x), 1 - \gamma_A(x) \rangle : x \in X \}$$

Definition 2.5: [15] A restricted neutrosophic topology (RN-topology in short) on a non empty set X is a family of restricted neutrosophic subsets in X satisfying the following axioms

- (i) $0_N, 1_N \in \tau$
- (ii) $\bigcup_i G_i \in \tau, \forall \{G_i : i \in J\} \subseteq \tau$
- (iii) $G_1 \cap G_2 \in \tau$ for any $G_1, G_2 \in \tau$.

The pair (X, τ) is called restricted neutrosophic topological space (RN-topological space in short). The members of τ are called restricted neutrosophic open sets. A RNS F is closed if and only if F^c is RN open set.

3 Restricted Interval Valued Neutrosophic Set

In this section we introduce the concept of restricted interval valued neutrosophic set along with some examples, operators and results.

Definition 3.1 Let X be a non empty set. A restricted interval valued neutrosophic set (RIVNS in short) A is an object having the form $A = \{ \langle x, \mu_A(x), \gamma_A(x), \delta_A(x) \rangle : x \in X \}$, where $\mu_A(x), \gamma_A(x), \delta_A(x) : X \rightarrow Int]-0, 1^+[$ are functions such that the condition: $\forall x \in X, \sup \mu_A(x) \wedge \sup \gamma_A(x) \wedge \sup \delta_A(x) \leq 0.5$ is satisfied.

Here $\mu_A(x), \gamma_A(x)$ and $\delta_A(x)$ represent truth-membership interval, indeterminacy-membership interval and falsity-membership interval respectively of the element $x \in X$. For the sake of simplicity, we shall use the symbol $A = \langle x, \mu_A, \gamma_A, \delta_A \rangle$ for the RIVNS $A = \{ \langle x, \mu_A(x), \gamma_A(x), \delta_A(x) \rangle : x \in X \}$.

Example 3.2 Let $X = \{x_1, x_2, x_3\}$, then the RIVNS $A = \{ \langle x, \mu_A(x), \gamma_A(x), \delta_A(x) \rangle : x \in X \}$ can be represent by the following table

X	$\mu_A(x)$	$\gamma_A(x)$	$\delta_A(x)$	$\sup \mu_A(x) \wedge \sup \gamma_A(x) \wedge \sup \delta_A(x)$
x_1	[.2, .3]	[0, .1]	[.4, .5]	.1
x_2	[.3, .5]	[.1, .4]	[.5, .6]	.4
x_3	[.4, .7]	[.2, .4]	[.6, .8]	.4

The RIVNSs $\tilde{0}$ and $\tilde{1}$ are defined as

$$\tilde{0} = \{ \langle x, [0, 0], [1, 1], [1, 1] \rangle : x \in X \}$$

and

$$\tilde{1} = \{ \langle x, [1, 1], [0, 0], [0, 0] \rangle : x \in X \}.$$

Definition 3.3 Let $J_1 = [inf J_1, sup J_1]$ and $J_2 = [inf J_2, sup J_2]$ be two intervals then

- (i) $J_1 \leq J_2$ iff $inf J_1 \leq inf J_2$ and $sup J_1 \leq sup J_2$.
- (ii) $J_1 \vee J_2 = [max(inf J_1, inf J_2), max(sup J_1, sup J_2)]$.
- (iii) $J_1 \wedge J_2 = [min(inf J_1, inf J_2), min(sup J_1, sup J_2)]$.

Definition 3.4 Let A and B be two RIVNSs on X defined by

$$A = \{ \langle x, \mu_A(x), \gamma_A(x), \delta_A(x) \rangle : x \in X \}$$

and

$B = \{ \langle x, \mu_B(x), \gamma_B(x), \delta_B(x) \rangle : x \in X \}$. Then we can define union, intersection, subset and complement in several ways.

(i) The RIVNunion of A and B is denoted by $A \cup B$ and is defined as

$$A \cup B = \{ \langle x, \mu_A(x) \vee \mu_B(x), \gamma_A(x) \wedge \gamma_B(x), \delta_A(x) \wedge \delta_B(x) \rangle : x \in X \}$$

or

$$A \cup B = \{ \langle x, \mu_A(x) \vee \mu_B(x), \gamma_A(x) \vee \gamma_B(x), \delta_A(x) \wedge \delta_B(x) \rangle : x \in X \}$$

We take first definition throughout the paper.

(ii) The RIVNintersection of A and B is denoted by $A \cap B$ and is defined as

$$A \cap B = \{ \langle x, \mu_A(x) \wedge \mu_B(x), \gamma_A(x) \vee \gamma_B(x), \delta_A(x) \vee \delta_B(x) \rangle : x \in X \}$$

or

$$A \cap B = \{ \langle x, \mu_A(x) \wedge \mu_B(x), \gamma_A(x) \wedge \gamma_B(x), \delta_A(x) \vee \delta_B(x) \rangle : x \in X \}$$

We take first definition throughout the paper.

(iii) A is called RIVN subset of B , denoted by $A \subseteq B$ if and only if

$$\mu_A(x) \leq \mu_B(x), \gamma_A(x) \geq \gamma_B(x), \delta_A(x) \geq \delta_B(x)$$

or

$$\mu_A(x) \leq \mu_B(x), \gamma_A(x) \leq \gamma_B(x), \delta_A(x) \geq \delta_B(x).$$

We take first definition throughout the paper.

(iv) The RIVN complement of A is denoted by A^c and is defined as

$$A^c = \{ \langle x, \delta_A(x), [1 - \sup \gamma_A(x), 1 - \inf \gamma_A(x)], \mu_A(x) \rangle : x \in X \}$$

or

$$A^c = \{ \langle x, \delta_A(x), \gamma_A(x), \mu_A(x) \rangle : x \in X \}$$

We take first definition throughout the paper.

Definition 3.5 Let $\{A_i : i \in J\}$ be an arbitrary family of RIVNSs in X , then $\bigcup A_i$ and $\bigcap A_i$ can be respectively defined as

$$\bigcup A_i = \{ \langle x, \bigvee_{i \in J} \mu_{A_i}(x), \bigwedge_{i \in J} \gamma_{A_i}(x), \bigwedge_{i \in J} \delta_{A_i}(x) \rangle : x \in X \}$$

or

$$\bigcup A_i = \{ \langle x, \bigvee_{i \in J} \mu_{A_i}(x), \bigvee_{i \in J} \gamma_{A_i}(x), \bigwedge_{i \in J} \delta_{A_i}(x) \rangle : x \in X \}$$

$$\bigcap A_i = \{ \langle x, \bigwedge_{i \in J} \mu_{A_i}(x), \bigvee_{i \in J} \gamma_{A_i}(x), \bigvee_{i \in J} \delta_{A_i}(x) \rangle : x \in X \}$$

or

$$\bigcap A_i = \{ \langle x, \bigwedge_{i \in J} \mu_{A_i}(x), \bigwedge_{i \in J} \gamma_{A_i}(x), \bigvee_{i \in J} \delta_{A_i}(x) \rangle : x \in X \}$$

Theorem 3.6 Let A, B and C be three RIVNSs then

- (1) $A \cup A = A$
- (2) $A \cap A = A$
- (3) $A \cup B = B \cup A$
- (4) $A \cap B = B \cap A$
- (5) $(A \cup B)^c = A^c \cap B^c$
- (6) $(A \cap B)^c = A^c \cup B^c$
- (7) $(A \cup B) \cup C = A \cup (B \cup C)$
- (8) $(A \cap B) \cap C = A \cap (B \cap C)$
- (9) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
- (10) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

Proof: Let $A = \langle x, [a_1, a_2], [a_3, a_4], [a_5, a_6] \rangle$,
 $B = \langle x, [b_1, b_2], [b_3, b_4], [b_5, b_6] \rangle$ and
 $C = \langle x, [c_1, c_2], [c_3, c_4], [c_5, c_6] \rangle$

(1) - (4) Straight forward.

$$(5) A \cup B = \langle x, [\max(a_1, b_1), \max(a_2, b_2)], [\min(a_3, b_3), \min(a_4, b_4)], [\min(a_5, b_5), \min(a_6, b_6)] \rangle$$

$$(A \cup B)^c = \langle x, [\min(a_5, b_5), \min(a_6, b_6)], [1 - \min(a_4, b_4), 1 - \min(a_3, b_3)], [\max(a_1, b_1), \max(a_2, b_2)] \rangle$$

Now

$$A^c = \langle x, [a_5, a_6], [1 - a_4, 1 - a_3], [a_1, a_2] \rangle$$

$$B^c = \langle x, [b_5, b_6], [1 - b_4, 1 - b_3], [b_1, b_2] \rangle$$

$$\begin{aligned} A^c \cap B^c &= \langle x, [\min(a_5, b_5), \min(a_6, b_6)], \\ &\quad [\max(1 - a_4, 1 - b_4), \max(1 - a_3, 1 - b_3)], \\ &\quad [\max(a_1, b_1), \max(a_2, b_2)] \rangle \\ &= \langle x, [\min(a_5, b_5), \min(a_6, b_6)], \\ &\quad [1 - \min(a_4, b_4), 1 - \min(a_3, b_3)], \\ &\quad [\max(a_1, b_1), \max(a_2, b_2)] \rangle \end{aligned}$$

(6) Same as (5).

$$\begin{aligned} (7) \quad A \cup B &= \langle x, [\max(a_1, b_1), \max(a_2, b_2)], \\ &\quad [\min(a_3, b_3), \min(a_4, b_4)], \\ &\quad [\min(a_5, b_5), \min(a_6, b_6)] \rangle \end{aligned}$$

$$\begin{aligned} (A \cup B) \cup C &= \langle x, [\max(\max(a_1, b_1), c_1), \\ &\quad \max(\max(a_2, b_2), c_2)], [\min(\min(a_3, b_3), c_3), \\ &\quad \min(\min(a_4, b_4), c_4)], [\min(\min(a_5, b_5), c_5), \\ &\quad \min(\min(a_6, b_6), c_6)] \rangle \\ &= \langle x, [\max(a_1, b_1, c_1), \max(a_2, b_2, c_2)], \\ &\quad [\min(a_3, b_3, c_3), \min(a_4, b_4, c_4)], \\ &\quad [\min(a_5, b_5, c_5), \min(a_6, b_6, c_6)] \rangle \end{aligned}$$

$$\begin{aligned} B \cup C &= \langle x, [\max(b_1, c_1), \max(b_2, c_2)], \\ &\quad [\min(b_3, c_3), \min(b_4, c_4)], \\ &\quad [\min(b_5, c_5), \min(b_6, c_6)] \rangle \end{aligned}$$

$$\begin{aligned} A \cup (B \cup C) &= \langle x, [\max(a_1, \max(b_1, c_1)), \\ &\quad \max(a_2, \max(b_2, c_2))], [\min(a_3, \min(b_3, c_3)), \\ &\quad \min(a_4, \min(b_4, c_4))], [\min(a_5, \min(b_5, c_5)), \\ &\quad \min(a_6, \min(b_6, c_6))] \rangle \\ &= \langle x, [\max(a_1, b_1, c_1), \max(a_2, b_2, c_2)], \\ &\quad [\min(a_3, b_3, c_3), \min(a_4, b_4, c_4)], \\ &\quad [\min(a_5, b_5, c_5), \min(a_6, b_6, c_6)] \rangle \end{aligned}$$

(8) Same as (7).

$$\begin{aligned} (9) \quad B \cap C &= \langle x, [\min(b_1, c_1), \min(b_2, c_2)], \\ &\quad [\max(b_3, c_3), \max(b_4, c_4)], \\ &\quad [\max(b_5, c_5), \max(b_6, c_6)] \rangle \\ A \cup (B \cap C) &= \langle x, [\max(a_1, \min(b_1, c_1)), \\ &\quad \max(a_2, \min(b_2, c_2))], [\min(a_3, \max(b_3, c_3)), \\ &\quad \min(a_4, \max(b_4, c_4))], [\min(a_5, \max(b_5, c_5)), \\ &\quad \min(a_6, \max(b_6, c_6))] \rangle \end{aligned}$$

$$\begin{aligned} A \cup B &= \langle x, [\max(a_1, b_1), \max(a_2, b_2)], \\ &\quad [\min(a_3, b_3), \min(a_4, b_4)], \\ &\quad [\min(a_5, b_5), \min(a_6, b_6)] \rangle \end{aligned}$$

$$\begin{aligned} A \cup C &= \langle x, [\max(a_1, c_1), \max(a_2, c_2)], \\ &\quad [\min(a_3, b_3), \min(a_4, c_4)], \\ &\quad [\min(a_5, c_5), \min(a_6, c_6)] \rangle \end{aligned}$$

$$\begin{aligned} (A \cup B) \cap (A \cup C) &= \langle x, [\min(\max(a_1, b_1), \max(a_1, c_1)), \\ &\quad \min(\max(a_2, b_2), \max(a_2, c_2))], \\ &\quad [\max(\min(a_3, b_3), \min(a_3, c_3)), \\ &\quad \max(\min(a_4, b_4), \min(a_4, c_4))], \\ &\quad [\max(\min(a_5, b_5), \min(a_5, c_5)), \\ &\quad \max(\min(a_6, b_6), \min(a_6, c_6))] \rangle \end{aligned}$$

Now let us consider a_1, b_1 and c_1 , six cases may arise as

$a_1 \geq b_1 \geq c_1$, for this

$$\begin{aligned} \max(a_1, \min(b_1, c_1)) &= \\ \min(\max(a_1, b_1), \max(a_1, c_1)) &= a_1 \end{aligned}$$

$a_1 \geq c_1 \geq b_1$, for this

$$\begin{aligned} \max(a_1, \min(b_1, c_1)) &= \\ \min(\max(a_1, b_1), \max(a_1, c_1)) &= a_1 \end{aligned}$$

$$\begin{aligned}
 & b_1 \geq a_1 \geq c_1, \text{ for this} \\
 & \max(a_1, \min(b_1, c_1)) = \\
 & \quad \min(\max(a_1, b_1), \max(a_1, c_1)) = a_1 \\
 & b_1 \geq c_1 \geq a_1, \text{ for this} \\
 & \max(a_1, \min(b_1, c_1)) = \\
 & \quad \min(\max(a_1, b_1), \max(a_1, c_1)) = c_1 \\
 & c_1 \geq a_1 \geq b_1, \text{ for this} \\
 & \max(a_1, \min(b_1, c_1)) = \\
 & \quad \min(\max(a_1, b_1), \max(a_1, c_1)) = a_1 \\
 & c_1 \geq b_1 \geq a_1, \text{ for this} \\
 & \max(a_1, \min(b_1, c_1)) = \\
 & \quad \min(\max(a_1, b_1), \max(a_1, c_1)) = b_1
 \end{aligned}$$

Similarly it can be shown that other results are true for $a_2, b_2, c_2 ; a_3, b_3, c_3 ; a_4, b_4, c_4 ; a_5, b_5, c_5 ; a_6, b_6, c_6$. Hence

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C) .$$

(10) Same as (9).

4. Restricted Interval Valued neutrosophic Topological Spaces

In this section we give the definition of restricted interval valued Neutrosophic topological spaces with some examples and results.

Definition 4.1A restricted interval valued neutrosophic topology (*RIVN*-topology in short) on a non empty set X is a family of restricted interval valued neutrosophic subsets in X satisfying the following axioms

- (iv) $\tilde{0}, \tilde{1} \in \tau$
- (v) $\bigcup_i G_i \in \tau, \forall \{G_i : i \in J\} \subseteq \tau$
- (vi) $G_1 \cap G_2 \in \tau$ for any $G_1, G_2 \in \tau$.

The pair (X, τ) is called restricted interval valued neutrosophic topological space (*RIVN*-topological space in short). The members of τ are called restricted interval valued neutrosophic open sets. A *RIVNSF* is closed if and only if F^c is *RIVN* open set.

Example 4.2 Let X be a non-empty set. Let us consider the following *RIVNS*

$$\begin{aligned}
 G_1 &= \{ \langle x, [.5, .8], [.2, .3], [.2, .5] \rangle : x \in X \} , \\
 G_2 &= \{ \langle x, [.6, .7], [.5, .6], [.3, .4] \rangle : x \in X \} , \\
 G_3 &= G_1 \cup G_2 = \{ \langle x, [.6, .8], [.2, .3], [.2, .4] \rangle : x \in X \} , \\
 G_4 &= G_1 \cap G_2 = \{ \langle x, [.5, .7], [.5, .6], [.3, .5] \rangle : x \in X \}
 \end{aligned}$$

The family $\tau_1 = \{ \tilde{0}, \tilde{1}, G_1, G_2, G_3, G_4 \}$ is a *RIVN*-topology in X and (X, τ_1) is called a *RIVN*-topological space. But $\tau_2 = \{ \tilde{0}, \tilde{1}, G_1, G_2 \}$ is not a *RIVN*-topology as $G_1 \cup G_2 = G_3 \notin \tau_2$.

Definition 4.3 The two *RIVN* subsets $\tilde{0}, \tilde{1}$ constitute a *RIVN*-topology on X , called indiscrete *RIVN*-topology. The family of all *RIVN* subsets of X constitutes a *RIVN*-topology on X , such topology is called discrete *RIVN*-topology.

Theorem 4.4 Let $\{ \tau_j : j \in J \}$ be a collection of *RIVN*-topologies on X . Then their intersection $\bigcap_{j \in J} \tau_j$ is also a *RIVS*-topology on X .

Proof: (i) Since $\tilde{0}, \tilde{1} \in \tau_j$ for each $j \in J$. Hence $\tilde{0}, \tilde{1} \in \bigcap_{j \in J} \tau_j$.

(ii) Let $\{ G_k : k \in K \}$ be an arbitrary family *RIVNS* where $G_k \in \bigcap_{j \in J} \tau_j$ for each $k \in K$. Then for each $j \in J$, $G_k \in \tau_j$ for $k \in K$ and since for each $j \in J$, τ_j is a *RIVN*-topology, therefore $\bigcup_{k \in K} G_k \in \tau_j$ for each $j \in J$. Hence $\bigcup_{k \in K} G_k \in \bigcap_{j \in J} \tau_j$.

(iii) Let $G_1, G_2 \in \bigcap_{j \in J} \tau_j$, then $G_1, G_2 \in \tau_j$ for each $j \in J$. Since for each $j \in J$, τ_j is an *RIVN*-topology, therefore $G_1, G_2 \in \tau_j$ for each $j \in J$. Hence $G_1 \cap G_2 \in \bigcap_{j \in J} \tau_j$.

Thus $\bigcap_{j \in J} \tau_j$ forms a *RIVN*-topology as it satisfies all the axioms of *RIVN*-topology. But union of *RIVN*-topologies need not be a *RIVN*-topology.

Let us show this with the following example.

Example 4.5 In example 4.2, let us consider two *RIVN*-topologies τ_3 and τ_4 on X as $\tau_3 = \{\tilde{0}, \tilde{1}, G_1\}$ and $\tau_4 = \{\tilde{0}, \tilde{1}, G_2\}$. Here their union $\tau_3 \cup \tau_4 = \{\tilde{0}, \tilde{1}, G_1, G_2\} = \tau_2$ is not a *RIVN*-topology on X .

Definition 4.6 Let (X, τ) be an *RIVN*-topological space over X . A *RIVN* subset G of X is called restricted interval valued neutrosophic closed set (in short *RIVN*-closed set) if its complement G^c is a member of τ .

Definition 4.7 Let (X, τ_1) and (X, τ_2) be two *RIVN*-topological spaces over X . If each $G \in \tau_2$ implies $G \in \tau_1$, then τ_1 is called restricted interval valued neutrosophic finer topology than τ_2 and τ_2 is called restricted interval valued neutrosophic coarser topology than τ_1 .

Example 4.8 In example 4.2 and 4.5, τ_1 is restricted interval valued neutrosophic finer topology than τ_3 and τ_3 is called restricted interval valued neutrosophic coarser topology than τ_1 .

Definition 4.9 Let τ be a *RIVN*-topological space on X and β be a subfamily of τ . If every element of τ can be express as the arbitrary restricted interval valued neutrosophic union of some elements of β , then β is called restricted interval valued neutrosophic basis for the *RIVN*-topology τ .

5 Some Properties of Restricted Interval Valued Neutrosophic Soft Topological Spaces

In this section some properties of *RIVN*-topological spaces are introduced. Some results on *RIVNInt* and *RIVNCl* are also introduced. Restricted interval valued neutrosophic subspace topology is also studied.

Definition 5.1 Let (X, τ) be a *RIVN*-topological space and A be a *RIVNS* in X . The restricted interval valued neutrosophic interior and restricted interval valued neutrosophic closer of A is denoted by *RIVNInt*(A) and *RIVNCl*(A) are defined as $RIVNInt(A) = \bigcup \{G : G \text{ is an } RIVN \text{ open set and } G \subseteq A\}$ and $RIVNCl(A) = \bigcap \{F : F \text{ is an } RIVN \text{ closed set and } F \supseteq A\}$ respectively.

Theorem 5.2 Let (X, τ) be a *RIVN*-topological space and G and H be two *RIVNS*s then the following properties hold

- (1) $RIVNInt(G) \subseteq G$
- (2) $G \subseteq H \Rightarrow RIVNInt(G) \subseteq RIVNInt(H)$
- (3) $RIVNInt(G) \in \tau$
- (4) $G \in \tau \Leftrightarrow RIVNInt(G) = G$
- (5) $RIVNInt(RIVNInt(G)) = RIVNInt(G)$
- (6) $RIVNInt(\tilde{0}) = \tilde{0}, RIVNInt(\tilde{1}) = \tilde{1}$

Proof:

(1) Straight forward.

(2) Let $G \subseteq H$, then all the *RIVN*-open sets Contained in G also contained in H .

$$\begin{aligned} \text{i.e. } \{G^* \in \tau : G^* \subseteq G\} &\subseteq \{H^* \in \tau : H^* \subseteq H\} \\ \text{i.e. } \bigcup \{G^* \in \tau : G^* \subseteq G\} &\subseteq \bigcup \{H^* \in \tau : H^* \subseteq H\} \\ \text{i.e. } RIVNInt(G) &\subseteq RIVNInt(H) \end{aligned}$$

(3) $RIVNInt(G) = \bigcup \{G^* \in \tau : G^* \subseteq G\}$

Now clearly $\bigcup \{G^* \in \tau : G^* \subseteq G\} \in \tau$

$$\therefore RIVNInt(G) \in \tau.$$

(4) Let $G \in \tau$, then by (1) $RIVNInt(G) \subseteq G$.

Now since $G \in \tau$ and $G \subseteq G$, therefore $G \subseteq \bigcup \{G^* \in \tau : G^* \subseteq G\} = RIVNInt(G)$

$$\text{i.e., } G \subseteq RIVNInt(G)$$

$$\text{Thus } RIVNInt(G) = G$$

Conversely, let $RIVNInt(G) = G$

Since by (3) $RIVNInt(G) \in \tau$

Therefore $G \in \tau$

(5) By (3) $RIVNInt(G) \in \tau$

∴ By (4)
 $RIVNInt(RIVNInt(f_A, E)) = RIVNInt(f_A, E)$

- (6) We know that $\tilde{0}, \tilde{1} \in \tau$
 ∴ By (4)
 $RIVNInt(\tilde{0}) = \tilde{0}, RIVNInt(\tilde{1}) = \tilde{1}$

Theorem 5.3 Let (X, τ) be a RIVN-topological space and G and H are two RIVNSs then the following properties hold

- (1) $G \subseteq RIVNCl(G)$
- (2) $G \subseteq H \Rightarrow RIVNCl(G) \subseteq RIVNCl(H)$
- (3) $(RIVNCl(G))^c \in \tau$
- (4) $G^c \in \tau \Leftrightarrow RIVNCl(G) = G$
- (5) $RIVNCl(RIVNCl(G)) = RIVNCl(G)$
- (6) $RIVNCl(\tilde{0}) = \tilde{0}, RIVNCl(\tilde{1}) = \tilde{1}$

Proof: straight forward.

Theorem 5.4 Let (X, τ) be an RIVN-topological space on X and A be a RIVNS of X and let $\tau_A = \{A \cap U : U \in \tau\}$. Then τ_A forms a RIVN-topology on A.

Proof:

- (i) Clearly $\tilde{0} = A \cap \tilde{0} \in \tau_A$ and $\tilde{1} = A \cap \tilde{1} \in \tau_A$.
- (ii) Let $G_j \in \tau_A, \forall j \in J$, then $G_j = A \cap U_j$ where $U_j \in \tau$ for each $j \in J$.
 Now $\bigcup_{j \in J} G_j = \bigcup_{j \in J} (A \cap U_j) = A \cap \left(\bigcup_{j \in J} U_j \right) \in \tau_A$ (since $\bigcup_{j \in J} U_j \in \tau$ as each $U_j \in \tau$).
- (iii) Let $G, H \in \tau_A$ then $G = A \cap U$ and $H = A \cap V$ where $U, V \in \tau$.
 Now
 $G \cap H = (A \cap U) \cap (A \cap V) = A \cap (U \cap V) \in \tau_A$
 (since $U \cap V \in \tau$ as $U, V \in \tau$).

Definition 5.5 Let (X, τ) be an RIVN-topological space on X and A be a RIVNS of X. Then $\tau_A = \{A \cap U : U \in \tau\}$ is called restricted interval valued neutrosophic subspace topology and (A, τ_A) is called restricted interval valued

neutrosophic subspace of RIVN-topological space (X, τ) .

Conclusion: In this paper we introduce the concept of restricted interval valued neutrosophic set which is the generalization of restricted neutrosophic set. We define some operators on RIVNS. We also introduce a topological structure based on this. RIVN interior and RIVN closer of a restricted

interval valued neutrosophic set are also defined. Restricted interval valued neutrosophic subspace topology is also studied. In future combining the ideas presented in this paper with concept of soft set one can define a new concept named restricted interval valued Neutrosophic soft set and can define a topological structure too.

References

- (1) H. Aktas, N. Cagman, Soft sets and soft groups, In-form. Sci., 177(2007), 2726-2735.
- (2) M. I. Ali, F. Feng, X. Liu, W. K. Min and M. Shabir, On some new operations in soft set theory, Computers and Mathematics with Applications 57 (9) (2009),1547-1553.
- (3) I. Arockiarani, I. R. Sumathi, J. Martina Jency, Fuzzy neutrosophic soft topological spaces, International Journal of Mathematical Archive, 4 (10) (2013), 225-238.
- (4) K. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems, 20 (1986), 87-96.
- (5) K. Atanassov, G. Gargov, Interval valued intuitionistic fuzzy sets, Fuzzy Sets and Systems, 31(1989), 343-349.
- (6) S. Broumi, I. Deli, and F. Smarandache, Relations on Interval Valued Neutrosophic Soft Sets, Journal of New Results in Science, 5 (2014), 1-20.
- (7) I. Deli, Interval-valued neutrosophic soft sets and its decision making, Kilis 7 Aralık University, 79000 Kilis, Turkey.
- (8) P. K. Maji, R. Biswas and A. R. Roy, Soft Set Theory, Computers and Mathematics with Applications, 45 (2003), 555-562.

- (9) P. K. Maji, Neutrosophic soft set, Annals of Fuzzy Mathematics and Information, 5(1) (2013), 157-168.
- (10) D. Molodtsov, Soft set theory-first results, Computers and Mathematics with Applications 37 (4-5) (1999), 19-31.
- (11) A. Mukherjee, A. K. Das, A. Saha, Interval valued intuitionistic fuzzy soft topological spaces, Annals of Fuzzy Mathematics and Informatics, 6 (3) (2013), 689-703.
- (12) A. Mukherjee, M. Datta, F. Smarandache, Interval Valued Neutrosophic Soft Topological Spaces, Neutrosophic Sets and Systems, 6 (2014), 18-27.
- (13) A. Mukherjee, M. Datta, A. Saha, Interval valued intuitionistic soft sets, The Journal of Fuzzy Mathematics, 23 (2) (2015), 283-294.
- (14) D. Pie, D. Miao, From soft sets to information systems, Proc. IEEE Int. Conf. Granular Comput. 2 (2005), 617-621.
- (15) A. A. Salma and S. A. Alblowi, Generalized Neutrosophic Set and Generalized Neutrosophic Topological Spaces, Computer Science and Engineering 2012, 2(7):129-132 DOI:10.5923/j.computer.20120207.01.
- (16) F. Smarandache, Neutrosophic Logic and Set, mss. , <http://fs.gallup.unm.edu/neutrosophy.htm>, 1995.
- (17) F. Smarandache, Neutrosophic set- a generalisation of the intuitionistic fuzzy sets, int. J. Pure Appl. Math., 24 (2005), 287-297.
- (18) H. Wang, F. Smarandache, Y.Q. Zhang, R. Sunderraman, Interval Neutrosophic Sets and logic: Theory and Applications in Computing, Hexis; Neutrosophic book series, No: 5, 2005.
- (19) L. A. Zadeh, Fuzzy sets, Information and Control, 8(1965), 338-353.

Received: March 26, 2016. Accepted: July 05, 2016.