

University of New Mexico

## UNM Digital Repository

---

Mathematics & Statistics ETDs

Electronic Theses and Dissertations

---

5-13-1971

# Convergence Rates for the Central Limit Theorem for Random Sums

Christopher E. Olson

Follow this and additional works at: [https://digitalrepository.unm.edu/math\\_etds](https://digitalrepository.unm.edu/math_etds)



Part of the [Applied Mathematics Commons](#), [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

---

### Recommended Citation

Olson, Christopher E.. "Convergence Rates for the Central Limit Theorem for Random Sums." (1971). [https://digitalrepository.unm.edu/math\\_etds/146](https://digitalrepository.unm.edu/math_etds/146)

This Dissertation is brought to you for free and open access by the Electronic Theses and Dissertations at UNM Digital Repository. It has been accepted for inclusion in Mathematics & Statistics ETDs by an authorized administrator of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).

UNIVERSITY OF NEW MEXICO-GENERAL LIBRARY



A14425 247311

LD

3781

N5640f8

cop. 2

COGNITIVE PSYCHOLOGY: ANSWERS FOR THE GENERAL LIMIT THEOREM

OLSSON



Embossed text, likely a name or title, centered on the page.

Embossed text, possibly a date or reference number, centered below the main text.

THE UNIVERSITY OF NEW MEXICO  
ALBUQUERQUE, NEW MEXICO 87106

POLICY ON USE OF THESES AND DISSERTATIONS

Unpublished theses and dissertations accepted for master's and doctor's degrees and deposited in the University of New Mexico Library are open to the public for inspection and reference work. *They are to be used only with due regard to the rights of the authors.* The work of other authors should always be given full credit. Avoid quoting in amounts, over and beyond scholarly needs, such as might impair or destroy the property rights and financial benefits of another author.

To afford reasonable safeguards to authors, and consistent with the above principles, anyone quoting from theses and dissertations must observe the following conditions:

1. Direct quotations during the first two years after completion may be made only with the written permission of the author.
2. After a lapse of two years, theses and dissertations may be quoted without specific prior permission in works of original scholarship provided appropriate credit is given in the case of each quotation.
3. Quotations that are complete units in themselves (e.g., complete chapters or sections) in whatever form they may be reproduced and quotations of whatever length presented as primary material for their own sake (as in anthologies or books of readings) ALWAYS require consent of the authors.
4. The quoting author is responsible for determining "fair use" of material he uses.

This thesis/dissertation by Christopher E. Olson has been used by the following persons whose signatures attest their acceptance of the above conditions. (A library which borrows this thesis/dissertation for use by its patrons is expected to secure the signature of each user.)

NAME AND ADDRESS

DATE

_____	_____
_____	_____
_____	_____
_____	_____
_____	_____

THE UNIVERSITY OF NEW BRUNSWICK  
ADMINISTRATIVE SERVICES DIVISION

Office of the Registrar  
100 University Avenue  
New Brunswick, N.B. A1B 3X9  
Canada

I hereby certify that the following information is true and correct to the best of my knowledge and belief. I am a duly qualified and licensed professional person in the Province of New Brunswick and I am duly qualified to perform the duties of the position of Registrar of the University of New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

I am a resident of the Province of New Brunswick and I am a resident of the City of Saint John, New Brunswick.

This dissertation, directed and approved by the candidate's committee, has been accepted by the Graduate Committee of The University of New Mexico in partial fulfillment of the requirements for the degree of

Doctor of Philosophy in Mathematics

CONVERGENCE RATES FOR THE  
CENTRAL LIMIT THEOREM FOR RANDOM SUMS

*Title*

Christopher E. Olson

*Candidate*

Department of Mathematics and Statistics

*Department*

David T. Bonchuk

*Dean*

May 13, 1971

*Date*

*Committee*

L.H. Koopmans

*Chairman*

Bernard Gasterin

Art Steger

THE UNIVERSITY OF CHICAGO  
LIBRARY

Division of Biological Sciences

Department of Zoology

Chicago, Illinois

Department of Zoology and Botany

Chicago, Illinois

Chicago, Illinois

Chicago, Illinois

Chicago, Illinois

Chicago, Illinois



CONVERGENCE RATES FOR THE  
CENTRAL LIMIT THEOREM FOR RANDOM SUMS

BY

CHRISTOPHER E. OLSON

DISSERTATION

Submitted in Partial Fulfillment of the  
Requirements for the Degree of  
Doctor of Philosophy in Mathematics  
in the Graduate School of  
The University of New Mexico  
Albuquerque, New Mexico

June, 1971

UNIVERSITY OF NEW MEXICO LIBRARY

LD  
3781  
N564 Ocl 8  
cop. 2

#### ACKNOWLEDGMENT

The author wishes to express his sincere appreciation to Professor L.H. Koopmans for his patience, guidance, and encouragement. The final form of several proofs benefited from his insight.

Acknowledgment is also due the Atomic Energy Commission and Sandia Laboratory for financial support and use of library facilities.

MEMORANDUM

The purpose of this memorandum is to provide information regarding the proposed changes to the existing policy. The proposed changes are intended to improve the efficiency of the process and to ensure that all relevant parties are kept informed of any developments. It is requested that you review the attached documents and provide your comments by the date indicated.

The proposed changes are outlined in the attached documents. The first document provides a detailed description of the changes, while the second document provides a summary of the key points. It is noted that the changes are subject to the approval of the relevant committees and the Board of Directors. Your input is valued and will be taken into account in the final decision.

Very truly yours,  
[Signature]  
[Title]

Enclosure

CONVERGENCE RATES FOR THE  
CENTRAL LIMIT THEOREM FOR RANDOM SUMS

BY

CHRISTOPHER E. OLSON

B.A., St. Mary's University (Texas), 1963

M.A., University of Kansas, 1965

ABSTRACT OF DISSERTATION

Submitted in Partial Fulfillment of the  
Requirements for the Degree of  
Doctor of Philosophy in Mathematics  
in the Graduate School of  
The University of New Mexico  
Albuquerque, New Mexico

June, 1971

CONFIDENTIAL - SECURITY INFORMATION  
DO NOT DISSEMINATE TO OTHERS

11

CONFIDENTIAL - SECURITY INFORMATION  
DO NOT DISSEMINATE TO OTHERS  
U.S. DEPARTMENT OF JUSTICE  
FEDERAL BUREAU OF INVESTIGATION

CONFIDENTIAL - SECURITY INFORMATION

CONFIDENTIAL

### Abstract

Let  $\{X_i\}$  be a sequence of independent, identically-distributed random variables with  $EX_i^2 < \infty$  and  $E(X_i - EX_i)^2 = 1$ . Let  $\bar{S}(n) = \left( \sum_{j=1}^n X_j - nEX_i \right) / \sqrt{n}$ . If  $\tau_n$  is a sequence of positive integer-valued random variables such that  $\tau_n/n \xrightarrow{p} \tau$ , a positive (possibly degenerate) random variable, and  $\bar{G}_n(x) \equiv P[\bar{S}(\tau_n) \leq x]$ , then by a theorem of Blum, Hanson, and Rosenblatt  $\sup_x |\bar{G}_n(x) - \bar{\Phi}(x)| \rightarrow 0$  as  $n \rightarrow \infty$ . This dissertation establishes sufficient conditions for series convergence rates of the form

$$\sum_n n^{-\alpha} \sup_x |\bar{G}_n(x) - \bar{\Phi}(x)| < \infty. \quad (1)$$

The exponent  $\alpha$  in (1) depends on the number of finite absolute moments of  $X_i$ . If  $E|X_i|^{2+\delta} < \infty$ ,  $0 < \delta < 1$ , then  $\alpha = 1 - \delta/2$ . If  $EX_i^2 < \infty$  but  $E|X_i|^{2+\delta} = \infty$ , for all  $\delta > 0$ , then  $\alpha = 1$ . In this case, unless  $EX_i^2 \log|X_i| < \infty$ , the term  $\bar{\Phi}(x)$  in (1) is replaced by  $E\bar{\Phi}(x/\sigma_{\tau_n})$ , where  $\sigma_n$  is the variance of  $X_i$  truncated at  $\sqrt{n}$ .

The problem is most naturally treated in two cases: In the first case, the sequences  $\{X_i\}$  and  $\{\tau_n\}$  are taken to be independent, and in the second, they are assumed to be dependent. In the independent case, a basic sufficient condition for (1) to hold is

$$\sum_n n^{-\alpha} \cdot P[\tau_n = k] = O(k^{-\alpha}). \quad (2)$$





It is shown that, if  $\tau_n$  is the nth epoch of a positive, integer-valued renewal process, then it satisfies (2). Another theorem gives a condition on the rate with which  $\tau_n/n \xrightarrow{P} c_0$ , a positive constant, which is sufficient for (2) to hold, and thus for (1).

In the dependent case, sufficient conditions are given in terms of the rate of decrease of

$$|P([\bar{S}(k) \leq x] \cap [\tau_n = k]) - P[\bar{S}(k) \leq x] \cdot P[\tau_n = k]|$$

and of

$$|P([\bar{S}(k) \leq x] \cap [\tau_n = k]) - P([\bar{S}(n) \leq x] \cap [\tau_n = k])| .$$

The sharpness of these sufficient conditions is indicated through simple examples whereas the conditions are violated and the convergence in (1) fails. A further example satisfying both of these conditions is then given.


THE UNIVERSITY OF CHICAGO  
DEPARTMENT OF CHEMISTRY  
5780 SOUTH CAMPUS DRIVE  
CHICAGO, ILLINOIS 60637  
TEL: 773-936-3700

### PROFESSOR OF CHEMISTRY

THE UNIVERSITY OF CHICAGO  
DEPARTMENT OF CHEMISTRY  
5780 SOUTH CAMPUS DRIVE  
CHICAGO, ILLINOIS 60637  
TEL: 773-936-3700

## TABLE OF CONTENTS

Section	Page
I. Introduction . . . . .	1
II. Convergence of the $A_j$ 's when the Sequences $\{\tau_n\}$ and $\{X_i\}$ are Independent . . . . .	10
III. Convergence of the $A_j$ 's when the Sequences $\{\tau_n\}$ and $\{X_i\}$ are Dependent . . . . .	24
Bibliography . . . . .	39



## I. Introduction

Throughout this dissertation, let  $X_1, X_2, \dots$  be a sequence of independent, identically-distributed random variables on a probability space  $(\Omega, \mathcal{A}, P)$ . The distribution function of  $X_1$  is  $F(x) = P(\{\omega | X_1(\omega) \leq x\})$ , or more simply  $F(x) = P[X_1 \leq x]$ .

The following conditions are assumed:

- (1)  $EX_1^2 < \infty$
- (2)  $E(X_1 - \mu)^2 = 1$ ,

where  $E$  denotes expectation and  $\mu \equiv EX_1$ . Let  $S(n) = \sum_{i=1}^n X_i$ ,  $\bar{S}(n) = (S(n) - n\mu)/\sqrt{n}$ . The distribution function of  $\bar{S}(n)$  is denoted by

$$\bar{F}_n(x) = P[\bar{S}(n) \leq x].$$

Let  $\Phi(x)$  and  $\varphi(x)$  be the distribution function and density function, respectively, of the normal probability distribution with zero mean and unit variance, i.e.,

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt,$$
$$\varphi(x) \equiv \Phi'(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$

A truncated variance is defined for  $X_1$  by

$$\sigma_n^2 = \int_{|x| \leq \sqrt{n}} x^2 dF - \left( \int_{|x| \leq \sqrt{n}} x dF \right)^2.$$



Note that  $\lim_{n \rightarrow \infty} \sigma_n = 1$ , the variance of  $X_1$ .

The central limit problem has been a principal line of research in probability theory since the inception of the theory. In its simplest form, the central limit problem consists of finding conditions under which

$$\Delta_n \equiv \sup_x |\bar{F}_n(x) - \Phi(x)| \rightarrow 0 \text{ as } n \rightarrow \infty .$$

The history of the problem begins with DeMoivre and Laplace, who proved that  $\Delta_n \rightarrow 0$  for random variables  $X_i$  with a binomial distribution. Lindeberg<sup>17</sup> and Feller<sup>8</sup> proved the necessity and sufficiency of the finiteness of  $E|X_i|^2$  for  $\Delta_n \rightarrow 0$ , for independent, identically distributed  $X_i$ . Further work in the central limit problem has extended these classic results to include sequences of non-identically distributed random variables and dependent random variables of certain types.

In many practical applications, the limiting distribution is used to approximate the distribution of the sum  $\bar{S}(n)$  for finite values of  $n$ . Such usage must be based on rigorous estimates of the remainder term or the speed of convergence. Liapounov<sup>15,16</sup> showed that, if  $E|X_i|^{2+\delta} < \infty$ ,  $0 < \delta < 1$ , then  $\Delta_n \leq cn^{-\delta/2} E|X_i|^{2+\delta}$ , where the constant  $c$  may depend on the distribution of the  $X_i$ . When  $E|X_i|^3 < \infty$ , Berry<sup>3</sup> and Esseen<sup>7</sup> proved that

$$\Delta_n \leq c_0 n^{-1/2} E|X_i|^3 ,$$





where  $c_0$  is a constant which does not depend on the distribution of  $X_1$ . This result was refined by Nagaev<sup>20</sup>, who showed that

$$|\bar{F}_n(x) - \Phi(x)| \leq \frac{c_0 E|X_1|^3}{n^{1/2}(1+|x|^3)},$$

where  $c_0$  is the Berry-Esseen constant. Another direction was taken by M. Katz<sup>12</sup>, who proved that, if  $g(x)$  is a real, non-negative, even function, non-decreasing on  $[0, \infty)$ , then  $EX_1^2 g(X_1) < \infty$  is sufficient for

$$\Delta_n \leq \frac{c_0 EX_1^2 g(X_1)}{g(n^{1/2})}.$$

The results of Berry and Esseen, and of Nagaev, are the best possible in the sense that the bound is achieved by some distribution satisfying the hypotheses.

In 1966, N. Friedman, M. Katz, and L. Koopmans<sup>9</sup> proved a series convergence rate theorem requiring only two moments. C.C. Heyde<sup>10</sup> and J.A. Davis<sup>6</sup> each noted improvements and simplifications which led to the following result:

Theorem: If  $E|X_1|^2 < \infty$ , say  $E|X_1|^2 = 1$  and  $EX_1 = 0$ , then

$\sum_{n=1}^{\infty} n^{-1} \sup_x |\bar{F}_n(x) - \Phi(x/\sigma_n)| < \infty$ . If, in addition,  $EX_1^2 \log|X_1| < \infty$ ,

then  $\sum_{n=1}^{\infty} n^{-1} \Delta_n < \infty$ . Further, if  $EX_1^{2+\delta} < \infty$ ,  $0 < \delta < 1$ , then

$\sum_{n=1}^{\infty} n^{-1+\delta/2} \Delta_n < \infty$ .



Suppose now that  $\{\tau_n\}$  is a sequence of positive, integer-valued random variables on  $(\Omega, \mathcal{A}, P)$ . Let  $\bar{G}_n(x) \equiv P(\{\omega/\bar{S}(\tau_n(\omega)) \leq x\})$ .

Then

$$\bar{G}_n(x) = E(I_{(-\infty, x]}(\bar{S}(\tau_n))) ,$$

where  $I_A$  is the set indicator function of the set  $A$ .

Proceeding in the same line,

$$\begin{aligned} \bar{G}_n(x) &= E(I_{(-\infty, x]}(\bar{S}(\tau_n))) \\ &= E_{\tau_n} [E(I_{(-\infty, x]}(\bar{S}(\tau_n)) | \tau_n)] \\ &= \sum_{k=1}^{\infty} E(I_{(-\infty, x]}(\bar{S}(k) | \tau_n=k)) \cdot P[\tau_n=k] \\ &= \sum_{k=1}^{\infty} P[\bar{S}(k) \leq x | \tau_n=k] \cdot P[\tau_n=k] . \end{aligned}$$

Note that in case  $\{\tau_n\}$  is independent of  $\{X_i\}$ ,

$$\bar{G}_n(x) = \sum_{k=1}^{\infty} \bar{F}_k(x) \cdot P[\tau_n=k] .$$


The question of when  $\sup_x |\bar{G}_n(x) - \Phi(x)|$  goes to zero is a natural extension of the classical central limit problem. For example, one might wish to approximate  $\bar{G}_n(x)$  when  $\tau_n$  is the first passage of the sequence  $S(n)$  past  $n$ , or when  $\tau_n$  is an integer-multiple of  $n$  determined by the value of  $S(n)$ . Such an approximation would be useful in sequential analysis for tests based on  $S(\tau_n)$  (see Wald<sup>11</sup>).



The case where the  $\tau_n$ 's are independent of the summands was dealt with by Robbins<sup>12</sup> in 1948. Anscombe<sup>1</sup> showed that, if  $\tau_n/n \xrightarrow{p} c$ , where  $c$  is a positive constant, then  $\Delta_{\tau_n} \equiv \sup_x |\bar{G}_n(x) - \bar{\Phi}(x)| \rightarrow 0$ , regardless of the fact that the  $\tau_n$ 's might depend on the  $X_i$ 's. This result was extended in 1960 by Renyi<sup>23</sup> to the case where  $\tau_n/n \xrightarrow{p} \tau$ , a positive discrete random variable; and by Blum, Hanson, and Rosenblatt<sup>4</sup>, and, independently, Mogyorodi<sup>19</sup>, in 1963, to the case where  $\tau$  is any positive random variable.

In 1964, Wittenberg<sup>27</sup> investigated, more generally, the conditions under which the Kolmogorov-Smirnov (K-S) distance between  $S(\tau_n)$  and  $S(n)$  tends to zero. If  $S(n)$ , when properly normed, converges in distribution to a random variable with non-degenerate law, and if the K-S distance between  $S(n)$  and  $S(\tau_n)$  tends to zero, then  $S(\tau_n)$  converges to a random variable with the same law. The theorem of Blum, Hanson and Rosenblatt becomes a particular case of Wittenberg's results.

This variety of results concerning the convergence of  $\bar{G}_n(x)$  to  $\bar{\Phi}(x)$  will be referred to as the random central limit theorem (RCLT). In particular, a pair of sequences  $\{X_i\}$ ,  $\{\tau_n\}$  will be said to satisfy the conditions of the RCLT if  $\{X_i\}$  is a sequence of independent, identically-distributed random variables with at least two finite moments, and if  $\{\tau_n\}$  is a sequence of positive, integer-valued random variables such that  $\tau_n/n \xrightarrow{p} \tau$ , a positive (possibly degenerate) random variable.



It is the purpose of this dissertation to investigate the rate of convergence of  $\bar{G}_n(x)$  to  $\bar{\Phi}(x)$ . In order to derive series convergence rates similar to those of the Friedman-Katz-Koopmans Theorem (F-K-K), one must treat functions similar to  $\bar{\Phi}(x/\sigma_n)$  where the normalizing factor is defined in terms of both  $\{X_i\}$  and  $\{\tau_n\}$ . To this end, consider

$$E\bar{\Phi}(x/\sigma_{\tau_n}) = \sum_k \bar{\Phi}(x/\sigma_k) \cdot P[\tau_n=k].$$

Since  $\sigma_k \rightarrow 1$ , in order to avoid trivial details, it will be assumed that  $\sigma_k > 0$  for all  $k$ . The series whose convergence properties are examined are the following:

$$A_1(x, \alpha) \equiv \sum_n n^{-\alpha} |\bar{G}_n(x) - E\bar{\Phi}(x/\sigma_{\tau_n})|$$

$$A_2(\alpha) \equiv \sum_n n^{-\alpha} \sup_x |\bar{G}_n(x) - E\bar{\Phi}(x/\sigma_{\tau_n})|$$

$$A_3(x, \alpha) \equiv \sum_n n^{-\alpha} |\bar{G}_n(x) - \bar{\Phi}(x)|$$

$$A_4(\alpha) = \sum_n n^{-\alpha} \sup_x |\bar{G}_n(x) - \bar{\Phi}(x)|.$$

The original statement of the F-K-K Theorem allowed for dropping the normalizing factor  $\sigma_n^{-1}$  of the argument of  $\bar{\Phi}$  provided that

$$\sum_{n=1}^{\infty} n^{-1}(1-\sigma_n^2) < \infty. \quad \text{J.A. Davis}^6 \text{ later pointed out that } \sum_n n^{-1}(1-\sigma_n^2) < \infty$$

if and only if  $EX_1^2 \log|X_1| < \infty$ . It will be valuable to have a

sufficient condition for dropping the normalizing factor  $\sigma_{\tau_n}^{-1}$  from  $A_1(x, \alpha)$  or  $A_2(\alpha)$ . The following lemma presents such a condition:





Lemma: If  $A_2(\alpha)$  (or  $A_1(x, \alpha)$ ) is finite and if

$$(1) \quad \sum_{n=1}^{\infty} n^{-\alpha} \sum_k P[\tau_n = k] \cdot |1 - \sigma_k^2| < \infty ,$$

then  $A_4(\alpha)$  (or  $A_3(x, \alpha)$ ) is finite.

Proof: Throughout this and following proofs,  $c, c', c'', \dots$  will denote positive constants that arise in the arguments, the values of which are unimportant to the discussion. For a sequence  $a_k \rightarrow 1$ , a straightforward computation yields

$$|\Phi(a_k x) - \Phi(x)| \leq \Phi'(x) |(a_k - 1)x| (1 + cx^2 |a_k - 1| \exp(-\frac{1}{2}\beta_k x^2)) ,$$

where  $\beta_k = 2 - a_k > 0$  as soon as  $a_k < 2$ . Thus, for  $k$  sufficiently large,

$$\sup_x |\Phi(a_k x) - \Phi(x)| \leq c' \sup_x \Phi'(x) |(a_k - 1)x| .$$

By taking  $c'$  sufficiently large, this inequality can be made valid for all  $k$ .

Let  $a_k = \sigma_k^{-1}$ . Then



$$\begin{aligned}
\sum_n n^{-\alpha} \sup_x |E\Phi(x/\sigma_{\tau_n}) - \Phi(x)| &\leq \sum_n n^{-\alpha} E(\sup_x |\Phi(x/\sigma_{\tau_n}) - \Phi(x)|) \\
&= \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \sup_x |\Phi(x/\sigma_k) - \Phi(x)| \\
&\leq c \cdot \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \cdot \sup_x \{|\Phi'(x)| \cdot |x/\sigma_k - x|\} \\
&= c \cdot \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \cdot \sup_x \{ |x\Phi'(x)| \cdot \left| \frac{1-\sigma_k}{\sigma_k} \right| \}.
\end{aligned}$$

Now,  $|x\Phi'(x)|$  is bounded, so

$$\begin{aligned}
\sum_n n^{-\alpha} \sup_x |E\Phi(x/\sigma_{\tau_n}) - \Phi(x)| &\leq c' \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \cdot \left| \frac{1-\sigma_k}{\sigma_k} \right| \\
&= c' \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \cdot \frac{|1-\sigma_k|^2}{\sigma_k(1+\sigma_k)} \\
&\leq c'' \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \cdot |1-\sigma_k|^2,
\end{aligned}$$

since  $\sigma_k$  and  $(1+\sigma_k)$  are bounded away from zero. But then

$$A_4(\alpha) \leq A_2(\alpha) + c'' \sum_n n^{-\alpha} \sum_k P[\tau_n=k] \cdot |1-\sigma_k|^2, \quad (2)$$

and the finiteness of  $A_4$  follows from the finiteness of the right-hand side of (2). Similar reasoning yields the result for  $A_3(x, \alpha)$ .  $\blacksquare$



Theorems giving sufficient conditions for the finiteness of  $A_1$  or  $A_2$  thus become theorems for  $A_3$  or  $A_4$  with the addition of condition 1 of the lemma. This fact will not usually be repeated in the following.



II. Convergence of the  $A_j$ 's when the Sequences  $\{\tau_n\}$  and  $\{X_i\}$  are Independent.

In this section, we assume that  $\{\tau_n\}$  is an independent sequence and is independent of  $\{X_n\}$ . Our basic theorem states, that in order for  $A_2$  (or  $A_4$ ) to be finite, it is sufficient that the probability that a given value  $k$  recurs in a sequence of  $\tau_n$ 's diminishes rapidly as  $n$  increases:

Theorem II A. Let  $E|X_i|^{2+\delta} < \infty$ ,  $0 \leq \delta < 1$ . If

$$\sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n=k] = O(k^{-1+\delta/2}), \quad (1)$$

then  $A_2(1-\delta/2) < \infty$ . If  $\delta > 0$  or  $\delta = 0$  but  $EX_i^2 \log|X_i| < \infty$ , then  $A_4(1) < \infty$ .

Proof:

$$\begin{aligned} & \sum_{n=1}^{\infty} n^{-1+\delta/2} \sup_x |\bar{G}_n(x) - E\bar{\Phi}(x/\sigma_{\tau_n})| \\ & \leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P[\tau_n=k] \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \\ & = \sum_{k=1}^{\infty} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \cdot \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n=k] \\ & \leq c \cdot \sum_{k=1}^{\infty} k^{-1+\delta/2} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)|. \end{aligned}$$





This final sum is finite by the F-K-K Theorem. If  $EX_1^2 \log|X_1| < \infty$ , then  $A_4(1) < \infty$  by the second part of the F-K-K Theorem. ■

The condition  $\sum_n n^{-1+\delta/2} P[\tau_n=k] = O(k^{-1+\delta/2})$  is a basic one. We first give an example in which it does not hold and the conclusion of Theorem A is false. This example provides information concerning the sharpness of Theorem A. Then we give examples of distributions for which the condition does hold.

Example II.1. Let  $\{X_i\}$  be any sequence of non-normal, independent, identically-distributed random variables satisfying  $E|X_1|^{2+\delta} < \infty$ ,  $0 < \delta < 1$ . Let

$$P[\tau_n=k] = \begin{cases} (\log n)^{-1} & \text{for } k = 1 \\ 1 - (\log n)^{-1} & \text{for } k = n, \end{cases}$$


and let  $x_0$  be chosen so that  $|\bar{F}_1(x_0) - \Phi(x_0)| > 0$ . Then

$$\sum_{n=1}^{\infty} n^{-1+\delta/2} \sup_x |\bar{G}_n(x) - \Phi(x)| \geq \left| \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} (\bar{F}_k(x_0) - \Phi(x_0)) \cdot P[\tau_n=k] \right|.$$

But

$$\begin{aligned} & \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} (\bar{F}_k(x_0) - \Phi(x_0)) \cdot P[\tau_n=k] \\ &= \sum_{n=1}^{\infty} \left\{ n^{-1+\delta/2} (\log n)^{-1} \cdot (\bar{F}_1(x_0) - \Phi(x_0)) \right. \\ & \quad \left. + n^{-1+\delta/2} \left( \frac{\log n - 1}{\log n} \right) (\bar{F}_n(x_0) - \Phi(x_0)) \right\}. \end{aligned}$$

By Liapounov's theorem,  $|\bar{F}_n(x_0) - \Phi(x_0)| \leq c \cdot n^{-\delta/2}$ . Thus, for large  $n$ , say  $n > N_0$ ,



$$\begin{aligned}
& \left| \sum_{n=N_0}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} (\bar{F}_k(x_0) - \Phi(x_0)) \cdot P[\tau_n = k] \right| \\
& \geq \left| \sum_{n=N_0}^{\infty} \left\{ n^{-1+\delta/2} (\log n)^{-1} (\bar{F}_1(x_0) - \Phi(x_0)) - c \left( \frac{\log n - 1}{\log n} \right) n^{-1} \right\} \right| \\
& = \left| \sum_{n=N_0}^{\infty} n^{-1} \left\{ c \frac{n^{\delta/2}}{\log n} - c^n \right\} \right|,
\end{aligned}$$

which diverges faster than  $\sum_n n^{-1+\gamma}$  for some  $\gamma > 0$ .

Ibragimov<sup>11</sup> has shown that, for random variables with characteristic function  $f(t)$  such that  $\overline{\lim}_{t \rightarrow \infty} |f(t)| < 1$ , the condition

$$\sup_x |\bar{F}_n(x) - \Phi(x)| = O(n^{-(k+\delta/2)}), \quad k = 1, 2, \dots,$$

$0 < \delta < 1$ , is equivalent to the joint conditions

$$(i) \quad \int_{|x| > z} |x|^{k+2} dF = O(z^{-\delta}), \quad z \rightarrow \infty; \text{ and}$$

(ii) the moments  $\mu_s$  of  $X_1$  up to order  $s = k+2$ , inclusive, coincide with the corresponding moments of the normal distribution.

Thus, no condition of the form  $\sum_n n^{-1+\delta/2} P[\tau_n = k] = O(k^\alpha)$ ,  $\alpha > 0$ , is necessary for  $A_4 < \infty$ ; while the sufficiency of conditions weaker than (1) of Theorem A depends on the specific distribution of  $X_1$ .

Some familiar distributions for  $\{\tau_n\}$  which satisfy the hypotheses of Theorem A are the following:



Example II.2. Let  $\tau_n$  be Poisson, with parameter  $n$ . Here

$$P[\tau_n = k] = e^{-n} \frac{n^k}{k!} .$$

Then

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] &= \frac{1}{k!} \sum_{n=1}^{\infty} n^{-1+\delta/2} n^k e^{-n} \\ &= \frac{1}{k!} \sum_{n=1}^{\infty} n^{k-1+\delta/2} e^{-n} \\ &= \frac{1}{k!} e \sum_{n=1}^{\infty} n^{k-1+\delta/2} e^{-n} . \end{aligned}$$

The summand  $n^{k-1+\delta/2} e^{-n}$  is everywhere positive and monotone decreasing for  $n > 1$ , so

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] &\leq \frac{1}{k!} e + \frac{1}{k!} \int_1^{\infty} t^{k-1+\delta/2} e^{-t} dt \\ &\leq \frac{1}{k!} e + \frac{1}{k!} \int_0^{\infty} t^{k-1+\delta/2} e^{-t} dt \\ &= \frac{1}{k!} e + \frac{1}{k!} \Gamma(k+\delta/2) \\ &= \frac{1}{k!} e + \frac{\Gamma(k+\delta/2)}{\Gamma(k+1)} \\ &= O(k^{-1+\delta/2}) . \quad (\text{See Ref. 19 for last equality.}) \end{aligned}$$



Example II.3. Let  $\tau_n$  be binomial with parameters  $n$  and  $p$ ,  $0 < p < 1$ . Then

$$P[\tau_n = k] = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, \quad \text{for } 0 \leq k < n.$$

We have

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] &= \sum_{n=k}^{\infty} n^{-1+\delta/2} P[\tau_n = k] \\ &= \frac{p^k}{k!} \sum_{n=k}^{\infty} \frac{n!}{(n-k)!} n^{-1+\delta/2} (1-p)^{n-k} \\ &\leq k^{-1+\delta/2} \cdot (p^k/k!) \cdot \sum_{n=0}^{\infty} (n+k)(n+k-1) \dots \\ &\quad (n+1) \cdot (1-p)^n \\ &= k^{-1+\delta/2} \cdot (p^k/k!) \cdot (k!/p^{k+1}) \quad (\text{Ref. 20}) \\ &= O(k^{-1+\delta/2}). \end{aligned}$$

Let  $\Theta$  be the class of real, non-increasing functions  $\theta(t)$  such that for all  $\epsilon > 0$ ,  $\sum_n n^{-1+\epsilon} \theta(n) = \infty$ , and let  $\Psi$  be the class of real, non-increasing functions  $\psi(t)$  such that  $\sum_{n=1}^{\infty} n^{-1} \psi(n) < \infty$ .

Theorem II.B. If the RCLT conditions hold for  $\{X_1\}$  and  $\{\tau_n\}$ , if

$$\sum_{n=1}^{\infty} n^{-1+\delta/2} E(\tau_n^{-\alpha} \psi(\tau_n)) < \infty \quad \text{for some } \psi \in \Psi \cap \Theta, \text{ and an arbitrary}$$

$\alpha$  such that  $\delta/2 \geq \alpha > 0$ ; and if  $a_k = \sum_n n^{-1+\delta/2} P[\tau_n = k]$  is monotone decreasing for large  $k$ ; then  $A_2(1-\delta/2) < \infty$ . If  $\delta > 0$  or  $\delta = 0$  and  $EX_1^2 \log|X_1| < \infty$ , then  $A_4(1) < \infty$ .





Proof:

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-1+\delta/2} E(\tau_n^{-\alpha} \psi(\tau_n)) &= \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} k^{-\alpha} \psi(k) \cdot P[\tau_n=k] \\ &= \sum_{k=1}^{\infty} k^{-\alpha} \psi(k) \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n=k] < \infty. \end{aligned}$$

(The final sum is finite by hypothesis.)

Now,  $\sum_{k=1}^{\infty} k^{-\alpha} \psi(k) a_k < \infty$  and  $a_k$  monotone imply<sup>21</sup> that

$$k^{-\alpha} a_k = O(k^{-1}), \text{ or}$$

$$a_k = \sum_n n^{-1+\delta/2} P[\tau_n=k] = O(k^{-1+\alpha}) = O(k^{-1+\delta/2}).$$


The result now follows from Theorem A. ■

Another approach is the following:

Theorem II.C. Suppose the RCLT conditions hold for  $\{X_i\}$  and  $\{\tau_n\}$  with  $\frac{\tau_n}{n} \xrightarrow{p} c_0$ , a positive constant. Suppose further that there is a positive monotone sequence  $\{\rho_n\}$ ,  $\rho_n = O(\frac{1}{n})$ , such that for all  $n$  and for some function  $\psi \in \Psi$ ,

$$P[|\tau_n - nc_0| > n\rho_n] = O(n^{-1} \psi(n)). \quad (1)$$

Then  $A_2(1-\delta/2) < \infty$ . If, in addition,  $EX_1^2 \log|X_1| < \infty$ , then  $A_4(1) < \infty$ .



Proof: Fix  $k$  sufficiently large so that  $\rho_k < c_0$  and let

$$K_1 = \left[ \frac{k}{c_0 + \rho_k} \right], \quad K_2 = \left[ \frac{k}{c_0 - \rho_k} \right],$$

where  $\rho_k \leq \rho < c_0$ . Here the symbol  $[a]$  denotes the largest integer not exceeding  $a$ . Then

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] &= \left\{ \sum_{n=1}^{K_1} + \sum_{n=K_1+1}^{K_2-1} + \sum_{n=K_2}^{\infty} \right\} (n^{-1+\delta/2} P[\tau_n = k]) \\ &\equiv I_1(k) + I_2(k) + I_3(k). \end{aligned}$$

We consider  $I_2(k)$  first:

$$\begin{aligned} I_2(k) &= \sum_{n=K_1+1}^{K_2-1} n^{-1+\delta/2} P[\tau_n = k] \leq (K_1+1)^{-1+\delta/2} \cdot \left( 2 + \frac{2k\rho_k}{c_0 - \rho_k} \right) \\ &= O(k^{-1+\delta/2}), \quad \text{since } \rho_k = O\left(\frac{1}{k}\right). \end{aligned}$$

$$\begin{aligned} I_3(k) &= \sum_{n=K_2}^{\infty} n^{-1+\delta/2} P[\tau_n = k] \leq K_2^{-1+\delta/2} \sum_{n=K_2}^{\infty} P[\tau_n = k] \\ &\leq K_2^{-1+\delta/2} \sum_{n=K_2}^{\infty} n^{-1} \psi(n) \\ &= O(k^{-1+\delta/2}), \quad \text{since } \sum_{n=1}^{\infty} n^{-1} \psi(n) < \infty. \end{aligned}$$

We next consider the term involving  $I_1(k) = \sum_{n=1}^{K_1} n^{-1+\delta/2} P[\tau_n = k]$ :



Let

$$\begin{aligned}
 J &= \sum_{k=1}^{\infty} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \cdot I_1(k) \\
 &= \sum_{k=1}^{\infty} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \sum_{n=1}^{\lfloor k/(c_0 + \rho_k) \rfloor} n^{-1+\delta/2} P(\tau_n = k) \dots
 \end{aligned}$$


For fixed  $n$ , let  $\hat{n} = \min\{j/j \geq n(c_0 + \rho_j)\}$ , and note that  $\hat{n}$  is well-defined because of the monotonicity of  $\rho_j$ . Then

$$\begin{aligned}
 J &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=\hat{n}}^{\infty} P(\tau_n = k) \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \\
 &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=\hat{n}}^{\infty} n^{-1} \psi(n) \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \\
 &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=\hat{n}}^{\infty} n^{-\delta/2} \psi(n) \left(\frac{k}{c_0 + \rho_n}\right)^{-1+\delta/2} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \\
 &\leq c \cdot \sum_{k=1}^{\infty} k^{-1+\delta/2} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \cdot \sum_{n=1}^{\infty} n^{-1} \psi(n) < \infty,
 \end{aligned}$$

since both  $\sum_n n^{-1} \psi(n)$  and  $\sum_{k=1}^{\infty} k^{-1+\delta/2} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)|$  are finite.

Thus,

$$\begin{aligned}
 A_2(1-\delta/2) &= \sum_{k=1}^{\infty} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| \cdot \{I_1(k) + I_2(k) + I_3(k)\} \\
 &\leq c' \cdot \sum_{k=1}^{\infty} k^{-1+\delta/2} \sup_x |\bar{F}_k(x) - \bar{\Phi}(x/\sigma_k)| < \infty. \quad \blacksquare
 \end{aligned}$$



It may not always be obvious that a given sequence of stopping times  $\{\tau_n\}$  satisfies the hypotheses of Theorems II.A, II.B, or II.C. In Theorem II.D below, a more easily recognizable class of random variables is treated.

Let  $\{T_n\}$  be a sequence of non-negative, independent, identically distributed random variables, and let  $\tau_n = \sum_{i=1}^n T_i$ . Consider a population of individuals such that when any individual "dies" or "fails", it is replaced by a new individual; and successive individuals live and die independently of one another. If the lifetime of the  $i$ th individual is  $T_i$ , then  $\tau_n$  represents the time of the  $n$ th replacement of a sequence of individuals. In such a situation,  $\{T_i\}$  is called an ordinary renewal process<sup>5</sup>, and  $\tau_n$  is called the  $n$ th renewal epoch.

Theorem II.D. Suppose  $\{X_i\}$  is a sequence of independent, identically-distributed random variables,  $E|X_i|^{2+\delta} < \infty$  for some  $\delta$ ,  $0 \leq \delta < 1$ . Let  $\{T_i\}$  be an integer-valued ordinary renewal process independent of the  $\{X_i\}$ , such that  $P\{T_i > 0\} = 1$ ,  $E|T_i|^{2+\delta} < \infty$  and  $ET_i = m > 0$ . Let  $\tau_n$  be the  $n$ th renewal epoch. Then

$$(i) \quad \sup_x |\bar{G}_n(x) - E\bar{\Phi}(x/\sigma_{\tau_n})| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

$$(ii) \quad A_2(1-\delta/2) < \infty.$$

If, in addition,  $EX_i^2 \log|X_i| < \infty$ , then  $\sup_x |\bar{G}_n(x) - \bar{\Phi}(x)| \rightarrow 0$  and  $A_4(1) < \infty$ .





Proof: Part (i) follows immediately from the strong law of large numbers and Anscombe's theorem, since the first implies  $\frac{\tau_n}{n} \xrightarrow{p} m$ .

(ii) Let  $\eta(k) = \min\{n/\tau_n \geq k\}$ . Then  $P[\eta(k) > n-1] = P[\tau_n \leq k]$ , since  $P[T_i > 0] = 1$  implies that  $P[\tau_n \leq k] = P[\tau_{n-1} < k]$ .

Let  $M = \lceil \frac{\lambda k}{m} \rceil$ , where  $\lambda$  is chosen arbitrarily between zero and one. Note that

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] &\leq \sum_{n=1}^{M-1} n^{-1+\delta/2} P[\tau_n > k] + \sum_{n=M}^{\infty} n^{-1+\delta/2} P[\tau_n \leq k] \\ &\leq \sum_{n=1}^{M-1} n^{-1+\delta/2} P[\tau_n > k-1] \\ &\quad + \sum_{n=M}^{\infty} n^{-1+\delta/2} P[\eta(k) > n-1] \\ &\equiv I_1(k) + I_2(k). \end{aligned}$$

Chebysheff's inequality is applied to  $I_1(k)$  to obtain the following inequality:

$$\begin{aligned} I_1(k) &= \sum_{n=1}^{M-1} n^{-1+\delta/2} P[\tau_n - nm > k-1-nm] \\ &\leq \sum_{n=1}^{M-1} n^{-1+\delta/2} \frac{n E T_1^{2+\delta}}{(k-1-nm)^{2+\delta}}. \end{aligned}$$

Write  $\alpha = \delta/2$  and note that  $M = \frac{\lambda k}{m^v}$ ,  $v \geq 1$ , and that for



$x > \frac{k-1}{2+\delta+m}$  ,  $x^{\alpha(k-1-mx)^{-2-\delta}}$  is monotone decreasing. Thus,

$$I_1(k) \leq ET_1^{2+\delta} \sum_{n=1}^{M-1} n^{\alpha(k-1-nm)^{-2-\delta}} = ET_1^{2+\delta} \sum_{n=1}^{\ell_1} n^{\alpha(k-1-nm)^{-2-\delta}} + ET_1^{2+\delta} \sum_{n=\ell_1}^{M-1} n^{\alpha(k-1-nm)^{-2-\delta}} ,$$

where  $\ell_1 = [\frac{k-1}{2+\delta+m}]$  . It follows that

$$I_1(k) \leq ET_1^{2+\delta} \left(\frac{k-1}{2+\delta+m}\right)^{\alpha} \cdot \frac{k-1}{2+\delta+m} \left\{ (k-1) \cdot \left(1 - \frac{m}{2+\delta+m}\right) \right\}^{-2-\delta} + ET_1^{2+\delta} \int_0^{\lambda k/m + 1} x^{\alpha(k-1-mx)^{-2-\delta}} dx .$$

The first term in the above inequality is of order  $O(k^{-1+\delta/2})$ . Thus,

$$I_1(k) \leq O(k^{-1+\delta/2}) + ET_1^{2+\delta} \cdot \frac{k^{\alpha}}{m^{\alpha} k^{2+\delta}} \int_0^{\frac{\lambda k}{m} + 1} \frac{m^{\alpha} (x/k)^{\alpha}}{\left(1 - \frac{1}{k-m(x/k)}\right)^{2+\delta}} dx = O(k^{-1+\delta/2}) + ET_1^{2+\delta} m^{-\alpha} k^{-1-\alpha} \int_0^{\frac{\lambda k+m}{k}} y^{\alpha} \left(1 - \frac{1}{k} - y\right)^{-2-\delta} dy ,$$

by the substitution of  $y = \frac{mx}{k}$  . It follows that



$$\begin{aligned}
I_1(k) &\leq O(k^{-1+\delta/2}) + ET_1^{2+\delta} m^{-\alpha} k^{-1-\alpha} \cdot c' \left(\frac{\lambda k+m}{k}\right)^{\alpha+1} \int_0^{\frac{\lambda k+m}{k}} \frac{dy}{\left(1-\frac{1}{k}y\right)^{2+\delta}} \\
&= O(k^{-1+\delta/2}) + c'' k^{-1-\alpha} \left\{ \left(1-\frac{1}{k}y\right)^{-1-\delta} \right\}_{y=0}^{y=\frac{\lambda k+m}{k}} \\
&= O(k^{-1+\delta/2}) + c'' k^{-1-\alpha} \left\{ \left(\frac{k(1-\lambda)+m-1}{k}\right)^{-1-\delta} - \left(\frac{k-1}{k}\right)^{-1-\delta} \right\} \\
&= O(k^{-1+\delta/2}) .
\end{aligned}$$

By Stein's theorem<sup>25</sup>, there exists  $p$ ,  $0 < p < 1$ , such that  $P[\eta(k) \geq n] \leq p^n$ . Thus

$$\begin{aligned}
I_2(k) &= \sum_{n=M}^{\infty} n^{-1+\delta/2} P[\eta(k) \geq n-1] \\
&\leq \left(\frac{\lambda k}{m}\right)^{-1-\delta/2} \sum_{n=M}^{\infty} p^{n-1} \leq c \cdot \left(\frac{\lambda}{m}\right)^{-1+\delta/2} k^{-1+\delta/2} \\
&= O(k^{-1+\delta/2}) .
\end{aligned}$$

Therefore,

$$\sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] \leq I_1(k) + I_2(k) = O(k^{-1+\delta/2}) ,$$

and the result follows from Theorem II.A.  $\blacksquare$



We note that in the preceding proof, the positivity of  $\{T_i\}$  was required in order to use the event  $[\eta(k) \geq n-1]$  and its known properties.


Theorems II.C and II.D are related to the following theorem of Baum and Katz<sup>2</sup>:

Theorem: For  $t > 1$ ,  $r > 1$ ,  $1/2 < r/t \leq 1$ , the following statements are equivalent:

- (a)  $E|X_1|^t < \infty$  and  $EX_1 = \mu$ .
- (b)  $\sum_{n=1}^{\infty} n^{r-2} P[|S(n) - n\mu| > n^{r/t} \epsilon] < \infty$  for all  $\epsilon > 0$ .

For  $r = t = 2$ , statement (b) of Katz's theorem is similar to condition (1) of II.C, with  $\tau_n$  replaced by a sum (see II.D), and with the monotone sequence  $p_n$  replaced by an arbitrary  $\epsilon > 0$ . A condition on  $\{\tau_n\}$  which required that it "resemble", in a suitable way, a sum of random variables might allow the sequence  $\{p_n\}$  in II.C to be replaced by any  $\epsilon > 0$ .

We close this section by considering analogues for random sums to the theorems of Berry and Esseen, and Nagaev. For an independent sequence of non-negative integer-valued random variables  $\{\tau_n\}$  with finite variances  $v_n$  and means  $a_n$ , Orazov and Hudai-Verenov<sup>21</sup> have shown that, if  $E|X_1|^3 < \infty$  and  $\frac{\tau_n}{n} \xrightarrow{P} c > 0$ , then





$$\sup_x |\bar{G}_n(x) - \Phi(x)| \leq B_0 E|X_1|^3 \max(a_n^{-1/2}, a_n^{-2\Delta_n}),$$

where  $\Delta_n = 1 - \frac{\log v_n}{\log a_n}$ , and  $B_0$  is the Berry Esseen constant. A

similar result is the following:

Theorem II.E. If  $E|X_1|^3 < \infty$  and the RCLT conditions hold for  $\{\tau_n\}$  and  $\{X_i\}$ , then

$$|\bar{G}_n(x) - \Phi(x)| \leq \frac{B_0 E|X_1|^3}{1 + |x|^3} \cdot E\left(\frac{1}{\sqrt{\tau_n}}\right).$$

Also,

$$\sup_x |\bar{G}_n(x) - \Phi(x)| \leq B_0 E|X_1|^3 E\left(\frac{1}{\sqrt{\tau_n}}\right).$$

Proof:

$$\begin{aligned} |\bar{G}_n(x) - \Phi(x)| &\leq \sum_{k=1}^{\infty} |\bar{F}_k(x) - \Phi(x)| \cdot P[\tau_n = k] \\ &\leq (1 + |x|^3)^{-1} B_0 E|X_1|^3 \cdot \sum_{k=1}^{\infty} k^{-1/2} P[\tau_n = k] \end{aligned}$$

(by Nagaev's theorem)

$$= B_0 E|X_1|^3 (1 + |x|^3)^{-1} \cdot E\left(\frac{1}{\sqrt{\tau_n}}\right). \quad \blacksquare$$

Note that the theorem of Orazov and Hudai-Verenov deals with moments of  $\tau_n$ , while Theorem II.E is expressed in terms of moments of  $\frac{1}{\tau_n}$ .



III. Convergence of the  $A_j$ 's when the Sequences  $\{\tau_n\}$  and  $\{X_i\}$  are Dependent.

The case where  $\{\tau_n\}$  is no longer required to be independent of  $\{X_i\}$  can involve deeper complications. It will no longer suffice to give conditions based solely on  $\{\tau_n\}$ . Rather, the conditions must be stated in terms of the interaction of  $\{\tau_n\}$  and  $\{X_i\}$ .


Suppose  $\tau_n$  depends only on  $X_1, \dots, X_n$ . Such a random variable is called a stopping time for the sequence  $\{S(n)\}$ . It follows from a Theorem of Renyi<sup>22</sup> on mixing sequences of random variable that for each  $x$  and every  $\epsilon > 0$ , there is a  $K$  such that for  $k > K$ ,

$$|P([\bar{S}(k) \leq x] \cap [\tau_n = k]) - P[\bar{S}(k) \leq x] \cdot P[\tau_n = k]| \leq \epsilon \cdot P[\tau_n = k]. \quad (1)$$

For future reference, let

$$D_{n,k}(x) = |P([\bar{S}(k) \leq x] \cap [\tau_n = k]) - P[\bar{S}(k) \leq x] \cdot P[\tau_n = k]|.$$

The phenomenon exhibited in (1), the asymptotic independence of  $\bar{S}(k)$  from each partial summand, suggests splitting the sums whose convergence we wish to investigate into an "independent" part and an "asymptotically independent" part. Sufficient conditions will be stated in terms of the rate with which  $D_{n,k}(x)$  goes to zero.



Theorem III.A. Let  $\{X_1\}, \{\tau_n\}$  satisfy the RCLT conditions. If

$$(i) \quad \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n = k] = O(k^{-1+\delta/2}),$$

and

$$(ii) \quad \sum_{\{k | P[\tau_n = k] > 0\}} D_{n,k}(x) = O(n^{-\delta/2} \psi(n))$$

for some  $\psi \in \Psi$ , then  $A_1(x, 1-\delta/2) < \infty$ . If, in addition,  $EX_1^2 \log |X_1| < \infty$ , then  $A_2(x, 1) < \infty$ .

Remark: Condition (ii) will certainly be true if

$$|P[\bar{S}(k) \leq x | \tau_n = k] - P[\bar{S}(k) \leq x]| = O(n^{-\delta/2} \psi(n)),$$

for all sufficiently large values of  $k$  for which  $P[\tau_n = k] > 0$ .

Proof: We proceed to split the sum  $A$ , as described above:

$$\begin{aligned} A_1(x, 1-\delta/2) &= \sum_{n=1}^{\infty} n^{-1+\delta/2} |P[\bar{S}(\tau_n) \leq x] - E\Phi(x/\sigma_{\tau_n})| \\ &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{\{k | P[\tau_n = k] > 0\}} |P([\bar{S}(k) \leq x] \cap [\tau_n = k]) \\ &\quad - P[\bar{S}(k) \leq x] \cdot P[\tau_n = k]| \\ &\quad + \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P[\tau_n = k] \cdot |P[\bar{S}(k) \leq x] - \Phi(x/\sigma_k)| \\ &= I_1(x) + I_2(x). \end{aligned}$$

1.  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = \lim_{x \rightarrow 0} \frac{\cos x}{1} = \frac{\cos 0}{1} = \frac{1}{1} = 1$$

2.  $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \lim_{x \rightarrow 0} \frac{\sin x}{2x} = \lim_{x \rightarrow 0} \frac{\cos x}{2} = \frac{\cos 0}{2} = \frac{1}{2}$$

3.  $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = \lim_{x \rightarrow 0} \frac{e^x}{1} = \frac{e^0}{1} = \frac{1}{1} = 1$$

4.  $\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2} = -\frac{1}{2}$

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2} = \lim_{x \rightarrow 0} \frac{e^x - 1}{2x} = \lim_{x \rightarrow 0} \frac{e^x}{2} = \frac{e^0}{2} = \frac{1}{2}$$

5.  $\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2}}{x^3} = \frac{1}{6}$

6.  $\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2} - \frac{x^3}{6}}{x^4} = \frac{1}{24}$

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2} - \frac{x^3}{6}}{x^4} = \lim_{x \rightarrow 0} \frac{e^x - 1}{4x^3} = \lim_{x \rightarrow 0} \frac{e^x}{12x^2} = \lim_{x \rightarrow 0} \frac{e^x}{24x} = \lim_{x \rightarrow 0} \frac{e^x}{24} = \frac{e^0}{24} = \frac{1}{24}$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2} - \frac{x^3}{6} - \frac{x^4}{24}}{x^5} = \lim_{x \rightarrow 0} \frac{e^x - 1}{120x^4} = \lim_{x \rightarrow 0} \frac{e^x}{480x^3} = \lim_{x \rightarrow 0} \frac{e^x}{1440x^2} = \lim_{x \rightarrow 0} \frac{e^x}{2880x} = \lim_{x \rightarrow 0} \frac{e^x}{2880} = \frac{e^0}{2880} = \frac{1}{2880}$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2} - \frac{x^3}{6} - \frac{x^4}{24} - \frac{x^5}{120}}{x^6} = \frac{1}{4320}$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2} - \frac{x^3}{6} - \frac{x^4}{24} - \frac{x^5}{120} - \frac{x^6}{720}}{x^7} = \frac{1}{5040}$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2} - \frac{x^3}{6} - \frac{x^4}{24} - \frac{x^5}{120} - \frac{x^6}{720} - \frac{x^7}{5040}}{x^8} = \frac{1}{40320}$$

Now

$$I_2(x) = \sum_{k=1}^{\infty} |P[\bar{S}(k) \leq x] - \Phi(x/\sigma_k)| \cdot \sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n=k]$$

$$\leq c \cdot \sum_{k=1}^{\infty} k^{-1+\delta/2} |P[\bar{S}(k) \leq x] - \Phi(x/\sigma_k)| ,$$

which is finite by the F-K-K Theorem. But  $I_1(x)$  is just

$$\sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{\{k | P[\tau_n=k] > 0\}} D_{n,k}(x) ,$$

so

$$I_1(x) \leq c \cdot \sum_{n=1}^{\infty} n^{-1} \psi(n) < \infty .$$

Thus

$$A_1(x, 1-\delta/2) \leq I_1(x) + I_2(x) < \infty .$$

The assertion about  $A_3(x, 1)$  follows from II.A. ■

It is not a hypothesis of Theorem III.A. that  $\tau_n$  depend only on  $X_1, \dots, X_n$ . Referring to (1), one can see that the rate of convergence of  $D_{n,k}$  to zero can be dominated either by rapidly decreasing values of  $P[\tau_n=k]$ , or by bounding  $D_{n,k}$  by an expression like  $\epsilon_{n,k} P[\tau_n=k]$ , where  $\epsilon_{n,k} \rightarrow 0$  with suitable speed. Thus, if  $\tau_n$  does in fact depend only on  $X_1, \dots, X_n$ , we need only a slightly more stringent condition on  $\{\tau_n\}$  to satisfy (ii) of Theorem III.A. The following example shows, however, that simple dependence on  $X_1, \dots, X_n$  is not enough.






Example III.1. Let  $X_i \geq 0$  be integer-valued,  $EX_i \geq 1$ ,  $EX_i^{2+\delta} < \infty$ ,  $\delta > 0$ , and let  $\tau_n = S(n)$ . Then

$$\begin{aligned}
 A_3(x,1) &= \sum_{n=1}^{\infty} n^{-1} |P[\bar{S}(\tau_n) \leq x] - \bar{\Phi}(x)| \\
 &= \sum_{n=1}^{\infty} n^{-1} \sum_{k=1}^{\infty} |P([\bar{S}(k) \leq x] \cap [\tau_n = k]) - P[\tau_n = k]| \cdot \bar{\Phi}(x) \\
 &= \sum_{n=1}^{\infty} n^{-1} \sum_{k=1}^{\infty} |P([\bar{S}(k) \leq x] \cap [S(n) = k]) - P[S(n) = k]| \cdot \bar{\Phi}(x) \\
 &= \sum_{n=1}^{\infty} n^{-1} \left\{ \sum_{k=1}^{n-1} |P([S(n-k) \leq x\sqrt{k} + k(\mu-1)] \cap [S(n) = k]) \right. \\
 &\quad \left. - P[S(n) = k] \cdot \bar{\Phi}(x) \right| \\
 &\quad + \sum_{k=n}^{\infty} |P([S(k-n) \leq x\sqrt{k} + k(\mu-1)] \cap [S(n) = k]) \\
 &\quad \left. - P[S(n) = k] \cdot \bar{\Phi}(x) \right\} \\
 &\geq \sum_{n=1}^{\infty} n^{-1} \sum_{k=n}^{\infty} P[S(n) = k] \cdot |P[S(k-n) \leq x\sqrt{k} + k(\mu-1)] - \bar{\Phi}(x)|.
 \end{aligned}$$

Now,  $P[S(k-n) \leq x\sqrt{k} + k(\mu-1)] \rightarrow 0$  as  $k \rightarrow \infty$  since

$$P[S(k-n) \leq x\sqrt{k} + k(\mu-1)] = P\left[\bar{S}(k-n) \leq x\sqrt{\frac{k}{k-n}} + \frac{n\mu-k}{\sqrt{k-n}}\right].$$



Let the quantity  $a(n,x)-1$  be the last integer for which  $|P[S(k-n) \leq x\sqrt{k} + k(\mu-1)] - \Phi(x)|$  is less than some constant  $d$  where  $\Phi(x) \geq d > 0$ . Then there is a constant  $c_1 > 0$  for which

$$\begin{aligned} A_3(x,1) &\geq c_1 \Phi(x) \sum_{n=1}^{\infty} n^{-1} \sum_{k=a(n,x)}^{\infty} P[S(n)=k] \\ &= c_1 \Phi(x) \sum_{n=1}^{\infty} n^{-1} P[S(n) \geq a(n,x)] . \end{aligned}$$

If  $a(n,x) \leq \mu n + c$ ,  $c > 0$ , then  $P[S(n) \geq a(n,x)] \rightarrow 1$ , and  $A_3(x,1) = \infty$ . To be more specific, then, let

$$P[X_1=m] = \begin{cases} \frac{1}{2} & \text{for } m = 2 \\ \frac{1}{2} & \text{for } m = 0 \\ 0 & \text{otherwise .} \end{cases}$$

Here,  $\mu = 1$ . Consider  $A_3(0,1)$  and note first that  $a(n,x)-1$  is the largest integer for which  $|P[S(k-n) \leq 0] - 1/2|$  is less than some constant  $d$ ,  $1/2 \geq d > 0$ . Now  $P[S(k-n) \leq 0] = (1/2)^{k-n}$ , so  $a(n,x) = n+2$ . From the above,  $A_3(0,1) = \infty$ . Notice that the sequences  $\{X_i\}$ ,  $\{\tau_n\}$  satisfy the RCLT conditions and condition (i) of Theorem III.A. For let  $k = 2l$  (otherwise  $P[S(n)=k]=0$ ). Then



$$\begin{aligned}
\sum_{n=1}^{\infty} n^{-1} P[S(n)=k] &= \sum_{n=l}^{\infty} n^{-1} \binom{n}{l} \left(\frac{1}{2}\right)^n \leq \left(\frac{k}{2}\right)^{-1} \sum_{n=l}^{\infty} \binom{n}{l} \left(\frac{1}{2}\right)^n \\
&= \left(\frac{k}{2}\right)^{-1} \frac{1}{l!} \sum_{n=0}^{\infty} (n+l)(n+l-1) \cdots (n-1) \left(\frac{1}{2}\right)^{n+l} \\
&\leq \left(\frac{k}{2}\right)^{-1} \frac{1}{l!} \frac{l!}{\left(\frac{1}{2}\right)^{l+1}} \left(\frac{1}{2}\right)^l \quad (\text{Ref. 20}) \\
&= O(k^{-1}).
\end{aligned}$$

For future convenience, let  $H_{n,k} = [\tau_n = k]$ . Now consider the case where  $[\bar{S}(\tau_n) \leq x]$  and  $[\bar{S}(n) \leq x]$  have, in some sense, the same asymptotic degree of dependence on  $H_{n,k}$ . One of the problems that must be faced is the question of convergence of

$$R \equiv \sum_{n=1}^{\infty} n^{-1+\delta/2} \sup_x |\Phi(x/\sigma_n) - E\Phi(x/\sigma_{\tau_n})|. \quad (2)$$

Recall that we assume  $\sigma_i \neq 0$  for all  $i > 0$ .

Lemma: Let  $\{X_i\}, \{\tau_n\}$  satisfy the RCLT conditions. If, for some  $\psi \in \Psi$ ,

$$\sum_{k=1}^{\infty} P(H_{n,k}) |\sigma_k^2 - \sigma_n^2| = O(n^{-\delta/2} \psi(n)), \quad (3)$$

then  $R$  is finite.



Proof:

$$\begin{aligned}
 R &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P(H_{n,k}) \sup_x |\Phi(x/\sigma_n) - \Phi(x/\sigma_k)| \\
 &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P(H_{n,k}) \sup_x \{ \Phi'(x/\sigma_n) \cdot (1+\theta_{n,k}) \cdot \left| \frac{x}{\sigma_n} - \frac{x}{\sigma_k} \right| \},
 \end{aligned}$$

as in the lemma of section I. Here,  $\theta_{n,k} \rightarrow 0$  as  $k \rightarrow \infty$  for every  $n$ . Thus

$$\begin{aligned}
 R &\leq c \cdot \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P(H_{n,k}) \sup_x |\Phi'(x/\sigma_n) \cdot \frac{x}{\sigma_n} \cdot \frac{\sigma_k - \sigma_n}{\sigma_k}| \\
 &\leq c' \cdot \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P(H_{n,k}) \left| \frac{\sigma_k - \sigma_n}{\sigma_k} \right|,
 \end{aligned}$$

because  $t\Phi'(t)$  is bounded. Since  $\sigma_k$  and  $\sigma_n$  are bounded and nonzero,

$$R \leq c'' \cdot \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{k=1}^{\infty} P(H_{n,k}) |\sigma_k^2 - \sigma_n^2| < \infty \quad (\text{by hypothesis}). \quad \blacksquare$$

Theorem III.B. Let  $\{X_1\}$ ,  $\{\tau_n\}$  satisfy the RCLT conditions and condition (3) of the lemma. If there is a function  $f(n,k)$  and a sequence  $\{a_n\}$  such that

$$(i) \quad \sum_n a_n n^{-1+\delta/2} < \infty,$$

$$(ii) \quad E f(n, \tau_n) = \sum_k f(n,k) P(H_{n,k}) \leq c_0 a_n \quad \text{for each } n,$$

when  $c_0$  is a fixed positive constant; and

$$(iii) \quad |P([\bar{S}(k) \leq x] \cap H_{n,k}) - P([\bar{S}(n) \leq x] \cap H_{n,k})| \leq f(n,k) \cdot P(H_{n,k}),$$

then  $A_2(1-\delta/2) < \infty$ . If, in addition,  $E X_1^2 \log |X_1| < \infty$ , then

$$A_4(1) < \infty.$$





Proof:

$$\begin{aligned}
 A_2(1-\delta/2) &= \sum_{n=1}^{\infty} n^{-1+\delta/2} \sup_x |P[\bar{S}(\tau_n) \leq x] - E\bar{\Phi}(x/\sigma_{\tau_n})| \\
 &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \left\{ \sup_x |P[\bar{S}(\tau_n) \leq x] - P[\bar{S}(n) \leq x]| \right. \\
 &\quad \left. + \sup_x |P[\bar{S}(n) \leq x] - \bar{\Phi}(x/\sigma_n)| + \sup_x |\bar{\Phi}(x/\sigma_n) - E\bar{\Phi}(x/\sigma_{\tau_n})| \right\} \\
 &= I_1 + I_2 + I_3 .
 \end{aligned}$$

$I_3 < \infty$  by the lemma, and  $I_2 < \infty$  by the F-K-K Theorem.


$$\begin{aligned}
 I_1 &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_{\{k | P(H_{n,k}) > 0\}} P(H_{n,k}) \sup_x |P[\bar{S}(k) \leq x] - P[\bar{S}(n) \leq x | H_{n,k}]| \\
 &\leq \sum_{n=1}^{\infty} n^{-1+\delta/2} \sum_k P(H_{n,k}) f(n,k) \leq c \cdot \sum_{n=1}^{\infty} n^{-1+\delta/2} a_n < \infty .
 \end{aligned}$$

Thus  $A_2(1-\delta/2) \leq I_1 + I_2 + I_3 < \infty$  . ■

One of the classical stopping times of sequential analysis is the first time the sum  $S(k)$  exceeds a given value; e.g.,

$$\tau_n = \min\{k | S(k) \geq n\} .$$

To see how this sequence  $\{\tau_n\}$  can lead to overdependence of  $[\bar{S}(k) \leq x]$  on  $H_{n,k}$  and thus to divergence of  $A_3$ , consider the following example.



Example III.2. Let  $X_i$  be zero with probability  $q$ ,  $1 > q > 0$ , and one with probability  $p, q = 1-p$ . Let

$$\tau_1 = \min\{i | X_i = 1\}, \quad \tau_n = \tau_{n-1} + \min\{i | X_{\tau_{n-1} + i} = 1\}.$$

It follows that

$$\begin{aligned} A_3(0,1) &= \sum_{n=1}^{\infty} n^{-1} \sum_{k=1}^{\infty} |P([S(k) \leq kp] \cap [\tau_n = k]) - \Phi(x) \cdot P[\tau_n = k]| \\ &= \sum_{n=1}^{\infty} n^{-1} \sum_{k=1}^{\infty} |P([S(k) \leq kp] \cap [S(k-1) = n-1] \cap [X_k = 1]) - \Phi(x) \\ &\quad \cdot P([S(k-1) = n-1] \cap [X_k = 1])|, \end{aligned}$$

since  $[\tau_n = k] = [S(k-1) = n-1] \cap [X_k = 1]$ . Thus

$$\begin{aligned} A_3(0,1) &= \sum_{n=1}^{\infty} n^{-1} \left\{ \sum_{k=n}^{\lfloor n/p \rfloor} p \Phi(x) \binom{k-1}{n-1} p^{n-1} q^{k-n} \right. \\ &\quad \left. + \sum_{k=\lfloor n/p \rfloor + 1}^{\infty} |p \binom{k-1}{n-1} p^{n-1} q^{k-n} - \Phi(x) p \binom{k-1}{n-1} p^{n-1} q^{k-n}| \right\} \\ &= \Phi(x) \cdot \sum_{n=1}^{\infty} n^{-1} \left\{ \sum_{k=n}^{\lfloor n/p \rfloor} \binom{k-1}{n-1} p^n q^{k-n} + \frac{1 - \Phi(x)}{\Phi(x)} \sum_{k=\lfloor n/p \rfloor + 1}^{\infty} \binom{k-1}{n-1} p^n q^{k-n} \right\} \\ &\geq c' \cdot \sum_{n=1}^{\infty} n^{-1} \sum_{k=n}^{\infty} \frac{n}{k} \binom{k}{n} p^n q^{k-n} = \sum_{k=1}^{\infty} k^{-1} \sum_{n=1}^{\infty} \binom{k}{n} p^n q^{k-n} \\ &= \sum_{k=1}^{\infty} k^{-1} (1-q^k) = \infty. \end{aligned}$$



The problem we have encountered in this example is that the sum  $S(\tau_n)$  is completely specified. In example III.1, a similar situation occurred--namely, the value of  $\tau_n$  was too closely tied to the value of the summand  $S(n)$ .

In the next example, we consider a sequence  $\{\tau_n\}$  which, though dependent on  $X_1, \dots, X_n$ , does not lead to a completely specified value of  $S(\tau_n)$  and whose dependence on  $S(n)$  is such that the hypothesis of Theorem III.A are satisfied.

Example III.3. Let  $\{X_i\}$  be an independent sequence of identically-distributed random variables with mean  $\mu$  and unit variance. Let  $\epsilon_0$  be a fixed positive constant. Define a sequence  $\{\tau_n\}$  as follows:

$$\tau_n = \begin{cases} n, & \text{if } S(n) \leq n(\mu + \epsilon_0) \\ 2n, & \text{if } S(n) > n(\mu + \epsilon_0) \end{cases}.$$

Note that the event  $[\tau_n = k]$  is the same as the event  $[S(n) \leq n(\mu + \epsilon_0)]$  for  $k = n$ , and the same as  $[S(n) > n(\mu + \epsilon_0)]$  for  $k = 2n$ . The sum  $\sum_{\{k | P[\tau_n = k] > 0\}} D_{n,k}(x)$  becomes, for  $x = 0$ , simply

$$\begin{aligned} \sum_{\{k | P[\tau_n = k] > 0\}} D_{n,k}(0) &\leq |P[S(n) \leq n\mu] \cap [S(n) \leq n(\mu + \epsilon_0)]| \\ &\quad - P[S(n) \leq n\mu] \cdot P[S(n) \leq n(\mu + \epsilon_0)] \\ &+ |P[[S(2n) \leq 2n\mu] \cap [S(n) > n(\mu + \epsilon_0)]]| \\ &\quad - P[S(2n) \leq 2n\mu] \cdot P[S(n) > n(\mu + \epsilon_0)]|. \end{aligned}$$



Consider the first term:

$$\begin{aligned}
 & |P([S(n) \leq n\mu] \cap [S(n) \leq n(\mu + \epsilon_0)]) - P[S(n) \leq n\mu] \cdot P[S(n) \leq n(\mu + \epsilon_0)]| \\
 &= |P[S(n) \leq n\mu] - P[S(n) \leq n\mu] \cdot P[S(n) \leq n(\mu + \epsilon_0)]| \\
 &= P[S(n) \leq n\mu] \cdot (1 - P[S(n) \leq n(\mu + \epsilon_0)]) \\
 &= P[S(n) \leq n\mu] \cdot P[S(n) > n(\mu + \epsilon_0)] .
 \end{aligned}$$

Let  $S^*(n) = S(2n) - S(n)$ . Then, for the second term,

$$\begin{aligned}
 & |P([S(2n) \leq 2n\mu] \cap [S(n) > n(\mu + \epsilon_0)]) - P[S(2n) \leq 2n\mu] \\
 & \qquad \qquad \qquad \cdot P[S(n) > n(\mu + \epsilon_0)]| \\
 &= |P([S^*(n) \leq 2n\mu - S(n)] \cap [S(n) > n(\mu + \epsilon_0)]) - P[S(2n) \leq 2n\mu] \\
 & \qquad \qquad \qquad \cdot P[S(n) > n(\mu + \epsilon_0)]| .
 \end{aligned}$$

It is certainly the case that

$$\begin{aligned}
 [S^*(n) \leq 2n\mu - S(n)] \cap [S(n) > n(\mu + \epsilon_0)] &\subset [S^*(n) \leq n(\mu - \epsilon_0)] \\
 &\cap [S(n) > n(\mu + \epsilon_0)] .
 \end{aligned}$$

Thus

$$\begin{aligned}
 P([S^*(n) \leq 2n\mu - S(n)] \cap [S(n) > n(\mu + \epsilon_0)]) &\leq P([S^*(n) \leq n(\mu - \epsilon_0)] \\
 &\cap [S(n) > n(\mu + \epsilon_0)]) .
 \end{aligned}$$





By the strong law of large numbers and the central limit theorem, for sufficiently large  $n$ ,  $P[S^*(n) \leq n(\mu - \epsilon_0)] < P[S(2n) \leq 2n\mu]$ , since the former goes to zero with  $n$ , and the latter to one-half.

Thus,

$$\begin{aligned} & P([S^*(n) \leq n(\mu - \epsilon_0)] \cap [S(n) > n(\mu + \epsilon_0)]) \\ &= P[S^*(n) \leq n(\mu - \epsilon_0)] \cdot P[S(n) > n(\mu + \epsilon_0)] \\ &< P[S(2n) \leq 2n\mu] \cdot P[S(n) > n(\mu + \epsilon_0)] . \end{aligned}$$

But, for sufficiently large  $n$ , it follows that

$$\begin{aligned} & |P([S^*(n) \leq 2n\mu - S(n)] \cap [S(n) > n(\mu + \epsilon_0)]) \\ & \quad - P[S(2n) \leq 2n\mu] \cdot P[S(n) > n(\mu + \epsilon_0)]| \\ &= P[S(2n) \leq 2n\mu] \cdot P[S(n) > n(\mu + \epsilon_0)] \\ & \quad - P([S^*(n) \leq 2n\mu - S(n)] \cap [S(n) > n(\mu + \epsilon_0)]) \\ &\leq P[S(2n) \leq 2n\mu] \cdot P[S(n) > n(\mu + \epsilon_0)] . \end{aligned}$$


Hence

$$\sum D_{n,k}(x) \leq 2 \cdot P[S(n) > n(\mu + \epsilon_0)] .$$

If  $E|X_1|^{2+\delta} < \infty$ ,  $\delta \geq 0$ , then Chebysheff's inequality yields

$$\begin{aligned} \sum D_{n,k}(x) &\leq 2 \cdot P[S(n) > n(\mu + \epsilon_0)] \\ &\leq \frac{2 \cdot n \cdot E|X_1|^{2+\delta}}{\epsilon_0^{2+\delta} n^{2+\delta}} = O(n^{-\delta/2} \psi(n)) , \end{aligned}$$

and condition (ii) of Theorem III.A is satisfied.



Condition (i) of Theorem III.A is easily verified, since

$$\sum_{n=1}^{\infty} n^{-1+\delta/2} P[\tau_n=k] = p_1 k^{-1+\delta/2} + p_2 \left(\frac{k}{2}\right)^{-1+\delta/2},$$

where  $p_1$  and  $p_2$  are the probabilities that  $\tau_n = n$  and  $2n$ , respectively. We also note that  $\{X_i\}$  and  $\{\tau_n\}$  satisfy the RCLT conditions, since  $\frac{\tau_n}{n} \rightarrow 1$  by the weak law of large numbers. Thus, by Theorem III.A,  $A_1(0, 1-\delta/2) < \infty$ .

This same example serves as an illustration of Theorem III.B. Referring to the theorem of Baum and Katz<sup>2</sup>, recall that if  $E|X_i|^{2+\delta} < \infty$ , then

$$\sum_{n=1}^{\infty} n^{\delta} \cdot P[|S(n) - n\mu| > nc_0] < \infty \text{ for } c_0 > 0.$$

For condition (iii) of III.B, we have for  $k = n$ ,

$$\begin{aligned} & |P([S(n) \leq n\mu] \cap [S(n) \leq n(\mu + \epsilon_0)]) - P([S(n) \leq n\mu] \\ & \cap [S(n) \leq n(\mu + \epsilon_0)])| = 0, \end{aligned}$$

and for  $k = 2n$ ,

$$\begin{aligned} & |P([S(2n) \leq 2n\mu] \cap [S(n) > n(\mu + \epsilon_0)]) - P([S(n) \leq n\mu] \\ & \cap [S(n) > n(\mu + \epsilon_0)])| \\ & \leq P[S^*(n) \leq n(\mu - \epsilon_0)] \cdot P[S(n) > n(\mu + \epsilon_0)]. \end{aligned}$$

For other values of  $k$ ,  $P[\tau_n=k] = 0$ . Using the notation of III.B,



$$f(n,k) = 0 \text{ for } k \neq 2n,$$

$$f(n,k) = P[S^*(n) \leq n(\mu + \epsilon_0)] \text{ for } k = 2n.$$

Thus

$$\sum_k f(n,k) P(H_{n,k}) \leq P[S^*(n) \leq n(\mu - \epsilon_0)] \cdot P[S(n) > n(\mu + \epsilon_0)],$$

and by the theorem of Baum and Katz,

$$\sum n^{-1+\delta/2} P[S(n) > n(\mu + \epsilon_0)] \cdot P[S^*(n) \leq n(\mu - \epsilon_0)] < \infty$$

with plenty to spare. The only condition of III.B not yet verified is the condition that

$$R = \sum_{k=1}^{\infty} P(H_{n,k}) \cdot |\sigma_k^2 - \sigma_n^2| = O(n^{-\delta/2} \psi(n)).$$

In this example,

$$R = |\sigma_{2n}^2 - \sigma_n^2| \cdot P[S(n) > n(\mu + \epsilon_0)].$$

Here  $|\sigma_{2n}^2 - \sigma_n^2|$  is bounded, and

$$P[S(n) > n(\mu + \epsilon_0)] = O(n^{-\delta/2} \psi(n))$$

by Theorem 3 and the lemma of the paper by Baum and Katz<sup>2</sup>. Thus the proof of Theorem III.B also yields  $A_1(0, 1-\delta/2) < \infty$ .

(1)  $\int_{-\infty}^{\infty} \delta(x) dx = 1$

(2)  $\int_{-\infty}^{\infty} f(x) \delta(x-a) dx = f(a)$

(3)  $\int_{-\infty}^{\infty} f(x) \delta(x) dx = f(0)$

(4)  $\int_{-\infty}^{\infty} f(x) \delta(x-a) dx = f(a)$

(5)  $\int_{-\infty}^{\infty} f(x) \delta(x) dx = f(0)$

$$\int_{-\infty}^{\infty} f(x) \delta(x-a) dx = f(a)$$

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

$$\int_{-\infty}^{\infty} f(x) \delta(x-a) dx = f(a)$$

$$\int_{-\infty}^{\infty} f(x) \delta(x) dx = f(0)$$

$$\int_{-\infty}^{\infty} f(x) \delta(x-a) dx = f(a)$$

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

The Dirac delta function is a distribution that is zero everywhere except at a single point, where it is infinite. It is used to model point charges, impulses, and other phenomena that are localized in space or time.

We have seen examples where the sufficiency conditions of Theorems III.A and III.B do not hold, and the series in question diverge, and an example where they do hold. This indicates that these conditions are reasonably sharp. However, the search for interesting necessary conditions has been fruitless. The openness of the necessity question for independent  $\{\tau_n\}$  suggests that no condition more stringent than

$$\sum_n n^{-1+\delta/2} P[\tau_n=k] < \infty$$

is likely to be, in general, necessary in the dependent case.






## BIBLIOGRAPHY

The following papers and books are referenced in the text:

1. Anscombe, F.J.: Large-sample Theory of Sequential Estimation. Proc. Cambridge Phil. Soc. 48, 1952, pp. 600-607.
2. Baum, L.E. and Katz, M.: Convergence Rates in the Law of Large Numbers. Trans. Am. Math. Soc. 120, 1965, pp. 103-123.
3. Berry, A.C.: The Accuracy of the Gaussian Approximation to the Sum of Independent Variates. Trans. Am. Math. Soc. 49, 1941, pp. 122-136.
4. Blum, J.R.; Hanson, D.L.; Rosenblatt, J.I.: On the Central Limit Theorem for the Sum of a Random Number of Independent Random Variables. Zeit. Wahrscheinlichkeitsth. verw. Geb. 1, 1962/63, pp. 389-393.
5. Cox, D.R.: Renewal Theory. Methuen and Co., Ltd., London, 1962.
6. Davis, J.A.: Convergence Rates for the Law of the Iterated Logarithm. Ann. Math. Stat., 39(5), 1968, pp. 1479-1485.
7. Esseen, C.G.: Fourier Analysis of Distribution Functions. A Mathematical Study of the Laplace-Gaussian Law. Acta Math., 77, 1945, pp. 1-25.
8. Feller, W.: Über den zentralen Grenzwertsatz der Wahrscheinlichkeitsrechnung. Math. Zeitschrift, 40, 1935, pp. 521-559.
9. Friedman, N.; Katz, M.; Koopmans, L.H.: Convergence Rates for the Central Limit Theorem. Proc. Nat. Acad. Sci. USA., 56, 1966, pp. 1062-1065.
10. Heyde, C.C.: On the Influence of Moments on the Rate of Convergence to the Normal Distribution. Zeit. Wahrscheinlichkeitsth. verw. Geb., 8, 1967, pp. 12-18.
11. Ibragimov, I.A.: On the Accuracy of the Gaussian Approximation to the Distribution Function of Sums of Independent Variables. Th. Prob. Appl. 11, 1966, pp. 559-576.
12. Katz, M.: Note on the Berry-Esseen Theorem. Ann. Math. Stat. 34, 1963, pp. 1107-1108.



13. Knopp, K.: Theory and Application of Infinite Series. Blackie and Sons, Ltd., London, 1928, p. 124.
14. Lebedev, N.N.: Special Functions and Their Applications. Prentice-Hall, Englewood Cliffs, 1965, p. 15.
15. Liapounov, A.M.: Sur une proposition de la théorie des probabilités. Bull. de l'Acad. Imp. des. Sci. de St. Petersbourg. (5)13, no. 4, 1900, pp. 359-386.
16. Liapounov, A.M.: Nouvelle forme du théorème sur la limite de probabilités. Mem. Acad. Sci. St. Petersbourg. (8) 12, no. 5, 1901, pp. 1-24.
17. Lindeberg, J.W.: Eine neue Herleitung des Exponential-gesetzes in der Wahrscheinlichkeitsrechnung. Math. Zeitschrift, 15, 1922, pp. 211-225.
18. Mangulis, V.: Handbook of Series for Scientists and Engineers. Academic Press, N.Y., 1965, p. 72.
19. Mogyoródi, J.: A Central Limit Theorem for the Sum of a Random Number of Independent Random Variables. Magyar Tud. Akad. Mat. Kutató Int. Közl., 7A, 1962, pp. 409-424.
20. Nagaev, S.V.: Some Theorems for Large Deviations. Th. Prob. Appl. 10, 1965, pp. 214-235.
21. Orazov, G. and Hudai-Verenov, M.: The Rapidity of Convergence to the Normal Law (Russian). Izv. Akad. Nauk. Turkmen. SSR Ser. Fiz.-Tehn. Him. Geol. Nauk, 1967, No. 3, pp. 3-9.
22. Renyi, A.: On the Central Limit Theorem for the Sum of a Random Number of Independent Random Variables. Acta Math. Hung. 8, 1957, pp. 97-102.
23. Renyi, A.: On Mixing Sequences of Sets. Acta Math. Hung. 9, 1958, pp. 215-228.
24. Robbins, H.: The Asymptotic Distribution of the Sum of a Random Number of Independent Random Variables. Bull. Am. Math. Soc. 54, 1948, pp. 1151-1161.
25. Stein, C.: A Note on Cumulative Sums. Ann. Math. Stat., 19, 1948, pp. 498-499.
26. Wald, A.: Sequential Analysis. Wiley, New York, 1947.
27. Wittenberg, H.: Limiting Distributions of Random Sums of Independent Random Variables. Zeit. Wahrscheinlichkeitsth. verw. Geb. 3, 1964, pp. 7-19.

1. The first part of the document discusses the general principles of the law of contract.

2. The second part of the document discusses the formation of a contract.

3. The third part of the document discusses the performance of a contract.

4. The fourth part of the document discusses the breach of a contract.

5. The fifth part of the document discusses the remedies available for breach of contract.

6. The sixth part of the document discusses the discharge of a contract.

7. The seventh part of the document discusses the assignment of a contract.

8. The eighth part of the document discusses the sub-contracting of a contract.

9. The ninth part of the document discusses the novation of a contract.

10. The tenth part of the document discusses the rescission of a contract.

11. The eleventh part of the document discusses the frustration of a contract.


12. The twelfth part of the document discusses the termination of a contract.

13. The thirteenth part of the document discusses the assignment of a contract.

14. The fourteenth part of the document discusses the sub-contracting of a contract.

In addition to the bibliography, the following works were consulted and provide a background for the material treated in the dissertation.

1. Chung, K.L.: A Course in Probability Theory. Harcourt, Brace, and Wald. New York, 1968. (Stein's Theorem, Berry-Esseen Theorem, random walks.)
2. Cramér, H.: Random Variables and Probability Distributions, 3rd Ed., Cambridge Univ. Press, 1970. (Classic results on the central limit problem.)
3. Feller, W.: The Fundamental Limit Theorems in Probability. Bull. AMS, vol. 51, 1945, pp. 800-832.
4. Feller, W.: On the Berry-Esseen Theorem. Z. Wahrscheinlichkeitsth. verw. Geb. 10, 1968, pp. 261-268. (Simple derivation of a generalized Berry-Esseen theorem not requiring moments.)
5. Heathcote, C.R.: Complete Exponential Convergence and Some Related Topics. J. Appl. Prob. 4, 1967, pp. 217-256. (Summary of convergence rate results for the central limit theorem and laws of large numbers.)
6. Hertz, Ellen S.: On Convergence Rates in the Central Limit Theorem. Ann. Math. Stat. 40, no. 2, 1969, pp. 475-479.  
(Convergence rates in terms of  $\sum_{|x| > c}^n \int x^2 dF_i(x)$ .)
7. Heyde, C.C.: Some Properties of Metrics in a Study on Convergence to Normality. Zeit. Wahrscheinlichkeitsth. verw. Geb. 11, 1969, pp. 181-192. (Use of  $\mathcal{L}_1$  and  $\mathcal{L}_\infty$  metrics in a study of convergence rates in the central limit theorem.)
8. Philipp, W.: The Remainder in the Central Limit Theorem for Mixing Stochastic Processes. Ann. Math. Stat., 40, no. 3, 1969, pp. 601-609. (Berry-Esseen-like rate theorems for mixing processes.)
9. Richter, W.: Limit Theorems for Sequences of Random Variables with Sequences of Random Indices. Theor. Prob. Appl. 10, no. 1, 1965, pp. 74-84.
10. Robbins, H.: Optimal Stopping. The Am. Math. Monthly, vol. 77, no. 4, April 1970, pp. 333-343.
11. Rosen, B.: On the Asymptotic Distribution of Sums of Independent Identically Distributed Random Variables. Ark. Mat. vol. 4, no. 24, 1961, pp. 323-332.



12. Sirashdinov, S.H.: Über Grenzwertsätze für unabhängige zufällige Summanden. Wiss. Zeit. Tech. Univ. Dresden, 15(5), 1966, pp. 915-921.
13. Zolotarev, V.M. A Sharpening of the Inequality of Berry-Esseen. Z. Wahrscheinlichkeitstheorie verw. Geb. 8, 1967, pp. 332-342.

THE UNIVERSITY OF CHICAGO  
DEPARTMENT OF CHEMISTRY  
5408 S. UNIVERSITY AVENUE  
CHICAGO, ILL. 60637

RECEIVED

NOV 15 1964

1964

NOV 15 1964

NOV 15 1964



## CURRICULUM VITAE

Christopher E. Olson was born in San Antonio, Texas on December 4, 1941. He attended public schools in San Antonio and graduated from Thomas Jefferson High School in 1958. He attended Washington University (St. Louis) for two years and received his B.A. magna cum laude from St. Mary's University (Texas) in 1963. Working as a teaching assistant, he received his M.A. in 1965 from the University of Kansas. He received his Ph.D. in 1971 from the University of New Mexico.





Embossed text, likely the brand name "Snooble Bond".

Embossed text, likely "25% COTTON FIBER".



Central Cotton Fibre

233 COTTON FIBRE

100% COTTON FIBRE

100% COTTON FIBRE

100% COTTON FIBRE

100% COTTON FIBRE

100% COTTON FIBRE

