

University of New Mexico

UNM Digital Repository

Project ECHO Bibliography

Project ECHO

12-1-2019

Impact of Project ECHO Models of Medical Tele-Education: a Systematic Review

Ryan K. McBain

Jessica L. Sousa

Adam J. Rose

Sangita M. Baxi

Laura J. Faherty

See next page for additional authors

Follow this and additional works at: https://digitalrepository.unm.edu/hsc_echo_bibliography

Authors

Ryan K. McBain, Jessica L. Sousa, Adam J. Rose, Sangita M. Baxi, Laura J. Faherty, Caroline Taplin, Andre Chappel, and Shira H. Fischer

REVIEWS

Impact of Project ECHO Models of Medical Tele-Education: a Systematic Review



Ryan K. McBain, PhD, MPH¹, Jessica L. Sousa, MPH, MSW¹, Adam J. Rose, MD, MS^{1,2}, Sangita M. Baxi, MAS³, Laura J. Faherty, MD, MPH, MS¹, Caroline Taplin, MS⁴, Andre Chappel, PhD⁴, and Shira H. Fischer, MD, PhD¹

¹RAND Corporation, Boston, MA, USA; ²Section of General Internal Medicine, Boston University School of Medicine, Boston, MA, USA; ³RAND Corporation, Santa Monica, CA, USA; ⁴Office of the Assistant Secretary for Planning and Evaluation, Washington, DC, USA.

BACKGROUND: Extension for Community Health care Outcomes (ECHO) and related models of medical tele-education are rapidly expanding; however, their effectiveness remains unclear. This systematic review examines the effectiveness of ECHO and ECHO-like medical tele-education models of healthcare delivery in terms of improved provider- and patient-related outcomes.

METHODS: We searched English-language studies in PubMed, Embase, and PsycINFO databases from 1 January 2007 to 1 December 2018 as well as bibliography review. Two reviewers independently screened citations for peer-reviewed publications reporting provider- and/or patient-related outcomes of technology-enabled collaborative learning models that satisfied six criteria of the ECHO framework. Reviewers then independently abstracted data, assessed study quality, and rated strength of evidence (SOE) based on Cochrane GRADE criteria.

RESULTS: Data from 52 peer-reviewed articles were included. Forty-three reported provider-related outcomes; 15 reported patient-related outcomes. Studies on provider-related outcomes suggested favorable results across three domains: satisfaction, increased knowledge, and increased clinical confidence. However, SOE was low, relying primarily on self-reports and surveys with low response rates. One randomized trial has been conducted. For patient-related outcomes, 11 of 15 studies incorporated a comparison group; none involved randomization. Four studies reported care outcomes, while 11 reported changes in care processes. Evidence suggested effectiveness at improving outcomes for patients with hepatitis C, chronic pain, dementia, and type 2 diabetes. Evidence is generally low-quality, retrospective, non-experimental, and subject to social desirability bias and low survey response rates.

DISCUSSION: The number of studies examining ECHO and ECHO-like models of medical tele-education has been modest compared with the scope and scale of implementation throughout the USA and internationally. Given the potential of ECHO to broaden access to healthcare in

rural, remote, and underserved communities, more studies are needed to evaluate effectiveness. This need for evidence follows similar patterns to other service delivery models in the literature.

J Gen Intern Med 34(12):2842–57

DOI: 10.1007/s11606-019-05291-1

© Society of General Internal Medicine 2019

INTRODUCTION

Technological innovations over the past decade have steadily reduced barriers to accessing healthcare¹ both in the USA² and internationally.³ Telemedicine holds the potential for patients to seek medical expertise more efficiently, reducing wait times and allowing specialists to direct their attention to individuals with the greatest health needs, regardless of geographic location.⁴ This is particularly true for the expertise of specialists, which may be unevenly distributed.⁵

Several mechanisms have arisen to enable increased access to specialist care, including e-consultations that allow specialists to consult remotely.⁶ However, with a growing physician and nursing shortage in settings ranging from the USA⁷ to large parts of sub-Saharan Africa,⁸ there is a fundamental need to equip front-line providers in rural areas with the specialized skills necessary to address community needs themselves. Such capacity-building is particularly relevant in the context of escalating health epidemics, such as the US opioid crisis or recent Ebola epidemic in West Africa, and for tackling increasingly common conditions like hepatitis C.

In 2016, the US Congress passed the ECHO (Expanding Capacity for Health Outcomes) Act, which aims to support and promote “technology-enabled collaborative learning and capacity building models.”⁹ Several such models have been developed, the most ubiquitous being Project ECHO (Extension for Community Health care Outcomes).¹⁰ Project ECHO involves pairing front-line clinicians, typically located in underserved areas, with specialist mentors at academic medical centers, or “hubs”, using videoconference and a case-based mode of pedagogy. Launched in 2003 out of the University of New Mexico to increase access to hepatitis C treatment in parts of the rural southwest, the program now operates at more

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11606-019-05291-1>) contains supplementary material, which is available to authorized users.

Received March 4, 2019

Revised May 31, 2019

Accepted July 31, 2019

Published online September 4, 2019

than over 100 academic and medical hubs across 48 states as well as multiple continents, and covers dozens of disease states and health conditions.¹¹

While Project ECHO has been successful at expanding its scope and scale, there remains a paucity of evidence regarding the impact of ECHO and ECHO-like models (EELM) on provider- and patient-related outcomes. An investigation of the evidence is particularly warranted, given the extent of human and financial capital invested in this model: thousands of trainers and trainees, and millions of dollars in financial support. While an earlier review examined the impact of ECHO through the middle of 2015,¹² this was prior to the ECHO Act and any experimental evidence, and did not extend beyond ECHO-affiliated programs. We present a systematic review of EELM that comprises peer-reviewed evidence of patient and provider outcomes between 2007 and 2018. We follow Cochrane Collaboration's GRADE framework¹³ to examine the strength of evidence (SOE) and use this review as a basis for highlighting potential next steps and future directions.

METHODS

Data Sources and Searches

We reviewed academic literature in accordance with PRISMA guidelines,¹⁴ targeting publications that evaluated EELM (2007–2018). In consultation with public health experts in the field of telehealth, we established an operational definition for EELM as “a technology-enabled educational model, in which a mentor with specialized knowledge provides interactive and case-based guidance to a group of mentees for the purpose of strengthening their skills and knowledge to provide high-quality healthcare.” We delimited our search according to six inclusion criteria: (1) using a technology-enabling platform, (2) having a health-focused objective, (3) leveraging specialists to train generalists, (4) using interactive mentorship, (5) using case-based learning, and (6) implementing a hub-spoke framework rather than 1:1 learning.

We implemented a Boolean search procedure based on key words defined under three domains: (i) a technology-enabling component, (ii) involvement of health providers, and/or (iii) terms denoting resource or geographic barriers, which EELM often address. As a complementary strategy, we searched for ECHO-specific terminology linked by “or” statements. A detailed list of search terms can be found in Appendix Table 1 (online). A total of six databases were searched: PubMed, Embase, PsycINFO, Google Scholar, the Cochrane Central Register (CENTRAL), and Scopus. Google Scholar was limited to the first 100 returns. Using seminal articles, including a 2016 review by Zhou and colleagues,¹² we also examined bibliographies.

Study Selection

We limited results to peer-reviewed articles reporting provider- or patient-related outcomes published in English between January 1, 2007 and December 1, 2018, including articles originating outside the USA. Returns were screened independently by two research team members for agreement with the six inclusion criteria. For situations in which agreement with criteria was unclear from the title and abstract, the full text was reviewed. Records that met inclusion criteria were flagged for full data abstraction (see Fig. 1). In the event a discrepancy arose, additional members of the research team were consulted.

Data Extraction and Quality Assessment

Articles meeting inclusion criteria were independently entered by an investigator into a data abstraction form. A second investigator was then tasked with reviewing abstracted data to ensure accuracy and completeness. Summaries of abstracted data can be found in Tables 1 and 2.

Data Synthesis and Analysis

We selected key features from each study to review and summarize, based on a hierarchy of evidence, with highest quality evidence the focus of synthesis. Provider-related outcomes included participation, satisfaction, knowledge, self-efficacy, and behavior change, following a core competencies model for implementation science⁶⁶ articulated by the National Implementation Research Network.⁶⁷

Patient-related outcomes were grouped according to health condition and involved careful examination of health condition-specific outcomes as stated in the literature. These were classified as either process or outcome measures, according to US Agency for Healthcare Research and Quality⁶⁸ definitions. Whenever possible, we report summary statistics—including means, standard deviations, odds ratios (ORs), and hazards ratios (HRs). We also report *p* values and 95% confidence intervals.

We rated SOE according to Cochrane GRADE criteria,¹³ following a two-step process. First, two research team members independently assigned a score to each article for the outcomes presented within, based on six GRADE characteristics: study design, risk of bias, inconsistency, indirectness, imprecision, and publication bias. The full research team collectively reviewed each score. Second, the research team deliberated weight of evidence across individual studies for each patient- and provider-related outcome. This systematic classification process is outlined in detail on the Cochrane website and involves, for example, evaluating the quantity of experimental versus observational evidence, and studying effect sizes and dose responses. SOE is assigned an ordinal score: very low (+), low (++) , medium (+++) , high (++++) (Table 3).




Figure 1 Study flow diagram.

Role of the Funder

This investigation was supported by the Office of the Assistant Secretary for Planning and Evaluation (ASPE), within the US Department of Health and Human Services (HHS).⁶⁹ In accordance with the ECHO Act of 2016, this investigation was commissioned as part of a Report to US Congress, released in 2019. Two staff members from ASPE are co-authors; the staff contributed to the selection of terms for the literature search and the criteria for study eligibility and provided edits of the final manuscript.

RESULTS

After implementation of search procedures, we reviewed 2970 records: 2965 from database searches, and an additional five from bibliographic reviews. There was an acceptable degree of inter-rater reliability: raters agreed 97.3% of the time, reflecting kappa coefficient of $\kappa = 0.46$. Following screening,

211 articles were identified for full-text review for eligibility. Of these, 52 met eligibility (see Fig. 1). Forty-three contained provider outcomes, 15 contained patient outcomes, and six contained both provider and patient outcomes.

The most common health topics addressed by EELM were hepatitis C, chronic pain management, and dementia and elderly care. Thirty-nine of 52 articles focused on EELM implemented in the USA, with Canada and Australia as the next most common countries. Year by year, there has been an overall increase in the number of published articles evaluating EELM (see Fig. 2).

The format of sessions ranged from weekly to monthly, lasting 60 to 180 min per session, with wide variation in number of sessions conducted—in part because of the continuous nature of intervention protocols. Similarly, there was variability in the number of trainees and number of patients served by trainees. In some instances, these numbers were not reported. Below, we present a topical synthesis of the articles, organized by provider-related outcomes and patient-related

Table 1 Included Studies Reporting Provider-Related Outcomes

Citation	Health content area	Number of trainees evaluated	Evaluation design	Main provider outcome measures	Main provider outcomes reported
Anderson et al., 2017 ¹⁵	Pain management	12 in the intervention group; 11 in the control group	Pre-post study design with comparison group	Pain-related knowledge and self-reported self-efficacy; frequency of formal assessment tool utilization; frequency of opioid agreements developed; patient concern about addiction to opioids	Increased pain knowledge in the intervention group ($p < 0.001$), not observed in the control group ($p = 0.11$); non-significant group difference in frequency of opioid agreement usage ($p = 0.05$); lower concern about patient addiction to opioids in the intervention group ($p = 0.006$).
Arora et al., 2011 ¹⁶	Hepatitis C	Varied by year: 17–52 providers	Pre-post study design without comparison group	Self-reported satisfaction with ECHO training; self-reported self-efficacy before versus after ECHO training; self-reported perceived benefits of ECHO training	Satisfaction with ECHO training ranged from 4.3 to 4.9 on 1–5 ordinal scale (2006); self-efficacy increased significantly across all categories ($p < 0.001$) (2006–2007); moderate-major benefits self-reported across eight categories 82–98% of time (2008).
Ball et al., 2018 ¹⁷	Pain management	25 (surveys); 14 (focus group discussions)	Pre-post study design without comparison group; focus group discussions	Self-reported confidence and knowledge treating patients with chronic pain before versus after ECHO training; barriers and facilitators to participation in ECHO	Increased provider confidence ($p < 0.01$) and increased provider knowledge ($p < 0.05$) on chronic pain; focus group discussions indicated increased provider self-efficacy and knowledge, as well as increased workload associated with participation.
Beste et al., 2017 ¹⁸	Hepatitis C	376	Retrospective cohort study with comparison group	Rate of PCPs who initiate hepatitis C treatment with antiviral treatment	Providers who received at least one SCAN-ECHO training were more likely to initiate antiviral treatment ($p < 0.01$), compared with those with no SCAN-ECHO training. This was attributable to more frequent initiation among those presented as cases during trainings.
Beste et al., 2016 ¹⁹	Infectious diseases; hepatitis C; pulmonology; nephrology	78	Participant survey	Self-reported benefits of ECHO participation, such as perceived impact on providers and perceived impact on care delivery; association between duration of participation and perceived benefits	Strong agreement with trainings' impact on providers ranged from 34.2 to 46.8% across questions; strong agreement with trainings' impact on care delivery ranged from 28.6 to 38.4% across questions; participation for more than 1 year was associated with greater perceived impact, particularly perceived patient access to specialty care ($p = 0.005$).
Carlin et al., 2018 ²⁰	Chronic pain management	37	Focus group discussions (6)	Qualitative feedback on barriers and facilitators to ECHO, as well as perceived benefits and drawbacks	Respondents reported insights defined under such themes as challenges of managing chronic pain; ECHO participation and improvement in patient-provider interaction and participant knowledge; ECHO participation generating a sense of community; and disadvantages associated with participating in ECHO.
Catic et al., 2014 ²¹	Dementia	Unknown	Prospective cohort study without comparison group	Self-reported adherence to recommendations of the ECHO-AGE expert team	Self-reported adherence to expert recommendations in 39 of 44 cases (89%) presented.
Chaple et al., 2018 ²²	Substance use disorder	20	Participant survey	Participant satisfaction in quality of training; self-reported enhancement in clinical skills	General participation satisfaction was 4.69 of 5; self-reported enhancement of clinical skills as a result of training was 4.45 of 5.
Cofa-Woerpel et al., 2018 ²³	Tobacco cessation	23	Participant survey	Self-reported confidence treating tobacco use; satisfaction with participation; tobacco-related knowledge survey	All respondents (22) reported moderate-to-high confidence to address tobacco use; a majority of knowledge questions yielded 69–85% correct answers; 77% agreed the program was satisfactory.

(continued on next page)

Table 1. (continued)

Citation	Health content area	Number of trainees evaluated	Evaluation design	Main provider outcome measures	Main provider outcomes reported
Cordasco et al., 2015 ²⁴	Women's health	Varied by type of survey, 18–53	Participant surveys; participant semi-structured interviews	Self-reported impact of training on care; self-reported satisfaction with participation	47 of 53 survey respondents (89%) reported that SCAN-ECHO information would influence their patient care; 18 of 18 interviewees (100%) reported SCAN-ECHO was useful for building and maintaining knowledge.
Covell et al., 2015 ²⁵	Co-occurring mental and substance use disorder	8 (provider-level); 11 (program-level)	Participant survey (provider level); prospective cohort study without comparison (program level)	Provider-level: self-reported satisfaction. Program-level: increased knowledge about integrated treatment; percent of charts with stage of treatment recorded.	All providers reported that the online learning collaborative was helpful, the implementation model was helpful, and strategies supporting implementation were helpful. At program-level, sites showed significant increase in dual disorder treatment knowledge survey ($p < 0.05$); sites showed increase in chart documentation ($p < 0.05$).
Eaton et al., 2018 ²⁶	Chronic pain management	41	Cluster randomized controlled trial (clinic participation in Tele-Pain sessions)	Pain management knowledge measured by KnowPain-12; self-reported knowledge and attitudes regarding pain; self-reported perceived competence	No significant change in knowledge scores or self-perceived competence when compared between intervention and control group PCPs ($p > 0.05$).
Farris et al., 2017 ²⁷	Dementia	12	Participant survey	Self-reported benefits of ECHO participation, including on patient treatment plans	Satisfaction on features of ECHO ranged from 3.25 to 3.58 on scale of 1 (strongly disagree) to 5 (strongly agree); providers demonstrated an average score of 3.64 on agreement that they incorporated training advice into treatment plans.
Fisher et al., 2017 ²⁸	Dementia	154 (cohort); 26 (qualitative interviews)	Semi-structured interviews; retrospective cohort study with comparison group	Semi-structured interviews explored participant perceptions and experiences in the program	Interviewees reported the program led to improvements in clinician geriatric mental healthcare knowledge and treatment practices.
Glass et al., 2017 ²⁹	Chronic liver disease	106	Retrospective cohort study with control comparison	Association between complexity of trainee cases presented and number of cases presented	Providers who presented more than ten SCAN-ECHO cases were more likely to present complex cases about a specific treatment or a procedure, compared with those presenting ten or fewer cases ($p < 0.001$).
Haozous et al., 2012 ³⁰	Cancer-related pain management	24 (education sessions); 32 (case conference sessions)	Retrospective cohort study with and without control comparison	Self-reported satisfaction survey on pain management educational sessions; self-reported perceived competence proceeding case conference calls	Providers who attended pain management sessions reported mean item-level satisfaction scores ranging from 2.75 to 3.47 on a 0–4 ordinal scale; providers who attended case conference calls reported significantly higher competence on pain management than a control comparison group ($p < 0.01$).
Jansen et al., 2018 ³¹	Pain management in end-stage dementia	18	Mixed-methods prospective cohort study	Participant self-efficacy and knowledge, based on KnowPain-50 and KnowPain-12 questionnaires; two focus group discussion interviews	Overall knowledge and self-efficacy scores were significantly higher post-ECHO than pre-ECHO for physicians ($p = 0.01$) and nurses ($p = 0.04$). Key themes that emerged were knowledge and skills development and dissemination, protected time, areas for improvement, and the future of ECHO.
Johnson et al., 2017 ³²	Multiple sclerosis	15 trainees participated in evaluation; 24 trainees total	Participant surveys	Self-reported confidence treating multiple sclerosis; self-reported satisfaction with program; self-reported feedback on program format	Mean self-reported confidence treating multiple sclerosis after training was 4.53 out of 5; 9 of 15 participants indicated the program met their expectations; 15 of 15 participants indicated that sessions expressed good value.

(continued on next page)

Table 1. (continued)

Citation	Health content area	Number of trainees evaluated	Evaluation design	Main provider outcome measures	Main provider outcomes reported
Katzman et al., 2014 ³³	Chronic pain	763 (surveys); 9 (focus group discussants)	Participant survey; focus group discussions	Percentage of providers who reported that trainings were “excellent” on five dimensions; exploratory feedback on utility of presentations and impact of participation	From 2010 to 2012, the percentage of providers reporting “excellent” increased significantly across categories ($p < 0.01$); provider feedback on utility of ECHO trainings and impact of participation were generally positive.
Kauth et al., 2015 ³⁴	Transgender health	13	Participant survey; pre-post study design with comparison group	Post-intervention self-reported satisfaction on training; pre- and post-intervention self-reported confidence providing care	92.3% of providers described the didactics as somewhat or very helpful. The majority (76.9%) reported that receiving consultation was somewhat or very helpful, and nearly everyone (92.3%) felt that they benefited from listening to other cases being discussed; 39.7% of providers increased in self-reported confidence to treat transgender veterans after SCAN-ECHO ($p = 0.007$).
Komaromy, Bartlett, et al., 2017 ³⁵	Integrated addictions and psychiatry	41	Participant survey	Percentage of participants who reported changing their patient care plan as a result of presenting a case; percentage who rated the value of expert input received as 5 on a scale of 1–5; percentage who reported training as useful in caring for their own patients	77% of case presenters reported that the case discussion changed their patient care plan; 86% reported the value of the input they received as a 5 out of 5; 93% reported training as useful in caring for their own patients.
Komaromy, Ceballos, et al., 2018 ³⁶	Community health worker training: obesity prevention and addiction recovery	16 (obesity prevention); 46 (addiction recovery)	Pre-post study design without comparison group; trainer-rated pre-post survey	Self-reported change in obesity prevention knowledge and abilities; trainer-reported change in motivational interviewing skills for addiction recovery	Self-reported obesity prevention knowledge and abilities increased on 12 of 13 dimensions ($p < 0.05$); trainer-reported provider performance on motivational interviewing improved ($p < 0.001$).
Lewiecki et al., 2017 ³⁷	Osteoporosis	16	Pre-post study design without comparison group	Pre-post intervention change in self-reported self-efficacy, based on self-efficacy questionnaire	Overall increase in reported self-efficacy among participants who completed the survey ($p = 0.005$). It uses a pre-post framework.
Marcjano et al., 2017 ³⁸	Hepatitis C	14	Pre-post study design without comparison group	Self-assessed provider knowledge on HCV	Increase in self-assessed knowledge on all ten aspects of HCV care from pre- to post-intervention ($p < 0.05$).
Masi et al., 2012 ³⁹	Hypertension	9 in the intervention group; 3 in the control group	Pre-post study design with comparison group	Knowledge surveys administered at baseline and endline; self-reported knowledge reported at baseline and endline	Tested knowledge of how to treat hypertension increased among intervention providers ($p < 0.01$) but not among controls. Self-assessed knowledge increased among intervention providers ($p < 0.01$) but not among controls.
Mazurek et al., 2017 ⁴⁰	Autism spectrum disorders	14	Pre-post study design without comparison group	Self-reported self-efficacy; self-reported use of M-CHAT or another screening tool; self-reported adherence to American Academy of Pediatrics autism spectrum disorder screening guidelines; self-reported use of 15 possible resources for autism; satisfaction with program	Self-efficacy improved significantly ($p = 0.002$); use of resources increased from 0.29 out of 15 to 4.07 out of 15, on average ($p = 0.003$); high satisfaction with ECHO trainings was reported.
Mehrotra et al., 2018 ⁴¹	Mental health	12	Pre-post study design without comparison group; participant survey	Self-reported responses to satisfaction survey; pre- and post-intervention knowledge test and pre- and post-intervention self-reported self-efficacy	Mean participant satisfaction was 4.5 or higher on a scale of 1–5 for five survey questions. Topical knowledge increased significantly ($p < 0.01$), as did self-reported self-efficacy ($p < 0.05$).

58

(continued on next page)

Table 1. (continued)

Citation	Health content area	Number of trainees evaluated	Evaluation design	Main provider outcome measures	Main provider outcomes reported
Meins et al., 2015 ⁴²	Chronic pain management		Participant survey; participant observation	Self-reported belief that participation enhanced knowledge of pain management; self-report that participant intends to use new knowledge gained	On scale of 1–4, mean score for statement that participation enhanced knowledge was 3.94; mean score for statement that participant intended to use new knowledge gained was 3.77.
Ni Cheallaigh et al., 2017 ⁴³	Hepatitis C	6	Participant semi-structured interviews	Self-reported care management skills following ECHO training	Respondents generally reported that ECHO participation increased their ability to manage HCV infection.
Oliveira, Branquinho, and Goncalves, 2012 ⁴⁴	Varied: e.g. dermatology, neurology, and gastroenterology	848	Participant survey	Overall participant satisfaction	Overall satisfaction was reported as medium, high, or very high (range: very low, low, medium, high, very high) by 90% of respondents in 2009 and 94% in 2010.
Parsons et al., 2017 ⁴⁵	Sleep medicine	39	Participant surveys	Self-reported comfort treating sleep disorders; self-reported clinical practice change	Increased provider comfort reported by 77% of respondents; a majority (85%) of respondents reported “some” or significant” practice change across practice domains.
Qaddoumi et al., 2007 ⁴⁶	Pediatric neuro-oncology	Unknown	Prospective cohort study without comparison group	Percentage of patients for whom expert recommendations differed from original care plan; percentage of patients for whom there was a significant change in the original care plan, conditional on recommendations	In 23 patients (36%), major changes from original plan were recommended on different aspects of the care; in 21 patients (91%), those recommendations were followed.
Rahman et al., 2012 ⁴⁷	Geriatric nutrition	Unknown	Participant survey; prospective cohort study without comparison group	Post-intervention participant satisfaction; pre-post intervention change in knowledge	89% of participants reported that they would participate in a similar project and recommend the course; knowledge scores on trainer-administered quiz improved significantly ($p < 0.05$).
Ray, Fried, and Lindsay, 2014 ⁴⁸	Palliative care	101	Pre-post study design without comparison group	Increased confidence to provide palliative care pre-versus post-intervention; post-intervention rating of content usefulness	Provider confidence increased significantly ($p < 0.05$); average rating of content usefulness was 3.50 on scale of 1–4.
Salgia et al., 2014 ⁴⁹	Hepatitis C	24	Participant survey	Self-reported change of care provision following the intervention	Most participants ($n = 20$; 83%) reported having encountered a case similar to the one presented in SCAN-ECHO; of these participants, 18 (90%) reported improvements in their perceived diagnostic approach, 16 (80%) reported having developed a better treatment plan, and 16 (80%) reported perceived improvements in follow-up plan development.
Shipherd et al., 2016 ⁵⁰	Transgender care	111	Participant surveys	Post-session knowledge test score; post-session self-reported satisfaction survey scores; post-intervention feedback survey; self-perceived confidence treating transgender veterans before versus after participation	Session participation ranged from 11 to 57, with 93% receiving a post-session knowledge survey score greater than 80%; average session satisfaction was 4.28 on 0–5 scale; participants rated all aspects of the intervention to be useful; 92% of participants increased in treatment confidence (p value unreported); 63% of participants expected to care for more transgender patients in the future.
Sockalingam et al., 2018 ⁵¹	Mental health	Varied by type of survey, 22–27	Pre-post study design without comparison group	Self-reported knowledge and self-efficacy	Increased mental health and additions knowledge ($p < 0.001$); increased provider self-efficacy approaching statistical significance ($p = 0.06$).

(continued on next page)

Table 1. (continued)

Citation	Health content area	Number of trainees evaluated	Evaluation design	Main provider outcome measures	Main provider outcomes reported
Swigert et al., 2014 ⁵²	Diabetes	Unknown	Pre-post study design without comparison group	Self-reported knowledge and confidence levels (including retrospective report of baseline knowledge and confidence); self-reported intention to change current clinical care practices	Self-reported increase in diabetes knowledge ($p < 0.001$) and increased confidence ($p < 0.001$) after individual ECHO sessions; a majority of participants (95%) reported an intention to change clinical practice after ECHO sessions.
Van Ast and Larson, 2007 ⁵³	Disability care	8	Semi-structured interviews	Perceived acceptability of technology; perceived benefits of participation	Participants generally reported favorable feedback about the technology platform; participants reported positive behavioral changes in caregiving.
Volpe, Boydell, and Pignatiello, 2014 ⁵⁴	Psychiatric services	Unknown	Focus group discussion; key informant interviews	Overall participant satisfaction; acceptability of tele-video technology	Focus group discussants and interviewees reported overall satisfaction; tele-video technology was regarded as an effective tool for learning.
White et al., 2015 ⁵⁵	Palliative care	28	Mixed-methods prospective cohort study	Provider knowledge score; self-reported self-efficacy scores, provider self-reported satisfaction with program	Mean knowledge score improved significantly (71.3% to 82.7%, $p < 0.001$); self-efficacy significantly improved ($p = 0.063$); 96% reported gains in learning; 90% felt ECHO had improved the care they provide; 83% would recommend ECHO to other healthcare providers; 70% said ECHO's technology gave them access to education they would have had difficulty accessing.
Wood et al., 2018 ⁵⁶	HIV/AIDS (PrEP)	45	Participant survey	Self-reported knowledge of PrEP, comfort level discussing PrEP, and prescribing practices	93.3% of survey respondents reported that the intervention helped them stay up to date on PrEP guidelines "extremely" or "moderately" well; 91.1% reported an "extremely" or "moderately" increased likelihood to prescribe PrEP; 40.0% reported that, without the intervention, they would have referred patients seeking PrEP to another provider.
Wood et al., 2016 ⁵⁷	HIV/AIDS	45	Prospective cohort study without comparison group	Self-assessed confidence to perform essential components of HIV care; self-reported feeling part of a community of practice; self-reported overall HIV care knowledge	Self-assessed confidence improved over time in several clinical skill areas on 14 of 18 dimensions of care provision ($p < 0.05$); feelings of professional isolation decreased while degree to which participants felt part of an HIV community of practice increased ($p < 0.05$); self-reported HIV care knowledge increased ($p = 0.004$).

outcomes; a full description of outcomes reported is found in the Online Appendix.

Provider-Related Outcomes

Between 2007 and 2018, 43 of 52 articles presented quantitative or qualitative evidence outlining provider-related outcomes of EELM. Studies most frequently measured outcomes in one of four areas: (i) provider satisfaction with quality and content of trainings ($n = 17$; 40%); (ii) provider knowledge acquired ($n = 18$; 42%); (iii) enhanced provider confidence or

self-efficacy associated with care delivery ($n = 18$; 42%); and (iv) changes in self-reported provider behaviors associated with patient care ($n = 7$; 16%). In terms of study design, 23 of 43 (53%) involved a counterfactual—either within- (pre vs. post) or between-subjects. While only one of the studies included an element of randomization, three studies involved both within- and between-subject comparisons.

Provider Satisfaction. Assessment of provider satisfaction largely entailed administration of post-intervention structured

Table 2 Included Studies Reporting Patient-Related Outcomes

Citation	Health content area	Patients in evaluation	Evaluation design	Main patient outcome measures	Main patient outcomes reported
Anderson et al., 2017 ¹⁵	Pain management	Exposure group: 1586 at baseline; 1485 at follow-up Control group: 2020 at baseline; 1695 at follow-up	Pre-post study design with comparison group	Percentage of patients with chronic pain treated with an opioid medication; average number of opioid prescriptions written per patient with pain; frequency of referrals for behavioral health and physical therapy	Greater reduction in the intervention group for percentage of patients with chronic pain treated with an opioid medication ($p = 0.002$); smaller increase in the intervention group for number of opioid prescriptions written per patient with pain ($p = 0.001$); frequency of referrals to behavioral health and physical therapy ($p < 0.001$)
Arora, Thornton, et al., 2011 ¹⁶	HCV	261 patients in exposure group 146 patients in control group	Prospective cohort study with comparison group	Percentage of patients with sustained viral response; percentage of patients among whom a serious adverse event occurred	No difference in percentage of patients with sustained viral response ($p = 0.89$); greater prevalence of serious adverse events reported in the control group ($p = 0.02$)
Beste et al., 2017 ¹⁸	HCV	6431 patients in exposure group 32,322 patients in control group	Retrospective cohort study with comparison group	Rate of patients with sustained virologic response	No significant difference in rates of sustained virologic response between providers with versus without SCAN-ECHO training ($p = 0.32$)
Carey et al., 2016 ⁵⁸	Pain management	371,646	Spatial reach analysis	Association between distance to specialty pain care and being seen in person at a specialty clinic; association between distance to specialty pain care and access to a Pain SCAN-ECHO participating PCP	Patient distance from home to specialty pain care associated with 22% lower odds of being seen in person at a specialty care clinic ($p < 0.001$); distance from home to specialty pain care associated with 2% lower odds of access to a Pain SCAN-ECHO participating PCP ($p = 0.01$)
Catic et al., 2014 ²¹	Dementia	47	Prospective cohort study without comparison group	Association between provider self-reported adherence to expert recommendations and provider self-reported (1) clinical improvement and (2) hospitalization of their patients	Clinical improvement among patients was self-reported as greater among those who adhered to expert recommendations ($p < 0.05$); hospitalization among patients was self-reported as lower among those who adhered to expert recommendations (p value unreported)
Fisher et al., 2017 ²⁸	Dementia	More than 70,000	Semi-structured interviews; retrospective cohort study with comparison group	Patient healthcare utilization and costs at participant practices, before and after enrollment in study	Reduction in emergency department costs (\$406 to \$311; $p < 0.05$) among those with mental disorder; increase in outpatient care utilization and costs among those without a mental disorder ($p < 0.05$)
Frank et al., 2015 ⁵⁹	Chronic pain	22,454 patients in exposure group 299,981 in non-exposure group	Prospective cohort study with comparison group	Association between case presentations and (1) delivery of outpatient care (physical medicine, mental health, SUD, and pain medicine) and (2) medication initiation (antidepressants, anticonvulsants, and opioid analgesics)	Patients whose case was presented during training sessions had greater likelihood of utilizing physical therapy ($p < 0.05$), but not care for mental health, SUD, or specialty pain medicine ($p > 0.05$), compared with patients whose cases were not discussed. Patients with presented cases also had greater likelihood of initiation on antidepressants and anticonvulsants ($p < 0.05$), but not an opioid analgesic ($p > 0.05$)
Glass et al., 2017 ²⁹	Chronic liver disease	582 in exposure group 1395 in comparison group	Retrospective cohort study with control comparison	Patient time to liver consultation; patient distance traveled to care	SCAN-ECHO liver consults were completed an average of 9.6 days sooner than in the liver clinic (p value unreported); average patient distance traveled to the liver clinic was 250 miles round-trip (p value unreported)

(continued on next page)

Table 2. (continued)

Citation	Health content area	Patients in evaluation	Evaluation design	Main patient outcome measures	Main patient outcomes reported
Gordon et al., 2016 ⁶⁰	Dementia	Unknown	2:1 matched cohort study	Percentage of patients receiving antipsychotic medications; percentage of patients physically restrained; nine other secondary outcomes	Patients at participant facilities were marginally less likely to be physically restrained than patients at nonparticipant facilities ($p=0.05$), and less likely to be prescribed antipsychotic medication ($p=0.07$). Patients at participant facilities were less likely to experience a urinary tract infection
Katzman et al., 2018 ⁶¹	Chronic pain management	Unknown	Prospective cohort study with comparison group	Prescription rates of opioid analgesics and co-prescribing of opioids and benzodiazepines	Clinics participating in the intervention (ECHO Pain) showed greater declines in opioid prescriptions than did comparison facilities (-23% versus -9% , $p < 0.001$); days of co-prescribed opioids and benzodiazepines also declined more ($p < 0.001$)
Mohsen et al., 2018 ⁶²	Hepatitis C	100 in exposure group 100 in comparison group	Retrospective cohort study with a comparison group	Percentage of patients with direct-acting antiviral therapy initiated; percentage of patients who complete their regimen; percentage of patients with sustained virologic response	Treatment was initiated among 78% of intervention patients versus 81% of those in a TLC; 89% of intervention participants completed treatment—of those, 87% had sustained virologic response compared with 86% and 96%, respectively, in the TLC group. Statistical significance not reported
Moore et al., 2017 ⁶³	Geriatric care	Exposure group: 213 at baseline, 148 at endline Comparison group: 220 at baseline, 214 at endline	Prospective cohort study with comparison group	30-day readmission rates; 30-day total cost of care; average length of stay at the skilled nursing facility; 30-day mortality rate	Readmission was lower in the intervention group ($p=0.04$); adjusted 30-day cost was lower in intervention group ($p < 0.001$); average length of stay at skilled nursing facility was shorter in intervention group ($p < 0.001$); 30-day mortality rate was not significantly different between groups ($p=0.11$)
Ní Cheallaigh et al., 2017 ⁴³	Hepatitis C	Unknown	Participant semi-structured interviews	Provider-reported benefits to patients	Respondents reported that patients attending their practice benefited from ECHO training
Su et al., 2018 ⁶⁴	Chronic liver disease	513 in VA SCAN-ECHO group; 62,237 in comparison group	Retrospective cohort study with comparison group	All-cause mortality among patients who received a SCAN-ECHO visit, propensity score matched to patients who received no visits	Propensity-adjusted mortality rates showed that a SCAN-ECHO visit was associated with a hazard ratio of 0.54 ($p=0.003$) compared with no visit
Watts et al., 2016 ⁶⁵	Diabetes	39	Pre-post study design without comparison group	Mean glycated HbA1c value (glycemic control) at intervention sites before and after intervention; comparative levels of HbA1c $> 9.0\%$ at intervention and comparison sites at baseline and endline	Mean HbA1c improved from $10.2 \pm 1.4\%$ to $8.4 \pm 1.8\%$ ($p < 0.001$) over the average follow-up period of five months, not explained by system-wide changes or improvements; comparative increase in HbA1c scores at comparison sites ($p < 0.05$)

surveys.^{19, 23–25, 27, 33, 34, 41, 44, 50, 70} The median response rate was low (under 50%); however, self-reports consistently indexed positive ratings, at both the item-level and survey-level. In several instances, satisfaction was framed in terms of participation benefits, such as “Because of [EELM], I have expanded my practice to include new skills.¹⁹” In addition to structured surveys, several authors conducted focus group discussions^{17, 20, 33, 54} and semi-structured interviews^{24, 28, 43, 53} to solicit feedback on aspects of EELM that worked well

or less well, often with a focus on acceptability of the technology platform utilized. Here, responses were also generally positive.

Provider Knowledge. In one study,⁴² authors evaluated provider knowledge by merely asking participants after training to self-report whether they perceived their knowledge to improve. More often, studies implemented a pre-post design, in which providers were asked to self-assess their

Table 3 Summary of SOE Scores for Patient- and Provider-Related Outcomes

Category	GRADE score	Quality of evidence
Provider measures		
Provider satisfaction (n = 15)	++	Low
Provider knowledge (n = 18)	++	Low
Provider confidence (n = 18)	++	Low
Provider behavior change (n = 7)	+	Very low
Patient measures		
Hepatitis C (n = 4)	++	Low
Chronic liver disease (n = 2)	++	Low
Chronic pain management/opioid use (n = 4)	++	Low
Gerontology, including mental health and dementia (n = 4)	+	Very low
Diabetes management (n = 1)	+	Very low

knowledge at baseline and again at endline, with significant changes observed.^{17, 36, 38, 39, 51, 52, 56, 71} In a subset of assessments, knowledge surveys were constructed by the authors and administered.^{15, 23, 26, 39, 41, 47} The authors found significant improvements in objectively measured content knowledge. By contrast, a cluster randomized controlled trial on chronic pain education did not show knowledge gains among ECHO participant clinics compared with non-participant clinics.²⁶

Provider Confidence. Change in confidence and self-efficacy focused on whether providers reported greater confidence in ability to diagnose and/or treat patients following EELM participation.^{17, 23, 26, 30, 32, 34, 41, 48, 51, 52, 56, 71} Metrics along these lines were reported in most studies, ranging from self-reported changes following participation,³² to within-subjects change from baseline to endline,⁴⁸ to between-subjects comparisons in perceived competence,³⁰ including in one randomized controlled trial (RCT).²⁶ In most instances, results were

positive and significant; a notable exception came from the RCT on chronic pain management.²⁶

Provider Behavior Change. Several studies administered surveys in which providers were asked to self-report behavior change as a result of participating in case presentations. For example, Komaromy and colleagues³⁵ found 77% of participants reported that case discussion changed their patient care plan. Likewise, Catic and colleagues²¹ observed that recommendations for treatment were incorporated by case presenters 89% of the time. Qaddoumi and colleagues⁴⁶ reported that 91% of case presenters followed recommendations. In other studies, providers were merely asked via survey whether EELM participation had or would alter care provision^{19, 43}; on such occasions, providers responded positively.

Patient-Related Outcomes

Fifteen of 52 identified studies (29%) discussed patient-related outcomes, including changes in care processes and outcomes of care. Few studies examined cost of care.

Hepatitis C. Four studies reviewed hepatitis C outcomes. Arora and colleagues¹⁶ compared sustained virologic response (SVR) between patients at training versus trainee sites and found no difference ($p > 0.05$), indicating trainees (generalists) performed at a level comparable to trainers (specialists). Similarly, Mohsen and colleagues⁶² compared 100 patients of providers who participated in an EELM to 100 patients who received care in a tertiary liver clinic (TLC). Initiation of direct-acting antiviral therapy was similar between groups (EELM, 78%; TLC, 81%), as was completion of treatment (EELM, 89%; TLC, 86%) and—to a lesser




Figure 2 Publications by topic, year, and outcome classification.

extent—SVR (EELM, 87%; TLC, 96%). Statistical significance was not reported.

Beste and colleagues¹⁸ identified providers trained via EELM, and compared likelihood of patient treatment initiation among EELM participants and non-participants. The authors found treatment initiation was higher among trainees (hazard ratio [HR], 1.20; $p < 0.01$), but this effect was a result of increased initiations among only those patients presented in case discussions (HR, 3.30; $p < 0.01$). Ni Cheallaigh and colleagues⁴³ conducted a series of semi-structured interviews with EELM trainees. Interviewees reported that patients attending their practice were beneficiaries of ECHO. For example, one trainee remarked, “Now, access to specialist clinics has improved. [The local specialist] has actually taken back some people he discharged. He’s also seen a couple of new people.”⁴³

Chronic Liver Disease. We identified two studies on chronic liver disease. The first, by Glass and colleagues,²⁹ found that EELM training allowed patients to access care an average of 9.6 days sooner and saved 250 miles of travel compared with those seeking in-clinic specialty care. A second study, by Su and colleagues,⁶⁴ examined the effect of receiving a virtual consultation through the VA’s SCAN-ECHO program. Between 2011 and 2015, 513 veterans with chronic liver disease received a virtual consultation from a SCAN-ECHO provider, while 62,237 did not. After propensity score matching on characteristics predictive of receiving a visit, researchers found hazard ratio of all-cause mortality among those receiving a virtual consultation to be 0.54 ($p = 0.003$), compared with no visit.

Chronic Pain Management and Opioid Addiction. Four studies examined chronic pain management. Anderson and colleagues¹⁵ compared providers at community health centers who participated in EELM trainings with those who did not. They found, among those who participated, the percent of patients with an opioid prescription declined from 56.2 to 50.5% ($p = 0.02$), with no decline observed in the comparison group. Conversely, referrals for behavioral health and physical therapy increased ($p < 0.001$). Two other studies, by Katzman and colleagues⁶¹ and Frank and colleagues,⁵⁹ examined prescription and referral rates, respectively. Katzman and colleagues⁶¹ inspected opioid prescription rates across 1382 clinics associated with the Army and Navy, 99 of which participated in an EELM. Compared with patients of providers who did not participate in EELM ($n = 1,187,945$), those with providers who did participate ($n = 52,941$) observed a much greater decline in prescriptions: from 23 to 9% ($p < 0.001$). Meanwhile, Frank and colleagues inspected likelihood of referral among patients presented as EELM cases versus those not presented as cases; cases were more likely to be referred to physical therapy (HR, 1.10; $p < 0.05$).⁵⁹ A final study, by Carey and colleagues,⁵⁸

performed a spatial reach analysis, concluding that patient travel distance to specialty pain care was associated with only slightly lower odds of access to an EELM-trained provider (OR, 0.98; $p = 0.01$), versus sizably lower odds of care receipt at a specialty care clinic (OR, 0.78; $p < 0.001$).

Geriatric Care. Three studies examined elderly care for those with mental health conditions, including dementia; one additional study examined transitional care. Catic and colleagues²¹ studied the effect of adhering to expert recommendations for residents with dementia, and found that providers who followed EELM recommendations were more likely to report “clinical improvement” among patients (74% vs. 20%; $p = 0.03$). Fisher and colleagues²⁸ examined the relative change in care utilization and costs among elderly patients with mental health conditions, compared with elderly patients without such conditions, before versus after providers participated in EELM training. Among patients with mental health conditions, there was a reduction in emergency department costs: from \$406 to \$311 ($p < 0.05$); this reduction was not observed in the comparison group. Gordon and colleagues⁶⁰ examined quality of care metrics among elderly patients at facilities with providers who were EELM-trained versus not. They observed non-significant differences on primary outcomes (restraint and antipsychotic medication use), but they did find lower rates of urinary tract infections (UTI) among patients seen at facilities with providers trained through EELM (OR for UTI, 0.77; $p < 0.05$).

Moore and colleagues⁶³ examined transitional care among elderly adults. Among patients with providers at a skilled nursing facility who had participated in EELM training, the authors found shorter lengths of inpatient stay ($p = 0.01$), lower 30-day hospital re-admission rates ($p = 0.03$), and lower 30-day care costs ($p < 0.001$) compared with providers who had not participated. This difference was significant even after adjusting for baseline differences in case mix.

Diabetes Management. Watts and colleagues⁶⁵ reported training two primary care physicians on diabetes management through EELM. Providers reported that—among patients with poorly controlled diabetes (i.e., all patients with HbA1c > 9%)—mean HbA1c levels decreased from 10.2% before training sessions to 8.4% after training ($p < 0.001$) 5 months later, a clinically significant difference.

Strength of Evidence

Provider-related outcomes have relied heavily on self-reports for providers who (i) self-select to participate in EELM, (ii) maintain participation in trainings over time, and (iii) complete feedback surveys. The one RCT examining provider-related outcomes concluded null results. Among studies that collected data before versus after EELM trainings, most offered no

control group, raising the question of what would have happened in the absence of training, or if EELM trainings were substituted with a different set of learning tools.

Quality of patient-related outcomes varied widely. While mental health and substance use disorders have been the most frequently implemented EELM in the USA, we found no literature describing the impact of EELMs on patient outcomes associated with these conditions apart from among elderly adults. For conditions like osteoporosis, for which there were provider-related outcomes, we identified no articles assessing patient-related outcomes. For hepatitis C, chronic pain management, dementia care, and diabetes, there was at least one article published in which a counterfactual was incorporated. Two studies, one by Anderson and colleagues¹⁵ and one by Katzman and colleagues,⁶¹ employed quasi-experimental approaches. In a majority of instances, authors identified statistically significant results in favor of EELM. With the exceptions of virologic suppression in the context of hepatitis C and A1C levels in the context of diabetes, reported outcomes were process measures rather than outcome measures.

DISCUSSION

We identified 52 studies between 2007 and 2018 that reported provider- and/or patient-related outcomes from EELM of medical tele-education. Based on our analysis, the empirical evidence for EELM's impact on patient and provider outcomes is low.

Regarding provider-related outcomes, 43 articles have been published in the past 11 years. Over three-quarters provide no between-subjects comparison group, raising a question as to what would be observed in the absence of intervention or under an alternative intervention. While measures like provider satisfaction and self-efficacy are inherently subjective and susceptible to social desirability bias, such biases could be addressed by the inclusion of an active control condition or a third-party evaluator. In 14 instances, no baseline data were recorded, meaning that change in outcomes due to the intervention was unobservable. For measures like provider knowledge, which can be more objectively measured through formal testing, over half of studies relied on subjective self-reports. The response rate was also as low as 7% of participants, suggesting self-selection bias.³⁷ The one recorded cluster RCT, by Eaton and colleagues,²⁶ did not find a benefit of EELM in terms of provider knowledge or perceived competence.

Arguably, the most important provider-related outcomes are behavior changes. To this end, three studies examined the effect of provider presentations on provider behavior. The authors found that providers who presented cases altered care 77–91% of the time.^{21, 35, 46} Beste and colleagues¹⁸ also found that presenting cases resulted in increased initiation of patient care for hepatitis C. However, increased treatment initiation was only observed among those cases presented during

trainings, and not among other patients treated by the same provider. This raises a question of whether EELM are truly building capacity to handle cases without assistance and constitutes an important area for further investigation.

With respect to patient-related outcomes, we identified 15 EELM studies published over the past 11 years. Eleven of these included a comparison group; however, none involved an element of randomization. For all but two measures, outcomes examined were process measures—for example, frequency of prescriptions or number of referrals. While process measures are likely associated with direct patient outcomes, the inferences drawn from these are indirect. However, for three conditions—hepatitis C, dementia care, and chronic pain management—studies showed improvements in different processes and outcomes of care, which suggests that EELM may be beneficial in those conditions.

In terms of direct patient outcome measures, three studies examined the rate of sustained virologic response among individuals treated for hepatitis C. While two articles found that SVR was similar among patients who sought care from EELM trainees and experts^{16, 62}—an indication of project success—another found that SVR did not differ between those seeing providers who received EELM training versus did not receive EELM training.¹⁸ In a separate study looking at the effect of EELM on diabetes care,⁶⁵ the authors found positive results that training led to reductions in patient mean HbA1c values within practitioner panels, a result not found in a comparison group. While results here show promise, the study was limited by sample size, with only 2 EELM providers and 2 control providers.

While existing evidence suggests the potential impact of EELM in improving patient outcomes, our SOE assessment underscores the need for more rigorous evaluation to substantiate the model. One of the main findings is that the quality of evidence for the effectiveness of EELM is generally rated as “low” or “very low” based on the GRADE system. However, it is important to note that this finding is by no means limited to EELM; in fact, many models of service delivery are supported by limited evidence, for example, the widescale initiative of Housing First, which provides rapid housing to improve client housing security and health. While it is a widely used model, only limited evidence exists for its impact on long-term health outcomes, despite four randomized controlled trials.⁷² Nevertheless, it is appropriate to continue to strive for higher quality evidence.

A few study limitations should be noted. First, while search criteria were meant to be broad, it is possible articles were overlooked, particularly if they did not contain key words in the Appendix Table online. Second, we were unable to include works in progress, though we identified several. Similarly, due to a limited subset of studies that were not ECHO affiliated, we were not equipped to make ECHO versus non-ECHO EELM comparisons, though this may be an interesting avenue to pursue as non-ECHO EELM proliferate.

In summary, we identified 52 articles over the past 10 years which outline provider- and patient-related outcomes from EELM implementations. One of these comprised a randomized controlled trial with non-significant findings, while a plurality were cross-sectional surveys with high risk of bias. Given the capacity-building orientation of EELM, it would be important for studies to include longer periods of follow-up to assess maintenance, as well as to compare the costs and outcomes of EELM to alternative forms of continuing medical education that rely on technology. As noted, models like EELM that are novel within the healthcare delivery landscape are liable to incrementally establish an evidence base. In this respect, our findings are not surprising. Rather, the assessment is meant to provide an inventory of the existing literature, which may establish a benchmark for EELM moving forward.

Acknowledgments: The authors thank the funders of this project at ASPE. We thank Caryn Marks, Nancy De Lew, and Rose Chu for their support. Among our colleagues at the RAND Corporation, we are grateful to Christine Eibner, Jody Larkin, Justin Timbie, Lisa Turner, Lori Uscher-Pines, Monique Martineau, Paul Koegel, and Tricia Soto.

Corresponding Author: Ryan K. McBain, PhD, MPH; RAND Corporation, Boston, MA, USA (e-mail: rmcbain@rand.org).

Funding Information: This work was supported by the Office of the Assistant Secretary for Planning and Evaluation, US Department of Health and Human Services, under master contract, Building Analytic Capacity for Monitoring and Evaluating the Implementation of the ACA, HHSP23320095649WC.

Compliance with Ethical Standards:

Conflict of Interest: Drs. McBain, Rose, Faherty, and Fischer, as well as Ms. Sousa and Ms. Baxi, report support from ASPE through a contract with the RAND Corporation during the conduct of the study. All remaining authors declare that they do not have a conflict of interest.

Disclaimer: The authors of this manuscript are responsible for its content. Statements in the manuscript should not be construed as endorsements of ASPE or HHS.

REFERENCES

1. **Kvedar J, Coye MJ, Everett W.** Connected health: A review of technologies and strategies to improve patient care with telemedicine and telehealth. *Health Affairs (Project Hope)* 2014;33(2):194-9.
2. **Dorsey ER, Topol EJ.** State of telehealth. *N Engl J Med* 2016;375(2):154-61.
3. **Alkmim MB, Figueira RM, Marcolino MS, Cardoso CS, Abreu MPD, Cunha LR, et al.** Improving patient access to specialized health care: The Telehealth Network of Minas Gerais, Brazil. *Bull World Health Organ* 2012;90:373-8.
4. **Terwiesch C, Asch DA, Volpp KG.** Technology and Medicine: Reimagining Provider Visits as the New Tertiary Care. *Ann Intern Med* 2017;167(11):814-5. doi:<https://doi.org/10.7326/m17-0597>
5. **Ricketts TC.** The migration of physicians and the local supply of practitioners: a five-year comparison. *Acad Med* 2013;88(12):1913-8.
6. **Liddy C, Rowan MS, Afkham A, Maranger J, Keely E.** Building access to specialist care through e-consultation. *Open Med* 2013;7(1):e1.
7. **Kirch DG, Petelle K.** Addressing the physician shortage: The peril of ignoring demography. *JAMA*. 2017;317(19):1947-8. doi:<https://doi.org/10.1001/jama.2017.2714>
8. **Kinфу Y, Dal Poz MR, Mercer H, Evans DB.** The health worker shortage in Africa: Are enough physicians and nurses being trained? : *SciELO Public Health*; 2009.
9. **ECHO Act, Pub L No. 114-270, 130 Stat 1395** (2016).
10. **University of New Mexico. Project ECHO.** Available at: <https://echo.unm.edu/>. Accessed June 16, 2019.
11. **Fischer SH, Rose AJ, McBain RK, Faherty LJ, Sousa J, Martineau M.** Evaluation of technology-enabled collaborative learning and capacity building models: Materials for a report to Congress. Santa Monica, CA: RAND Corporation, 2019. Available at: https://www.rand.org/pubs/research_reports/RR2934.html.
12. **Zhou C, Crawford A, Serhal E, Kurdyak P, Sockalingam S.** The impact of Project ECHO on participant and patient outcomes: A systematic review. *Academic medicine : journal of the Association of American Medical Colleges* 2016;91(10):1439-61. doi:<https://doi.org/10.1097/acm.0000000000001328>
13. **Ryan R, Hill S.** How to GRADE the quality of the evidence. *Cochrane Consumers and Communication Group*. 2016. Available at: <http://cccg.cochrane.org/author-resources>. Version 3.0. Accessed June 16, 2019.
14. **Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group.** Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Ann Intern Med* 2009;151(4):264-9.
15. **Anderson D, Zlateva I, Davis B, Bifulco L, Giannotti T, Coman E, et al.** Improving pain care with Project ECHO in community health centers. *Pain Med (Malden, Mass.)*. 2017;18(10):1882-9. doi:<https://doi.org/10.1093/pm/pnx187>
16. **Arora S, Thornton K, Murata G, Deming P, Kalishman S, Dion D, et al.** Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med* 2011;364(23):2199-207. doi:<https://doi.org/10.1056/NEJMoa1009370>
17. **Ball S, Wilson B, Ober S, McHaourab A.** SCAN-ECHO for Pain Management: Implementing a Regional Telementoring Training for Primary Care Providers. *Pain Med (Malden, Mass.)*. 2018;19(2):262-8. doi:<https://doi.org/10.1093/pm/pnx122>
18. **Beste LA, Glorioso TJ, Ho PM, Au DH, Kirsh SR, Todd-Stenberg J, et al.** Telemedicine specialty support promotes hepatitis C treatment by primary care providers in the Department of Veterans Affairs. *Am J Med*. 2017;130(4):432-8.e3. doi:<https://doi.org/10.1016/j.amjmed.2016.11.019>
19. **Beste LA, Mattox EA, Pichler R, Young BA, Au DH, Kirsh SF, et al.** Primary care team members report greater individual benefits from long-versus short-term specialty telemedicine mentorship. *Telemedicine journal and e-health : the official journal of the American Telemedicine Association* 2016;22(8):699-706. doi:<https://doi.org/10.1089/tmj.2015.0185>
20. **Carlin L, Zhao J, Dubin R, Taenzer P, Sidrak H, Furlan A.** Project ECHO telementoring intervention for managing chronic pain in primary care: Insights from a qualitative study. *Pain Med (Malden, Mass.)*. 2018;19(6):1140-6. doi:<https://doi.org/10.1093/pm/pnx233>
21. **Catic AG, Mattison ML, Bakaev I, Morgan M, Monti SM, Lipsitz L.** ECHO-AGE: an innovative model of geriatric care for long-term care residents with dementia and behavioral issues. *J Am Med Dir Assoc* 2014;15(12):938-42. doi:<https://doi.org/10.1016/j.jamda.2014.08.014>
22. **Chaple MJ, Freese TE, Rutkowski BA, Krom L, Kurtz AS, Peck JA, Warren P, Garrett S.** Using ECHO Clinics to Promote Capacity Building in Clinical Supervision. *American Journal of Preventive Medicine*. 54:S275-S280. 2018.
23. **Cofta-Woerpel L, Lam C, Reitzel LR, Wilson W, Karam-Hage M, Beneventi D, et al.** A tele-mentoring tobacco cessation case consultation and education model for healthcare providers in community mental health centers. *Cogent Med*. 2018;5(1). doi:<https://doi.org/10.1080/2331205X.2018.1430652>
24. **Cordasco KM, Zuchowski JL, Hamilton AB, Kirsh S, Veet L, Saavedra JO, et al.** Early lessons learned in implementing a women's health educational and virtual consultation program in VA. *Med Care* 2015;53(4 Suppl 1):S88-92. doi:<https://doi.org/10.1097/mlr.0000000000000313>
25. **Covell NH, Foster FP, Margolies PJ, Lopez LO, Dixon LB.** Using distance technologies to facilitate a learning collaborative to implement stage-wise treatment. *Psychiatric services (Washington, D.C.)*. 2015;66(6):645-8. doi:<https://doi.org/10.1176/appi.ps.201400155>
26. **Eaton LH, Godfrey DS, Langford DJ, Rue T, Tauben DJ, Doorenbos AZ.** Telementoring for improving primary care provider knowledge and competence in managing chronic pain: A randomised controlled trial. *J Telemed Telecare*. 2018;1357633x18802978. doi:<https://doi.org/10.1177/1357633x18802978>

- novel video-conferencing program: ECHO-Care Transitions. *Am J Med* 2017;130(10):1199–204. doi:<https://doi.org/10.1016/j.amjmed.2017.04.041>
64. **Su GL, Glass L, Tapper EB, Van T, Waljee AK, Sales AE.** Virtual Consultations through the Veterans Administration SCAN-ECHO Project Improves Survival for Veterans with Liver Disease. *Hepatology*. 2018. doi:<https://doi.org/10.1002/hep.30074>
65. **Watts SA, Roush L, Julius M, Sood A.** Improved glycemic control in veterans with poorly controlled diabetes mellitus using a Specialty Care Access Network-Extension for Community Healthcare Outcomes model at primary care clinics. *J Telemed Telecare* 2016;22(4):221–4. doi:<https://doi.org/10.1177/1357633x15598052>
66. **Moore JE, Rashid S, Park JS, Khan S, Straus SE.** Longitudinal evaluation of a course to build core competencies in implementation practice. *Implementation Sci* : IS 2018;13(1):106.
67. **Metz A, Louison L, Ward C, Burke K.** Global Implementation Specialist Practice Profile: Skills and competencies for implementation practitioners. Chapel Hill, N.C.: National Implementation Research Network and Centre for Effective Services, 2017. Available at: <https://fpg.unc.edu/node/9178>.
68. Agency for Healthcare Research and Quality. Types of health care quality measures. 2011. Available at: <http://www.ahrq.gov/talkingquality/measures/types.html>. Accessed June 16, 2019.
69. U.S. Department of Health & Human Services. Office of the Assistant Secretary for Planning and Evaluation. Available at: <https://aspe.hhs.gov/>. Accessed June 16, 2019.
70. **Rahman AN, Schnelle JF, Yamashita T, Patry G, Prasauskas R.** Distance learning: A strategy for improving incontinence care in nursing homes. *The Gerontologist* 2010;50(1):121–32. doi:<https://doi.org/10.1093/geront/gnp126>
71. **Ricketts EJ, Goetz AR, Capriotti MR, Bauer CC, Brei NG, Himle MB,** et al. A randomized waitlist-controlled pilot trial of voice over Internet protocol-delivered behavior therapy for youth with chronic tic disorders. *J Telemed Telecare* 2016;22(3):153–62. doi:<https://doi.org/10.1177/1357633x15593192>
72. **Baxter AJ, Tweed EJ, Katikireddi SV, Thomson H.** Effects of Housing First approaches on health and well-being of adults who are homeless or at risk of homelessness: Systematic review and meta-analysis of randomised controlled trials. *J Epidemiol Community Health*. 2019;jech-2018-210981.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.