

University of New Mexico

UNM Digital Repository

Mathematics and Statistics Faculty and Staff
Publications

Academic Department Resources

4-28-2018

On Rugina's system of thought

Florentin Smarandache

University of New Mexico, smarand@unm.edu

Follow this and additional works at: https://digitalrepository.unm.edu/math_fsp



Part of the [Logic and Foundations Commons](#), [Number Theory Commons](#), and the [Other Mathematics Commons](#)

Recommended Citation

Florentin Smarandache, (2001) "On Rugina's system of thought", *International Journal of Social Economics*, Vol. 28 Issue: 8, pp.623-647, <https://doi.org/10.1108/EUM0000000005543>

This Article is brought to you for free and open access by the Academic Department Resources at UNM Digital Repository. It has been accepted for inclusion in Mathematics and Statistics Faculty and Staff Publications by an authorized administrator of UNM Digital Repository. For more information, please contact amywinter@unm.edu, lsloane@salud.unm.edu, sarahrk@unm.edu.



International Journal of Social Economics

On Rugina's system of thought
Florentin Smarandache,

Article information:

To cite this document:

Florentin Smarandache, (2001) "On Rugina's system of thought", International Journal of Social Economics, Vol. 28 Issue: 8, pp.623-647, <https://doi.org/10.1108/EUM0000000005543>

Permanent link to this document:

<https://doi.org/10.1108/EUM0000000005543>

Downloaded on: 27 April 2018, At: 08:11 (PT)

References: this document contains references to 13 other documents.

To copy this document: permissions@emeraldinsight.com

The fulltext of this document has been downloaded 171 times since 2006*

Access to this document was granted through an Emerald subscription provided by

For Authors

If you would like to write for this, or any other Emerald publication, then please use our Emerald for Authors service information about how to choose which publication to write for and submission guidelines are available for all. Please visit www.emeraldinsight.com/authors for more information.

About Emerald www.emeraldinsight.com

Emerald is a global publisher linking research and practice to the benefit of society. The company manages a portfolio of more than 290 journals and over 2,350 books and book series volumes, as well as providing an extensive range of online products and additional customer resources and services.

Emerald is both COUNTER 4 and TRANSFER compliant. The organization is a partner of the Committee on Publication Ethics (COPE) and also works with Portico and the LOCKSS initiative for digital archive preservation.

*Related content and download information correct at time of download.



On Rugina's system of thought

Florentin Smarandache

University of New Mexico, Gallup, New Mexico, USA

On Rugina's
system of
thought

Keywords *Language, Mathematics, Philosophy*

623

Abstract *This article investigates Rugina's orientation table and gives particular examples for several of its seven models. Leon Walras's Economics of Stable Equilibrium and Keynes's Economics of Disequilibrium are combined in Rugina's orientation table in systems which are s percent stable and $100 - s$ percent unstable, where s may be 100, 95, 65, 50, 35, 5, and 0. Classical logic and modern logic are united in Rugina's integrated logic, and then generalized in neutrosophic logic.*

I. Introduction

Coming across Rugina's system of thoughts, in his published books and articles (Rugina, 1984, 1989, 1994, 1998), I learned about the connection between classic and modern. It is not a contradiction, but a complementarity from the part of modern with respect to the classic; and always the new "modern" will have something to bring to the old knowledge. In a similar way we may talk on the complementarity between theory and practice, rather than their contradiction.

In economics, Rugina negated Marx's social justice for the mass and Keynes's involuntary unemployment. His methodology in science tries to unite all scientific fields, preserving, however, independence in thinking and judgement. Einstein worked in the last period of his life on the unified field theory (a single general theory in physics), but didn't succeed. At the present, his supposition that the speed of light is a barrier in the universe is also being denied.

The economical systems are characterized by free market or centrally-planned and controlled economy. I think each system has a mixture of the previous, where a part of the market is free and another is centrally-planned and controlled.

Rugina's universal hypothesis of duality states that the physical universe is composed of stable and unstable elements arranged in various proportions, may be completed with unknown elements, a strip border between stable and unstable, which are continuously changing from the state of equilibrium to disequilibrium and vice-versa, and which therefore are giving the dynamics of the universe.

Unknowns may be:

- anomalies;
- relativities;
- uncertainties;
- revolution risks; and
- hidden parameters.

The internal parameters are involved in Rugina's universal law of natural parameter (NaPa):

Any system in order to reach and maintain a position of stable equilibrium must have a very strong natural parameter (center of weight).

Whereas the external parameters are involved in Rugina's universal law of general consistency:

Any system produces and maintains a position of stable equilibrium if there is a suitable space-time frame work.

II. Theory of paradoxes

How did I get to the theory of paradoxes? I have observed that: what's good for someone, may be bad for others – and reciprocally. There are people who are considered terrorists by their enemies, and patriots by their friends. All of them are right and wrong at the same time. If one changes the referential system, the result is different.

Let's see a few nice paradoxes:

Social paradox. In a democracy should the nondemocratic ideas be allowed?

- (a) If yes, i.e. the nondemocratic ideas are allowed, then one has not a democracy anymore. (The nondemocratic ideas may overturn the society.)
- (b) If no, i.e. other ideas are not allowed – even those nondemocratic – then one has not a democracy either, because the freedom of speech is restricted.

The sets' paradox. The notion of “set of all sets”, introduced by Georg Cantor, does not exist.

Let all sets be noted by $\{S_a\}_a$, where a indexes them. But the set of all sets is itself another set, say T_1 ; and then one constructs again another “set of ⟨all sets⟩”, but ⟨all sets⟩ are this time $\{S_a\}$ and T_1 , and then the “set of all sets” is now T_2 , different from T_1 ; and so on. Even the notion of “all sets” can not exactly be defined (like the largest number of an open interval, which doesn't exist), as one was just seeing above (we can construct a new set as the “set of all sets”) and reunites it to “all sets”.

A paradoxist psychological complex (with the accent on the first syllable). A collection of fears stemming from previous unsuccessful experience or from unconscious feelings that, wanting to do something ⟨S⟩, the result would be ⟨Anti-S⟩, which give rise to feelings, attitudes, and ideas pushing the subject towards a deviation of action ⟨S⟩ eventually towards an ⟨Anti-S⟩ action. (From the positive and negative brain's electrical activities.) For example, a shy boy, attempting to invite a girl to dance, inhibits himself through fear that she would turn him down. How to manage this phobia? To dote and anti-dote! By transforming it into an opposite one, thinking differently, and being fear in our mind that we would pass our expectancies but we shouldn't. People who do not try for fear of being rejected: they lose by not competing!

Auto-suggestion

If an army leaves for war with anxiety to lose, that army is half-defeated before starting the confrontation.

Paradoxist psychological behavior. How can we explain contrary behaviors of a person: in the same conditions, without any reason, cause? Because our deep unconsciousness is formed of contraries.

Ceaseless anxiety. What you want is, normally, what you don't get. And this is for eternity. Like a chain, because, when you get it (if ever), something else will be your next desire. Man can't live without a new hope.

Inverse desire. The wish to purposely have bad luck, to suffer, to be pessimistic as stimulating factors for more and better creation or work. (Applies to some artists, poets, painters, sculptors, spiritualists.)

All is possible, the impossible too! Is this an optimistic or pessimistic paradox?

- (a) It is an optimistic paradox, because it shows that all is possible.
- (b) It is a pessimistic paradox, because it shows that the impossible is possible.

Mathematician's paradox. Let M be a mathematician who may not be characterized by his mathematical work.

- (a) To be a mathematician, M should have some mathematical work done, therefore M should be characterized by that work.
- (b) The reverse judgement: if M may not be characterized by his mathematical work, then M is not a mathematician.

Divine paradox (I). Can God commit suicide?

- (a) If God cannot, then it appears that there is something God cannot do, therefore God is not omnipotent.
- (b) If God can commit suicide, then God dies – because He has to prove it, therefore God is not immortal.

Divine paradox (II). Can God be atheist, governed by scientific laws?

- (a) If God can be atheist, then God doesn't believe in Himself, therefore why should we believe in Him?
- (b) If God cannot, then again He's not omnipotent.

Religion is full of god-ism and evil-ism. God and evil in the same being. Man is a bearer of good and bad simultaneously. Man is enemy to himself. God and Magog!

Expect the unexpected. If we expect someone to do the unexpected, then:

- Is it possible for him to do the unexpected?
- Is it possible for him to do the expected?

If he does the unexpected, then that's what we expected. If he doesn't do the expected, then he did the unexpected.

The ultimate paradox. Living is the process of dying.

Reciprocally: death of one is the processes of somebody else's life (an animal eating another one).

Exercises for readers:

- If China and Japan are in the Far East, why from the USA do we go west to get there?
- Are humans inhuman, because they committed genocides?

The invisible paradoxes. Our visible world is composed of a totality of invisible particles. Things with mass result from atoms with quasi-null mass. Infinity is formed of finite part(icle)s.

Look at these Sorites paradoxes (associated with Eubulides of Miletus, fourth century BC):

- (a) An invisible particle does not form a visible object, nor do two invisible particles, three invisible particles, etc. However, at some point, the collection of invisible particles becomes large enough to form a visible object, but there is apparently no definite point where this occurs.
- (b) A similar paradox is developed in an opposite direction. It is always possible to remove an atom from an object in such a way that what is left is still a visible object. However, repeating and repeating this process, at some point, the visible object is decomposed so that the part left becomes invisible, but there is no definite point where this occurs.

Between $\langle A \rangle$ and $\langle \text{Non-}A \rangle$ there is no clear distinction, no exact frontier. Where does $\langle A \rangle$ really end and $\langle \text{Non-}A \rangle$ begin? We extend Zadeh's fuzzy set term to fuzzy concept.

Paradoxist existentialism

Life's value consists in its lack of value; life's sense consists in its lack of sense.

Semantic Paradox (I): "I am who I am not". If I am not Socrates, and since I am who I am not, it results that I am Socrates. If I am Socrates, and since I am who I am not, it results that I am not Socrates. Generally speaking: "I am X" if and only if "I am not X". Who am I?

In a similar pattern one constructs the paradoxes: "I am myself when I am not myself"; and "I exist when I don't exist". And, for the most part, "I {verb} when I don't {verb}" (Smarandache, 1997).

What is a dogma? Is it an idea that makes you have no other idea. How can we get rid of such authoritative tenet? (To un-read and un-study it!)

Semantic Paradox (II): "I don't think". This can not be true for, in order to even write this sentence, I needed to think (otherwise I was writing with mistakes, or was not writing it at all). Whence "I don't think" is false, which means "I think".

Unsolved mysteries:

- (a) Is it true that for each question there is at least an answer?
- (b) Is any statement the result of a question?
- (c) Let $P(n)$ be the following assertion: "If $S(n)$ is true, then $S(n + 1)$ is false", where $S(n)$ is a sentence relating on parameter n . Can we prove by mathematical induction that $P(n)$ is true?
- (d) " $\langle A \rangle$ is true if and only if $\langle A \rangle$ is false". Is this true or false?
- (e) How can this assertion "Living without living" be true? Find a context. Explain.

On Rugina's
system of
thought

627

Other examples of such paradoxes are:

- $\langle \text{Anti-}A \rangle$ of $\langle A \rangle$ (anti-literature of literature);
- $\langle \text{Non-}A \rangle$ of $\langle A \rangle$ (language of non-language); and
- $\langle A \rangle$ of $\langle \text{Non-}A \rangle$ (artistic of the non-artistic).

Tautologies. I want because I want (showing will, ambition) ($\langle A \rangle$ because of $\langle A \rangle$) (Smarandache, 1997).

Other examples include:

- Our axiom is to break down all axioms.
- Be patient without patience.
- Non-existence exists.
- Culture exists through its non-existence.
- Our culture is our lack of culture.
- Style without style.
- The rule we apply: there is no rule.

Paradox of the paradoxes

Is "this is a paradox" a paradox? I mean is it true or false? Examples are:

- To speak without speaking (without words (body language)).
- To communicate without communicating.
- To do the impossible.
- To know nothing about everything, and everything about nothing.
- I do only what I can't! If I can't do something, of course "I can do" is false. And, if I can do, it's also false because I can do only what I am not able to do.
- I cannot for I can.
- Paradoxal sleep, from a French "*Larousse*" dictionary (1989), is a phase of the sleep when the dreams occur. Sleep, sleep, but why paradoxal? How do the dreams put up with reality?

- Is O.J. Simpson's crime trial an example of: justice of injustice, or injustice of justice? However, his famous release is a victory against the system!
- Corrupt the incorruptible!

Everything which is not paradoxist, is, however, paradoxist. This is the great universal paradox. A superparadox (as a superman in a hyperspace).

Facts exist in isolation from other facts (the analytic philosophy), and in connection as well with each other (Whitehead's and Bergson's thoughts). The neutrosophic philosophy unifies contradictory and noncontradictory ideas in any human field.

The antagonism doesn't exist. Or, if the antagonism does exist, this becomes (by neutrosophic view) a non- (or un-)antagonism: a normal thought. I don't worry about it as well as Wordsworth.

Platonism is the observable of unobservable, the thought of the non-thought.

The essence of a thing may never be reached. It is a symbol, a pure and abstract and absolute notion.

An action may be considered g percent good (or right) and b percent bad (or wrong), where $0 \leq g + b \leq 100$ – the remainder being indeterminacy, not only ⟨good⟩ or only ⟨bad⟩ – with rare exceptions, if its consequence is g percent happiness (pleasure). In this case the action is g percent-useful (in a semi-utilitarian way). Utilitarianism shouldn't work with absolute values only!

Verification has a pluri-sense because we have to demonstrate or prove that something is t percent true, and f percent false, where $0 \leq t, f \leq 100$ and $t + f \leq 100$, not only $t = 0$ or 100 – which occurs in rare/absolute exceptions, by means of formal rules of reasoning of this neutrosophic philosophy.

The logical cogitation's structure is discordant. Scientism and empiricism are strongly related. They can't run one without other, because one exists in order to complement the other and to differentiate it from its opponent. Plus doesn't work without minus, and both of them supported by zero. They all are cross-penetrating sometimes up to confusion. The non-understandable is understandable. If vices wouldn't exist, the virtues will not be seen (T. Muşatescu).

Any new born theory (notion, term, event, phenomenon) automatically generates its non-theory – not necessarily anti-(notion, term, event, phenomenon). Generally speaking, for any ⟨A⟩ a ⟨Non-A⟩ (not necessarily ⟨Anti-A⟩) will exist for compensation.

The neutrosophy is a theory of theories, because at any moment new ideas and conceptions are appearing and implicitly their negative and neutral senses are highlighted.

The non-important is important, because the first one is the second one's shadow that makes it grow its value. The important things would not be so without any unimportant comparison.

The neutrosophic philosophy accepts *a priori* and *a posteriori* any philosophical idea, but associates it with adverse and neutral ones, as a

summum. This is to be neutrosophic without being! Its schemes are related to the neutrality of everything.

On Rugina's
system of
thought

III. On Rugina's orientation table

Starting from a new viewpoint in philosophy, the neutrosophy, one extends the classical "probability theory", "fuzzy set" and "fuzzy logic" to ⟨neutrosophic probability⟩, ⟨neutrosophic set⟩ and ⟨neutrosophic logic⟩ respectively.

They are useful in artificial intelligence, neural networks, evolutionary programming, neutrosophic dynamic systems, quantum theory, and decision making in economics.

With the neutrosophic logic help one explores Rugina's orientation table, a remarkable tool of study, at the micro- and macro-level, of problems in all sciences.

(1) *Neutrosophy: a new branch of mathematical philosophy*

(A) *Etymology*. Neutro-sophy (French *neutre* < Latin *neuter*, neutral, and Greek *sophia*, skill/wisdom) means knowledge of neutral thought.

(B) *Definition*. Neutrosophy is a new branch of philosophy which studies the origin, nature, and scope of neutralities, as well as their interactions with different ideational spectra.

(C) *Characteristics*. This mode of thinking:

- proposes new philosophical theses, principles, laws, methods, formulas, movements;
- interprets the uninterpretable;
- regards, from many different angles, old concepts, systems: showing that an idea, which is true in a given referential system, may be false in another one, and vice versa;
- measures the stability of unstable systems, and instability of stable systems.

(D) *Methods of neutrosophic study*. The methods of neutrosophic study are mathematization (neutrosophic logic, neutrosophic probability and statistics, duality), generalization, complementarity, contradiction, paradox, tautology, analogy, reinterpretation, combination, interference, aphoristic, linguistic, and multidisciplinary.

(E) *Formalization*. Let's note by ⟨A⟩ an idea or theory or concept, by ⟨Non-A⟩ what is not ⟨A⟩, and by ⟨Anti-A⟩ the opposite of ⟨A⟩. Also, ⟨Neut-A⟩ means what is neither ⟨A⟩, nor ⟨Anti-A⟩, i.e. neutrality in between the two extremes. And ⟨A'⟩ a version of ⟨A⟩. ⟨Non-A⟩ is different from ⟨Anti-A⟩.

For example, if ⟨A⟩ = white, then ⟨Anti-A⟩ = black (antonym), but ⟨Non-A⟩ = green, red, blue, yellow, black, etc. (any color, except white), while ⟨Neut-A⟩ = green, red, blue, yellow, etc. (any color, except white and black), and ⟨A'⟩ = dark white, etc. (any shade of white). ⟨Neut-A⟩ ≡ ⟨Neut-(Anti-A)⟩, neutralities of ⟨A⟩ are identical with neutralities of ⟨Anti-A⟩.

$\langle \text{Non-A} \rangle \supset \langle \text{Anti-A} \rangle$, and $\langle \text{Non-A} \rangle \supset \langle \text{Neut-A} \rangle$ as well, also $\langle A \rangle \cap \langle \text{Anti-A} \rangle = \phi$, $\langle A \rangle \cap \langle \text{Non-A} \rangle = \phi$. $\langle A \rangle$, $\langle \text{Neut-A} \rangle$, and $\langle \text{Anti-A} \rangle$ are disjoint two by two. $\langle \text{Non-A} \rangle$ is the complementitude of $\langle A \rangle$ with respect to the universal set.

(F) *Main principle*. Between an idea $\langle A \rangle$ and its opposite $\langle \text{Anti-A} \rangle$, there is a continuum-power spectrum of neutralities $\langle \text{Neut-A} \rangle$.

(G) *Fundamental thesis*.

Any idea $\langle A \rangle$ is t percent true, i percent indeterminate, and f percent false, where $t + i + f = 100$.

(H) *Main laws*. Let $\langle \alpha \rangle$ be an attribute, and $(a, i, b) \in [0, 100]^3$, with $a + i + b = 100$. Then:

- There is a proposition $\langle P \rangle$ and a referential system $\langle R \rangle$, such that $\langle P \rangle$ is a percent $\langle \alpha \rangle$, i percent indeterminate or $\langle \text{Neut-}\alpha \rangle$, and b percent $\langle \text{Anti-}\alpha \rangle$.
- For any proposition $\langle P \rangle$, there is a referential system $\langle R \rangle$, such that $\langle P \rangle$ is a percent $\langle \alpha \rangle$, i percent indeterminate or $\langle \text{Neut-}\alpha \rangle$, and b percent $\langle \text{Anti-}\alpha \rangle$.
- $\langle \alpha \rangle$ is at some degree $\langle \text{Anti-}\alpha \rangle$, while $\langle \text{Anti-}\alpha \rangle$ is at some degree $\langle \alpha \rangle$.

(2) *Neutrosophic probability and neutrosophic statistics*

Let's first generalize the classical notions of "probability" and "statistics" for practical reasons.

(A) *Definitions*. Neutrosophic probability studies the chance that a particular event E will occur, where that chance is represented by three coordinates (variables): t percent true, i percent indeterminate, and f percent false, with $t + i + f = 100$ and $f, i, t \in [0, 100]$. Neutrosophic statistics is the analysis of such events.

(B) *Neutrosophic probability space*. The universal set, endowed with a neutrosophic probability defined for each of its subsets, forms a neutrosophic probability space.

(C) *Applications*.

- (1) The probability that candidate C will win an election is say 25 percent true (percentage of people voting for him), 35 percent false (percentage of people voting against him), and 40 percent indeterminate (percentage of people not coming to the ballot box, or giving a blank vote – not selecting anyone, or giving a negative vote – cutting all candidates on the list). Dialectic and dualism don't work in this case anymore.
- (2) Another example, the probability that tomorrow it will rain is say 50 percent true according to meteorologists who have investigated the past years' weather, 30 percent false according to today's very sunny and droughty summer, and 20 percent undecided (indeterminate).

(3) *Neutrosophic set*

Let's second generalize, in the same way, the fuzzy set.

(A) *Definition.* Neutrosophic set is a set such that an element belongs to the set with a neutrosophic probability, i.e. t percent is true that the element is in the set, f percent false, and i percent indeterminate.

(B) *Neutrosophic set operations.* Let M and N be two neutrosophic sets. One can say, by language abuse, that any element neutrosophically belongs to any set, due to the percentage of truth/indeterminacy/falsity which varies between 0 and 100. For example: $x(50, 20, 30) \in M$ (which means, with a probability of 50 percent x is in M , with a probability of 30 percent x is not in M , and the rest is undecidable), or $y(0, 0, 100) \in M$ (which normally means y is not for sure in M), or $z(0, 100, 0) \in M$ (which means one doesn't know absolutely anything about z 's affiliation with M).

Let $0 \leq t_1, t_2, t' \leq 1$ represent the truth-probabilities, $0 \leq i_1, i_2, i' \leq 1$ the indeterminacy-probabilities, and $0 \leq f_1, f_2, f' \leq 1$ the falsity-probabilities of an element x to be in the set M and in the set N respectively, and of an element y to be in the set N , where $t_1 + i_1 + f_1 = 1$, $t_2 + i_2 + f_2 = 1$, and $t' + i' + f' = 1$.

One notes, with respect to the given sets, $x = x(t_1, i_1, f_1) \in M$ and $x = x(t_2, i_2, f_2) \in N$, by mentioning x 's neutrosophic probability appurtenance. And, similarly, $y = y(t', i', f') \in N$.

Also, for any $0 \leq x \leq 1$ one notes $1 - x = \bar{x}$. Let $W(a, b, c) = (1 - a)/(b + c)$ and $W(R) = W(R(t), R(i), R(f))$ for any tridimensional vector $R = (R(t), R(i), R(f))$.

Complement of M : Let $N(x) = 1 - x = \bar{x}$. Therefore: if $x(t_1, i_1, f_1) \in M$, then $x(N(t_1), N(i_1), N(f_1)) \in C(M)$.

Intersection. Let $C(x, y) = xy$, and $C(z_1, z_2) = C(z)$ for any bidimensional vector $z = (z_1, z_2)$. Therefore: if $x(t_1, i_1, f_1) \in M$, $x(t_2, i_2, f_2) \in N$, then $x(C(t), C(i), C(f)) \in M \cap N$.

Union. Let $D1(x, y) = x + y - xy = x + \bar{x}y = y + x\bar{y}$, and $D1(z_1, z_2) = D1(z)$ for any bidimensional vector $z = (z_1, z_2)$. Therefore: if $x(t_1, i_1, f_1) \in M$, $x(t_2, i_2, f_2) \in N$, then $x(D1(t), D1(i), D1(f)) \in M \cup N$.

Cartesian product. If $x(t_1, i_1, f_1) \in M$, $y(t', i', f') \in N$, then $(x(t_1, i_1, f_1), y(t', i', f')) \in M \times N$.

Difference. Let $D(x, y) = x - xy = x\bar{y}$, and $D(z_1, z_2) = D(z)$ for any bidimensional vector $z = (z_1, z_2)$. Therefore: if $x(t_1, i_1, f_1) \in M$, $x(t_2, i_2, f_2) \in N$, then $x(D(t), D(i), D(f)) \in M/N$, because $M/N = M \cap C(N)$.

(C) *Applications.* From a pool of refugees, waiting in a political refugee camp to get the American visa of emigration, a percent are accepted, r percent rejected, and p percent in pending (not yet decided), $a + r + p = 100$. The chance for someone in the pool to emigrate to USA is not a percent as in classical probability, but a percent true and p percent pending (therefore normally bigger than a percent) – because later, the p percent pending refugees will be distributed into the first two categories, either accepted or rejected.

Another example, a cloud is a neutrosophic set, because its borders are ambiguous, and each element (water drop) belongs with a neutrosophic probability to the set (e.g. there are separated water drops, around a compact mass of water drops, that we don't know how to consider them: in or out of the

cloud). We are not sure where the cloud ends nor where it begins, neither if some elements are or are not in the set. That's why the percent of indeterminacy is required: for a more organic, smooth, and especially accurate estimation.

(4) Neutrosophic logic: a generalization of fuzzy logic

(A) *Introduction.* One passes from the classical $\{0, 1\}$ bivalent logic of George Boole, to the three-valued logic of Reichenbach (leader of the logical empiricism), then to the $\{0, a_1, \dots, a_n, 1\}$ plurivalent one of Lukasiewicz (and Post's m -valued calculus), and finally to the $[0, 1]$ infinite logic as in mathematical analysis and probability: a transcendental logic (with values of the power of continuum), or fuzzy logic.

Falsehood is infinite, and truthhood quite alike; in between, at different degrees, indeterminacy as well. Everything is G percent good, I percent indeterminate, and B percent bad, where $G + I + B = 100$.

Besides Diderot's dialectics on good and bad ("Rameau's Nephew", 1772), any act has its percentage of "good", "indeterminate", and of "bad" as well incorporated.

Rodolph Carnap said:

Metaphysical propositions are neither true nor false, because they assert nothing, they contain neither knowledge nor error. . .

Hence, there are infinitely many statuses in between "good" and "bad", and generally speaking in between "A" and "Anti-A", like on the real number segment:

[0,	1]
False	True
Bad	Good
Non-sense	Sense
Anti-A	A

0 is the absolute falsity, 1 the absolute truth. In between each opposite pair, normally in a vicinity of 0.5, are being set up the neutralities.

There exist as many states in between "true" and "false" as in between "good" and "bad". Irrational and transcendental standpoints belong to this interval.

Even if an act apparently looks to be only good, or only bad, the other headed side should be sought. The following ratios vary indefinitely

$$\frac{\text{Anti-A}}{A}, \frac{\text{Non-A}}{A}.$$

They are transfinite.

If a statement is 30 percent T (true) and 60 percent I (indeterminate), then it is 10 percent F (false). This is somehow alethic, meaning pertaining to

truthhood and falsehood in the same time. In opposition to fuzzy logic, if a statement is 30 percent T doesn't involve it is 70 percent F . We have to study its indeterminacy as well.

(B) *Definition of neutrosophic logic.* This is a generalization (for the case of null indeterminacy) of the fuzzy logic. Neutrosophic logic is useful in the real-world systems for designing control logic, and may work in quantum mechanics.

If a proposition P is t percent true, doesn't necessarily mean it is $100 - t$ percent false as in fuzzy logic. There should also be a percentage of indeterminacy on the values of P . A better approach of the logical value of P is f percent false, i percent indeterminate, and t percent true, where $t + i + f = 100$ and $t, i, f \in [0, 100]$, called neutrosophic logical value of P , and noted by $n(P) = (t, i, f)$.

Neutrosophic logic means the study of neutrosophic logical values of the propositions. There exist, for each individual event, PRO parameters, CONTRA parameters, and NEUTER parameters which influence the above values. Indeterminacy results from any hazard which may occur, from unknown parameters, or from new arising conditions. This resulted from practice.

(C) *Applications.*

- (1) "The candidate C , who runs for election in a metropolis M of p people with right to vote, will win". This proposition is, say, 25 percent true (percentage of people voting for him), 35 percent false (percentage of people voting against him), and 40 percent indeterminate (percentage of people not coming to the ballot box, or giving a blank vote – not selecting anyone, or giving a negative vote – cutting all candidates on the list).
- (2) "Tomorrow it will rain". This proposition is, say, 50 percent true according to meteorologists who have investigated the past years' weather, 30 percent false according to today's very sunny and droughty summer, and 20 percent undecided.
- (3) "This is a heap". As an application to the sorites paradoxes, we may now say this proposition is t percent true, f percent false, and i percent indeterminate (the neutrality comes for we don't know exactly where is the difference between a heap and a non-heap; and, if we approximate the border, our "accuracy" is subjective).

We are not able to distinguish the difference between yellow and red as well if a continuum spectrum of colors is painted on a wall imperceptibly changing from one into another.

(D) *Definition of neutrosophic logical connectors.* One uses the definitions of neutrosophic probability and neutrosophic set. Let $0 \leq t_1, t_2 \leq 1$ represent the truth-probabilities, $0 \leq i_1, i_2 \leq 1$ the indeterminacy-probabilities, and $0 \leq f_1, f_2 \leq 1$ the falsity-probabilities of two events P_1 and P_2 respectively, where $t_1 + i_1 + f_1 = 1$ and $t_2 + i_2 + f_2 = 1$. One notes the neutrosophic logical

values of P_1 and P_2 by:

$$n(P_1) = (t_1, i_1, f_1) \text{ and } n(P_2) = (t_2, i_2, f_2).$$

Also, for any $0 \leq x \leq 1$ one notes $1 - x = \bar{x}$. Let $W(a, b, c) = (1 - a)/(b + c)$ and $W(R) = W(R(t), R(i), R(f))$ for any tridimensional vector $R = (R(t), R(i), R(f))$.

Negation. Let $N(x) = 1 - x = \bar{x}$. Then:

$$n(\neg P_1) = (N(t_1), N(i_1)W(N), N(f_1)W(N)).$$

Conjunction. Let $C(x, y) = xy$, and $C(z_1, z_2) = C(z)$ for any bidimensional vector $z = (z_1, z_2)$. Then:

$$n(P_1 \wedge P_2) = (C(t), C(i)W(C), C(f)W(C)).$$

(And, in a similar way, generalized for n propositions.)

Weak or inclusive disjunction. Let $D1(x, y) = x + y - xy = x + \bar{x}y = y + x\bar{y}$, and $D1(z_1, z_2) = D1(z)$ for any bidimensional vector $z = (z_1, z_2)$. Then:

$$n(P_1 \vee P_2) = (D1(t), D1(i)W(D1), D1(f)W(D1)).$$

(And, in a similar way, generalized for n propositions.)

Strong or exclusive disjunction. Let $D2(x, y) = x(1 - y) + y(1 - x) - xy(1 - x)(1 - y) = \bar{x}y + \bar{y}x - \bar{x}\bar{y}xy$, and $D2(z_1, z_2) = D2(z)$ for any bidimensional vector $z = (z_1, z_2)$. Then:

$$n(P_1 \underline{\vee} P_2) = (D2(t), D2(i)W(D2), D2(f)W(D2)).$$

(And, in a similar way, generalized for n propositions.)

Material conditional (implication). Let $I(x, y) = 1 - x + xy = \bar{x} + xy = 1 - \bar{x}\bar{y}$, and $I(z_1, z_2) = I(z)$ for any bidimensional vector $z = (z_1, z_2)$. Then:

$$n(P_1 \rightarrow P_2) = (I(t), I(i)W(I), I(f)W(I)).$$

Material biconditional (equivalence). Let $E(x, y) = (1 - x + xy)(1 - y + xy) = (\bar{x} + xy)(\bar{y} + xy) = (1 - \bar{x}\bar{y})(1 - \bar{y}\bar{x})$, and $E(z_1, z_2) = E(z)$ for any bidimensional vector $z = (z_1, z_2)$:

$$n(P \leftrightarrow Q) = (E(t), E(i)W(E), E(f)W(E)).$$

Sheffer's connector. Let $S(x, y) = 1 - xy$, and $S(z_1, z_2) = S(z)$ for any bidimensional vector $z = (z_1, z_2)$:

$$n(P | Q) = n(\neg P \vee \neg Q) = (S(t), S(i)W(S), S(f)W(S)).$$

Peirce's connector. Let $P(x, y) = (1 - x)(1 - y) = \bar{x}\bar{y}$, and $P(z_1, z_2) = P(z)$ for any bidimensional vector $z = (z_1, z_2)$.

$$n(P \downarrow Q) = n(\neg P \wedge \neg Q) = (P(t), P(i)W(P), P(f)W(P)).$$

(E) *Properties of neutrosophic logical connectors.*

Let's note by $t(P)$ the truth-component of the neutrosophic value $n(P)$, and $t(P) = p, t(Q) = q$.

On Rugina's
system of
thought

(a) *Conjunction.* $t(P \wedge Q) \leq \min\{p, q\}$:

$$\bigwedge_{k=1}^{\infty} (t(P) = 0 \text{ if } t(P) \neq 1).$$

(b) *Weak disjunction.* $t(P \vee Q) \geq \max\{p, q\}$:

$$\bigvee_{k=1}^{\infty} (t(P) = 1 \text{ if } t(P) \neq 0).$$

(c) *Implication.* $t(P \rightarrow Q) = 1$ if $t(P) = 0$ or 1 , and $> p$ otherwise:

$$\lim_{t(P) \rightarrow 0} t(P \rightarrow Q) = 1$$

$$\lim_{t(Q) \rightarrow 1} t(P \rightarrow Q) = 1$$

$$\lim_{t(P) \rightarrow 1} t(P \rightarrow Q) = q$$

$$\lim_{t(Q) \rightarrow 0} t(P \rightarrow Q) = 1 - p$$

(d) *Equivalence.* $t(P \leftrightarrow Q) = t(Q \leftrightarrow P) = t(\neg P \leftrightarrow \neg Q)$:

$$\lim_{\substack{t(P) \rightarrow 0 \\ t(Q) \rightarrow 0}} t(P \leftrightarrow Q) = 1$$

$$\lim_{\substack{t(P) \rightarrow 1 \\ t(Q) \rightarrow 1}} t(P \leftrightarrow Q) = 1$$

$$\lim_{\substack{t(P) \rightarrow 0 \\ t(Q) \rightarrow 1}} t(P \leftrightarrow Q) = 0$$

$$\lim_{\substack{t(P) \rightarrow 1 \\ t(Q) \rightarrow 0}} t(P \leftrightarrow Q) = 0$$

$$\lim_{t(P) \rightarrow 0} t(P \leftrightarrow Q) = 1 - q$$

$$\lim_{t(P) \rightarrow 1} t(P \leftrightarrow Q) = q$$

Let $q \neq 0, 1$ be constant, and one notes $p_{\max}(q) = (q^2 - 3q + 1)/(2q^2 - 2q)$. Then $\max t(P \leftrightarrow Q)$ occurs when $0 \leq t(P) \leq 1$. $p = p_{\max}(q)$ if $p_{\max}(q) \in [0, 1]$,

or $p = 0$ if $p_{\max}(q) < 0$, or $p = 1$ if $p_{\max}(q) > 1$, because the equivalence connector is described by a parabola of equation:

$$e_q(p) = (q^2 - q)p^2 + (-q^2 + 3q - 1)p + (1 - q).$$

This equation is concave down.

(5) *Neutrosophic topology*

(A) *Definition.* Let's construct a neutrosophic topology on $NT = [0, 1]$, considering the associated family of subsets $(0, p)$, for $0 \leq p \leq 1$, the whole set $[0, 1]$, and the empty set $\phi = (0, 0)$, called open sets, which is closed under set union and finite intersection. The union is defined as $(0, p) \cup (0, q) = (0, d)$, where $d = p + q - pq$, and the intersection as $(0, p) \cap (0, q) = (0, c)$, where $c = pq$. The complementary of $(0, p)$ is $(0, n)$, where $n = 1 - p$, which is a closed set.

(B) *Neutrosophic topological space.* The interval NT, endowed with this topology, forms a neutrosophic topological space.

(C) *Isomorphism.* Neutrosophic logical space, neutrosophic topological space, and neutrosophic probability space are all isomorphic.

A method of neutrosophy is described below.

(6) *Transdisciplinarity*

(A) *Introduction.* Transdisciplinarity means to find common features to uncommon entities: $\langle A \rangle \cap \langle \text{Non-A} \rangle \neq \phi$, even if they are disjunct.

(B) *Multi-structure and multi-space.* I consider that life and practice do not deal with "pure" spaces, but with a group of many spaces, with a mixture of structures, a "mongrel", a heterogeneity – the ardently preoccupation is to reunite them, to constitute a multi-structure.

I thought to a multi-space also: fragments (potsherds) of spaces put together, say as an example: Banach, Hausdorff, Tikhonov, compact, paracompact, Fock symmetric, Fock antisymmetric, path-connected, simply connected, discrete metric, indiscrete pseudo-metric, etc. spaces that work together as a whole mechanism. The difficulty is to be the passage over "frontiers" (borders between two disjoint spaces); i.e. how can we organically tie a point $P1$ from a space $S1$ with a point $P2$ from a structurally opposite space $S2$?

Does the problem become more complicated when the spaces' sets are not disjoint?

Let S_1 and S_2 be two distinct structures, induced by the group of laws L which verify the axiom groups A_1 and A_2 respectively, such that A_1 is strictly included in A_2 .

One says that the set M , endowed with the properties is called an S_1 -structure with respect to the S_2 -structure:

- M has an S_1 -structure;
- there is a proper subset P (different from the empty set, from the unitary element, and from M) of the initial set M which has an S_2 -structure; and
- M doesn't have an S_2 -structure.

Let S_1, S_2, \dots, S_k be distinct space-structures. We define the multi-space (or k -structured-space) as a set M such that for each structure $S_i, 1 \leq i \leq k$, there is a proper (different from ϕ and from M) subset M_i of it which has that structure. The M_1, M_2, \dots, M_k proper subsets are different two by two.

Let's introduce new terms.

(C) *Psychomathematics*. A discipline which studies psychological processes in connection with mathematics.

(D) *Mathematical modeling of psychological processes*. Weber's law and Fechner's law on sensations and stimuli are improved.

(E) *Psychoneutrosophy*. Psychology of neutral thought, action, behavior, sensation, perception, etc. This is a hybrid field deriving from theology, philosophy, economics, psychology, etc. For example, to find the psychological causes and effects of individuals supporting neutral ideologies (neither capitalists, nor communists), politics (not in the left, not in the right), etc.

(F) *Socioneutrosophy*. Sociology of neutralities. For example, the sociological phenomena and reasons which determine a country or group of people or class to remain neutral in a military, political, ideological, cultural, artistic, scientific, economical, etc. international or internal war (dispute).

(G) *Econoneutrosophy*. Economics of non-profit organizations, groups, such as: churches, philanthropic associations, charities, emigrating foundations, artistic or scientific societies, etc. How they function, how they survive, who benefits and who loses, why are they necessary, how they improve, how they interact with for-profit companies.

These terms are in the process of development.

(7) *Rugina's orientation table*

In order to clarify the anomalies in science, Rugina (1989, 1998) proposes an original method, starting first from an economic point of view but generalizing it to any science, to study the equilibrium and disequilibrium of systems. His table comprises seven basic models:

- (1) Model M_1 (which is 100 percent stable)
- (2) Model M_2 (which is 95 percent stable, and 5 percent unstable);
- (3) Model M_3 (which is 65 percent stable, and 35 percent unstable);
- (4) Model M_4 (which is 50 percent stable, and 50 percent unstable);
- (5) Model M_5 (which is 35 percent stable, and 65 percent unstable);
- (6) Model M_6 (which is 5 percent stable, and 95 percent unstable); and
- (7) Model M_7 (which is 100 percent unstable)

He gives orientation tables for physical sciences and mechanics (Rugina, 1989, p. 18), for the theory of probability, for logic, and generally for any natural or social science (Rugina, 1989, pp. 286-88):

An anomaly can be simply defined as a deviation from a position of stable equilibrium represented by Model M_1 (Rugina, 1989, p. 17).

Rugina proposes the universal hypothesis of duality:

The physical universe in which we are living, including human society and the world of ideas, all are composed in different and changeable proportions of stable (equilibrium) and unstable (disequilibrium) elements, forces, institutions, behavior and value.

He also proposes the general possibility theorem:

...there is an unlimited number of possible combinations or systems in logic and other sciences.

According to the last assertions one can extend Rugina's orientation table in the way that any system in each science is s percent stable and u percent unstable, with $s + u = 100$ and both parameters $0 \leq s, u \leq 100$, somehow getting to a fuzzy approach.

But, because each system has hidden features and behaviors, and there would always be unexpected occurring conditions we are not able to control – we mean the indeterminacy plays a role as well, a better approach would be the neutrosophic model:

Any system in each science is s percent stable, i percent indeterminate, and u percent unstable, with $s + i + u = 100$ and all three parameters $0 \leq s, i, u \leq 100$.

Examples of Rugina's orientation table are given in Appendices 1-5.

References and further reading

- Bonola, R. (1938), *Non-Euclidean Geometry: a Critical and Historical Study of its Development*, Dover, New York, NY.
- Chimienti, S.P. and Bencze, M. (1998), "Smarandache paradoxist geometry", *Bulletin of Pure and Applied Sciences*, Vol. 17E No. 1, pp. 123-4.
- Chimienti, S.P. and Bencze, M. (1998), "Smarandache non-geometry", *Bulletin of Pure and Applied Sciences*, Vol. 17E No. 1, pp. 115-16.
- Chimienti, S.P. and Bencze, M. (1998), "Smarandache anti-geometry", *Bulletin of Pure and Applied Sciences*, Vol. 17E No. 1, pp. 103-14.
- Chimienti, S.P. and Bencze, M. (1998), "Smarandache counter-projective geometry", *Bulletin of Pure and Applied Sciences*, Vol. 17E No. 1, pp. 117-8.
- Hilbert, D. (1950), *Foundations of Geometry*, E.J. Townsend (trans), Kegan Paul & Co., London.
- McNeil, F.M. and Thro, E. (1994), *Fuzzy Logic/A Practical Approach*, Academic Press, Orlando, FL.
- Rugina, A.N. (1984), "The problem of values and value-judgements in science and a positive solution", *International Journal of Sociology and Social Policy*, Vol. 4 No. 3.
- Rugina, A.N. (1989), *Principia Methologica 1: A Bridge from Economics to all Other Natural Sciences / Towards a Methodological Unification of all Sciences*, MCB University Press, Bradford.
- Rugina, A.N. (1994), "A long journey to the third revolution in economic thinking", *International Journal of New Ideas*, Vol. 3 No. 1/2.
- Rugina, A.N. (1998), "Prolegomena to any future study in economics, finance and other social sciences: the road to a third revolution in economic, financial, social, ethical, logical and political thinking", *International Journal of Social Economics*, Vol. 25 No. 5.
- Smarandache, F. (1997), *Collected Papers*, Vol. 2, University of Kishinev, Kishinev.
- Webster's New World Dictionary of American English* (1998), 3rd College ed. (edited by Neufeldt, V. and Guralnik, D.B.), Simon & Schuster, New York, NY.

Appendix 1. Example of model M_3 in Rugina's orientation table

The paradoxist geometry (actually the percentage of instability is between 20-35)

In 1969, intrigued by geometry, I simultaneously constructed a partially Euclidean and partially non-Euclidean space by a strange replacement of the Euclid's fifth postulate (axiom of parallels) with the following five-statement proposition:

- (1) there are at least a straight line and a point exterior to it in this space for which only one line passes through the point and does not intersect the initial line (1 parallel);
- (2) there are at least a straight line and a point exterior to it in this space for which only a finite number of lines $l_1, \dots, l_k (k \geq 2)$ pass through the point and do not intersect the initial line (two or more (in a finite number) parallels);
- (3) there are at least a straight line and a point exterior to it in this space for which any line that passes through the point intersects the initial line (0 parallels);
- (4) there are at least a straight line and a point exterior to it in this space for which an infinite number of lines that pass through the point (but not all of them) do not intersect the initial line (an infinite number of parallels, but not all lines passing through); and
- (5) there are at least a straight line and a point exterior to it in this space for which any line that passes through the point does not intersect the initial line (an infinite number of parallels, all lines passing through the point).

This geometry unites all together: Euclid, Lobachevsky/Bolyai, and Riemann geometries. And separates them as well!

Appendix 2. First example of model M_7 in Rugina's orientation table

The non-geometry (the percentage of instability is 100)

It's a lot easier to deny the Euclid's five postulates than Hilbert's 20 thorough axioms:

- (1) It is not always possible to draw a line from an arbitrary point to another arbitrary point. For example this axiom can be denied only if the model's space has at least a discontinuity point (in our model bellow, MD, one takes an isolated point I in between f_1 and f_2 , the only one which will not verify the axiom).
- (2) It is not always possible to extend by continuity a finite line to an infinite line. For example, consider the model bellow, and the segment AB , where both A and B lie on f_1 , A in between P and N , while B on the left side of N ; one can not at all extend AB either beyond A or beyond B , because the resulted curve, noted say $A'-A-B-B'$, would not be a geodesic (i.e. line in our Model) anymore.
If A and B lie in δ_1-f_1 , both of them closer to f_1 , A in the left side of P , while B in the right side of P , then the segment AB , which is in fact $A-P-B$, can be extended beyond A and also beyond B only up to f_1 (therefore one gets a finite line too, $A'-A-P-B'$, where A', B' are the intersections of PA, PB respectively with f_1). If A, B lie in δ_1-f_1 , far enough from f_1 and P , such that AB is parallel to f_1 , then AB verifies this postulate.
- (3) It is not always possible to draw a circle from an arbitrary point and of an arbitrary interval. For example, same as for the first axiom, the isolated point I , and a very small interval not reaching f_1 neither f_2 , will deny this axiom.
- (4) Not all the right angles are congruent. (See example of the anti-geometry, explained below.)
- (5) If a line, cutting two other lines, forms the interior angles of the same side of it strictly less than two right angles, then not always the two lines extended towards infinite cut each other in the side where the angles are strictly less than two right angles. For example, let h_1, h_2 and l be three lines in $\delta_1-\delta_2$, where h_1 intersects f_1 in A , and h_2 intersects f_1 in B , with A, B, P different from each other, such that h_1 and h_2 do not intersect, but l cuts h_1 and h_2 and forms the interior angles of one of its sides (towards

f_1) strictly less than two right angles; the assumption of the fifth postulate is fulfilled, but the consequence does not hold, because h_1 and h_2 do not cut each other (they may not be extended beyond A and B respectively, because the lines would not be geodesics anymore).

Appendix 3. Second example of model M_7 in Rugina's orientation table

The counter-projective geometry (the percentage of instability is 100)

Let P, L be two sets, and r a relation included in $P \times L$. The elements of P are called points, and those of L lines. When (p, l) belongs to r , we say that the line l contains the point p . For these, one imposes the following counter-axioms:

- There exist: either at least two lines, or no line, that contains two given distinct points.
- Let p_1, p_2, p_3 be three non-collinear points, and q_1, q_2 two distinct points. Suppose that $\{p_1, q_1, p_3\}$ and $\{p_2, q_2, p_3\}$ are collinear triples. Then the line containing p_1, p_2 , and the line containing q_1, q_2 do not intersect.
- Every line contains at most two distinct points.

Does the duality principle hold in a counter-projective space? What about Desargues's theorem, fundamental theorem of projective geometry/theorem of Pappus, and Staudt algebra? Or Pascal's theorem, Brianchon's theorem? (I think none of them will hold!) However, Rugina's hypothesis of duality does hold (although the this geometry is formed by unstable elements only!).

Appendix 4. Third example of model M_7 in Rugina's orientation table

The anti-geometry (the percentage of instability is 100 – even... more, this is the geometry of total chaos!)

It is possible to entirely de-formalize Hilbert's groups of axioms of the Euclidean geometry, and to construct a model such that none of his fixed axioms holds.

Let's consider the following things:

- a set of (points): A, B, C, \dots ;
- a set of (lines): h, k, l, \dots ; and
- a set of (planes): $\alpha, \beta, \gamma, \dots$

Let us also consider a set of relationships among these elements: "are situated", "between", "parallel", "congruent", "continuous", etc.

Then, we can deny all Hilbert's 20 axioms (see Hilbert, 1950; Binola, 1938).

There exist cases, within a geometric model, when the same axiom is verified by certain points/lines/planes and denied by others.

Group I. Anti-axioms connection

I.1. Two distinct points A and B do not always completely determine a line.

Let's consider the following model MD: get an ordinary plane δ , but with an infinite hole as shown in Figure A1.

Plane δ is a reunion of two disjoint planar semi-planes; f_1 lies in MD, but f_2 does not; P, Q are two extreme points on f that belong to MD.

One defines a LINE l as a geodesic curve: if two points A, B that belong to MD lie in l , then the shortest curve lying in MD between A and B lies in l also. If a line passes twice through the same point, then it is called double point (KNOT).

One defines a PLANE α as a surface such that for any two points A, B that lie in α and belong to MD there is a geodesic which passes through A, B and lies in α also.

Now, let's have two strings of the same length: one ties P and Q with the first string s_1 such that the curve s_1 is folded in two or more different planes and s_1 is under the plane




Figure A1.
A model of connection

delta; next, do the same with string s_2 , tie Q with P , but over the plane delta and such that s_2 has a different form from s_1 ; and a third string s_3 , from P to Q , much longer than s_1 . s_1, s_2, s_3 belong to MD.

Let I, J, K be three isolated points – as some islands, i.e. not joined with any other point of MD, exterior to the plane delta. This model has a measure, because the (pseudo-)line is the shortest way (length) to go from a point to another (when possible). Of course, this model is not perfect, and is far from the best. Readers are asked to improve it, or to make up a new one that is better. (Let A, B be two distinct points in delta1-f1 . P and Q are two points on s_1 , but they do not completely determine a line, referring to the first axiom of Hilbert, because $A-P-s_1-Q$ are different from $B-P-s_1-Q$.)

- I.2. There is at least a line l and at least two distinct points A and B of l , such that A and B do not completely determine the line l . (Line $A-P-s_1-Q$ are not completely determined by P and Q in the previous construction, because $CVB-P-s_1-Q$ is another line passing through P and Q too.)
- I.3. Three points A, B, C not situated in the same line do not always completely determine a plane alpha. (Let A, B be two distinct points in delta1-f1 , such that A, B, P are not co-linear. There are many planes containing these three points: delta1 extended with any surface s containing s_1 , but not cutting s_2 in between P and Q , for example.)
- I.4. There is at least a plane, alpha, and at least three points A, B, C in it not lying in the same line, such that A, B, C do not completely determine the plane alpha. (See the previous example.)
- I.5. If two points A, B of a line l lie in a plane alpha, it doesn't mean that every point of l lies in alpha. (Let A be a point in delta1-f1 , and B another point on s_1 in between P and Q . Let alpha be the following plane: delta1 extended with a surface s containing s_1 , but not cutting s_2 in between P and Q , and tangent to delta2 on a line QC , where C is a point in delta2-f2 . Let D be point in delta2-f2 , not lying on the line QC . Now, A, B, D are lying on the same line $A-P-s_1-Q-D$, A, B are in the plane alpha, but D is not.)

- I.6. If two planes α , β have a point A in common, it doesn't mean they have at least a second point in common. (Construct the following plane α : a closed surface containing s_1 and s_2 , and intersecting δ_1 in one point only, P . Then α and δ_1 have a single point in common.)
- I.7. There exist lines where lie only one point, or planes where lie only two points, or space where lie only three points. (Hilbert's I.7 axiom may be contradicted if the model has discontinuities. Let's consider the isolated points area. The point I may be regarded as a line, because it's not possible to add any new point to I to form a line. One constructs a surface that intersects the model only in the points I and J .)

Group II. Anti-axioms of order

- II.1. If A, B, C are points of a line and B lies between A and C , it doesn't mean that always B lies also between C and A .

[Let T lie in s_1 , and V lie in s_2 , both of them closer to Q , but different from it. Then: P, T, V are points on the line $P-s_1-Q-s_2-P$ (i.e. the closed curve that starts from the point P and lies in s_1 and passes through the point Q and lies back to s_2 and ends in P), and T lies between P and V – because PT and TV are both geodesics – but T doesn't lie between V and P because from V the line goes to P and then to T , therefore P lies between V and T .]

[By definition: a segment AB is a system of points lying upon a line between A and B (the extremes are included). Warning, AB may be different from BA ; for example: the segment PQ formed by the system of points starting with P , ending with Q , and lying in s_1 , is different from the segment QP formed by the system of points starting with Q , ending with P , but belonging to s_2 . Worse, AB may be sometimes different from AB ; for example, the segment PQ formed by the system of points starting with P , ending with Q , and lying in s_1 , is different from the segment PQ formed by the system of points starting with P , ending with Q , but belonging to s_2 .]

- II.2. If A and C are two points of a line, then: there does not always exist a point B lying between A and C , or there does not always exist a point D such that C lies between A and D .

[For example, let F be a point on f_1 , F different from P , and G a point in δ_1 , G doesn't belong to f_1 ; draw the line l which passes through G and F ; then there exists a point B lying between G and F – because GF is an obvious segment – but there is no point D such that F lies between G and D – because GF is right bounded in F (GF may not be extended to the other side of F , because otherwise the line will not remain a geodesic anymore).]

- II.3. There exist at least three points situated on a line such that one point lies between the other two, and another point lies also between the other two.

[For example, let R, T be two distinct points, different from P and Q , situated on the line $P-s_1-Q-s_2-P$, such that the lengths PR, RT, TP are all equal; then R lies between P and T , and T lies between R and P ; also P lies between T and R .]

- II.4. Four points A, B, C, D of a line can not always be arranged such that B lies between A and C and also between A and D , and such that C lies between A and D and also between B and D .

[For example:

- let R, T be two distinct points, different from P and Q , situated on the line $P-s_1-Q-s_2-P$ such that the lengths PR, RQ, QT, TP are all equal, therefore R belongs to s_1 , and T belongs to s_2 ; then P, R, Q, T are situated on the same line: such that R lies between P and Q , but not between P and T – because the geodesic PT does not pass through R – and such that Q does not lie between P and T – because the geodesic PT does not pass through Q – but lies between R and T .
- Let A, B be two points in δ_2-f_2 such that A, Q, B are colinear, and C, D two points on s_1, s_2 respectively, all of the four points being different from P and Q ; then A, B, C, D are points situated on the same line $A-Q-s_1-P-s_2-Q-B$, which is the same with line $A-Q-$

s_2 - P - s_1 - Q - B , therefore we may have two different orders of these four points in the same time: A, C, D, B and A, D, C, B .]

- II.5. Let A, B, C be three points not lying in the same line, and l a line lying in the same plane ABC and not passing through any of the points A, B, C . Then, if the line l passes through a point of the segment AB , it doesn't mean that always the line l will pass through either a point of the segment BC or a point of the segment AC .
[For example, let AB be a segment passing through P in the semi-plane δ_1 , and C a point lying in δ_1 too on the left side of the line AB ; thus A, B, C do not lie on the same line; now, consider the line Q - s_2 - P - s_1 - Q - D , where D is a point lying in the semi-plane δ_2 not on f_2 : therefore this line passes through the point P of the segment AB , but does not pass through any point of the segment BC , nor through any point of the segment AC .]

Group III. Anti-axiom of parallels

In a plane α there can be drawn through a point A , lying outside of a line l , either no line, or only one line, or a finite number of lines, or an infinite number of lines which do not intersect the line l . (At least two of these situations should occur.) The line(s) is (are) called the parallel(s) to l through the given point A .

[For example:

- Let l_0 be the line N - P - s_1 - Q - R , where N is a point lying in δ_1 not on f_1 , and R is a similar point lying in δ_2 not on f_2 , and let A be a point lying on s_2 , then: no parallel to l_0 can be drawn through A (because any line passing through A , hence through s_2 , will intersect s_1 , hence l_0 , in P and Q).
- If the line l_1 lies in δ_1 such that l_1 does not intersect the frontier f_1 , then through any point lying on the left side of l_1 one and only one parallel will pass.
- Let B be a point lying in f_1 , different from P , and another point C lying in δ_1 , not on f_1 ; let A be a point lying in δ_1 outside of BC ; then: an infinite number of parallels to the line BC can be drawn through the point A .

Theorem. There are at least two lines l_1, l_2 of a plane, which do not meet a third line l_3 of the same plane, but they meet each other, (i.e. if l_1 is parallel to l_3 , and l_2 is parallel to l_3 , and all of them are in the same plane, it's not necessary that l_1 is parallel to l_2).

[For example: consider three points A, B, C lying in f_1 , and different from P , and D a point in δ_1 not on f_1 ; draw the lines AD, BE and CE such that E is a point in δ_1 not on f_1 and both BE and CE do not intersect AD ; then: BE is parallel to AD , CE is also parallel to AD , but BE is not parallel to CE because the point E belongs to both of them.]

Group IV. Anti-axioms of congruence

IV.1. If A, B are two points on a line l , and A' is a point upon the same or another line l' , then upon a given side of A' on the line l' , we can not always find only one point B' so that the segment AB is congruent to the segment $A'B'$.

[For example:

- Let AB be segment lying in δ_1 and having no point in common with f_1 , and construct the line C - P - s_1 - Q - s_2 - P (noted by l') which is the same with C - P - s_2 - Q - s_1 - P , where C is a point lying in δ_1 not on f_1 nor on AB ; take a point A' on l' , in between C and P , such that $A'P$ is smaller than AB ; now, there exist two distinct points B_1' on s_1 and B_2' on s_2 , such that $A'B_1'$ is congruent to AB and $A'B_2'$ is congruent to AB , with $A'B_1'$ different from $A'B_2'$.
- But if we consider a line l' lying in δ_1 and limited by the frontier f_1 on the right side (the limit point being noted by M), and take a point A' on l' , close to M , such that $A'M$ is less than $A'B'$, then there is no point B' on the right side of l' so that $A'B'$ is congruent to AB .]

A segment may not be congruent to itself!

[For example, let A be a point on s_1 , closer to P , and B a point on s_2 , closer to P also; A and B are lying on the same line $A-Q-B-P-A$ which is the same with line $A-P-B-Q-A$, but AB measured on the first representation of the line is strictly greater than AB measured on the second representation of their line.]

- IV.2. If a segment AB is congruent to the segment $A'B'$ and also to the segment $A''B''$, then not always the segment $A'B'$ is congruent to the segment $A''B''$.

[For example, let AB be a segment lying in δ_1 , and consider the line $C-P-s_1-Q-s_2-P-D$, where C, D are two distinct points in δ_1 such that C, P, D are colinear. Suppose that the segment AB is congruent to the segment CD (i.e. $C-P-s_1-Q-s_2-P-D$). Get also an obvious segment $A'B'$ in δ_1 , different from the preceding ones, but congruent to AB . Then the segment $A'B'$ is not congruent to the segment CD (considered as $C-P-D$, i.e. not passing through Q).

- IV.3. If AB, BC are two segments of the same line l which have no points in common aside from the point B , and $A'B', B'C'$ are two segments of the same line or of another line l' having no point other than B' in common, such that AB is congruent to $A'B'$ and BC is congruent to $B'C'$, then not always the segment AC is congruent to $A'C'$.

[For example, let l be a line lying in δ_1 , not on f_1 , and A, B, C three distinct points on l , such that AC is greater than s_1 ; let l' be the following line: $A'-P-s_1-Q-s_2-P$ where A' lies in δ_1 , not on f_1 , and get B' on s_1 such that $A'B'$ is congruent to AB , get C' on s_2 such that BC is congruent to $B'C'$ (the points A, B, C are thus chosen); then the segment $A'C'$ which is first seen as $A'-P-B'-Q-C'$ is not congruent to AC , because $A'C'$ is the geodesic $A'-P-C'$ (the shortest way from A' to C' does not pass through B') which is strictly less than AC .]

Definitions. Let h, k be two lines having a point O in common. Then the system (h, O, k) is called the angle of the lines h and k in the point O . (Because some of our lines are curves, we take the angle of the tangents to the curves in their common point.)

The angle formed by the lines h and k situated in the same plane, noted by $\angle(h, k)$, is equal to the arithmetic mean of the angles formed by h and k in all their common points.

- IV.4. Let an angle (h, k) be given in the plane α , and let a line h' be given in the plane β . Suppose that in the plane β a definite side of the line h' be assigned, and a point O' . Then in the plane β there are one, or more, or even no half-line(s) k' emanating from the point O' such that the angle (h, k) is congruent to the angle (h', k') , and at the same time the interior points of the angle (h', k') lie upon one or both sides of h' .

For example:

- Let A be a point in δ_1 , and B, C two distinct points in δ_2 ; let h be the line $A-P-s_1-Q-B$, and k be the line $A-P-s_2-Q-C$; because h and k intersect in an infinite number of points (the segment AP), where they normally coincide – i.e. in each such point their angle is congruent to zero, the angle (h, k) is congruent to zero. Now, let A' be a point in δ_1 , different from A , and B' a point in δ_2 , different from B , and draw the line h' as $A'-P-s_1-Q-B'$; there exist an infinite number of lines k' , of the form $A'-P-s_2-Q-C'$ (where C' is any point in δ_2 , not on the line QB'), such that the angle (h, k) is congruent to (h', k') , because (h', k') is also congruent to zero, and the line $A'-P-s_2-Q-C'$ is different from the line $A'-P-s_2-Q-D'$ if D' is not on the line QC' .
- If h, k , and h' are three lines in δ_1 , which intersect the frontier f_1 in at most one point, then there exists only one line k' on a given part of h' such that the angle (h, k) is congruent to the angle (h', k') .
- Is there any case when, with these hypotheses, no k' exists?
- Not every angle is congruent to itself; for example, $\angle(s_1, s_2)$ is not congruent to $\angle(s_1, s_2)$ [because one can construct two distinct lines: $P-s_1-Q-A$ and $P-s_2-Q-A$, where A is a

point in δ_2 , for the first angle, which becomes equal to zero; and $P-s_1-Q-A$ and $P-s_2-Q-B$, where B is another point in δ_2 , B different from A , for the second angle, which becomes strictly greater than zero!].

- IV. 5. If the angle (h, k) is congruent to the angle (h', k') and the angle (h'', k'') , then the angle (h', k') is not always congruent to the angle (h'', k'') . (A similar construction to the previous one.)
- IV. 6. Let ABC and $A'B'C'$ be two triangles such that AB is congruent to $A'B'$, AC is congruent to $A'C'$, $\sphericalangle BAC$ is congruent to $\sphericalangle B'A'C'$. Then not always $\sphericalangle ABC$ is congruent to $\sphericalangle A'B'C'$ and $\sphericalangle ACB$ is congruent to $\sphericalangle A'C'B'$.

[For example, Let M, N be two distinct points in δ_2 , thus obtaining the triangle PMN ; Now take three points R, M', N' in δ_1 , such that RM' is congruent to PM , RN' is congruent to PN , and the angle (RM', RN') is congruent to the angle (PM, PN) . $RM'N'$ is an obvious triangle. Of course, the two triangles are not congruent, because for example PM and PN cut each other twice – in P and Q – while RM' and RN' only once – in R . (These are geodesical triangles.)]

Definitions. Two angles are called supplementary if they have the same vertex, one side in common, and the other sides not common form a line. A right angle is an angle congruent to its supplementary angle. Two triangles are congruent if its angles are congruent two by two, and its sides are congruent two by two.

Propositions. A right angle is not always congruent to another right angle.

For example: Let $A-P-s_1-Q$ be a line, with A lying in δ_1 , and $B-P-s_1-Q$ another line, with B lying in δ_1 and B not lying in the line AP ; we consider the tangent t at s_1 in P , and B chosen in a way that $\sphericalangle(AP, t)$ is not congruent to $\sphericalangle(BP, t)$; let A', B' be other points lying in δ_1 such that $\sphericalangle APA'$ is congruent to $\sphericalangle A'P-s_1-Q$, and $\sphericalangle BPB'$ is congruent to $\sphericalangle B'P-s_1-Q$.

Then the angle APA' is right, because it is congruent to its supplementary (by construction), and the angle BPB' is also right, because it is congruent to its supplementary (by construction). But $\sphericalangle APA'$ is not congruent to $\sphericalangle BPB'$, because the first one is half of the angle $A-P-s_1-Q$, i.e. half of $\sphericalangle(AP, t)$, while the second one is half of the $B-P-s_1-Q$, i.e. half of $\sphericalangle(BP, t)$.

The theorems of congruence for triangles [side, side, and angle in between; angle, angle, and common side; side, side, side] may not hold either in the critical zone (s_1, s_2, f_1, f_2) of the model.

Property: The sum of the angles of a triangle can be:

- 180 degrees, if all its vertexes A, B, C are lying, for example, in δ_1 ;
- strictly less than 180 degrees (any value in the interval $(0, 180)$). For example, let R, T be two points in δ_2 such that Q does not lie in RT , and S another point on s_2 ; then the triangle SRT has $\sphericalangle(SR, ST)$ congruent to 0 because SR and ST have an infinite number of common points (the segment SQ), and $\sphericalangle QTR + \sphericalangle TRQ$ congruent to $180 - \sphericalangle TQR$ [by construction we may vary $\sphericalangle TQR$ in the interval $(0, 180)$], even 0 degrees! Let A be a point in δ_1 , B a point in δ_2 , and C a point on s_3 , very close to P , then ABC is a non-degenerate triangle (because its vertexes are non-collinear), but $\sphericalangle(A-P-s_1-Q-B, A-P-s_3-C) = \sphericalangle(B-Q-s_1-P-A, B-Q-s_1-P-s_3-C) = \sphericalangle(C-s_3-P-A, C-s_3-P-s_1-Q-B) = 0$ (one considers the length $C-s_3-P-s_1-Q-B$ strictly less than $C-s_3-B$); the area of this triangle is also 0!
- more than 180 degrees. For example; let A, B be two points in δ_1 , such that $\sphericalangle PAB + \sphericalangle PBA + \sphericalangle(s_1, s_2; \text{in } Q)$ is strictly greater than 180 degrees; then the triangle ABQ , formed by the intersection of the lines $A-P-s_2-Q, Q-s_1-P-B, AB$ will have the sum of its angles strictly greater than 180 degrees.

Definition. A circle of center M is a totality of all points A for which the segments MA are congruent to one another.

For example, if the center is Q , and the length of the segments MA is chosen greater than the length of s_1 , then the circle is formed by the arc of circle centered in Q , of radius MA , and lying in δ_2 , plus another arc of circle centered in P , of radius MA -length of s_1 , lying in δ_1 .

Group V. Anti-axiom of continuity (anti-Archimedean axiom)

Let A, B be two points. Take the points $A_1, A_2, A_3, A_4, \dots$ so that A_1 lies between A and A_2 , A_2 lies between A_1 and A_3 , A_3 lies between A_2 and A_4 , etc. and the segments $AA_1, A_1A_2, A_2A_3, A_3A_4, \dots$ are congruent to one another.

Then, among this series of points, not always there exists a certain point A_n such that B lies between A and A_n .

[For example, let A be a point in δ_1 , and B a point on f_1 , B different from P ; on the line AB consider the points $A_1, A_2, A_3, A_4, \dots$ in between A and B , such that $AA_1, A_1A_2, A_2A_3, A_3A_4$, etc. are congruent to one another; then we find that there is no point behind B (considering the direction from A to B), because B is a limit point (the line AB ends in B).]

Bolzano's (intermediate value) theorem may not hold in the critical zone of the model.

Appendix 5. Fourth example of model M_7 in Rugina's orientation table

The inconsistent system of axioms, and the contradictory theory (the percentage of instability is 100 – even... more, this is the system of chaos!)

Let $(a_1), (a_2), \dots, (a_n), (b)$ be $n + 1$ independent axioms, with $n \geq 1$; and let (b') be another axiom contradictory to (b) . We construct a system of $n + 2$ axioms:

$$[I] \quad (a_1), (a_2), \dots, (a_n), (b), (b'),$$

which is inconsistent. But this system may be shared into two consistent systems of independent axioms:

$$[C] \quad (a_1), (a_2), \dots, (a_n), (b),$$

and

$$[C'] \quad (a_1), (a_2), \dots, (a_n), (b').$$

We also consider the partial system of independent axioms:

$$[P] \quad (a_1), (a_2), \dots, (a_n).$$

Developing $[P]$, we find many propositions (theorems, lemmas):

$$(p_1), (p_2), \dots, (p_m),$$

by combinations of its axioms.

Developing $[C]$, we find all propositions of $[P]$:

$$(p_1), (p_2), \dots, (p_m),$$

resulting by combinations of $(a_1), (a_2), \dots, (a_n)$, plus other propositions:

$$(r_1), (r_2), \dots, (r_t),$$

resulting by combinations of (b) with any of $(a_1), (a_2), \dots, (a_n)$.

Similarly for $[C']$, we find the propositions of $[P]$:

$$(p_1), (p_2), \dots, (p_m),$$

plus other propositions

$$(r'_1), (r'_2), \dots, (r'_t),$$

resulting by combinations of (b') with any of $(a1), (a2), \dots, (an)$, where $(r'1)$ is an axiom contradictory to $(r1)$, and so on.

Now, developing $[I]$, we'll find all the previous resulted propositions:

$$\begin{aligned} &(p1), (p2), \dots, (pm), \\ &(r1), (r2), \dots, (rt), \\ &(r'1), (r'2), \dots, (r't). \end{aligned}$$

Therefore, $[I]$ is equivalent to $[C]$ reunited to $[C']$.

From one pair of contradictory propositions $\{(b) \text{ and } (b')\}$ in its beginning, $[I]$ adds t more such pairs, where $t \geq 1$, $\{(r1) \text{ and } (r'1), \dots, (rt) \text{ and } (r't)\}$, after a complete step. The further we go, the more pairs of contradictory propositions are accumulating in $[I]$.

It is interesting to study the case when $n = 0$.

Why do people avoid thinking about the contradictory theory?

As you know, nature is not perfect: opposite phenomena occur together, and opposite ideas are simultaneously asserted and, ironically, proved that both of them are true! How is that possible?

A statement may be true in a referential system, but false in another one. The truth is subjective. The proof is relative. (In philosophy there is a theory that "knowledge is relative to the mind, or things can be known only through their effects on the mind, and consequently there can be no knowledge of reality as it is in itself", called "the Relativity of Knowledge"; see *Webster's New World Dictionary of American English*, 1988, p. 1133)

You know? . . . sometimes is good to be wrong!