

10-12-2019

An Outranking Approach for MCDM-Problems with Neutrosophic Multi-Sets

Vakkas Uluçay

Adil Kılıç

Ismet Yıldız

Memet Şahin

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Uluçay, Vakkas; Adil Kılıç; Ismet Yıldız; and Memet Şahin. "An Outranking Approach for MCDM-Problems with Neutrosophic Multi-Sets." *Neutrosophic Sets and Systems* 30, 1 (). https://digitalrepository.unm.edu/nss_journal/vol30/iss1/17

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu, lsloane@salud.unm.edu, sarahrk@unm.edu.



An Outranking Approach for MCDM-Problems with Neutrosophic Multi-Sets

Vakkas Uluçay^{1*}, Adil Kılıç², İsmet Yıldız³ and Memet Şahin⁴

¹Kokluce neighborhood, Gaziantep, 27650, Turkey. E-mail: vulucay27@gmail.com

²Department of Mathematics, Gaziantep University, Gaziantep, 27310, Turkey. E-mail: adilkilic@gantep.edu.tr

³Department of Mathematics, Duzce University, Duzce, 81620, Turkey. E-mail: ismetyildiz@duzce.edu.tr

⁴Department of Mathematics, Gaziantep University, Gaziantep, 27310, Turkey. E-mail: mesahin@gantep.edu.tr

* Correspondence: Vakkas Uluçay (vulucay27@gmail.com)

Abstract: In this paper, we introduced a new outranking approach for multi-criteria decision making (MCDM) problems to handle uncertain situations in neutrosophic multi environment. Therefore, we give some outranking relations of neutrosophic multi sets. We also examined some desired properties of the outranking relations and developed a ranking method for MCDM problems. Moreover, we describe a numerical example to verify the practicality and effectiveness of the proposed method.

Keywords: Single valued neutrosophic sets, neutrosophic multi-sets, outranking relations, decision making.

1. Introduction

Fuzzy set theory, intuitionistic fuzzy set theory and neutrosophic set theory is introduced by Zadeh [59], Atanassov [1] and Smarandache [28] to handle the uncertain, incomplete, indeterminate and inconsistent information, respectively. The above set theories have been applied to many different areas including real decision making problems [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 18, 19, 21, 22, 23, 24, 25, 26, 27, 32, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 49, 58]. Also, several generalizations of the set theories made such as fuzzy multi-set theory [34, 35, 48], intuitionistic fuzzy multi-set theory [16, 31, 36, 37, 57] and n-valued refined neutrosophic set theory [29].

Another generalization of above theories that is relevant for our work is single valued neutrosophic refined (multi) set theory introduced by Ye [53, 56] which contain a few different values. A single valued neutrosophic multi set theory has truth-membership sequence $(\mu_{\mathcal{A}}^1(t), \mu_{\mathcal{A}}^2(t), \dots, \mu_{\mathcal{A}}^p(t))$, indeterminacy membership sequence $(\nu_{\mathcal{A}}^1(t), \nu_{\mathcal{A}}^2(t), \dots, \nu_{\mathcal{A}}^p(t))$ and falsity-membership sequence $(\omega_{\mathcal{A}}^1(t), \omega_{\mathcal{A}}^2(t), \dots, \omega_{\mathcal{A}}^p(t))$ of element $t \in T$. Recently, the single valued neutrosophic multi set theory have attracted widely attention in [20, 33, 50, 51, 52, 54, 55]. The paper is organized as follows; In Section 2 we give some basic notions of neutrosophic sets and neutrosophic multi-sets. In Section 3, we first introduce outranking relations of neutrosophic multi-sets with proprieties. In Section 4, we propose an outranking approach for to solving the multi-criteria decision making problems based on neutrosophic multi-set information. In Section 5, we propose a selection example to validate the practicality. Finally, in Section 6, we conclude the paper.

2. Preliminaries

In this section, we present the basic definitions and results of neutrosophic set theory [28, 33] and neutrosophic multi (or refined) set theory [12, 53] that are useful for subsequent discussions.

Definition 1 [28] let T be a universe. A neutrosophic set \mathcal{A} over T is defined by

$$\mathcal{A} = \left\{ \langle t, (\mu_{\mathcal{A}}(t), \nu_{\mathcal{A}}(t), \omega_{\mathcal{A}}(t)) \rangle, t \in T \right\}.$$

where $\mu_{\mathcal{A}}(t)$, $\nu_{\mathcal{A}}(t)$ and $\omega_{\mathcal{A}}(t)$ are called truth-membership function, indeterminacy-membership function and falsity-membership function, respectively. They are respectively defined by

$$\mu_{\mathcal{A}}(t) : T \rightarrow]^{-}0, 1^{+}[, \nu_{\mathcal{A}}(t) : T \rightarrow]^{-}0, 1^{+}[, \omega_{\mathcal{A}}(t) : T \rightarrow]^{-}0, 1^{+}[$$

such that $^{-}0 \leq \mu_{\mathcal{A}}(t) + \nu_{\mathcal{A}}(t) + \omega_{\mathcal{A}}(t) \leq 3^{+}$.

Definition 2 [33] Let T be a universe. An single valued neutrosophic set (SVN-set) over T is a neutrosophic set over T , but the truth-membership function, indeterminacy-membership function and falsity-membership function are respectively defined by

$$\mu_{\mathcal{A}}(t) : T \rightarrow [0, 1], \nu_{\mathcal{A}}(t) : T \rightarrow [0, 1], \omega_{\mathcal{A}}(t) : T \rightarrow [0, 1]$$

such that $0 \leq \mu_{\mathcal{A}}(t) + \nu_{\mathcal{A}}(t) + \omega_{\mathcal{A}}(t) \leq 3$.

Definition 3 [53] Let T be a universe. A neutrosophic multiset set (Nms) \mathcal{A} on T can be defined as follows:

$$\mathcal{A} = \{ \langle t, (\mu_{\mathcal{A}}^1(t), \mu_{\mathcal{A}}^2(t), \dots, \mu_{\mathcal{A}}^p(t)), (v_{\mathcal{A}}^1(t), v_{\mathcal{A}}^2(t), \dots, v_{\mathcal{A}}^p(t)), (w_{\mathcal{A}}^1(t), w_{\mathcal{A}}^2(t), \dots, w_{\mathcal{A}}^p(t)) \rangle : t \in T \}$$

Where,

$$\mu_{\mathcal{A}}^1(t), \mu_{\mathcal{A}}^2(t), \dots, \mu_{\mathcal{A}}^p(t) : T \rightarrow [0, 1],$$

$$v_{\mathcal{A}}^1(t), v_{\mathcal{A}}^2(t), \dots, v_{\mathcal{A}}^p(t) : T \rightarrow [0, 1],$$

and

$$w_{\mathcal{A}}^1(t), w_{\mathcal{A}}^2(t), \dots, w_{\mathcal{A}}^p(t) : T \rightarrow [0, 1]$$

such that

$$0 \leq \sup \mu_{\mathcal{A}}^i(t) + \sup v_{\mathcal{A}}^i(t) + \sup w_{\mathcal{A}}^i(t) \leq 3$$

($i = 1, 2, \dots, P$) and $(\mu_{\mathcal{A}}^1(t), \mu_{\mathcal{A}}^2(t), \dots, \mu_{\mathcal{A}}^p(t)), (v_{\mathcal{A}}^1(t), v_{\mathcal{A}}^2(t), \dots, v_{\mathcal{A}}^p(t))$ and $(w_{\mathcal{A}}^1(t), w_{\mathcal{A}}^2(t), \dots, w_{\mathcal{A}}^p(t))$ Is the truth-membership sequence, indeterminacy-membership sequence and falsity-membership sequence of the element u , respectively. Also, P is called the dimension (cardinality) of Nms \mathcal{A} , denoted $d(\mathcal{A})$. We arrange the truth-membership sequence in decreasing order but the corresponding indeterminacy-membership and falsity-membership sequence may not be in decreasing or increasing order.

The set of all Neutrosophic multisets on T is denoted by $NMS(T)$.

Definition 4 [12, 53, 56] Let $A, B \in NMS(T)$. Then,

- (1) \mathcal{A} is said to be Nm-subset of \mathcal{B} is denoted by $\mathcal{A} \subseteq \mathcal{B}$ if $\mu_{\mathcal{A}}^i(t) \leq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) \geq v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) \geq w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, \dots, P$.
- (2) \mathcal{A} is said to be neutrosophic equal of \mathcal{B} is denoted by $\mathcal{A} = \mathcal{B}$ if $\mu_{\mathcal{A}}^i(t) = \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, \dots, P$.
- (3) The complement of \mathcal{A} denoted by \mathcal{A}^c and is defined by

$$\mathcal{A}^c = \{ \langle t, (w_{\mathcal{A}}^1(t), w_{\mathcal{A}}^2(t), \dots, w_{\mathcal{A}}^p(t)), (v_{\mathcal{A}}^1(t), v_{\mathcal{A}}^2(t), \dots, v_{\mathcal{A}}^p(t)), (\mu_{\mathcal{A}}^1(t), \mu_{\mathcal{A}}^2(t), \dots, \mu_{\mathcal{A}}^p(t)) \rangle : t \in T \}$$

(4) If $\mu_{\mathcal{A}}^i(t) = 0$ and $\nu_{\mathcal{A}}^i(t) = w_{\mathcal{A}}^i(t) = 1$ for all $t \in T$ and $i = 1, 2, \dots, P$, then \mathcal{A} is called null ns-set and denoted by Φ .

(5) If $\mu_{\mathcal{A}}^i(t) = 1$ and $\nu_{\mathcal{A}}^i(t) = w_{\mathcal{A}}^i(t) = 0$ for all $t \in T$ and $i = 1, 2, \dots, P$, then \mathcal{A} is called universal ns-set and denoted by \tilde{T} .

(6) The union of \mathcal{A} and \mathcal{B} is denoted by $\mathcal{A} \tilde{\cup} \mathcal{B} = \mathcal{C}$ and is defined by

$$\mathcal{C} = \{ \langle t, (\mu_{\mathcal{C}}^1(t), \mu_{\mathcal{C}}^2(t), \dots, \mu_{\mathcal{C}}^p(t)), (v_{\mathcal{C}}^1(t), v_{\mathcal{C}}^2(t), \dots, v_{\mathcal{C}}^p(t)), (w_{\mathcal{C}}^1(t), w_{\mathcal{C}}^2(t), \dots, w_{\mathcal{C}}^p(t)) \rangle : t \in T \}$$

Where $\mu_{\mathcal{C}}^i = \mu_{\mathcal{A}}^i(t) \vee \mu_{\mathcal{B}}^i(t)$, $\nu_{\mathcal{C}}^i = \nu_{\mathcal{A}}^i(t) \wedge \nu_{\mathcal{B}}^i(t)$, $w_{\mathcal{C}}^i = w_{\mathcal{A}}^i(t) \wedge w_{\mathcal{B}}^i(t)$, $\forall t \in T$ and $i = 1, 2, \dots, P$.

(7) The intersection of \mathcal{A} and \mathcal{B} is denoted by $\mathcal{A} \tilde{\cap} \mathcal{B} = \mathcal{D}$ and is defined by

$$\mathcal{D} = \{ \langle t, (\mu_{\mathcal{D}}^1(t), \mu_{\mathcal{D}}^2(t), \dots, \mu_{\mathcal{D}}^p(t)), (v_{\mathcal{D}}^1(t), v_{\mathcal{D}}^2(t), \dots, v_{\mathcal{D}}^p(t)), (w_{\mathcal{D}}^1(t), w_{\mathcal{D}}^2(t), \dots, w_{\mathcal{D}}^p(t)) \rangle : t \in T \}$$

where $\mu_{\mathcal{D}}^i = \mu_{\mathcal{A}}^i(t) \wedge \mu_{\mathcal{B}}^i(t)$, $\nu_{\mathcal{D}}^i = \nu_{\mathcal{A}}^i(t) \vee \nu_{\mathcal{B}}^i(t)$, $w_{\mathcal{D}}^i = w_{\mathcal{A}}^i(t) \vee w_{\mathcal{B}}^i(t)$, $\forall t \in T$ and $i = 1, 2, \dots, P$.

(8) The addition of \mathcal{A} and \mathcal{B} is denoted by $\mathcal{A} \tilde{+} \mathcal{B} = \mathcal{U}_1$ and is defined by

$$\mathcal{U}_1 = \{ \langle t, (\mu_{\mathcal{U}_1}^1(t), \mu_{\mathcal{U}_1}^2(t), \dots, \mu_{\mathcal{U}_1}^p(t)), (v_{\mathcal{U}_1}^1(t), v_{\mathcal{U}_1}^2(t), \dots, v_{\mathcal{U}_1}^p(t)), (w_{\mathcal{U}_1}^1(t), w_{\mathcal{U}_1}^2(t), \dots, w_{\mathcal{U}_1}^p(t)) \rangle : t \in T \}$$

where $\mu_{\mathcal{U}_1}^i = \mu_{\mathcal{A}}^i(t) + \mu_{\mathcal{B}}^i(t) - \mu_{\mathcal{A}}^i(t) \cdot \mu_{\mathcal{B}}^i(t)$, $\nu_{\mathcal{U}_1}^i = \nu_{\mathcal{A}}^i(t) \cdot \nu_{\mathcal{B}}^i(t)$, $w_{\mathcal{U}_1}^i = w_{\mathcal{A}}^i(t) \cdot w_{\mathcal{B}}^i(t)$ $\forall t \in T$ and $i = 1, 2, \dots, P$.

(9) The multiplication of \mathcal{A} and \mathcal{B} is denoted by $\mathcal{A} \tilde{\times} \mathcal{B} = \mathcal{U}_2$ and is defined by

$$\mathcal{U}_2 = \{ \langle t, (\mu_{\mathcal{U}_2}^1(t), \mu_{\mathcal{U}_2}^2(t), \dots, \mu_{\mathcal{U}_2}^p(t)), (v_{\mathcal{U}_2}^1(t), v_{\mathcal{U}_2}^2(t), \dots, v_{\mathcal{U}_2}^p(t)), (w_{\mathcal{U}_2}^1(t), w_{\mathcal{U}_2}^2(t), \dots, w_{\mathcal{U}_2}^p(t)) \rangle : t \in T \}$$

where $\mu_{\mathcal{U}_2}^i = \mu_{\mathcal{A}}^i(t) \cdot \mu_{\mathcal{B}}^i(t)$, $\nu_{\mathcal{U}_2}^i = \nu_{\mathcal{A}}^i(t) + \nu_{\mathcal{B}}^i(t) - \nu_{\mathcal{A}}^i(t) \cdot \nu_{\mathcal{B}}^i(t)$, $w_{\mathcal{U}_2}^i = w_{\mathcal{A}}^i(t) + w_{\mathcal{B}}^i(t) - w_{\mathcal{A}}^i(t) \cdot w_{\mathcal{B}}^i(t)$ $\forall t \in T$ and $i = 1, 2, \dots, P$.

Here $\vee, \wedge, +, \cdot, -$ denotes maximum, minimum, addition, multiplication, subtraction of real numbers respectively.

Definition 5 [13] Let

$$\mathcal{A} = \{ \langle t, (\mu_{\mathcal{A}}^1(t), \mu_{\mathcal{A}}^2(t), \dots, \mu_{\mathcal{A}}^p(t)), (v_{\mathcal{A}}^1(t), v_{\mathcal{A}}^2(t), \dots, v_{\mathcal{A}}^p(t)), (w_{\mathcal{A}}^1(t), w_{\mathcal{A}}^2(t), \dots, w_{\mathcal{A}}^p(t)) \rangle : t \in T \}$$

and

$$\mathcal{B} = \{ \langle t, (\mu_{\mathcal{B}}^1(t), \mu_{\mathcal{B}}^2(t), \dots, \mu_{\mathcal{B}}^p(t)), (v_{\mathcal{B}}^1(t), v_{\mathcal{B}}^2(t), \dots, v_{\mathcal{B}}^p(t)), (w_{\mathcal{B}}^1(t), w_{\mathcal{B}}^2(t), \dots, w_{\mathcal{B}}^p(t)) \rangle : t \in T \}$$

and be two NMSs, then the normalized hamming distance between \mathcal{A} and \mathcal{B} can be defined as follows:

$$d_{NHD}(\mathcal{A}, \mathcal{B}) = \frac{1}{3n \cdot P} \sum_{j=1}^P \sum_{i=1}^n (|\mu_{\mathcal{A}}^j(t_i) - \mu_{\mathcal{B}}^j(t_i)| + |\nu_{\mathcal{A}}^j(t_i) - \nu_{\mathcal{B}}^j(t_i)| + |w_{\mathcal{A}}^j(t_i) - w_{\mathcal{B}}^j(t_i)|).$$

3. The Outranking Relations of Neutrosophic Multi-Sets

In this section, the binary relations between two neutrosophic refined sets that are based on ELECTRE by extending the studies in [22]. Some of it is quoted from [13, 22, 35, 49].

Definition 6 Let $\mathcal{A} = \{ \langle t, (\mu_{\mathcal{A}}^i(t), \nu_{\mathcal{A}}^i(t), w_{\mathcal{A}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ and

$\mathcal{B} = \{ \langle t, (\mu_{\mathcal{B}}^i(t), \nu_{\mathcal{B}}^i(t), w_{\mathcal{B}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ be two NMS on T . Then, the strong dominance relation, weak dominance relation, and indifference relation of NMS can be defined as follows:

1. If $\mu_{\mathcal{A}}^i(t) \geq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) < v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) < w_{\mathcal{B}}^i(t)$ or $\mu_{\mathcal{A}}^i(t) > \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. Then \mathcal{A} strongly dominates \mathcal{B} (\mathcal{B} is strongly dominated by \mathcal{A}), denoted by $\mathcal{A} \succ_s \mathcal{B}$.
2. If $\mu_{\mathcal{A}}^i(t) \geq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) \geq v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) < w_{\mathcal{B}}^i(t)$ or $\mu_{\mathcal{A}}^i(t) \geq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) < v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) \geq w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. Then \mathcal{A} weakly dominates \mathcal{B} (\mathcal{B} is weakly dominated by \mathcal{A}), denoted by $\mathcal{A} \succ_w \mathcal{B}$.
3. If $\mu_{\mathcal{A}}^i(t) = \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. Then \mathcal{A} indifferent to \mathcal{B} , denoted by $\mathcal{A} \sim_l \mathcal{B}$.
4. If none of the relations mentioned above exist between \mathcal{A} and \mathcal{B} for any $t \in T$, then \mathcal{A} and \mathcal{B} are incomparable, denoted by $\mathcal{A} \perp \mathcal{B}$.

Proposition 7 Let $\mathcal{A} = \{ \langle t, (\mu_{\mathcal{A}}^i(t), v_{\mathcal{A}}^i(t), w_{\mathcal{A}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ and

$\mathcal{B} = \{ \langle t, (\mu_{\mathcal{B}}^i(t), v_{\mathcal{B}}^i(t), w_{\mathcal{B}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ be two NMS on T , then the following properties can be obtained:

1. If $\mathcal{B} \subset \mathcal{A}$, then $\mathcal{A} \succ_s \mathcal{B}$;
2. If $\mathcal{A} \succ_s \mathcal{B}$, then $\mathcal{B} \subseteq \mathcal{A}$;
3. $\mathcal{A} \sim_l \mathcal{B}$ if and only if $\mathcal{A} = \mathcal{B}$.

Proof:

1. If $\mathcal{B} \subset \mathcal{A}$, then $\mu_{\mathcal{B}}^i(t) \leq \mu_{\mathcal{A}}^i(t), v_{\mathcal{B}}^i(t) \geq v_{\mathcal{A}}^i(t), w_{\mathcal{B}}^i(t) \geq w_{\mathcal{A}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. $\mathcal{A} \succ_s \mathcal{B}$ is definitely validated according to the strong dominance relation in Definition 6.

2. $\mathcal{A} \succ_s \mathcal{B}$ then based on Definition 6, $\mu_{\mathcal{A}}^i(t) \geq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) < v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) < w_{\mathcal{B}}^i(t)$ or $\mu_{\mathcal{A}}^i(t) > \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. are realized. Then we have $\mathcal{B} \subseteq \mathcal{A}$.

3. Necessity: $\mathcal{A} \sim_l \mathcal{B} \Rightarrow \mathcal{A} = \mathcal{B}$. According to the indifference relation in Definition 6 it is known that $\mu_{\mathcal{A}}^i(t) = \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. Clearly $\mathcal{A} \subseteq \mathcal{A}$ and $\mathcal{B} \subseteq \mathcal{A}$ are achieved, then $\mathcal{A} = \mathcal{B}$.

Sufficiency: $\mathcal{A} = \mathcal{B} \Rightarrow \mathcal{A} \sim_l \mathcal{B}$. If $\mathcal{A} = \mathcal{B}$, then it is known that $\mathcal{A} \subseteq \mathcal{B}$ and $\mathcal{B} \subseteq \mathcal{A}$, which means

$\mu_{\mathcal{B}}^i(t) \leq \mu_{\mathcal{A}}^i(t), v_{\mathcal{B}}^i(t) \geq v_{\mathcal{A}}^i(t), w_{\mathcal{B}}^i(t) \geq w_{\mathcal{A}}^i(t)$ or $\mu_{\mathcal{A}}^i(t) = \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$. are obtained. Due to the indifference relation in Definition 6, $\mathcal{A} \sim_l \mathcal{B}$ is definitely obtained.

Proposition 8 Let $\mathcal{A} = \{ \langle t, (\mu_{\mathcal{A}}^i(t), v_{\mathcal{A}}^i(t), w_{\mathcal{A}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$,

$\mathcal{B} = \{ \langle t, (\mu_{\mathcal{B}}^i(t), v_{\mathcal{B}}^i(t), w_{\mathcal{B}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ and $\mathcal{C} = \{ \langle t, (\mu_{\mathcal{C}}^i(t), v_{\mathcal{C}}^i(t), w_{\mathcal{C}}^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ be three NMS on T , if $\mathcal{A} \succ_s \mathcal{B}$ and $\mathcal{B} \succ_s \mathcal{C}$, then $\mathcal{A} \succ_s \mathcal{C}$.

Proof: According to the strong dominance relation in Definition 6, if $\mathcal{A} \succ_s \mathcal{B}$, then $\mu_{\mathcal{A}}^i(t) \geq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) < v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) < w_{\mathcal{B}}^i(t)$ or $\mu_{\mathcal{A}}^i(t) > \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) = v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) = w_{\mathcal{B}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$.

if $\mathcal{B} \succ_s \mathcal{C}$, then $\mu_{\mathcal{B}}^i(t) \geq \mu_{\mathcal{C}}^i(t), v_{\mathcal{B}}^i(t) < v_{\mathcal{C}}^i(t), w_{\mathcal{B}}^i(t) < w_{\mathcal{C}}^i(t)$ or $\mu_{\mathcal{B}}^i(t) > \mu_{\mathcal{C}}^i(t), v_{\mathcal{B}}^i(t) = v_{\mathcal{C}}^i(t), w_{\mathcal{B}}^i(t) = w_{\mathcal{C}}^i(t), \forall t \in T$ and $i = 1, 2, 3, \dots, p$.

Therefore the further derivations are: If

$$\mu_{\mathcal{A}}^i(t) \geq \mu_{\mathcal{B}}^i(t), v_{\mathcal{A}}^i(t) < v_{\mathcal{B}}^i(t), w_{\mathcal{A}}^i(t) < w_{\mathcal{B}}^i(t), \dots(1)$$

$$\mu_B^i(t) \geq \mu_C^i(t), v_B^i(t) < v_C^i(t), w_B^i(t) < w_C^i(t), \dots (2)$$

from (1) and (2)

$$\mu_A^i(t) \geq \mu_C^i(t), v_A^i(t) < v_C^i(t), w_A^i(t) < w_C^i(t),$$

then based on Definition 6 $\mathcal{A} \succ_s C$ is realized. If

$$\mu_A^i(t) \geq \mu_B^i(t), v_A^i(t) < v_B^i(t), w_A^i(t) < w_B^i(t), \dots (3)$$

$$\mu_B^i(t) > \mu_C^i(t), v_B^i(t) = v_C^i(t), w_B^i(t) = w_C^i(t), \dots (4)$$

from (3) and (4)

$$\mu_A^i(t) \geq \mu_C^i(t), v_A^i(t) < v_C^i(t), w_A^i(t) < w_C^i(t),$$

then based on Definition 6 $\mathcal{A} \succ_s C$ is achieved. If

$$\mu_A^i(t) > \mu_B^i(t), v_A^i(t) = v_B^i(t), w_A^i(t) = w_B^i(t), \dots (5)$$

$$\mu_B^i(t) \geq \mu_C^i(t), v_B^i(t) < v_C^i(t), w_B^i(t) < w_C^i(t), \dots (6)$$

from (5) and (6)

$$\mu_A^i(t) > \mu_C^i(t), v_A^i(t) = v_C^i(t), w_A^i(t) = w_C^i(t),$$

then based on Definition 6 $\mathcal{A} \succ_s C$ is obtained. If

$$\mu_A^i(t) > \mu_B^i(t), v_A^i(t) = v_B^i(t), w_A^i(t) = w_B^i(t), \dots (7)$$

$$\mu_B^i(t) > \mu_C^i(t), v_B^i(t) = v_C^i(t), w_B^i(t) = w_C^i(t), \dots (8)$$

from (7) and (8)

$$\mu_A^i(t) > \mu_C^i(t), v_A^i(t) = v_C^i(t), w_A^i(t) = w_C^i(t),$$

then based on Definition 6 $\mathcal{A} \succ_s C$ is realized. Therefore, if $\mathcal{A} \succ_s B$ and $B \succ_s C$, then $\mathcal{A} \succ_s C$.

Proposition 9 Let $\mathcal{A} = \{ \langle t, (\mu_A^i(t), v_A^i(t), w_A^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$,

$\mathcal{B} = \{ \langle t, (\mu_B^i(t), v_B^i(t), w_B^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ and $\mathcal{C} = \{ \langle t, (\mu_C^i(t), v_C^i(t), w_C^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ be three NMS on T , if $\mathcal{A} \sim_l B$ and $B \sim_l C$, then $\mathcal{A} \sim_l C$.

Proof: Clearly, if $\mathcal{A} \sim_l B$ and $B \sim_l C$, then $\mathcal{A} \sim_l C$ is surely validated.

Proposition 10 Let $\mathcal{A} = \{ \langle t, (\mu_A^i(t), v_A^i(t), w_A^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$,

$\mathcal{B} = \{ \langle t, (\mu_B^i(t), v_B^i(t), w_B^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ and $\mathcal{C} = \{ \langle t, (\mu_C^i(t), v_C^i(t), w_C^i(t)) \rangle : t \in T, (i = 1, 2, 3, \dots, p) \}$ be three NMS on $T = \{t_1, t_2, \dots, t_n\}$, then the following results can be obtained.

- 1 – irreflexivity : $\forall \mathcal{A} \in \text{NMSs}, \mathcal{A} \not\succeq_s \mathcal{A}$;
- 2 – asymmetry : $\forall \mathcal{A}, \mathcal{B}$ on NMSs; $\mathcal{A} \succ_s \mathcal{B} \Rightarrow \mathcal{B} \not\succeq_s \mathcal{A}$;
- 3 – transitivity: $\forall \mathcal{A}, \mathcal{B}, \mathcal{C}$ on NMSs; $\mathcal{A} \succ_s \mathcal{B}, \mathcal{B} \succ_s \mathcal{C}$, then $\mathcal{A} \succ_s \mathcal{C}$.
- 4 – irreflexivity : $\forall \mathcal{A} \in \text{NMSs}, \mathcal{A} \not\succeq_w \mathcal{A}$;
- 5 – asymmetry : $\forall \mathcal{A}, \mathcal{B}$ on NMSs; $\mathcal{A} \succ_w \mathcal{B} \Rightarrow \mathcal{B} \not\succeq_w \mathcal{A}$;
- 6 – non – transitivity: $\exists \mathcal{A}, \mathcal{B}, \mathcal{C}$ on NMSs; $\mathcal{A} \succ_s \mathcal{B}, \mathcal{B} \succ_s \mathcal{C}$, then $\mathcal{A} \not\succeq_s \mathcal{C}$.
- 7 – reflexivity : $\forall \mathcal{A} \in \text{NMSs}, \mathcal{A} \sim_l \mathcal{A}$;
- 8 – symmetry : $\forall \mathcal{A}, \mathcal{B}$ on NMSs; $\mathcal{A} \sim_l \mathcal{B} \Rightarrow \mathcal{B} \sim_l \mathcal{A}$;
- 9 – transitivity: $\exists \mathcal{A}, \mathcal{B}, \mathcal{C}$ on NMSs; $\mathcal{A} \sim_l \mathcal{B}, \mathcal{B} \sim_l \mathcal{C}$, then $\mathcal{A} \sim_l \mathcal{C}$.

Example 11 1,2,4,5 and 6 are exemplified as follows.

1. If $\mathcal{A} = \langle (0.8, 0.5, \dots, 0.6), (0.3, 0.1, \dots, 0.5), (0.2, 0.3, \dots, 0.4) \rangle$ is a NMSs, then $\mathcal{A} \not\succeq_s \mathcal{A}$ can be obtained.
2. If $\mathcal{A} = \langle (0.5, 0.7, \dots, 0.6), (0.2, 0.3, \dots, 0.4), (0.1, 0.3, \dots, 0.2) \rangle$ and $\mathcal{B} = \langle (0.4, 0.6, \dots, 0.5), (0.3, 0.4, \dots, 0.5), (0.2, 0.5, \dots, 0.3) \rangle$ are two NMSs, then $\mathcal{A} \succ_s \mathcal{B}$, but $\mathcal{B} \not\succeq_s \mathcal{A}$ is realized.

3. If $\mathcal{A} = \langle (0.7, 0.4, \dots, 0.5), (0.4, 0.2, \dots, 0.6), (0.3, 0.3, \dots, 0.2) \rangle$ is a NMSs, then $\mathcal{A} \not>_w \mathcal{A}$ can be obtained.
4. If $\mathcal{A} = \langle (0.5, 0.7, \dots, 0.6), (0.5, 0.6, \dots, 0.4), (0.1, 0.3, \dots, 0.2) \rangle$ and $\mathcal{B} = \langle (0.3, 0.5, \dots, 0.6), (0.2, 0.3, \dots, 0.1), (0.2, 0.5, \dots, 0.3) \rangle$ are two NMSs, then $\mathcal{A} >_w \mathcal{B}$, however $\mathcal{B} \not>_w \mathcal{A}$.
5. If $\mathcal{A} = \langle (0.5, 0.7, \dots, 0.6), (0.3, 0.2, \dots, 0.4), (0.1, 0.3, \dots, 0.2) \rangle$,
6. $\mathcal{B} = \langle (0.5, 0.6, \dots, 0.4), (0.5, 0.4, \dots, 0.6), (0.2, 0.5, \dots, 0.3) \rangle$ and $\mathcal{C} = \langle (0.4, 0.3, \dots, 0.2), (0.6, 0.5, \dots, 0.7), (0.3, 0.6, \dots, 0.8) \rangle$ are three NMSs, then $\mathcal{A} >_w \mathcal{B}$ and $\mathcal{B} >_w \mathcal{C}$ are obtained, $\mathcal{A} >_w \mathcal{C}$.

Proposition 11 [22] Let t_1 and t_2 be two actions, the performances for actions t_1 and t_2 be in the form of NMSs, and $P = s \cup w \cup l$ mean that “ t_1 is at least as good as t_2 ”, then four situations may arise:

1. t_1Pt_2 and not t_2Pt_1 , that is $t_1 >_s t_2$ or $t_1 >_w t_2$;
2. t_2Pt_1 and not t_1Pt_2 , that is $t_2 >_s t_1$ or $t_2 >_w t_1$;
3. t_1Pt_2 and t_2Pt_1 , that is $t_1 \sim_l t_2$;
4. not t_1Pt_2 and not t_2Pt_1 , that is $t_1 \perp t_2$.

4. An outranking approach for MCDM with simplified neutrosophic multi-set information

In this section, we introduced an approach for a MCDM problem with neutrosophic multi-set information. Some of it is quoted from [22, 35, 49].

Definition 12 [15] Let $X = (x_1, x_2, \dots, x_n)$ be a set of alternatives, $C = (c_1, c_2, \dots, c_n)$ be the set of criteria, $w = (w_1, w, \dots, w_n)^T$ be the weight vector of the criterions $C_j (j = 1, 2, \dots, n)$ such that $w_j \geq 0$ and $\sum_{j=1}^n w_j = 1$ and $Z_{ij} = \langle (\mu_{ij}^1, \mu_{ij}^2, \dots, \mu_{ij}^n), (v_{ij}^1, v_{ij}^2, \dots, v_{ij}^n), (w_{ij}^1, w_{ij}^2, \dots, w_{ij}^n) \rangle$ be the decision matrix in which the rating values of the alternatives in for NMSs. Then,

$$[Z_{ij}]_{m \times n} = \begin{matrix} & c_1 & c_2 & \dots & c_n \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{matrix} & \begin{pmatrix} Z_{11} & Z_{12} & \dots & Z_{1n} \\ Z_{21} & Z_{22} & \dots & Z_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ Z_{m1} & Z_{m2} & \dots & Z_{mn} \end{pmatrix} \end{matrix}$$

is called an NMS-multi-criteria decision making matrix of the decision maker.

Definition 13 [22, 35] In multi-criteria decision making problems;

1. The cost-type criterion values can be transformed into benefit-type criterion values as follows:

$$\alpha_{ij} = \begin{cases} Z_{ij} & \text{for benefit criterion } C_j, \\ (Z_{ij})^c & \text{for benefit criterion } C_j, \end{cases} \quad (i = 1, 2, \dots, m; j = 1, 2, \dots, n) \quad (9)$$

where $(Z_{ij})^c$ is complement of Z_{ij} as defined in Definition 4.

2. The concordance set of subscripts, which should satisfy the constraint $Z_{ij}PZ_{kj}$, is represented as:

$$O_{ik} = \{j: Z_{ij}PZ_{kj}\} \quad (i, k = 1, 2, \dots, m).$$

$Z_{ij}PZ_{kj}$ represents $Z_{ij} >_s Z_{kj}$ or $Z_{ij} >_w Z_{kj}$ or $Z_{ij} \sim_l Z_{kj}$.

3. The concordance index h_{ik} between x_i and x_k is thus defined as follows:

$$h_{ik} = \sum_{j \in O_{ik}} w_j \quad (10)$$

Thus, the concordance matrix C is:

$$H = h_{ik} = \begin{pmatrix} - & h_{12} & \cdots & h_{1n} \\ h_{21} & - & \cdots & h_{2n} \\ \vdots & & \ddots & \vdots \\ h_{n1} & h_{n2} & \cdots & - \end{pmatrix}$$

In H ; h_{ik} ($i \neq k$) denote the degree to which the evaluations of x_i are at least as good as those of the competitor x_k , and the degree to which x_i is inferior to x_k decreases with increasing h_{ik} .

4. The discordance set of subscripts for criteria is given as;

$$G_{ik} = J - O_{ik}.$$

5. The discordance index $G(x_i; x_k)$ is represented as:

$$G_{ik} = \frac{\max_{j \in G_{ik}} \{d(Z_{ij}, Z_{kj})\}}{\max_{j \in J} \{d(Z_{ij}, Z_{kj})\}} \tag{11}$$

here $d(Z_{ij}, Z_{kj})$ denotes the normalized Hamming distance between Z_{ij} and Z_{kj} as defined in Definition 5.

Thus, the discordance matrix D is:

$$g_{ik} = \begin{pmatrix} - & g_{12} & \cdots & g_{1n} \\ g_{21} & - & \cdots & g_{2n} \\ \vdots & & \ddots & \vdots \\ g_{n1} & g_{n2} & \cdots & - \end{pmatrix}$$

In G ; g_{ik} ($i \neq k$) denote the degree to which the evaluations of x_i are at least as good as those of the competitor x_k , and the degree to which x_i is inferior to x_k decreases with increasing g_{ik} .

6. To rank all alternatives, the net dominance index of x_k

$$h_{ik} = \sum_{i=1, i \neq k}^n h_{ik} - \sum_{i=1, i \neq k}^n h_{ki} \tag{12}$$

and the net disadvantage index of x_k is

$$g_{ik} = \sum_{i=1, i \neq k}^n g_{ik} - \sum_{i=1, i \neq k}^n g_{ki} \tag{13}$$

In here, h_k is the sum of the concordance indices between x_k and x_k ($i \neq k$) minus the sum of the concordance indices between x_k ($i \neq k$) and x_k , and reflects the dominance degree of the alternative x_k among the relevant alternatives. Meanwhile, g_k reflects the disadvantage degree of the alternative x_k among the relevant alternatives. Therefore, x_k obtains a greater dominance over the other alternatives that are being compared as h_k increases and g_k decreases.

Definition 14 [35] The ranking rules of two alternatives are

- i. If $h_i < h_k$ and $g_i > g_k$ then x_k is superior to x_i , as denoted by $x_k > x_i$;
- ii. If $h_i = h_k$ and $g_i = g_k$ then x_k is indifferent to x_i , as denoted by $x_k \sim x_i$;
- i. if the relation between x_k and x_i does not belong to (i) or (ii); then x_k and x_i are incomparable; as denoted by $x_k \perp x_i$.

Now, we give an algorithm to develop a new approach as

Algorithm:

Step 1 Give the decision-making matrix $[Z_{ij}]_{m \times n}$; for decision;

Step 2 Compute the weighted normalized matrix as;

$$[\gamma_{ij}]_{m \times n} = \alpha_{ij} w_j \quad i = 1, 2, \dots, m; j = 1, 2, \dots, n.$$

where w_j is the weight of the j th criterion with $\sum_{j=1}^n w_j = 1$.

Step 3 Find the concordance set of subscripts;

Step 4 Find the discordance set of subscripts;

Step 5 Compute the concordance matrix $H = (h_{ik})_{n \times n}$

Step 6 Compute the discordance matrix $G = (g_{ik})_{n \times n}$

Step 7. Compute the net dominance index of each alternative h_i ($i=1,2,3,\dots,m$)

Step 8. Compute the net disadvantage index of each alternative g_i ($i=1,2,\dots,m$)

Step 9. Rank all alternatives and select the best alternative.

5 Illustrative examples

In this section, we introduced an example for a MCDM problem with neutrosophic refined information. Some of it is quoted from [22, 35, 49].

Example 15 Assume that $X = (x_1, x_2, x_3, x_4)$ be a set of alternatives and $C = (c_1, c_2, c_3, c_4)$ be the set of criteria, $w = (0.1, 0.3, 0.2, 0.4)^T$ be the weight vector of the criteria C_j ($j = 1, 2, \dots, n$). The four alternatives are to be evaluated under the above four criteria in the form of NMSs. Then,

Step 1. The decision matrix $[Z_{ij}]_{m \times n}$ is given as;

$$\begin{pmatrix} \langle(0:1; 0:2; 0:4; 0:5); (0:6; 0:3; 0:5; 0:2); (0:2; 0:4; 0:5; 0:6)\rangle \\ \langle(0:3; 0:4; 0:6; 0:7); (0:2; 0:5; 0:1; 0:8); (0:3; 0:4; 0:6; 0:8)\rangle \\ \langle(0:1; 0:2; 0:5; 0:6); (0:1; 0:3; 0:5; 0:2); (0:1; 0:5; 0:7; 0:9)\rangle \\ \langle(0:2; 0:3; 0:4; 0:5); (0:3; 0:2; 0:4; 0:6); (0:2; 0:3; 0:5; 0:7)\rangle \\ \\ \langle(0:3; 0:5; 0:7; 0:8); (0:4; 0:3; 0:6; 0:2); (0:1; 0:3; 0:5; 0:2)\rangle \\ \langle(0:2; 0:3; 0:4; 0:5); (0:1; 0:4; 0:3; 0:6); (0:2; 0:3; 0:4; 0:5)\rangle \\ \langle(0:1; 0:2; 0:6; 0:7); (0:3; 0:2; 0:5; 0:4); (0:1; 0:2; 0:5; 0:6)\rangle \\ \langle(0:3; 0:4; 0:6; 0:8); (0:2; 0:1; 0:3; 0:6); (0:4; 0:3; 0:2; 0:5)\rangle \\ \\ \langle(0:2; 0:4; 0:5; 0:6); (0:3; 0:5; 0:2; 0:6); (0:1; 0:2; 0:5; 0:6)\rangle \\ \langle(0:4; 0:5; 0:7; 0:8); (0:1; 0:6; 0:2; 0:3); (0:1; 0:4; 0:3; 0:6)\rangle \\ \langle(0:3; 0:6; 0:8; 0:9); (0:2; 0:4; 0:1; 0:5); (0:2; 0:1; 0:3; 0:6)\rangle \\ \langle(0:1; 0:2; 0:4; 0:6); (0:1; 0:3; 0:7; 0:4); (0:3; 0:4; 0:6; 0:7)\rangle \\ \\ \langle(0:1; 0:2; 0:4; 0:5); (0:2; 0:3; 0:5; 0:4); (0:1; 0:3; 0:7; 0:4)\rangle \\ \langle(0:3; 0:4; 0:5; 0:6); (0:3; 0:1; 0:2; 0:5); (0:3; 0:6; 0:8; 0:9)\rangle \\ \langle(0:1; 0:3; 0:4; 0:5); (0:1; 0:4; 0:6; 0:7); (0:1; 0:2; 0:6; 0:7)\rangle \\ \langle(0:2; 0:4; 0:5; 0:7); (0:2; 0:3; 0:5; 0:6); (0:3; 0:2; 0:4; 0:6)\rangle \end{pmatrix}$$

Step 2. The weighted normalized matrix $[\gamma_{ij}]_{m \times n}$ is computed as;

$$\begin{pmatrix} (0:7943; 0:8513; 0:9124; 0:9330); (0:0875; 0:0350; 0:0669; 0:0220); (0:0220; 0:0104; 0:0669; 0:0875) \\ (0:6968; 0:7596; 0:8579; 0:8985); (0:0647; 0:1877; 0:0311; 0:3829); (0:1014; 0:1420; 0:2403; 0:3829) \\ (0:6309; 0:7247; 0:8705; 0:9028); (0:2080; 0:0688; 0:1294; 0:0436); (0:2080; 0:1294; 0:2140; 0:3690) \\ (0:5253; 0:6178; 0:6931; 0:7578); (0:1329; 0:0853; 0:1848; 0:3068); (0:0853; 0:1329; 0:2421; 0:3822) \\ \\ (0:8865; 0:9330; 0:9649; 0:9779); (0:0498; 0:0350; 0:0875; 0:0620); (0:0104; 0:0350; 0:0669; 0:0220) \\ (0:6170; 0:6968; 0:7596; 0:8122); (0:0311; 0:1420; 0:1014; 0:2403); (0:0647; 0:1014; 0:1420; 0:1877) \\ (0:6309; 0:7247; 0:9028; 0:9311); (0:0188; 0:0436; 0:1294; 0:0971); (0:0208; 0:0436; 0:1294; 0:1674) \\ (0:6178; 0:6931; 0:8151; 0:9146); (0:0853; 0:0412; 0:1329; 0:3068); (0:1848; 0:1329; 0:0853; 0:2421) \\ \\ (0:8513; 0:9124; 0:9330; 0:9502); (0:0350; 0:0669; 0:0720; 0:0875); (0:0104; 0:0220; 0:0669; 0:0875) \\ (0:7596; 0:8122; 0:8985; 0:9352); (0:0311; 0:0203; 0:0647; 0:1014); (0:0311; 0:1420; 0:1014; 0:2403) \\ (0:7860; 0:9028; 0:9563; 0:9791); (0:0436; 0:0971; 0:0208; 0:1294); (0:0436; 0:0208; 0:0688; 0:1674) \\ (0:3981; 0:5253; 0:6931; 0:8151); (0:0412; 0:1329; 0:3822; 0:1848); (0:0412; 0:1329; 0:3822; 0:6018) \end{pmatrix}$$

(0: 7943; 0: 8513; 0: 9124; 0: 9330); (0: 0220; 0: 0350; 0: 0669; 0: 0498); (0: 0104; 0: 0350; 0: 1134; 0: 0498)
 (0: 6968; 0: 7596; 0: 8122; 0: 8579); (0: 1014; 0: 0311; 0: 0647; 0: 1877); (0: 1014; 0: 2403; 0: 2403; 0: 4988)
 (0: 6309; 0: 7860; 0: 8325; 0: 8705); (0: 0228; 0: 0971; 0: 1674; 0: 2140); (0: 0208; 0: 0436; 0: 1674; 0: 2140)
 (0: 5253; 0: 6931; 0: 7578; 0: 8670); (0: 1853; 0: 1329; 0: 2421; 0: 3068); (0: 0329; 0: 0853; 0: 1848; 0: 3068)

Step 3. The concordance set is found as;

$$O_{12} = \{ \}; O_{21} = \{4\}; O_{31} = \{ \}; O_{41} = \{ \}; O_{13} = \{1, 2\}; O_{23} = \{ \};$$

$$O_{32} = \{ \}; O_{42} = \{ \}; O_{14} = \{4\}; O_{24} = \{1, 3\}; O_{34} = \{1, 2\}; O_{43} = \{ \}.$$

Step 4. The discordance set is found as;

$$G_{12} = \{1, 2, 3, 4\}; G_{21} = \{1, 2, 3\}; G_{31} = \{1, 2, 3, 4\}; G_{41} = \{1, 2, 3, 4\}; O_{13} = \{1, 2\}; G_{23} = \{1, 2, 3, 4\};$$

$$G_{32} = \{1, 2, 3, 4\}; G_{42} = \{1, 2, 3, 4\}; G_{14} = \{1, 2, 3\}; G_{24} = \{2, 4\}; G_{34} = \{3, 4\}; G_{43} = \{1, 2, 3, 4\}.$$

where $\{ \}$ denotes "empty".

Step 5. The concordance is computed as;

$$H = \begin{pmatrix} - & 0 & 0.4 & 0.4 \\ 0.4 & - & 0.4 & 0.3 \\ 0 & 0 & - & 0.4 \\ 0 & 0 & 0 & - \end{pmatrix}$$

Step 6. The discordance matrix is computed as;

$$G = \begin{pmatrix} - & 1 & 0.6612 & 1 \\ 0.9958 & - & 1 & 0.5778 \\ 1 & 1 & - & 1 \\ 1 & 1 & 1 & - \end{pmatrix}$$

Step 7. The net dominance index of each alternative h_i ($i=1,2,3,4$) is computed as;

$$h_1 = 0.4, h_2 = 1.1, h_3 = -0.4 \text{ and } h_4 = -1.1, \Rightarrow h_4 < h_3 < h_1 < h_2;$$

Step 8. The net disadvantage index of each alternative g_i ($i=1,2,3,4$) is computed as;

$$g_1 = -0.3346, g_2 = -0.428, g_3 = 0.3388 \text{ and } g_4 = 0.4242, \Rightarrow g_4 > g_3 > g_1 > g_2.$$

Step 9. The final ranking is and the best alte $x_2 > x_1 > x_3 > x_4$ rnative is x_2 .

6. Conclusions

This paper developed a multi-criteria decision making method for neutrosophic multi-sets based on these given the outranking relations. In further research, we will develop different methods and compare the different methods on neutrosophic multi-sets. The contribution of this study is that the proposed approach is simple and convenient with regard to computing, and effective in decreasing the loss of evaluative information. More effective decision methods of this proposes a new outranking approach will be investigated in the near future and applied these concepts to engineering, game theory, multi-agent systems, decision-making and so on.

Funding: This research received no external funding

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Atanassov, K. Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, **1986**, 20 87–96.
2. Abdel-Basset, M., Mohamed, R., Zaied, A. E. N. H., Smarandache, F. (2019). A Hybrid Plithogenic Decision-Making Approach with Quality Function Deployment for Selecting Supply Chain Sustainability Metrics. *Symmetry*, **2019**, 11(7), 903.
3. Abdel-Basset, M., Nabeeh, N. A., El-Ghareeb, H. A., Aboelfetouh, A. Utilising neutrosophic theory to solve transition difficulties of IoT-based enterprises. *Enterprise Information Systems*, **2019**, 1-21.

4. Nabeeh, N. A., Abdel-Basset, M., El-Ghareeb, H. A., Aboelfetouh, A. Neutrosophic multi-criteria decision making approach for iot-based enterprises. *IEEE Access*, **2019**, 7, 59559-59574.
5. Abdel-Baset, M., Chang, V., Gamal, A. Evaluation of the green supply chain management practices: A novel neutrosophic approach. *Computers in Industry*, **2019**, 108, 210-220.
6. Abdel-Basset, M., Saleh, M., Gamal, A., Smarandache, F. An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number. *Applied Soft Computing*, **2019**, 77, 438-452.
7. Abdel-Baset, M., Chang, V., Gamal, A., Smarandache, F. An integrated neutrosophic ANP and VIKOR method for achieving sustainable supplier selection: A case study in importing field. *Computers in Industry*, **2019**, 106, 94-110.
8. Abdel-Basset, M., Manogaran, G., Gamal, A., Smarandache, F. A group decision making framework based on neutrosophic TOPSIS approach for smart medical device selection. *Journal of medical systems*, **2019**, 43(2), 38
9. Athar, K. A neutrosophic multi-criteria decision making method. *New Mathematics and Natural Computation*, **2014**, 10(02), 143-162.
10. S. Broumi and F. Smarandache, (2013). Several similarity measures of neutrosophic sets, *Neutrosophic Sets and Systems*, **2013**, 1(1) 54-62.
11. Chen, N., Xu, Z. Hesitant fuzzy ELECTRE II approach: a new way to handle multi-criteria decision making problems. *Information Sciences*, **2015**, 292, 175-197.
12. Deli, I., Broumi, S., Ali, M. Neutrosophic Soft Multi-Set Theory and Its Decision Making. *Neutrosophic Sets and Systems*, **2014**, 5, 65-76.
13. Deli, I. Refined Neutrosophic Sets and Refined Neutrosophic Soft Sets: Theory and Applications. *Handbook of Research on Generalized and Hybrid Set Structures and Applications for Soft Computing*, **2016**, 321-343.
14. Deli, I., Broumi S. Neutrosophic Soft Matrices and NSM-decision Making. *Journal of Intelligent and Fuzzy Systems*, **2015**, 28: 2233-2241.
15. Deli, I., S. Broumi, F. Smarandache, On neutrosophic refined sets and their applications in medical diagnosis, *Journal of New Theory*, **2015**, 6, 88-98.
16. Devi, K., S.P. Yadav, A multicriteria intuitionistic fuzzy group decision making for plant location selection with ELECTRE method, *Int. J. Adv. Manuf. Technol.* **2013**, 66 (912), 1219-1229.
17. Figueira, J.R., S. Greco, B. Roy, R. Slowinski, ELECTRE methods: main features and recent developments, *Handbook of Multicriteria Analysis*, vol. 103, Springer-Verlag, Berlin/Heidelberg, **2010**, pp. 51-89.
18. Hashemi, S. S., Hajiagha, S. H. R., Zavadskas, E. K., Mahdiraji, H. A. Multicriteria group decision making with ELECTRE III method based on interval-valued intuitionistic fuzzy information. *Applied Mathematical Modelling*, **2016**, 40(2), 1554-1564.
19. Karaaslan, F. Correlation Coefficient between Possibility Neutrosophic Soft Sets. *Math. Sci. Lett.* **2016**, 5/1, 71-74.
20. Karaaslan, F. Correlation coefficients of single-valued neutrosophic refined soft sets and their applications in clustering analysis. *Neural Computing and Applications*, **2016**, 1-13.
21. Mondal, K., Pramanik, S. Neutrosophic tangent similarity measure and its application to multiple attribute decision making. *Neutrosophic Sets and Systems*, **2015**, 9, 85-92.
22. Peng, J. J., Wang, J. Q., Zhang, H. Y., Chen, X. H. An outranking approach for multi-criteria decision-making problems with simplified neutrosophic sets. *Applied Soft Computing*, **2014**, 25, 336-346.
23. Peng, J. J., Wang, J. Q., Wang, J., Yang, L. J., Chen, X. H. An extension of ELECTRE to multi-criteria decision-making problems with multi-hesitant fuzzy sets. *Information Sciences*, **2015**, 307, 113-126.
24. Peng, J. J., Wang, J. Q., Wu, X. H. An extension of the ELECTRE approach with multi-valued neutrosophic information. *Neural Computing and Applications*, **2016**, 1-12.
25. Pramanik, S., Biswas, P., Giri, B. C. Hybrid vector similarity measures and their applications to multi-attribute decision making under neutrosophic environment. *Neural computing and Applications*, **2015**, 1-14.
26. Pramanik, S., Mondal, K. Cosine similarity measure of rough neutrosophic sets and its application in medical diagnosis. *Global Journal of Advanced Research*, **2015**, 2(1), 212-220.
27. Roy, B. The outranking approach and the foundations of ELECTRE methods. *Theory and decision*, **1991**, 31(1), 49-73.

28. Smarandache F. A Unifying Field in Logics Neutrosophy: Neutrosophic Probability, Set and Logic. Rehoboth: American Research Press. **1998**.
29. Smarandache, F. n-Valued Refined Neutrosophic Logic and Its Applications in Physics, Progress in Physics, **2013**, 4; 143–146.
30. Smarandache F. Neutrosophic set, a generalisation of the intuitionistic fuzzy sets. Int J Pure Appl Math , **2005**, 24:287-297.
31. Shen, F., Xu, J., Xu, Z. An outranking sorting method for multi-criteria group decision making using intuitionistic fuzzy sets. Information Sciences, **2016**, 334, 338–353.
32. Sahin, M., I. Deli, V. Ulucay, Jaccard Vector Similarity Measure of Bipolar Neutrosophic Set Based on Multi-Criteria Decision Making, International Conference on Natural Science and Engineering, **2016**, (ICNASE'16), March 19–20, Kilis.
33. Wang H, Smarandache FY, Q. Zhang Q, Sunderraman R (2010). Single valued neutrosophic sets. Multispace and Multistructure **2010**, 4:410–413.
34. Wang, J.Q., J.T. Wu, J. Wang, H.Y. Zhang, X.H. Chen, Interval-valued hesitant fuzzy linguistic sets and their applications in multi-criteria decision-making problems Original, Information Sciences, **2014**, 288/20 , 55–72.
35. Wang, J., Wang, J. Q., Zhang, H. Y., Chen, X. H. Multi-criteria decision-making based on hesitant fuzzy linguistic term sets: an outranking approach. Knowledge-Based Systems, **2015**, 86, 224–236.
36. M.C. Wu, T.Y. Chen, The ELECTRE multicriteria analysis approach based on Atanassovs intuitionistic fuzzy sets, Expert Syst. Appl. **2011**, 38 (10) , 12318-12327.
37. , J., Shen, F. A new outranking choice method for group decision making under Atanassovs interval-valued intuitionistic fuzzy environment. Knowledge-Based Systems, **2014**, 70, 177–188.
38. Ulucay, V., Deli, I., and Sahin, M. Similarity measures of bipolar neutrosophic sets and their application to multiple criteria decision making. Neural Computing and Applications, **2018**, 29(3), 739-748.
39. Ulucay, V., Deli, I., and Sahin, M. Intuitionistic trapezoidal fuzzy multi-numbers and its application to multi-criteria decision-making problems. Complex and Intelligent Systems, **2019**, 1-14.
40. Ulucay, V., Deli, I., and Sahin, M. Trapezoidal fuzzy multi-number and its application to multi-criteria decisionmaking problems. Neural Computing and Applications, **2018**, 30(5), 1469-1478.
41. Sahin, M., Olgun, N., Ulucay, V., Kargn, A., and Smarandache, F. A new similarity measure based on falsity value between single valued neutrosophic sets based on the centroid points of transformed single valued neutrosophic numbers with applications to pattern recognition. **2017**, Infinite Study.
42. Ulucay, V., Sahin, M., Olgun, N., and Kilicman, A. (2017). On neutrosophic soft lattices. Afrika Matematika, **2017**, 28(3-4), 379-388.
43. Ulucay, V., Kilic, A., Sahin, M., Deniz, H. A New Hybrid Distance-Based Similarity Measure for Refined Neutrosophic sets and its Application in Medical Diagnosis. MATEMATIKA: Malaysian Journal of Industrial and Applied Mathematics, **2019**, 35(1), 83-94.
44. Bakbak, D., Ulucay, V. Chapter Eight Multiple Criteria Decision Making in Architecture Based on Q-Neutrosophic Soft Expert Multiset. NEUTROSOPHIC TRIPLET STRUCTURES, **2019**, 90.
45. Bakbak, D., Ulucay, V., Sahin, M. Neutrosophic soft expert multiset and their application to multiple criteria decision making. Mathematics, **2019**, 7(1), 50.
46. Ulucay, V., Sahin, M. Neutrosophic Multigroups and Applications. MATHEMATICS, **2019**, 7(1).
47. Ulucay, V., Kilic, A., Yildiz, I., Sahin, M. A new approach for multi-attribute decision-making problems in bipolar neutrosophic sets. Neutrosophic Sets Syst, **2018**, 23(1), 142-159.
48. Yang, W. E., Wang, J. Q., Wang, X. F. An outranking method for multi-criteria decision making with duplex linguistic information. Fuzzy Sets and Systems, **2012**, 198, 20–33.
49. Ye, J. A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets, J. Intell. Fuzzy Syst. **2014**, 26 (5) 2459-2466.
50. Ye, J. Vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making. International Journal of Fuzzy Systems, **2014**, 16(2), 204–215.
51. Ye J. Improved correlation coefficients of single valued neutrosophic sets and interval neutrosophic sets for multiple attribute decision making. J Intell Fuzzy Syst **2014**, 27:2453–2462.
52. Ye, J., Zhang, Q. S. Single valued neutrosophic similarity measures for multiple attribute decision making. Neutrosophic Sets and Systems, **2014**, 2, 48–54.
53. Ye, S., and J. Ye, Dice Similarity Measure between Single Valued Neutrosophic Multisets and Its Application in Medical Diagnosis, Neutrosophic Sets and Systems, **2014**, 6, 49–54.

54. J., and J. Fub, Multi-period medical diagnosis method using a single-valued neutrosophic similarity measure based on tangent function, computer methods and programs in biomedicine doi:10.1016/j.cmpb.2015.10.002.
55. Ye,J., Single-valued neutrosophic similarity measures based on cotangent function and their application in the fault diagnosis of steam trbine, Soft Computing, DOI 10.1007/s00500-015- 1818-y.
56. Ye, S., Fu, J., Ye, J. Medical Diagnosis Using Distance-Based Similarity Measures of Single Valued Neutrosophic Multisets. Neutrosophic Sets and Systems, **2015**, 7, 47–52.
57. Wu, Y., Zhang, J., Yuan, J., Geng, S.,Zhang, H. Study of decision framework of offshore wind power station site selection based on ELECTRE-III under intuitionistic fuzzy environment: A case of China. Energy Conversion and Management, **2016**, 113, 66–81.
58. Zhang, H., Wang, J., Chen, X. An outranking approach for multi-criteria decision-making problems with intervalvalued neutrosophic sets. Neural Computing and Applications, **2015**,1–13.
59. L.A. Zadeh, (1965). Fuzzy Sets, Inform. and Control, **1965**,8: 338–353.

Received: Mar 15, 2019. Accepted: Nov 28, 2019