

10-15-2019

Vague –Valued Possibility Neutrosophic Vague Soft Expert Set Theory and Its Applications

Anjan Mukherjee

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Mukherjee, Anjan. "Vague –Valued Possibility Neutrosophic Vague Soft Expert Set Theory and Its Applications." *Neutrosophic Sets and Systems* 29, 1 (2020). https://digitalrepository.unm.edu/nss_journal/vol29/iss1/12

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu, lsloane@salud.unm.edu, sarahrk@unm.edu.



Vague –Valued Possibility Neutrosophic Vague Soft Expert Set Theory and Its Applications

Anjan Mukherjee

Department of Mathematics, Tripura University, Suryamaninagar, Agartala-799022, Tripura, India,

anjan2011_m@tripurauniv.in

*Correspondence: Anjan Mukherjee, anjan2011_m@tripurauniv.in

Abstract: In this paper, we first propose the concept of Vague-valued possibility Neutrosophic vague soft expert sets (VPNVSEsets in short). It is a combination of vague-valued possibility neutrosophic vague sets and soft expert sets. We also define its basic operations and study some related properties. Lastly an algorithm is proposed applied to the concept of vague-valued possibility Neutrosophic vague soft expert sets in hypothetical decision making problem. Here we associate the degree of belongingness degree of indeterminacy and non-belongingness of the elements of universe set with the vague –valued possibility set.

Keywords: Soft set, Neutrosophic soft expert set, Neutrosophic Vague soft set.

1. Introduction

Most real life problems involve data with a high level of uncertainty and imprecision. Traditionally, classical mathematical theories such as fuzzy mathematics, probability theories and interval mathematics are used to deal with uncertain and fuzziness. But all these theories have their difficulties and weakness as pointed out by Molodstov [14]. This led to the introduction of the theory of soft sets by Molodstov [14] in 1999. However, in order to handle the indeterminate and inconsistent information, neutrosophic set is defined [18]. The theory of vague set was first proposed by Gau and Buehrer [12]. It is an extension of fuzzy set theory. In 2010, W. Xu J. Ma, S. Wang and G. Hao, introduced Vague soft sets and their properties as a generalization of [12]. G. Selvachandran and A.R. sallah [19], introduced Possibility vague soft expert theory and its application in decision making.

In, [18] Smarandache talked about neutrosophic set theory. It is an important new mathematical tools for handling problems involving imprecise, indeterminacy and inconsistent data. Neutrosophic vague set was defined by S. Allehezaleh [2] in 2015. The concept of neutrosophic vague soft expert set was first introduced by Ashraf Al-Qurn and N. Hassan in 2016 [16]. It is the combination of neutrosophic vague sets and soft expert sets. In 2016, [19] G.Selvachandran and Abdul Razak Salleh introduced the concept of Possibility Intuitionistic Fuzzy Soft Expert and Its Application in Decision Making . In [15], Mukherjee and Sarkar introduced the concept of possibility interval

valued intuitionistic fuzzy soft expert theory which is a generalization of [20]. N. Hassan and A.Al-Quran [13], introduced Possibility Neutrosophic Vague soft expert set for decision under uncertainty. For further applications we refer the papers{[4],[5],[6],[7],[8],[9],[10],[11]}.

We first introduce the concept of vague-valued possibility neutrosophic vague soft expert set. It is a combination of vague-valued possibility neutrosophic vague set and soft expert set. The concept is to improve the reasonability of decision making in reality. Next we define its basic operation as a generalization of [13]. Finally we present an application of this concept in solving a decision making problem.

2. Preliminaries

We give some basic notions in neutrosophic vague set, neutrosophic vague soft set, soft expert set and neutrosophic soft expert set.

Definition 2.1. [2] A neutrosophic vague set A_{NV} (NVS in short) on the universe of discourse X written as $A_{NV} = \{ \langle x; \hat{T}A_{NV}(x); \hat{I}A_{NV}(x); \hat{F}A_{NV}(x) \rangle; x \in X \}$ whose truth-membership, indeterminacy-membership, and falsity-membership functions is defined as $\hat{T}A_{NV}(x) = [T^-, T^+]$, $\hat{I}A_{NV}(x) = [I^-, I^+]$ and $\hat{F}A_{NV}(x) = [F^-, F^+]$, where (1) $T^+ = 1 - F^-$, (2) $F^+ = 1 - T^-$ and (3) $-0 \leq T^- + I^- + F^- \leq 2^+$.

Definition 2.2. [2] If Ψ_{NV} is a NVS of the universe U , where $\forall u_i \in U$, $\hat{T}\Psi_{NV}(x) = [1, 1]$, $\hat{I}\Psi_{NV}(x) = [0, 0]$, $\hat{F}\Psi_{NV}(x) = [0, 0]$, then Ψ_{NV} is called a unit NVS, where $1 \leq i \leq n$. If Φ_{NV} is a NVS of the universe U , where $\forall u_i \in U$, $\hat{T}\Phi_{NV}(x) = [0, 0]$, $\hat{I}\Phi_{NV}(x) = [1, 1]$, $\hat{F}\Phi_{NV}(x) = [1, 1]$, then Φ_{NV} is called a zero NVS, where $1 \leq i \leq n$.

Definition 2.3. [2] Let A_{NV} and B_{NV} be two NVSs of the universe U . If $\forall u_i \in U$, (1) $\hat{T}A_{NV}(u_i) = \hat{T}B_{NV}(u_i)$, (2) $\hat{I}A_{NV}(u_i) = \hat{I}B_{NV}(u_i)$ and (3) $\hat{F}A_{NV}(u_i) = \hat{F}B_{NV}(u_i)$, then the NVS A_{NV} is equal to B_{NV} , denoted by $A_{NV} = B_{NV}$, where $1 \leq i \leq n$.

Definition 2.4. [2] Let A_{NV} and B_{NV} be two NVSs of the universe U . If $\forall u_i \in U$, (1) $\hat{T}A_{NV}(u_i) \leq \hat{T}B_{NV}(u_i)$, (2) $\hat{I}A_{NV}(u_i) \geq \hat{I}B_{NV}(u_i)$ and (3) $\hat{F}A_{NV}(u_i) \geq \hat{F}B_{NV}(u_i)$, then the NVS A_{NV} is included by B_{NV} , denoted by $A_{NV} \subseteq B_{NV}$, where $1 \leq i \leq n$.

Definition 2.5. [2] The complement of a NVS A_{NV} is denoted by A^c and is defined by

$$\begin{aligned} \hat{T}^c A_{NV}(x) &= [1 - T^+, 1 - T^-], \\ \hat{I}^c A_{NV}(x) &= [1 - I^+, 1 - I^-], \text{ and} \\ \hat{F}^c A_{NV}(x) &= [1 - F^+, 1 - F^-]. \end{aligned}$$

Definition 2.6. [2] The union of two NVSs A_{NV} and B_{NV} is a NVS C_{NV} , written as $C_{NV} = A_{NV} \cup B_{NV}$, whose truth-membership, indeterminacy-membership and false-membership functions are related to those of A_{NV} and B_{NV} given by

$$\begin{aligned} T_{C_{NV}}(x) &= [\max(T_{A_{NV}x}^-, T_{B_{NV}x}^-), \max(T_{A_{NV}x}^+, T_{B_{NV}x}^+)] \\ I_{C_{NV}}(x) &= [\min(I_{A_{NV}x}^-, I_{B_{NV}x}^-), \min(I_{A_{NV}x}^+, I_{B_{NV}x}^+)] \text{ and} \\ F_{C_{NV}}(x) &= [\min(F_{A_{NV}x}^-, F_{B_{NV}x}^-), \min(F_{A_{NV}x}^+, F_{B_{NV}x}^+)] \end{aligned}$$

Definition 2.7. [2] The intersection of two NVSs A_{NV} and B_{NV} is a NVS C_{NV} , written as $H_{NV} = A_{NV} \cap B_{NV}$, whose truth-membership, indeterminacy-membership and false-membership functions are related to those of A_{NV} and B_{NV} given by

$$\begin{aligned} T_{H_{NV}}(x) &= [\min(T_{A_{NV}x}^-, T_{B_{NV}x}^-), \min(T_{A_{NV}x}^+, T_{B_{NV}x}^+)] \\ I_{H_{NV}}(x) &= [\max(I_{A_{NV}x}^-, I_{B_{NV}x}^-), \max(I_{A_{NV}x}^+, I_{B_{NV}x}^+)] \text{ and} \end{aligned}$$

$$F_{HNV}(x) = [\max(F_{ANV_x}^-, F_{BNV_x}^-), \max(F_{ANV_x}^+, F_{BNV_x}^+)]$$

Definition 2.8. [17] Let U be an initial universal set. Let E be a set of parameters. Let $NV(U)$ denote the power set of all neutrosophic vague subsets of U and let $A \subseteq E$. A collection of pairs (\hat{F}, E) is called a neutrosophic vague soft set $\{NVSset\}$ over U , where \hat{F} is a mapping given by $\hat{F} : A \rightarrow NV(U)$.

Let U be a universe. E a set of parameters. X a set of experts (agents), and O a set of opinions. Let $Z = E \times X \times O$ and $A \subseteq Z$.

Definition 2.9. [3] A pair (F, A) is called a soft expert set over U , where F is a mapping given by $F : A \rightarrow P(U)$, where $P(U)$ denotes the power set of U .

Let U be a universe, E a set of parameters, X a set of experts (agents), and $O = \{1 = \text{agree}, 0 = \text{disagree}\}$ a set of opinions. Let $Z = E \times X \times O$ and $A \subseteq Z$.

Definition 2.10. [16] A pair (F, A) is called a neutrosophic soft expert set (NSES in short) over U , where F is a mapping given by $F : A \rightarrow PN(U)$, where $PN(U)$ denotes the power neutrosophic set of U .

Let U be a universe, E a set of parameters, X a set of experts (agents), and $O = \{1 = \text{agree}, 0 = \text{disagree}\}$ a set of opinions. Let $Z = E \times X \times O$ and $A \subseteq Z$.

Definition 2.11. [16] A pair (F, A) is called a neutrosophic vague soft expert set over U , where F is a mapping given by $F : A \rightarrow NV^u$, where NV^u denotes the power neutrosophic vague set of U .

Suppose $F : A \rightarrow NV^u$ is a function defined as $F(a) = F(a)(u), \forall u \in U$. For each $a_i \in A, F(a_i) = F(a_i)(u)$, where $F(a_i)$ represents the degree of belongingness, degree of indeterminacy and non-belongingness of the elements of U in $F(a_i)$. Hence $F(a_i)$ can be written as:

$$F(a_i) = \left\{ \frac{u_i}{F(a_i)(u_i)} \right\}, \text{ for } i = 1, 2, 3, \dots$$

Where $F(a_i)(u_i) = \langle [T_{F(a_i)}^-(u_i), T_{F(a_i)}^+(u_i)], [I_{F(a_i)}^-(u_i), I_{F(a_i)}^+(u_i)], [F_{F(a_i)}^-(u_i), F_{F(a_i)}^+(u_i)] \rangle$ and $T_{F(a_i)}^+(u_i) = 1 - F_{F(a_i)}^-(u_i), F_{F(a_i)}^+(u_i) = 1 - T_{F(a_i)}^-(u_i)$ with $[T_{F(a_i)}^-(u_i), T_{F(a_i)}^+(u_i)], [I_{F(a_i)}^-(u_i), I_{F(a_i)}^+(u_i)]$ and $[F_{F(a_i)}^-(u_i), F_{F(a_i)}^+(u_i)]$ representing the truth-membership function, indeterminacy-membership function and falsity-membership function of each of the elements $u_i \in U$, respectively.

Example 2.12 [16]. Suppose that a company produced new types of its products and wishes to take the opinion of some experts concerning these products. Let $U = \{u_1, u_2, u_3, u_4\}$ be a set of products. $E = \{e_1, e_2\}$ a set of decision parameters where $e_i (i = 1, 2)$ denotes the decision ‘‘easy to use,’’ and ‘‘quality,’’ respectively. Let $X = \{p, q\}$ be a set of experts. Suppose that the company has distributed a questionnaire to the two experts to make decisions on the company’s products, and we get the following:

$$F(e_1, p, 1) = \left\{ \frac{u_1}{\langle [0.2, 0.8]; [0.1, 0.3]; [0.2, 0.8] \rangle}, \frac{u_2}{\langle [0.1, 0.7]; [0.2, 0.5]; [0.3, 0.9] \rangle}, \frac{u_3}{\langle [0.5, 0.6]; [0.3, 0.7]; [0.4, 0.5] \rangle}, \frac{u_4}{\langle [0.8, 1]; [0.1, 0.2]; [0.0, 2] \rangle} \right\}$$

$$F(e_1, q, 1) = \left\{ \frac{u_1}{\langle [0.8, 0.9]; [0.3, 0.4]; [0.1, 0.2] \rangle}, \frac{u_2}{\langle [0.2, 0.4]; [0.2, 0.4]; [0.6, 0.8] \rangle}, \frac{u_3}{\langle [0.0, 5]; [0.5, 0.7]; [0.5, 1] \rangle}, \frac{u_4}{\langle [0.6, 0.7]; [0.2, 0.4]; [0.3, 0.4] \rangle} \right\}$$

$$F(e_2, p, 1) = \left\{ \frac{u_1}{\langle [0.3, 0.9]; [0.1, 0.3]; [0.1, 0.7] \rangle}, \frac{u_2}{\langle [0.2, 0.5]; [0.2, 0.5]; [0.5, 0.8] \rangle}, \frac{u_3}{\langle [0.6, 0.9]; [0.1, 0.7]; [0.1, 0.4] \rangle}, \frac{u_4}{\langle [0.2, 0.4]; [0.2, 0.2]; [0.6, 0.8] \rangle} \right\}$$

$$F(e_2, q, 1) = \left\{ \frac{u_1}{\langle [0.4, 0.6]; [0.1, 0.4]; [0.4, 0.6] \rangle}, \frac{u_2}{\langle [0.1, 0.3]; [0.2, 0.4]; [0.7, 0.9] \rangle}, \frac{u_3}{\langle [0.1, 0.5]; [0.5, 0.7]; [0.5, 0.9] \rangle}, \frac{u_4}{\langle [0.2, 0.7]; [0.2, 0.4]; [0.3, 0.8] \rangle} \right\}$$

$$F(e_1, p, 0) = \left\{ \frac{u_1}{\langle [0.2, 0.8]; [0.7, 0.9]; [0.2, 0.8] \rangle}, \frac{u_2}{\langle [0.3, 0.9]; [0.5, 0.8]; [0.1, 0.7] \rangle}, \frac{u_3}{\langle [0.4, 0.5]; [0.3, 0.7]; [0.5, 0.6] \rangle}, \frac{u_4}{\langle [0, 0.2]; [0.8, 0.9]; [0.8, 1] \rangle} \right\}$$

$$F(e_1, q, 0) = \left\{ \frac{u_1}{\langle [0.1, 0.7]; [0.7, 0.9]; [0.3, 0.9] \rangle}, \frac{u_2}{\langle [0.5, 0.8]; [0.5, 0.8]; [0.2, 0.5] \rangle}, \frac{u_3}{\langle [0.5, 1]; [0.3, 0.5]; [0, 0.5] \rangle}, \frac{u_4}{\langle [0.3, 0.4]; [0.6, 0.8]; [0.6, 0.7] \rangle} \right\}$$

$$F(e_2, p, 0) = \left\{ \frac{u_1}{\langle [0.2, 0.8]; [0.7, 0.9]; [0.2, 0.8] \rangle}, \frac{u_2}{\langle [0.3, 0.9]; [0.5, 0.8]; [0.1, 0.7] \rangle}, \frac{u_3}{\langle [0.1, 0.4]; [0.3, 0.9]; [0.6, 0.9] \rangle}, \frac{u_4}{\langle [0.6, 0.8]; [0.8, 0.8]; [0.2, 0.4] \rangle} \right\}$$

$$F(e_2, q, 0) = \left\{ \frac{u_1}{\langle [0.4, 0.6]; [0.6, 0.9]; [0.4, 0.6] \rangle}, \frac{u_2}{\langle [0.7, 0.9]; [0.6, 0.8]; [0.1, 0.3] \rangle}, \frac{u_3}{\langle [0.5, 0.9]; [0.3, 0.5]; [0.1, 0.5] \rangle}, \frac{u_4}{\langle [0.3, 0.8]; [0.6, 0.8]; [0.2, 0.7] \rangle} \right\}$$

The neutrosophic vague soft expert set (F, Z) is a parameterized family $\{F(e_i), i = 1, 2, 3, \dots\}$ of all neutrosophic vague sets of U and describes a collection of approximation of an object.

Definition 2.13. [16] The complement of a NVSE set (F, A) is denoted by $(F, A)^c$ and is defined by $(F, A)^c = (F^c, A)$ where $F^c: A \rightarrow NV^u$ is a mapping given by $F^c(\alpha) = \tilde{c}(F(\alpha)), \forall \alpha \in A$.

Where \tilde{c} is a neutrosophic vague complement.

Definition 2.14. [15] The union of two NVSE sets (F, A) and (G, B) over U , denoted by $(F, A) \tilde{\cup} (G, B)$, is a neutrosophic vague soft expert set (H, C) , where $C = A \cup B$ and $\forall \varepsilon \in C$,

$$(H, C) = \begin{cases} F(\varepsilon), & \text{if } \varepsilon \in A - B, \\ G(\varepsilon), & \text{if } \varepsilon \in B - A, \\ F(\varepsilon) \tilde{\cup} G(\varepsilon), & \text{if } \varepsilon \in A \cap B \end{cases} \quad \text{where } \tilde{\cup} \text{ denotes the union of the neutrosophic vague set}$$

Definition 2.15. [16] The intersection of two neutrosophic vague soft expert sets (F, A) and (G, B) over a universe U , is a neutrosophic vague soft expert set (H, C) , denoted by $(F, A) \tilde{\cap} (G, B)$ such that $C = A \cap B$ and $\forall \varepsilon \in C$

$$(H, C) = \begin{cases} F(\varepsilon), & \text{if } \varepsilon \in A - B, \\ G(\varepsilon), & \text{if } \varepsilon \in B - A, \\ F(\varepsilon) \tilde{\cap} G(\varepsilon), & \text{if } \varepsilon \in A \cap B \end{cases} \quad \text{where } \tilde{\cap} \text{ denotes the intersection of neutrosophic vague set.}$$

Definition 2.16 [16]. Let (F, A) and (G, B) be any two NVSE sets over a soft universe (U, Z) .

Then “ (F, A) AND (G, B) ” denoted $(F, A) \tilde{\wedge} (G, B)$ is defined by $(F, A) \tilde{\wedge} (G, B) = (H, A \times B)$, where $(H, A \times B) = H(\alpha, \beta)$, such that $H(\alpha, \beta) = F(\alpha) \cap G(\beta)$, for all $(\alpha, \beta) \in A \times B$, where \cap represents the basic intersection.

Definition 2.17 [16] Let (F, A) and (G, B) be any two neutrosophic vague soft expert sets over a soft universe (U, Z) .

Then “ (F, A) OR (G, B) ” denoted $(F, A) \tilde{\vee} (G, B)$ is defined by $(F, A) \tilde{\vee} (G, B) = (H, A \times B)$, where $(H, A \times B) = H(\alpha, \beta)$, such that $H(\alpha, \beta) = F(\alpha) \cup G(\beta)$, for all $(\alpha, \beta) \in A \times B$, where \cup represents the basic union.

Definition 2.18 [16]. Let U be a Universe. E a set of parameters, X a set of experts. $Q = \{1 = \text{agree}, 0 = \text{disagree}\}$ a set of opinions. Let $Z = E \times X \times Q$ and $A \subseteq Z$.

Let $U = \{u_1, u_2, \dots, u_n\}$ be a universal set of elements, let $E = \{e_1, e_2, e_3, \dots, e_m\}$ be a universal set of parameters. Let $X = \{x_1, x_2, \dots, x_i\}$ be a set of experts and let $Q = \{1 = \text{agree}, 0 = \text{disagree}\}$ be a set of opinions. Let $Z = E \times X \times Q$ and $A \subseteq Z$. Then the pair (U, Z) is called a soft universe. Let $F: Z \rightarrow NVSs(V)$, and p be a fuzzy subset of Z define by $p: Z \rightarrow I^U$, where I^U is the collection of all fuzzy subsets of U . Suppose $F_p: Z \rightarrow NVSs(U) \times I^U$ be a function define by $F_p = \{(F(Z)(u_i), P(Z)(u_i))\}$, for all $u_i \in U$. Then F_p is called a possibility neutrosophic vague soft expert set (denoted by PNVSES) over the soft universe (U, Z) . For each $z_i \in Z$, $F_p(z_i) = (F(z_i)(u_i), P(z_i)(u_i))$ where $F(z_i)$ represent the degree of belongingness degree of indeterminacy and non-belongingness of the elements of U in $F(z_i)$ and $P(z_i)$ represents the degree of possibility of belongingness of the elements of U in $F(z_i)$.

3. Vague-valued possibility neutrosophic vague soft expert set

In this section we introduce the definition of a vague-valued Possibility neutrosophic vague soft expert set (VPNVSE set).

Let U be a Universe. E a set of parameters. X a set of experts and $Q = \{1 = \text{agree}, 0 = \text{disagree}\}$ a set of opinions. Let $Z = E \times X \times Q$ and $A \subseteq Z$.

Definition 3.1. Let $U = \{u_1, u_2, \dots, u_n\}$ be a universal set of elements, let $E = \{e_1, e_2, e_3, \dots, e_m\}$ be a universal set of parameters. Let $X = \{x_1, x_2, \dots, x_i\}$ be a set of experts and let $Q = \{1 = \text{agree}, 0 = \text{disagree}\}$ be a set of opinions. Let $Z = E \times X \times Q$ and $A \subseteq Z$. Then the pair (U, Z) is called a soft universe. Let $F: Z \rightarrow NVSs(V)$, and p be a vague-valued subset of Z define by $p: Z \rightarrow V(U)$.

Suppose $F_p: Z \rightarrow NVSs(U) \times V(U)$ be a function define by $F_p = \{(F(Z)(u_i), P(Z)(u_i))\}$, $\forall u_i \in V$. Then F_p is called a vague-valued possibility neutrosophic vague soft expert set (denoted by VPNVSEs) over the soft universe (U, Z) . For each $z_i \in Z$, $F_p(z_i) = (F(z_i)(u_i), P(z_i)(u_i))$ where $F(z_i)$ represent the degree of belongingness degree of indeterminacy and non-belongingness of the elements of U in $F(z_i)$.

$$\text{So } F(z_i)(u_i) = \{[T_{F(z_i)}^-(u_i), T_{F(z_i)}^+(u_i)], [I_{F(z_i)}^-(u_i), I_{F(z_i)}^+(u_i)], [F_{F(z_i)}^-(u_i), F_{F(z_i)}^+(u_i)]\}$$

and $T_{F(z_i)}^+(u_i) = 1 - F_{F(z_i)}^-(u_i)$, $F_{F(z_i)}^+(u_i) = 1 - T_{F(z_i)}^-(u_i)$ with $[T_{F(z_i)}^-(u_i), T_{F(z_i)}^+(u_i)]$, $[I_{F(z_i)}^-(u_i), I_{F(z_i)}^+(u_i)]$, $[F_{F(z_i)}^-(u_i), F_{F(z_i)}^+(u_i)]$ representing the truth membership function indeterminacy membership function and fails membership function of each of the elements

$u_i \in U$ respectively. $P(z_i)$ represents the vague -value $[t_A(x), 1-f_A(x)]$, indicates that the exact grade of membership of x to A (which may be unknown but it is bounded by $t_A(x)$ and $1-f_A(x)$). Hence $F_p(z_i)$

can be written as $F_p(z_i) = \left\{ \left(\frac{u_i}{F(z_i)(u_i)} \right), P(z_i)(u_i) \right\}$ for $i = 1, 2, 3, \dots$. The $VPNVSEs(F_p, z)$ can be written simply as F_p . If $A \subseteq Z$, it is also possible to have a $VPNVSEs(F_p, A)$. For simplicity we take the set of opinion contains of only two values namely agree and disagree.

Suppose that a company produced new types of its products & wishes to take the opinion of some experts corresponding those products. Let $U = \{u_1, u_2, u_3\}$ be a set of products. $E = \{e_1, e_2\}$ a set of decision parameters. Here, e_i ($i=1,2$) denote the decision "easy to use" and "equality". Let $X = \{p, q\}$ be a set of experts. Suppose that the company has distributed questionnaire to, the two experts to make decisions on the company products. Then we have to following.

$F_p: Z \rightarrow VNVs(U) \times V(U)$ is a function then

$$F_p(e_1, p, 1) = \left\{ \left(\frac{u_1}{[0.2,0.8]; [0.1,0.3]; [0.2,0.8]}, [0.3,0.5], \left(\frac{u_2}{[0.1,0.7]; [0.2,0.5]; [0.3,0.9]}, [0.5,0.7], \left(\frac{u_3}{[0.5,0.6]; [0.3,0.7]; [0.4,0.5]}, [0.7,0.9] \right) \right\}$$

$$F_p(e_1, q, 1) = \left\{ \left(\frac{u_1}{[0.8,0.9]; [0.3,0.4]; [0.1,0.2]}, [0.4,0.6], \left(\frac{u_2}{[0.2,0.4]; [0.2,0.4]; [0.6,0.8]}, [0.6,0.8], \left(\frac{u_3}{[0.0,0.5]; [0.5,0.7]; [0.5,1]}, [0.8,1] \right) \right\}$$

$$F_p(e_2, p, 1) = \left\{ \left(\frac{u_1}{[0.3,0.9]; [0.1,0.3]; [0.1,0.7]}, [0.5,0.7], \left(\frac{u_2}{[0.2,0.5]; [0.2,0.5]; [0.5,0.8]}, [0.6,0.8], \left(\frac{u_3}{[0.6,0.9]; [0.1,0.7]; [0.1,0.4]}, [0.3,0.5] \right) \right\}$$

$$F_p(e_2, q, 1) = \left\{ \left(\frac{u_1}{[0.4,0.6]; [0.1,0.4]; [0.4,0.6]}, [0.2,0.4], \left(\frac{u_2}{[0.1,0.3]; [0.2,0.4]; [0.7,0.9]}, [0.4,0.6], \left(\frac{u_3}{[0.1,0.3]; [0.5,0.7]; [0.7,0.9]}, [0.7,0.9] \right) \right\}$$

$$F_p(e_1, p, 0) = \left\{ \left(\frac{u_1}{[0.2,0.8]; [0.7,0.9]; [0.2,0.8]}, [0.1,0.3], \left(\frac{u_2}{[0.3,0.9]; [0.5,0.8]; [0.1,0.7]}, [0.3,0.6], \left(\frac{u_3}{[0.4,0.5]; [0.3,0.7]; [0.5,0.6]}, [0.5,0.7] \right) \right\}$$

$$F_p(e_1, q, 0) = \left\{ \left(\frac{u_1}{[0.1,0.2]; [0.6,0.7]; [0.8,0.9]}, [0.8,0.9], \left(\frac{u_2}{[0.6,0.8]; [0.6,0.8]; [0.2,0.4]}, [0.6,0.8], \left(\frac{u_3}{[0.5,1]; [0.3,0.5]; [0.0,5]}, [0.3,0.7] \right) \right\}$$

$$F_p(e_2, p, 0) = \left\{ \left(\frac{u_1}{[0.1,0.7]; [0.7,0.9]; [0.3,0.9]}, [0.2,0.5], \left(\frac{u_2}{[0.5,0.8]; [0.5,0.8]; [0.2,0.5]}, [0.3,0.6], \left(\frac{u_3}{[0.1,0.4]; [0.3,0.9]; [0.6,0.9]}, [0.3,0.7] \right) \right\}$$

$$F_p(e_2, q, 0) = \left\{ \left(\frac{u_1}{[0.4,0.6]; [0.8,0.9]; [0.4,0.6]}, [0.2,0.5], \left(\frac{u_2}{[0.7,0.9]; [0.6,0.8]; [0.1,0.3]}, [0.4,0.6], \left(\frac{u_3}{[0.5,0.9]; [0.3,0.5]; [0.1,0.5]}, [0.5,0.7] \right) \right\}$$

Thus we have the VPNVSE set (F_p, Z) as follows:

$$(F_p, Z) = \{(e_1, p, 1) = \left\{ \left(\frac{u_1}{[0.2,0.8]; [0.1,0.3]; [0.2,0.8]}, [0.3,0.5], \left(\frac{u_2}{[0.1,0.7]; [0.2,0.5]; [0.3,0.9]}, [0.5,0.7], \left(\frac{u_3}{[0.5,0.6]; [0.3,0.7]; [0.4,0.5]}, [0.7,0.9] \right) \right\}$$

$$(e_2, p, 1) = \left\{ \left(\frac{u_1}{[0.3,0.9]; [0.1,0.3]; [0.1,0.7]}, [0.5,0.7], \left(\frac{u_2}{[0.2,0.5]; [0.2,0.5]; [0.5,0.8]}, [0.6,0.8], \left(\frac{u_3}{[0.6,0.9]; [0.1,0.7]; [0.1,0.4]}, [0.3,0.5] \right) \right\}$$

$$(e_1, q, 1) = \left\{ \left(\frac{u_1}{[0.8,0.9]; [0.3,0.4]; [0.1,0.2]} , [0.4,0.6], \left(\frac{u_2}{[0.2,0.4]; [0.2,0.4]; [0.6,0.8]} , [0.6,0.8], \left(\frac{u_3}{[0.0,0.5]; [0.5,0.7]; [0.5,1]} , [0.8,1] \right) \right) \right\}$$

$$(e_2, q, 1) = \left\{ \left(\frac{u_1}{[0.4,0.6]; [0.1,0.4]; [0.4,0.6]} , [0.2,0.4], \left(\frac{u_2}{[0.1,0.3]; [0.2,0.4]; [0.7,0.9]} , [0.4,0.6], \left(\frac{u_3}{[0.1,0.3]; [0.5,0.7]; [0.7,0.9]} , [0.7,0.9] \right) \right) \right\}$$

$$(e_1, p, 0) = \left\{ \left(\frac{u_1}{[0.2,0.8]; [0.7,0.9]; [0.2,0.8]} , [0.1,0.3], \left(\frac{u_2}{[0.3,0.9]; [0.5,0.8]; [0.1,0.7]} , [0.3,0.5], \left(\frac{u_3}{[0.4,0.5]; [0.3,0.7]; [0.5,0.6]} , [0.5,0.7] \right) \right) \right\}$$

$$(e_2, p, 0) = \left\{ \left(\frac{u_1}{[0.1,0.7]; [0.7,0.9]; [0.3,0.9]} , [0.2,0.5], \left(\frac{u_2}{[0.5,0.8]; [0.5,0.8]; [0.2,0.5]} , [0.3,0.6], \left(\frac{u_3}{[0.1,0.4]; [0.3,0.9]; [0.6,0.9]} , [0.3,0.7] \right) \right) \right\}$$

$$(e_1, q, 0) = \left\{ \left(\frac{u_1}{[0.1,0.2]; [0.6,0.7]; [0.8,0.9]} , [0.8,0.9], \left(\frac{u_2}{[0.6,0.8]; [0.6,0.8]; [0.2,0.4]} , [0.6,0.8], \left(\frac{u_3}{[0.5,1]; [0.3,0.5]; [0.0,5]} , [0.3,0.6] \right) \right) \right\}$$

$$(e_2, q, 0) = \left\{ \left(\frac{u_1}{[0.4,0.6]; [0.8,0.9]; [0.4,0.6]} , [0.2,0.3], \left(\frac{u_2}{[0.7,0.9]; [0.6,0.8]; [0.1,0.3]} , [0.4,0.6], \left(\frac{u_3}{[0.5,0.9]; [0.3,0.5]; [0.1,0.5]} , [0.5,0.7] \right) \right) \right\}$$

The collection (F_p, Z) is a VPNVSE set over the soft inverse (U, Z) .

Definition 3.3: Let (F_p, A) and (G_q, B) be two VPNVSE sets over the soft inverse (U, Z) then (F_p, A) is a VPNVSE sub set of (G_q, B) if $A \subseteq B$ and for all $\varepsilon \in A$ the following conditions are satisfied.

- (i) $p(\varepsilon)$ is a vague sub set of $q(\varepsilon)$.
- (ii) $F(\varepsilon)$ is a neutrosophic vague soft set of $G(\varepsilon)$.

It is denoted by $(F_p, A) \subseteq (G_q, A)$. Then (G_q, A) is called a vague-valued possibility neutrosophic soft expert superset of (F_p, A) .

Definition 3.4. Let (F_p, A) and (G_q, B) be two VPNVSE sets over the soft inverse (U, Z) then (F_p, A) equal to (G_q, B) if for all $\varepsilon \in A$ the following holds

- (i) $p(\varepsilon) = q(\varepsilon)$.
- (ii) $F(\varepsilon) = G(\varepsilon)$.

In other words $(F_p, A) = (G_q, B)$ if (F_p, A) is a subset of (G_q, B) and (G_q, B) is a subset of (F_p, A) .

4. Basic Operations On Vague-Valued Possibility Neutrosophic Soft Expert Sets.

Now we introduce some basic operations on PNVSE sets. These are ‘complement’ Union & intersection. Then we study some of the properties related to these operations.

Definition 4.1 Let (F_p, A) be a VPNVSE set over the soft universe (U, Z) then the complement of (F_p, A) denoted by $(F_p, A)^c$ is defined as

$$(F_p, A)^c = (\bar{c}(F\alpha), c(P(\alpha))) \forall \alpha \in A.$$

Where \bar{c} a neutrosophic vague complement and c is a Vague-valued set complement.

If A be a vague set over the universe U , then

$A = \{[x, t_A(x), 1-f_A(x)]: x \in V\}$ in this definition $t_A(x)$ is a lower bound on the grade of membership of x to A derived from the evidence for x and $f_A(x)$ is a lower bound on the negation of x to A derived from the evidence against x . The vague value $[t_A(x), 1-f_A(x)]$ indicates that the exact grade of membership of x to A may be unknown, but it is bounded by $t_A(x)$ & $1-f_A(x)$. It is to be noted that every fuzzy set α correspondence to the following vague set: $\alpha = \{(x, [\alpha(x), 1-\alpha(x)]: x \in U\}$ thus the notion of vague sets is a generalization of fuzzy sets. The complement of the vague set A is $A^c = \{[x, f_A(x), 1-t_A(x)]: x \in U\}$.

Example 4.2: Consider the VPNVSE (F_p, A) over a soft universe (U, Z) as an example 3.2. Now by definition 4.1 $(F_p, A)^c$ is given as follows:

$$\begin{aligned} (F_p, z)^c = \{(e_1, p, 1) = \{(\frac{u_1}{[0.2,0.8]; [0.7,0.9]; [0.2,0.8]}, [0.5, 0.7]), (\frac{u_2}{[0.3,0.9]; [0.5,0.8]; [0.1,0.7]}, [0.3, 0.5]), \\ (\frac{u_3}{[0.4,0.5]; [0.3,0.7]; [0.5,0.6]}, [0.1, 0.3])\}, (e_2, p, 1) = \{(\frac{u_1}{[0.1,0.7]; [0.7,0.9]; [0.3,0.9]}, [0.3, 0.5]), (\frac{u_2}{[0.5,0.8]; [0.5,0.8]; [0.2,0.5]}, \\ [0.2, 0.4]), (\frac{u_3}{[0.1,0.4]; [0.3,0.9]; [0.6,0.9]}, [0.5, 0.7])\}, (e_1, q, 1) = \{(\frac{u_1}{[0.1,0.2]; [0.6,0.7]; [0.8,0.9]}, [0.4, 0.6]), \\ (\frac{u_2}{[0.6,0.8]; [0.6,0.8]; [0.2,0.4]}, [0.2, 0.4]), (\frac{u_3}{[0.5,1]; [0.3,0.5]; [0.0,0.5]}, [0.0, 0.2])\}, (e_2, q, 1) = \{(\frac{u_1}{[0.4,0.6]; [0.6,0.9]; [0.4,0.6]}, [0.6, \\ 0.8]), (\frac{u_2}{[0.7,0.9]; [0.6,0.8]; [0.1,0.3]}, [0.4, 0.6]), (\frac{u_3}{[0.7,0.9]; [0.3,0.5]; [0.1,0.3]}, [0.1, 0.3])\}, (e_1, p, 0) = \\ \{(\frac{u_1}{[0.2,0.8]; [0.1,0.3]; [0.2,0.8]}, [0.7, 0.9]), (\frac{u_2}{[0.1,0.7]; [0.2,0.5]; [0.3,0.9]}, [0.5, 0.7]), (\frac{u_3}{[0.5,0.6]; [0.3,0.7]; [0.4,0.5]}, [0.3, 0.5])\}, \\ (e_2, p, 0) = \{(\frac{u_1}{[0.3,0.9]; [0.1,0.3]; [0.1,0.7]}, [0.5, 0.8]), (\frac{u_2}{[0.2,0.5]; [0.2,0.5]; [0.5,0.8]}, [0.4, 0.7]), (\frac{u_3}{[0.6,0.9]; [0.1,0.7]; [0.1,0.4]}, \\ [0.3, 0.7])\}, (e_1, q, 0) = \{(\frac{u_1}{[0.8,0.9]; [0.3,0.4]; [0.1,0.2]}, [0.1, 0.2]), (\frac{u_2}{[0.2,0.4]; [0.2,0.4]; [0.6,0.8]}, [0.2, 0.4]), \\ (\frac{u_3}{[0.0,5]; [0.5,0.7]; [0.5,1]}, [0.4, 0.7])\}, (e_2, q, 0) = \{(\frac{u_1}{[0.4,0.6]; [0.1,0.2]; [0.4,0.6]}, [0.5, 0.8]), (\frac{u_2}{[0.1,0.3]; [0.2,0.4]; [0.7,0.9]}, [0.4, \\ 0.6]), (\frac{u_3}{[0.1,0.5]; [0.5,0.7]; [0.5,0.9]}, [0.3, 0.5])\} \end{aligned}$$

Proposition 4.3: Let (F_p, A) be a VPNVSE set over the soft universe (U, Z) . Here, $(F_p, A) = (F(e), p(e))$ then $((F_p, A)^c)^c = (F_p, A)$.

Proof: Let $(F_p, A)^c = (G_q, B)$ then by definition $(G_q, B) = (G(e), q(e))$

$G(e) = \bar{C}(F(e))$ and $q(e) = C(p(e))$. Where \bar{c} a neutrosophic vague complement and c is a Vague-valued set complement.

So it follows that

$$\begin{aligned} (G_q, B)^c &= \{\bar{C}(G(e)), C(q(e))\} \\ &= \{\bar{C}(\bar{C}(F(e))), C(C(p(e)))\} \\ &= (F(e), p(e)) = (F_p, A) \\ ((F(e), p(e))^c)^c &= (F_p, A) \end{aligned}$$

Definition 4.4: Let (F_p, A) and (G_q, B) be two V PNVSE set over a soft universe (U, Z) then the intersection of (F_p, A) and (G_q, B) denoted by $(F_p, A) \bar{\cap} (G_q, B)$ is a VPNVSE set defined as $(F_p, A) \bar{\cap} (G_q, B) = (H_r, C)$,

where $C = A \cap B$ and

$$r(\alpha) = p(\alpha) \cap q(\alpha) \quad \forall \alpha \in C$$

$$H(\alpha) = F(\alpha) \bar{\cap} G(\alpha) \quad \forall \alpha \in C$$

$$\text{And } H(\alpha) = \begin{cases} F(\alpha) & \text{if } \alpha \in A - B \\ G(\alpha) & \text{if } \alpha \in B - A \\ F(\alpha) \cap G(\alpha) & \text{if } \alpha \in A \cap B \end{cases}$$

Definition 4.5 Let (F_p, A) and (G_q, B) be two VPNVSE sets over a soft universe (U, Z) . Then the union of (F_p, A) and (G_q, B) denoted by $(F_p, A) \bar{\cup} (G_q, B)$ is a PNVSE set defined as $(F_p, A) \bar{\cup} (G_q, B) = (H_r, C)$, where $C = A \cup B$ and

$$r(\alpha) = p(\alpha) \cup q(\alpha) \quad \forall \alpha \in C$$

$$H(\alpha) = F(\alpha) \bar{\cup} G(\alpha) \quad \forall \alpha \in C$$

$$\text{And } H(\alpha) = \begin{cases} F(\alpha) & \text{if } \alpha \in A - B \\ G(\alpha) & \text{if } \alpha \in B - A \\ F(\alpha) \bar{\cup} G(\alpha) & \text{if } \alpha \in A \cup B \end{cases}$$

5. Application of vague-valued possibility neutrosophic vague soft expert in a decision making problem

A company is looking to have a person to fill the vacancy for a position in their company. Out of all the candidates were short listed - The three candidates form the universe of the element $U = \{u_1, u_2, u_3\}$ were short listed out of all candidates. The hiring committee consists of hiring manager, head of the department and HR director of the firm . The committee is represented by the set $X = \{x, y, z\}$ (a set of experts), while the set $Q = \{1 = \text{agree}, 0 = \text{disagree}\}$ represents the set of opinions of the hiring committee members. The hiring committee consider a set of parameters $E = \{e_1, e_2, e_3, e_4\}$. The

parameters e_i ($i = 1, 2, 3, 4$) represents the characteristic or qualities that the candidates are assessed on namely “experience”, “academic qualifications”, “attitude towards the professionalism” and “technical knowledge” respectively. After finishing the interview of all the candidates and going through their certificates and other supporting papers. The hire committee constitutes the VPNVSE set (F_p, z) as follows:

$$\begin{aligned}
 (F_p, z) = & \{(e_1, x, 1) = \left\{ \left(\frac{u_1}{[0.2,0.8]; [0.1,0.3]; [0.2,0.8]} , [0.3, 0.5] \right), \left(\frac{u_2}{[0.1,0.7]; [0.2,0.5]; [0.3,0.9]} , [0.5, 0.7] \right), \right. \\
 & \left. \left(\frac{u_3}{[0.5,0.6]; [0.3,0.7]; [0.4,0.5]} , [0.7, 0.9] \right) \right\}, (e_2, x, 1) = \left\{ \left(\frac{u_1}{[0.3,0.9]; [0.1,0.3]; [0.1,0.7]} , [0.5, 0.7] \right), \left(\frac{u_2}{[0.2,0.5]; [0.2,0.5]; [0.5,0.8]} , \right. \right. \\
 & \left. \left. [0.6, 0.8] \right), \left(\frac{u_3}{[0.6,0.9]; [0.1,0.7]; [0.1,0.4]} , [0.3, 0.5] \right) \right\}, (e_3, x, 1) = \left\{ \left(\frac{u_1}{[0.2,0.7]; [0.5,0.7]; [0.3,0.8]} , [0.3, 0.5] \right), \right. \\
 & \left. \left(\frac{u_2}{[0.1,0.7]; [0.4,0.5]; [0.3,0.9]} , [0.2, 0.5] \right), \left(\frac{u_3}{[0.2,0.6]; [0.3,0.5]; [0.4,0.8]} , [0.4, 0.6] \right) \right\}, (e_4, x, 1) = \left\{ \left(\frac{u_1}{[0.2,0.3]; [0.4,0.6]; [0.7,0.8]} , \right. \right. \\
 & \left. \left. [0.5, 0.7] \right), \left(\frac{u_2}{[0.1,0.3]; [0.2,0.5]; [0.7,0.9]} , [0.3, 0.6] \right), \left(\frac{u_3}{[0.3,0.4]; [0.4,0.6]; [0.6,0.7]} , [0.6, 0.8] \right) \right\}, (e_1, y, 1) = \\
 & \left\{ \left(\frac{u_1}{[0.8,0.9]; [0.3,0.4]; [0.1,0.2]} , [0.4, 0.6] \right), \left(\frac{u_2}{[0.2,0.4]; [0.2,0.4]; [0.6,0.8]} , [0.6, 0.8] \right), \left(\frac{u_3}{[0.0,0.5]; [0.5,0.7]; [0.5,1]} , [0.8, 1] \right) \right\}, (e_2, \\
 & y, 1) = \left\{ \left(\frac{u_1}{[0.4,0.6]; [0.1,0.4]; [0.4,0.6]} , [0.2, 0.4] \right), \left(\frac{u_2}{[0.1,0.3]; [0.2,0.4]; [0.7,0.9]} , [0.4, 0.6] \right), \left(\frac{u_3}{[0.1,0.3]; [0.5,0.7]; [0.7,0.9]} , [0.7, \right. \right. \\
 & \left. \left. 0.9] \right) \right\}, (e_3, y, 1) = \left\{ \left(\frac{u_1}{[0.2,0.5]; [0.4,0.6]; [0.5,0.8]} , [0.3, 0.5] \right), \left(\frac{u_2}{[0.1,0.3]; [0.2,0.4]; [0.7,0.9]} , [0.6, 0.8] \right), \right. \\
 & \left. \left(\frac{u_3}{[0.4,0.5]; [0.8,0.9]; [0.5,0.6]} , [0.5, 0.7] \right) \right\}, (e_4, y, 1) = \left\{ \left(\frac{u_1}{[0.3,0.5]; [0.5,0.7]; [0.5,0.7]} , [0.5, 0.7] \right), \left(\frac{u_2}{[0.5,0.7]; [0.9,1]; [0.3,0.5]} , \right. \right. \\
 & \left. \left. [0.2, 0.5] \right), \left(\frac{u_3}{[0.6,0.9]; [0.2,0.3]; [0.1,0.4]} , [0.3, 0.6] \right) \right\}, (e_1, z, 1) = \left\{ \left(\frac{u_1}{[0.1,0.4]; [0.3,0.6]; [0.6,0.9]} , [0.4, 0.7] \right), \right. \\
 & \left. \left(\frac{u_2}{[0.5,0.7]; [0.2,0.5]; [0.3,0.5]} , [0.3, 0.5] \right), \left(\frac{u_3}{[0.2,0.5]; [0.4,0.7]; [0.5,0.8]} , [0.7, 0.9] \right) \right\}, (e_2, z, 1) = \left\{ \left(\frac{u_1}{[0.1,0.5]; [0.4,0.6]; [0.5,0.9]} , \right. \right. \\
 & \left. \left. [0.4, 0.5] \right), \left(\frac{u_2}{[0.6,0.7]; [0.3,0.5]; [0.3,0.4]} , [0.3, 0.5] \right), \left(\frac{u_3}{[0.0,0.1]; [0.2,0.4]; [0.9,1]} , [0.7, 0.9] \right) \right\}, (e_3, z, 1) = \\
 & \left\{ \left(\frac{u_1}{[0.3,0.5]; [0.5,0.7]; [0.5,0.7]} , [0.1, 0.3] \right), \left(\frac{u_2}{[0.3,0.4]; [0.5,0.6]; [0.6,0.7]} , [0.3, 0.7] \right), \left(\frac{u_3}{[0.4,0.6]; [0.3,0.5]; [0.4,0.6]} , [0.7, 0.9] \right) \right\}, \\
 & (e_4, z, 1) = \left\{ \left(\frac{u_1}{[0.1,0.5]; [0.3,0.7]; [0.5,0.9]} , [0.4, 0.7] \right), \left(\frac{u_2}{[0.5,0.7]; [0.1,0.2]; [0.3,0.5]} , [0.3, 0.7] \right), \left(\frac{u_3}{[0.4,0.5]; [0.7,0.8]; [0.5,0.6]} , \right. \right. \\
 & \left. \left. [0.7, 1] \right) \right\}, (e_1, x, 0) = \left\{ \left(\frac{u_1}{[0.2,0.8]; [0.7,0.9]; [0.2,0.8]} , [0.1, 0.3] \right), \left(\frac{u_2}{[0.3,0.9]; [0.5,0.8]; [0.1,0.7]} , [0.3, 0.5] \right), \right. \\
 & \left. \left(\frac{u_3}{[0.4,0.5]; [0.3,0.7]; [0.5,0.6]} , [0.5, 0.7] \right) \right\}, (e_2, x, 0) = \left\{ \left(\frac{u_1}{[0.1,0.7]; [0.7,0.9]; [0.3,0.9]} , [0.2, 0.5] \right), \left(\frac{u_2}{[0.5,0.8]; [0.5,0.8]; [0.2,0.5]} , \right. \right. \\
 & \left. \left. [0.3, 0.6] \right), \left(\frac{u_3}{[0.1,0.4]; [0.3,0.9]; [0.6,0.9]} , [0.3, 0.7] \right) \right\}, (e_3, x, 0) = \left\{ \left(\frac{u_1}{[0.2,0.4]; [0.3,0.6]; [0.6,0.8]} , [0.3, 0.5] \right), \right. \\
 & \left. \left(\frac{u_2}{[0.5,0.8]; [0.3,0.6]; [0.2,0.5]} , [0.2, 0.5] \right), \left(\frac{u_3}{[0.4,0.7]; [0.7,0.8]; [0.3,0.6]} , [0.6, 0.9] \right) \right\}, (e_4, x, 0) = \left\{ \left(\frac{u_1}{[0.3,0.5]; [0.7,0.9]; [0.5,0.7]} , \right. \right. \\
 & \left. \left. [0.3, 0.5] \right), \left(\frac{u_2}{[0.5,0.7]; [0.8,0.9]; [0.3,0.5]} , [0.4, 0.7] \right), \left(\frac{u_3}{[0.5,1]; [0.3,0.5]; [0.5]} , [0.3, 0.6] \right) \right\}, (e_1, y, 0) =
 \end{aligned}$$

$$\{(\frac{u_1}{[0.3,0.5]; [0.5,0.8]; [0.5,0.7]}, [0.1, 0.3]), (\frac{u_2}{[0.1,0.3]; [0.4,0.6]; [0.7,0.9]}, [0.3, 0.5]), (\frac{u_3}{[0.4,0.6]; [0.7,0.9]; [0.4,0.6]}, [0.4, 0.7])\},$$

$$(e_2, y, 0) = \{(\frac{u_1}{[0.1,0.2]; [0.6,0.7]; [0.8,0.9]}, [0.8, 0.9]), (\frac{u_2}{[0.6,0.8]; [0.6,0.8]; [0.2,0.4]}, [0.6, 0.8]), (\frac{u_3}{[0.5,1]; [0.3,0.5]; [0,0.5]}, [0.3,$$

$$0.6])\}, (e_3, y, 0) = \{(\frac{u_1}{[0.4,0.6]; [0.8,0.9]; [0.4,0.6]}, [0.2, 0.5]), (\frac{u_2}{[0.7,0.9]; [0.6,0.8]; [0.1,0.3]}, [0.4, 0.6]),$$

$$(\frac{u_3}{[0.5,0.9]; [0.3,0.5]; [0.1,0.5]}, [0.5, 0.7])\}, (e_4, y, 0) = \{(\frac{u_1}{[0.3,0.5]; [0.6,0.8]; [0.5,0.7]}, [0.4, 0.6]), (\frac{u_2}{[0.1,0.3]; [0.4,0.6]; [0.7,0.9]}, [0.4,$$

$$0.7]), (\frac{u_3}{[0.2,0.6]; [0.5,0.7]; [0.4,0.8]}, [0.1, 0.4])\}, (e_1, z, 0) = \{(\frac{u_1}{[0.2,0.5]; [0.4,0.7]; [0.5,0.8]}, [0.4, 0.8]),$$

$$(\frac{u_2}{[0.2,0.4]; [0.3,0.5]; [0.6,0.8]}, [0.4, 0.7]), (\frac{u_3}{[0.2,0.7]; [0.5,0.7]; [0.3,0.8]}, [0.1, 0.5])\}, (e_2, z, 0) = \{(\frac{u_1}{[0.4,0.5]; [0.5,1]; [0.5,0.6]}, [0.5,$$

$$0.8]), (\frac{u_2}{[0.1,0.4]; [0.3,0.5]; [0.6,0.9]}, [0.5, 0.8]), (\frac{u_3}{[0.3,0.5]; [0.7,0.9]; [0.5,0.7]}, [0.2, 0.5])\}, (e_3, z, 0) =$$

$$\{(\frac{u_1}{[0.5,0.9]; [0.6,0.8]; [0.1,0.5]}, [0.4, 0.7]), (\frac{u_2}{[0.2,0.5]; [0.5,1]; [0.5,0.8]}, [0.2, 0.4]), (\frac{u_3}{[0.4,0.7]; [0.6,0.8]; [0.3,0.6]}, [0.6, 0.8])\}, (e_4,$$

$$z, 0) = \{(\frac{u_1}{[0.3,0.5]; [0.6,0.8]; [0.5,0.7]}, [0.4, 0.6]), (\frac{u_2}{[0.1,0.4]; [0.3,0.5]; [0.6,0.9]}, [0.6, 0.8]), (\frac{u_3}{[0.6,0.8]; [0.5,0.7]; [0.2,0.4]}, [0.5,$$

$$0.7])\}.$$

The collection (F_p, z) is a VPNVSE set over the soft universe (U, Z) . The VPNVSE set (F_p, Z) is used together with an algorithm to solve the decision making problem. The algorithm given below is taken by the committee to determine the most suitable candidate to be hired for the position. The sets of algorithm are as follows:

Step 1: Input the VPNVSE set (F_p, Z) .

Step 2: Calculate the value of $\alpha_{F(a_i)}(u_i) = T_{F(a_i)}^-(u_i) - F_{F(a_i)}^-(u_i)$ for interval truth-membership part $[T_{F(a_i)}^-(u_i), T_{F(a_i)}^+(u_i)]$, where $T_{F(a_i)}^+(u_i) = 1 - F_{F(a_i)}^-(u_i)$, for each element $u_i \in U$.

Step 3: Calculate the arithmetic overage $\beta_{F(a_i)}(u_i)$ of the end points of the interval indeterminacy membership part $[I_{F(a_i)}^-(u_i), I_{F(a_i)}^+(u_i)]$, for each element $u_i \in U$.

Step 4: Find the value of $\gamma_{F(a_i)}(u_i) = F_{F(a_i)}^-(u_i) - T_{F(a_i)}^-(u_i)$ for interval falsity-membership part $[F_{F(a_i)}^-(u_i), F_{F(a_i)}^+(u_i)]$, where $F_{F(a_i)}^+(u_i) = 1 - T_{F(a_i)}^-(u_i)$, for each element $u_i \in U$.

Step 5: Find $\alpha_{F(a_i)}(u_i) - \beta_{F(a_i)}(u_i) - \gamma_{F(a_i)}(u_i)$ for each element $u_i \in U$.

Step 6: Find the higher numerical grade from the agree-PNVSE set & disagree-PNVSE set.

Step 7: Take the arithmetic average of $[t_A, 1-f_A]$ of the set corresponding vague set associated with the Neutrosophic vague soft set.

Step 8: Find the higher numerical grade for the average vague set value for the highest agree-VPNVSE set & disagree-VPNVSE set

Step 9: Compute the score of each element $u_i \in U$ by taking the sum of the product of the maximum numerical grade (λ_i) with the corresponding average numerical value of vague set μ_i for the agree-VPNVSE set and disagree-VPNVSE set by A_i & D_i respectively.

Step 10: Find the value $r_i = A_i - D_i$, for each element $u_i \in U$.

Step 11: Determine the values of highest scores = $\max u_i \in U \{r_i\}$. Then the decision is to choose element u_i as optimal or best solution if there are more than one element.

Table-1 Value of $\alpha_{F(a_i)}(u_i)$, $\beta_{F(a_i)}(u_i)$, $\gamma_{F(a_i)}(u_i)$ The value of $\alpha_{F(a_i)}(u_i) - \beta_{F(a_i)}(u_i) - \gamma_{F(a_i)}(u_i)$ & the average of the vague set corresponding to the highest numerical grade

	u1	u2	u3		u1	u2	u3
(e1, x, 1)	0, 0.2, 0	-0.2, 0.35, 0.2	0.1, 0.5, -0.1	(e1, x, 0)	0, 0.8, 0	0.2, 0.65, -0.2	-0.1, 0.5, 0.1
	-0.2, (0.4)	-0.75, (0.6)	-0.3, (0.8)		-0.8, (0.2)	-0.25, (0.4)	-0.7, (0.6)
(e2, x, 1)	0.2, 0.2, -0.2	-0.3, 0.35, 0.3	0.5, 0.4, -0.5	(e2, x, 0)	-0.2, 0.8, 0.2	0.3, 0.65, -0.3	-0.5, 0.6, 0.5
	0.2, (0.6)	-0.85, (0.35)	0.6, (0.4)		-1.2, (0.35)	-0.05, (0.45)	-1.6, (0.5)
(e3, x, 1)	-0.5, 0.5, 0.5	-0.6, 0.35, 0.6	-0.3, 0.5, 0.3	(e3, x, 0)	-0.4, 0.45, 0.4	0.3, 0.45, -0.3	0.1, 0.75, -0.1
	-1.5, (0.6)	-1.55, (0.45)	-0.8, (0.5)		-1.25, (0.4)	0.15, (0.35)	0.55, (0.75)
(e4, x, 1)	-0.5, 0.5, 0.5	-0.6, 0.35, 0.6	-0.3, 0.5, 0.3	(e4, x, 0)	-0.2, 0.8, 0.2	0.2, 0.85, -0.2	0.5, 0.4, -0.5
	-1.5, (0.6)	-1.55, (0.45)	-1.1, (0.7)		-1.20, (0.4)	-0.45, (0.55)	0.6, (.45)
(e1, y, 1)	0.7, 0.35, -0.7	-0.4, 0.3, 0.4	-0.5, 0.6, 0.5	(e1, y, 0)	-0.2, 0.8, 0.2	-0.6, 0.5, 0.6	0, 0.8, 0
	1.05, (0.5)	-1.1, (0.7)	-0.6, (0.9)		-1.05, (0.2)	-1.7, (0.4)	-0.8, (0.55)
(e2, y, 1)	0, 0.25, 0	-0.6, 0.3, 0.6	-0.6, 0.6, 0.6	(e2, y, 0)	-0.7, 0.65, 0.7	0.4, 0.7, -0.4	0.5, 0.4, -0.5
	-0.25, (0.3)	-1.5, (0.5)	-1.8, (0.8)		-2.05, (0.85)	0.1, (0.7)	0.6, (0.45)
(e3, y, 1)	-0.3, 0.5, 0.3	-0.6, 0.3, 0.6	-0.1, 0.85, 0.1	(e3, y, 0)	0, 0.85, 0	0.6, 0.7, -0.6	0.5, 0.4, -0.5

	-1.1, (0.4)	-1.5, (0.7)	-1.05, (0.6)		-0.85,(0.35)	0.5,(0.5)	0.6,(0.45)
(e ₄ , y, 1)	-0.2,0.6,0.2	0.2,0.95,-0.2	0.5,0.25,-0.5	(e ₄ , y, 0)	-0.2,0.7,0.2	-0.6,0.5,0.6	-0.2,0.6,0.2
	-1.0,(0.6)	-0.55,(0.35)	0.75,(0.45)		-1.1,(0.5)	-1.7,(0.55)	-1.0,(0.25)
(e ₁ , z, 1)	-0.5,0.45,0.5	0.2,0.35,-0.2	-0.3,0.55,0.3	(e ₁ , z, 0)	-0.3,0.55,0.3	-0.4,0.35,0.4	-0.1,0.6,0.1
	-1.45,(0.55)	0.5,(0.4)	-1.1,(0.8)		-1.15,(0.6)	-1.15,(0.55)	-0.8,(0.3)
(e ₂ , z, 1)	-0.4,0.5,0.4	0.3,0.4,-0.3	-0.9,0.3,0.9	(e ₂ , z, 0)	-0.1,0.75,0.1	-0.5,0.4,0.5	-0.2,0.8,0.2
	-1.3,(0.45)	0.2,(0.4)	-2.1,(0.8)		-0.95,(0.65)	-1.4,(0.65)	-1.2,(0.35)
(e ₃ , z, 1)	-0.2,0.6,0.2	-0.3,0.55,0.3	0,0.4,0	(e ₃ , z, 0)	0.4,0.7,-0.4	-0.3,0.75,0.3	0.1,0.7,-0.1
	-1.0,(0.2)	-1.15,(0.5)	-0.4,(0.8)		0.1,(0.55)	-1.65,(0.3)	-0.5,(0.7)
(e ₄ , z, 1)	-0.4,0.5,0.4	0.2,0.15,-0.2	-0.1,0.75,0.1	(e ₄ , z, 0)	-0.2,0.7,0.2	-0.5,0.4,0.5	0.4,0.6,-0.4
	-1.3,(0.55)	0.25,(0.5)	-0.95,(0.85)		-1.1,(0.5)	-1.4,(0.7)	0.2,(0.6)

Table-2

	High numerical grad for agree PNVSE set (λ_i)	High numerical average value of the vague set (μ_i) corresponding to highest numerical grad	$\lambda_i \times \mu_i$		High numerical grad for disagree PNVSE set (λ_i)	High numerical average value of the vague set (μ_i) corresponding to highest numerical grad	$\lambda_i \times \mu_i$
(e ₁ , x, 1)	u ₁ (-0.2)	0.4	-0.08	(e ₁ , x, 0)	u ₂ (-0.25)	0.4	-0.1
(e ₂ , x, 1)	u ₃ (0.6)	0.4	0.24	(e ₂ , x, 0)	u ₂ (-0.05)	0.45	-0.0225
(e ₃ , x, 1)	u ₃ (0.8)	0.5	-0.40	(e ₃ , x, 0)	u ₃ (0.55)	0.75	0.4125

$(e_4, x, 1)$	$u_3(-1.1)$	0.7	-0.77	$(e_4, x, 0)$	$u_3(0.6)$	0.45	0.27
$(e_1, y, 1)$	$u_1(1.05)$	0.5	0.525	$(e_1, y, 0)$	$u_3(-0.8)$	0.55	-0.44
$(e_2, y, 1)$	$u_1(-0.25)$	0.3	-0.075	$(e_2, y, 0)$	$u_3(0.6)$	0.45	0.27
$(e_3, y, 1)$	$u_3(-1.05)$	0.6	-0.63	$(e_3, y, 0)$	$u_3(0.6)$	0.6	0.36
$(e_4, y, 1)$	$u_3(0.75)$	0.45	0.3375	$(e_4, y, 0)$	$u_3(-1.0)$	0.25	-0.25
$(e_1, z, 1)$	$u_2(0.05)$	0.4	0.02	$(e_1, z, 0)$	$u_3(-0.8)$	0.3	-0.24
$(e_2, z, 1)$	$u_2(0.2)$	0.4	0.08	$(e_2, z, 0)$	$u_1(-0.95)$	0.65	-0.6175
$(e_3, z, 1)$	$u_3(-0.4)$	0.8	-0.32	$(e_3, z, 0)$	$u_1(0.1)$	0.55	0.055
$(e_4, z, 1)$	$u_2(0.25)$	0.5	0.125	$(e_4, z, 0)$	$u_3(0.2)$	0.6	0.12

For agree

Score $u_1 = -0.08+0.525+(-0.075) = 0.370$

Score $u_2 = 0.02+0.08+0.125 = 0.225$

Score $u_3 = 0.24+(-0.40)+(-0.75)+(-0.63) +0.3375+(-0.32) = -1.5225$

For disagree

Score $u_1 = -0.6175+0.055 = -0.5625$

Score $u_2 = -0.1+(-0.0225) = -0.1225,$

Score $u_3 = 0.4125+0.27+0.27+0.36 +(-0.25)+(-0.24)+0.12 = 0.9425$

Table 3. The score $r_i = A_i-D_i$

A_i	D_i	r_i
Score $u_1 = 0.37$	Score $u_1 = -0.5625$	0.9325
Score $u_2 = 0.225$	Score $u_2 = -0.1225$	0.3475
Score $u_3 = -1.5225$	Score $u_3 = 0.9425$	-2.465

Thus $S = \max u_i \in U \{r_i\} = r_1$. So, the committee is advised to hire candidate u_1 to fill the vacant position.

6. Conclusions

We give the advances of our proposal method using VPNVSE set as compared to that PVSE set as proposed by [19]. The VPNVSE set is a generalization of PVSE set. The VPNVSE set each examine the universal U in never detail with three membership functions, especially when there are many parameters involved, where PVSE set can tell us limited information about the universal U. It can

only handle the incomplete information comparing both the truth-membership value and falsity-membership values with corresponding vague set. But VPNVSE set can handle problems involving imprecise, indeterminacy and incomplete data with corresponding vague set. Thus it makes more accurate and realistic than PVSE set (PNVSE set [13]). In future many applications in decision making problems can be solved with VPNSE sets- especially in medical sciences.

References

1. K. Alhazaymeh and N. Hassan, *Possibility vague soft set and its application in decision making*, International Journal of pure and Applied Mathematics 2012, 77(4), 549-563.
2. S. Alkhazuleh, Neutrosophic vague set theory, critical review 2015, 29-39.
3. S. Allkhazaleh, A.R.Salleh, Soft Expert Sets, Advance in Sciences, Volume 2011, Article ID757868, 12 pages.
4. Abdel-Basset, M., El-hoseny, M., Gamal, A., & Smarandache, F. (2019), A Novel Model for Evaluation Hospital Medical Care Systems Based on Plithogenic Sets. Artificial Intelligence in Medicine, 101710.
5. Abdel-Basset, M., Manogaran, G., Gamal, A., & Chang, V. (2019), A Novel Intelligent Medical Decision Support Model Based on Soft Computing and IoT. IEEE Internet of Things Journal.
6. Abdel-Basset, M., Mohamed, R., Zaied, A. E. N. H., & Smarandache, F. , A hybrid plithogenic decision-making approach with quality function deployment for selecting supply chain sustainability metrics. Symmetry, 2019 ,11(7), 903.
7. Abdel-Basset, M., & Mohamed, M. ,A novel and powerful framework based on neutrosophic sets to aid patients with cancer. Future Generation Computer Systems, 2019, 98, 144-153.
8. Abdel-Basset, M., Mohamed, M., & Smarandache, F., Linear fractional programming based on triangular neutrosophic numbers. International Journal of Applied Management Science, 2019,11(1), 1-20.
9. Abdel-Basset, M., Atef, A., & Smarandache, F., A hybrid Neutrosophic multiple criteria group decision making approach for project selection. Cognitive Systems Research, 2019,57, 216-227.
10. Abdel-Basset, M., Gamal, A., Manogaran, G., & Long, H. V., A novel group decision making model based on neutrosophic sets for heart disease diagnosis. Multimedia Tools and Applications, 2019, 1-26.
11. Abdel-Basset, M., Chang, V., Mohamed, M., & Smarandache, F., A Refined Approach for Forecasting Based on Neutrosophic Time Series. Symmetry, 2019,11(4), 457.
12. W. L. Gau and D. J. Buehrer, Vague sets, IEEE Transaction on System, Man and Cybernetics 1993, 23(2), 610-614.
13. N. Hassan and A. Al-Quran, Possibility Neutrosophic Vague soft expert set for decision under uncertainty, The 4th International Conference on Mathematical Sciences, AIP conference Proc. 1830,070007-1-070007-7;doi.10.1063/1.4980956 Published by AIP publishing,978-0-7354-1498-3/\$30.00, 2017
14. D. Molodtsov, Soft set theory first result, Computers and Mathematics with Applications, 1999, 37(4-5), 19-31.
15. Anjan Mukherjee and Sadhan Sarkar, Possibility interval valued intuitionistic fuzzy soft expert set theory in complex phenomena and its application in decision making, Bull. Cal. Math. Soc. 2017, 109(6), 501-524.

16. A. Al-Quran and N. Hassan, Neutrosophic Vague soft expert set theory, *Journal of Intelligent and Fuzzy System* 2016, 30, 3691-3702.
17. A. Al-Quran and N. Hassan, Neutrosophic Vague Soft Set and its Applications, *Malaysian Journal of Mathematical Sciences* 2017,11(2),141-163.
18. F. Smarandache, Neutrosophic set – A generalization of the intuitionistic fuzzy sets, *International Journal of pure and Applied Mathematics* 2005, 24(3), 287-297.
19. G. Selvachandran and A.R. Salleh, *Possibility vague soft expert theory and its application in decision making*, proc. 1st int. conf. on soft computing in Data Science (SCDS 2015), communication in Computer and Information Science 545, edited by M.W. Berry, A.Hj. Mohamed and B.W. Yap, Springer 2015, 77-87.
20. Ganeshsree Selvachandran and Abdul RazakSalleh, Possibility Intuitionistic Fuzzy Soft Expert and Its Application in Decision Making, Hindawi Publishing Corporation *International Journal of Mathematics and Mathematical Sciences*, 2015, Article ID 314285, 11 pages <http://dx.doi.org/10.1155/2015/314285>.
21. W. Xu J. Ma, S. Wang and G. Hao, Vague soft sets and their properties, *Computer and Mathematics with Applications*, 2010, 59(2), 7876-794.

Received: June 19, 2019. Accepted: October 19, 2019