

5-1-2020

Structures on Doubt Neutrosophic Ideals of BCK/BCI-Algebras under (S, T)-Norms

Anas Al-Masarwah

Abd Ghafur Ahmad

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Al-Masarwah, Anas and Abd Ghafur Ahmad. "Structures on Doubt Neutrosophic Ideals of BCK/BCI-Algebras under (S, T)-Norms." *Neutrosophic Sets and Systems* 33, 1 (2020).
https://digitalrepository.unm.edu/nss_journal/vol33/iss1/17

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu, lsloane@salud.unm.edu, sarahrk@unm.edu.



Structures on Doubt Neutrosophic Ideals of BCK/BCI -Algebras under (S, T) -Norms

Anas Al-Masarwah^{1,*} and Abd Ghafur Ahmad²

^{1,2}School of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi 43600, Selangor, Malaysia.

E-mail: ¹ P90674@siswa.ukm.edu.my, almasarwah85@gmail.com, ² ghafur@ukm.edu.my

*Correspondence: almasarwah85@gmail.com

Abstract. Smarandache implemented the idea of neutrosophic set theory as a method for dealing undetermined data. Neutrosophic set theory is commonly used in various algebraic structures, such as groups, rings and BCK/BCI -algebras. At present, there exist no results on doubt neutrosophic ideals of BCK/BCI -algebras using t -conorm and t -norm. First, the notions of (S, T) - normed doubt neutrosophic subalgebras and ideals of BCK/BCI -algebras are introduced and the characteristic properties are described. Then, images and preimages of (S, T) - normed doubt neutrosophic ideals under homomorphism are considered. Moreover, the direct product and (S, T) - product of (S, T) - normed doubt neutrosophic ideals of BCK/BCI -algebras are also discussed.

Keywords: BCK/BCI -algebra; doubt neutrosophic subalgebra (ideal); (S, T) -normed doubt neutrosophic subalgebra (ideal).

1. Introduction

BCK -algebras entered into mathematics in 1966 through the work of Imai and Iséki [1], and were applied to various mathematical fields, such as group theory, topology, functional analysis and probability theory, etc. In the same way, the concept of a BCI -algebra, which is a generalization of a BCK -algebra, was proposed by Iséki [2]. Zadeh [3] introduced the idea of fuzzy set theory in 1965, where the degree of membership is discussed, and Xi [4] introduced fuzzy subalgebras and ideals in BCK/BCI -algebras in 1991. Later on, fuzzy sets have been generalized to intuitionistic fuzzy sets [5] by adding a non-membership function by Atanassov in 1986 and this concept has been applied to BCK/BCI -algebras by Jun and Kim [6].

As anew idea and based on the concept defined by Xi [4], Jun [7] in 1994 introduced the notions of doubt fuzzy subalgebras and ideals in BCK/BCI -algebras. Bej and Pal [8] introduced the concepts of doubt intuitionistic fuzzy subalgebras and ideals in BCK/BCI -algebras.

Al-Masarwah and Ahmad [9] introduced the concepts of doubt bipolar fuzzy subalgebras and ideals in BCK/BCI -algebras. After that, many other researchers used these ideas and published numerous articles in different branches of algebraic structures [10–14].

Triangular norms were formulated by Schweizer and Sklar [15] to model the distances in probabilistic metric spaces. Triangular norms play an important role in many fields of mathematics, statistics, cooperative games, decision making and artificial intelligence [16]. In particular, in fuzzy set theory, t -conorm (S) and t -norm (T) have been widely used for fuzzy logic, fuzzy relation equations and fuzzy operations. In algebraic structures, Senapati [17] proposed the idea of (imaginable) T -fuzzy subalgebras and (imaginable) T -fuzzy closed ideals of BG -algebras. Kim [18] presented the intuitionistic (S, T) -normed fuzzy subalgebras in BCK/BCI -algebras using triangular norms. Also, Kutukcu and Tuna [19], presented a new classification of intuitionistic fuzzy subalgebras, ideals and implicative ideals in BCK/BCI -algebras.

Neutrosophy, [20, 21] a new branch of science that deals with indeterminacy, was launched by Smarandache in 1998. This concept is a generalization of the classical set, fuzzy set and intuitionistic fuzzy set. Neutrosophic set theory has been applied to several fields of mathematics including decision making [22–24], pattern recognition and medical diagnosis [25] and others [26–31]. In the aspect of algebraic structures, the papers [32–38] address neutrosophic algebraic structures in BCK/BCI -algebras.

As no studies have been reported so far to generalize the above mentioned concepts, so the aim of this present article is:

(1) To propose the concept of (S, T) -normed doubt neutrosophic subalgebras and (S, T) -normed doubt neutrosophic ideals of BCK/BCI -algebras as a generalization of (S, T) -normed intuitionistic fuzzy subalgebras and ideals of BCK/BCI -algebras.

(2) To consider images and preimages of (S, T) - normed doubt neutrosophic ideals under homomorphism.

(3) To define and discuss the direct product and (S, T) - product of (S, T) - normed doubt neutrosophic ideals of BCK/BCI -algebras.

To do so, the rest of the article is structured as follows: In Section 2, we review some basic notions. In Section 3, we introduce the notions of (S, T) - normed doubt neutrosophic subalgebras and ideals of BCK/BCI -algebras and then describe some of the characteristic properties. Furthermore, we consider images and preimages of (S, T) - normed doubt neutrosophic ideals under homomorphism. In Section 4, we discuss the direct product and (S, T) - product of (S, T) - normed doubt neutrosophic ideals of BCK/BCI -algebras. Finally, in Section 5, we present the conclusion and future works of the study.

2. Preliminaries

In the current section, we remember some of the basic notions of *BCK/BCI*-algebras which will be very helpful in further study of the paper. Let X be a *BCK/BCI*-algebra in what follows, unless otherwise stated.

By a *BCI*-algebra, we mean a set X with a special element 0 and a binary operation $*$, for all $p, q, s \in X$, that satisfies the following axioms:

- (I) $[(p * q) * (p * s)] * (s * q) = 0$,
- (II) $[p * (p * q)] * q = 0$,
- (III) $p * p = 0$,
- (IV) $p * q = 0$ and $q * p = 0$ imply $p = q$.

If a *BCI*-algebra X satisfies $0 * p = 0$, then X is called a *BCK*-algebra. In a *BCK/BCI*-algebra, $p * 0 = p$ holds. A partial ordering \leq on a *BCK/BCI*-algebra X can be defined by $p \leq q$ if and only if $p * q = 0$. A non-empty subset K of a *BCK/BCI*-algebra X is called a subalgebra of X if $p * q \in K, \forall p, q \in X$, and an ideal of X if $\forall p, q \in X$,

- (1) $0 \in K$,
- (2) $p * q \in K$ and $q \in K$ imply $p \in K$.

Definition 2.1. A neutrosophic set in a non-empty set X (see [14]) is a structure of the form:

$$B = \{ \langle p; B_T(p), B_I(p), B_F(p) \rangle | p \in X \},$$

where $B_T, B_I, B_F : X \rightarrow [0, 1]$. We shall use the symbol $B = (B_T, B_I, B_F)$, for the neutrosophic set $B = \{ \langle p; B_T(p), B_I(p), B_F(p) \rangle | p \in X \}$.

If $B = (B_T, B_I, B_F)$ is a neutrosophic set in X , then $\square B = (B_T, B_I, B_T^c)$ and $\diamond B = (B_F^c, B_I, B_F)$ are also neutrosophic sets in X .

Definition 2.2 ([15]). A function $T : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is called a triangular norm, if it satisfies the following conditions: $\forall p, q, s \in [0, 1]$,

- (1) $T(0, 0) = 0, T(1, 1) = 1$,
- (2) $T(p, T(q, s)) = T(T(p, q), s)$,
- (3) $T(p, q) = T(q, p)$,
- (4) $T(p, q) \leq T(p, s)$ if $q \leq s$.

If $T(p, 0) = p$ and $T(p, 1) = p$ for all $p \in [0, 1]$, then T is called a t -conorm and a t -norm, respectively. Throughout this paper, denote S and T as a t -conorm and a t -norm, respectively.

Some examples of t -conorms and t -norms are:

- (1) $S_M(p, q) = \max\{p, q\}$ and $T_M(p, q) = \min\{p, q\}$.
- (2) $S_L(p, q) = \min\{p + q, 1\}$ and $T_L(p, q) = \max\{p + q - 1, 0\}$.

$$(3) S_P(p, q) = p + q - pq \text{ and } T_P(p, q) = pq.$$

A t -conorm S and a t -norm T are associated [39], i.e., $S(p, q) = 1 - T(1-p, 1-q)$, $\forall p, q \in [0, 1]$.

Lemma 2.3 ([40]). For any $p, q \in [0, 1]$, we have $0 \leq \max\{p, q\} \leq S(p, q) \leq 1$ and $0 \leq T(p, q) \leq \min\{p, q\} \leq 1$.

Definition 2.4 ([7]). A fuzzy set λ of X is called a doubt fuzzy subalgebra of X if $\lambda(p * q) \leq \max\{\lambda(p), \lambda(q)\} \forall p, q \in X$, and a doubt fuzzy ideal of X if $\lambda(0) \leq \lambda(p) \leq \max\{\lambda(p * q), \lambda(q)\} \forall p, q \in X$.

Definition 2.5 ([41]). A neutrosophic set $B = (B_T, B_I, B_F)$ of X is called a neutrosophic subalgebra of X if for all $p, q \in X$,

- (1) $B_T(p * q) \geq \min\{B_T(p), B_T(q)\}$,
- (2) $B_I(p * q) \geq \min\{B_I(p), B_I(q)\}$,
- (3) $B_F(p * q) \leq \max\{B_F(p), B_F(q)\}$.

Definition 2.6 ([41]). A neutrosophic set $B = (B_T, B_I, B_F)$ of X is called a neutrosophic ideal of X if for all $p, q \in X$,

- (1) $B_T(0) \geq B_T(p) \geq \min\{B_T(p * q), B_T(q)\}$,
- (2) $B_I(0) \geq B_I(p) \geq \min\{B_I(p * q), B_I(q)\}$,
- (3) $B_F(0) \leq B_F(p) \leq \max\{B_F(p * q), B_F(q)\}$.

3. (S, T) -Normed doubt neutrosophic ideals

Definition 3.1. A neutrosophic set $B = (B_T, B_I, B_F)$ of X is called a doubt neutrosophic subalgebra of X if for all $p, q \in X$,

- (1) $B_T(p * q) \leq \max\{B_T(p), B_T(q)\}$,
- (2) $B_I(p * q) \leq \max\{B_I(p), B_I(q)\}$,
- (3) $B_F(p * q) \geq \min\{B_F(p), B_F(q)\}$.

Definition 3.2. A neutrosophic set $B = (B_T, B_I, B_F)$ of X is called a doubt neutrosophic subalgebra of X with respect to a t -conorm S and a t -norm T (or simply, an (S, T) -normed doubt neutrosophic subalgebra of X) if for all $p, q \in X$,

- (1) $B_T(p * q) \leq S(B_T(p), B_T(q))$,
- (2) $B_I(p * q) \leq S(B_I(p), B_I(q))$,
- (3) $B_F(p * q) \geq T(B_F(p), B_F(q))$.

Definition 3.3. A neutrosophic set $B = (B_T, B_I, B_F)$ of X is called a doubt neutrosophic ideal of X if for all $p, q \in X$,

- (1) $B_T(0) \leq B_T(p) \leq \max\{B_T(p * q), B_T(q)\}$,

- (2) $B_I(0) \leq B_I(p) \leq \max\{B_I(p * q), B_I(q)\}$,
- (3) $B_F(0) \geq B_F(p) \geq \min\{B_F(p * q), B_F(q)\}$.

Definition 3.4. A neutrosophic set $B = (B_T, B_I, B_F)$ of X is called a doubt neutrosophic ideal of X with respect to a t -conorm S and a t -norm T (or simply, an (S, T) -normed doubt neutrosophic ideal of X) if for all $p, q \in X$,

- (1) $B_T(0) \leq B_T(p) \leq S(B_T(p * q), B_T(q))$,
- (2) $B_I(0) \leq B_I(p) \leq S(B_I(p * q), B_I(q))$,
- (3) $B_F(0) \geq B_F(p) \geq T(B_F(p * q), B_F(q))$.

Example 3.5. Consider a given BCK -algebra $X = \{0, k, l, m\}$ in Table 1:

TABLE 1. Tabular representation of a BCK -algebra $X = \{0, k, l, m\}$.

*	0	k	l	m
0	0	0	0	0
k	k	0	0	k
l	l	k	0	l
m	m	m	m	0

Define a neutrosophic set $B = (B_T, B_I, B_F)$ of X by Table 2:

TABLE 2. Neutrosophic set $B = (B_T, B_I, B_F)$.

X	$B_T(p)$	$B_I(p)$	$B_F(p)$
0	0	0	1
k	0.50	0.40	0.33
l	0.50	0.40	0.33
m	1	0.90	0

Clearly, $B_T(0) \leq B_T(p) \leq S_M(B_T(p * q), B_T(q))$, $B_I(0) \leq B_I(p) \leq S_M(B_I(p * q), B_I(q))$ and $B_F(0) \geq B_F(p) \geq T_L(B_F(p * q), B_F(q))$ for all $p, q \in X$. Hence, $B = (B_T, B_I, B_F)$ is an (S_M, T_L) -normed doubt neutrosophic ideal of X . Also, note that a t -conorm S_M and a t -norm T_L are not associated.

Remark 3.6. Example 3.5 holds even with the t -conorm S_M and t -norm T_M . Hence, $B = (B_T, B_I, B_F)$ is an (S_M, T_M) -normed doubt neutrosophic ideal of X .

Remark 3.7. Every doubt neutrosophic ideal of X is an (S, T) -normed doubt neutrosophic ideal of X , but the converse is not true.

Example 3.8. Consider a given *BCK*-algebra $X = \{0, 1, 2, 3, 4\}$ in Table 3:

TABLE 3. Tabular representation of a *BCK*-algebra $X = \{0, 1, 2, 3, 4\}$.

*	0	1	2	3	4
0	0	0	0	0	0
1	1	0	1	0	0
2	2	2	0	0	0
3	3	2	1	0	0
4	4	4	4	4	0

Define a neutrosophic set $B = (B_T, B_I, B_F)$ of X by Table 4:

TABLE 4. Neutrosophic set $B = (B_T, B_I, B_F)$.

X	$B_T(p)$	$B_I(p)$	$B_F(p)$
0	0.50	0.50	0.33
1	0.50	0.50	0.33
2	0.50	0.50	0.33
3	0.75	0.75	0.25
4	0.75	0.75	0.25

Clearly, $B_T(0) \leq B_T(p) \leq S_L(B_T(p * q), B_T(q))$, $B_I(0) \leq B_I(p) \leq S_L(B_I(p * q), B_I(q))$ and $B_F(0) \geq B_F(p) \geq T_P(B_F(p * q), B_F(q))$ for all $p, q \in X$. Hence, $B = (B_T, B_I, B_F)$ is an (S_L, T_P) -normed doubt neutrosophic ideal of X , but it is not a doubt neutrosophic ideal of X .

Lemma 3.9. *If $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic ideal of X , then so is $\square B = (B_T, B_I, B_T^c)$, where a t -conorm S and a t -norm T are associated.*

Proof. Let $B = (B_T, B_I, B_F)$ be an (S, T) -normed doubt neutrosophic ideal of X . Then, $B_T(0) \leq B_T(p) \forall p \in X$ and so $1 - B_T^c(0) \leq 1 - B_T^c(p)$. Hence, $B_T^c(0) \geq B_T^c(p)$. Also, for all $p, q \in X$, we have $B_T(p) \leq S(B_T(p * q), B_T(q))$ and so $1 - B_T^c(p) \leq S(1 - B_T^c(p * q), 1 - B_T^c(q))$ which implies $B_T^c(p) \geq 1 - S(1 - B_T^c(p * q), 1 - B_T^c(q))$. Since S and T are associated, we have $B_T^c(p) \geq T(B_T^c(p * q), B_T^c(q))$. Thus, $\square B = (B_T, B_I, B_T^c)$ is an (S, T) -normed doubt neutrosophic ideal of X . \square

Lemma 3.10. *If $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic ideal of X , then so is $\diamond B = (B_F^c, B_I, B_F)$, where a t -conorm S and a t -norm T are associated.*

Proof. The proof is similar to the proof of Lemma 3.9. \square

Combining Lemmas 3.9 and 3.10, we deduce that:

Theorem 3.11. *A neutrosophic set $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic ideal of X if and only if $\square B$ and $\diamond B$ are (S, T) -normed doubt neutrosophic ideals of X , where a t -conorm S and a t -norm T are associated.*

Lemma 3.12. *Every (S, T) -normed doubt neutrosophic ideal $B = (B_T, B_I, B_F)$ of X satisfies: for all $p, q \in X$,*

$$p \leq q \Rightarrow B_T(p) \leq B_T(q), B_I(p) \leq B_I(q) \text{ and } B_F(q) \geq B_F(p).$$

Proof. Let $p, q \in X$ be such that $p \leq q$. Then, $p * q = 0$ and so

$$B_T(p) \leq S(B_T(p * q), B_T(q)) = S(B_T(0), B_T(q)) \leq B_T(q),$$

$$B_I(p) \leq S(B_I(p * q), B_I(q)) = S(B_I(0), B_I(q)) \leq B_I(q),$$

and

$$B_F(p) \geq T(B_F(p * q), B_F(q)) = T(B_F(0), B_F(q)) \geq B_F(q).$$

This completes the proof. \square

Theorem 3.13. *Every (S, T) -normed doubt neutrosophic ideal of X is an (S, T) -normed doubt neutrosophic subalgebra of X .*

Proof. Let $B = (B_T, B_I, B_F)$ be an (S, T) -normed doubt neutrosophic ideal of X . Since $p * q \leq p \forall p, q \in X$, it follows from Lemma 3.12 that $B_T(p * q) \leq B_T(p)$, $B_I(p * q) \leq B_I(p)$ and $B_F(p * q) \geq B_F(p)$. Then,

$$B_T(p * q) \leq B_T(p) \leq S(B_T(p * q), B_T(q)) \leq S(B_T(p), B_T(q)),$$

$$B_I(p * q) \leq B_I(p) \leq S(B_I(p * q), B_I(q)) \leq S(B_I(p), B_I(q)),$$

and

$$B_F(p * q) \geq B_F(p) \geq T(B_F(p * q), B_F(q)) \geq T(B_F(p), B_F(q)).$$

Hence, $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic subalgebra of X . \square

Remark 3.14. The converse of Theorem 3.13 is not hold in general.

Example 3.15. Reconsider the BCK -algebra X given in Example 3.5. Define a neutrosophic set $B = (B_T, B_I, B_F)$ of X by Table 5:

TABLE 5. Neutrosophic set $B = (B_T, B_I, B_F)$.

X	$B_T(p)$	$B_I(p)$	$B_F(p)$
0	0	0	1
k	0.50	0.50	0.33
l	1	1	0
m	1	1	0

Clearly, $B = (B_T, B_I, B_F)$ is an (S_M, T_M) -normed doubt neutrosophic subalgebra of X , but it is not (S_M, T_M) -normed doubt neutrosophic ideal of X , since

$$B_T(l) = 1 > \max\{B_T(l * k), B_T(k)\},$$

$$B_I(l) = 1 > \max\{B_I(l * k), B_I(k)\},$$

and

$$B_F(l) = 0 < \min\{B_F(l * k), B_F(k)\}.$$

Definition 3.16. A mapping $\theta : X \rightarrow Y$ of BCK/BCI -algebras is said to be a homomorphism if $\theta(p * q) = \theta(p) * \theta(q) \forall p, q \in X$. If $\theta : X \rightarrow Y$ is a homomorphism, then $\theta(0) = 0$.

Let $\theta : X \rightarrow Y$ be a homomorphism of BCK/BCI -algebras. For any neutrosophic set $B = (B_T, B_I, B_F)$ in Y , we define a new neutrosophic set $B[\theta] = (B_T[\theta], B_I[\theta], B_F[\theta])$ such that for all $p \in X$,

$$B_T[\theta] : X \rightarrow [0, 1], B_T[\theta](p) = B_T(\theta(p)),$$

$$B_I[\theta] : X \rightarrow [0, 1], B_I[\theta](p) = B_I(\theta(p)),$$

$$B_F[\theta] : X \rightarrow [0, 1], B_F[\theta](p) = B_F(\theta(p)).$$

Theorem 3.17. Let $\theta : X \rightarrow Y$ be a homomorphism of BCK/BCI -algebras. If $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic ideal of Y , then $B[\theta] = (B_T[\theta], B_I[\theta], B_F[\theta])$ is an (S, T) -normed doubt neutrosophic ideal of X .

Proof. We first have

$$B_T[\theta](0) = B_T(\theta(0)) = B_T(0) \leq B_T(\theta(p)) = B_T[\theta](p),$$

$$B_I[\theta](0) = B_I(\theta(0)) = B_I(0) \leq B_I(\theta(p)) = B_I[\theta](p),$$

$$B_F[\theta](0) = B_F(\theta(0)) = B_F(0) \geq B_F(\theta(p)) = B_F[\theta](p)$$

for all $p, q \in X$. Let $p, q \in X$. Then,

$$\begin{aligned} B_T[\theta](p) &= B_T(\theta(p)) \leq S(B_T(\theta(p) * \theta(q)), B_T(\theta(q))) \\ &= S(B_T(\theta(p * q)), B_T(\theta(q))) \\ &= S(B_T[\theta](p * q), B_T[\theta](q)), \end{aligned}$$

$$\begin{aligned} B_I[\theta](p) &= B_I(\theta(p)) \leq S(B_I(\theta(p) * \theta(q)), B_I(\theta(q))) \\ &= S(B_I(\theta(p * q)), B_I(\theta(q))) \\ &= S(B_I[\theta](p * q), B_I[\theta](q)) \end{aligned}$$

and

$$\begin{aligned} B_F[\theta](p) &= B_F(\theta(p)) \geq T(B_F(\theta(p) * \theta(q)), B_F(\theta(q))) \\ &= T(B_F(\theta(p * q)), B_F(\theta(q))) \\ &= T(B_F[\theta](p * q), B_F[\theta](q)). \end{aligned}$$

Therefore, $B[\theta] = (B_T[\theta], B_I[\theta], B_F[\theta])$ is an (S, T) -normed doubt neutrosophic ideal of X . \square

Theorem 3.18. *Let $\theta : X \rightarrow Y$ be an onto homomorphism of BCK/BCI -algebras and let $B = (B_T, B_I, B_F)$ be a neutrosophic set of Y . If $B[\theta] = (B_T[\theta], B_I[\theta], B_F[\theta])$ is an (S, T) -normed doubt neutrosophic ideal of X , then $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic ideal of Y .*

Proof. For any $b \in Y$, there exists $a \in X$ such that $\theta(a) = b$. Then,

$$\begin{aligned} B_T(0) &= B_T(\theta(0)) = B_T[\theta](0) \leq B_T[\theta](a) = B_T(\theta(a)) = B_T(b), \\ B_I(0) &= B_I(\theta(0)) = B_I[\theta](0) \leq B_I[\theta](a) = B_I(\theta(a)) = B_I(b), \\ B_F(0) &= B_F(\theta(0)) = B_F[\theta](0) \geq B_F[\theta](a) = B_F(\theta(a)) = B_F(b). \end{aligned}$$

Let $p, q \in Y$. Then, $\theta(a) = p$ and $\theta(b) = q$ for some $a, b \in X$. It follows that

$$\begin{aligned} B_T(p) &= B_T(\theta(a)) = B_T[\theta](a) \\ &\leq S(B_T[\theta](a * b), B_T[\theta](b)) \\ &= S(B_T(\theta(a * b)), B_T(\theta(b))) \\ &= S(B_T(\theta(a) * \theta(b)), B_T(\theta(b))) \\ &= S(B_T(p * q), B_T(q)), \end{aligned}$$

$$\begin{aligned}
B_I(p) &= B_I(\theta(a)) = B_I[\theta](a) \\
&\leq S(B_I[\theta](a * b), B_I[\theta](b)) \\
&= S(B_I(\theta(a * b)), B_I(\theta(b))) \\
&= S(B_I(\theta(a) * \theta(b)), B_I(\theta(b))) \\
&= S(B_I(p * q), B_I(q))
\end{aligned}$$

and

$$\begin{aligned}
B_F(p) &= B_F(\theta(a)) = B_F[\theta](a) \\
&\geq T(B_F[\theta](a * b), B_F[\theta](b)) \\
&= T(B_F(\theta(a * b)), B_F(\theta(b))) \\
&= T(B_F(\theta(a) * \theta(b)), B_F(\theta(b))) \\
&= T(B_F(p * q), B_F(q)),
\end{aligned}$$

Therefore, $B = (B_T, B_I, B_F)$ is an (S, T) -normed doubt neutrosophic ideal of Y . \square

4. Product of (S, T) -normed doubt neutrosophic ideals

In this section, we discuss the direct product and (S, T) - product of (S, T) - normed doubt neutrosophic ideals of BCK/BCI -algebras.

Lemma 4.1 ([16]). *Let S and T be a t -conorm and t -norm, respectively. Then,*

$$\begin{aligned}
S(S(p, q), S(a, b)) &= S(S(p, a), S(q, b)), \\
T(T(p, q), T(a, b)) &= T(T(p, a), T(q, b))
\end{aligned}$$

for all $p, q, a, b \in [0, 1]$.

Theorem 4.2. *Let $X = P_1 \times P_2$ be the direct product BCK/BCI -algebra of two BCK/BCI -algebras P_1 and P_2 . If $B = (B_T, B_I, B_F)$ (resp., $C = (C_T, C_I, C_F)$) is an (S, T) -normed doubt neutrosophic ideal of P_1 (resp., P_2), then $D = (D_T, D_I, D_F)$ is an (S, T) -normed doubt neutrosophic ideal of X defined by $D_T = B_T \times C_T$, $D_I = B_I \times C_I$, and $D_F = B_F \times C_F$ such that*

$$\begin{aligned}
D_T(p_1, p_2) &= (B_T \times C_T)(p_1, p_2) = S(B_T(p_1), C_T(p_2)), \\
D_I(p_1, p_2) &= (B_I \times C_I)(p_1, p_2) = S(B_I(p_1), C_I(p_2)), \\
D_F(p_1, p_2) &= (B_F \times C_F)(p_1, p_2) = T(B_F(p_1), C_F(p_2))
\end{aligned}$$

for all $(p_1, p_2) \in X$.

Proof. Let $(p_1, p_2), (q_1, q_2) \in P_1 \times P_2$. Since $X = P_1 \times P_2$ is a BCK/BCI -algebra, we have

$$\begin{aligned} D_T(0, 0) &= (B_T \times C_T)(0, 0) = S(B_T(0), C_T(0)) \\ &\leq S(B_T(p_1), C_T(p_2)) \\ &= S((B_T \times C_T)(p_1, p_2)) \\ &= D_T(p_1, p_2), \end{aligned}$$

$$\begin{aligned} D_I(0, 0) &= (B_I \times C_I)(0, 0) = S(B_I(0), C_I(0)) \\ &\leq S(B_I(p_1), C_I(p_2)) \\ &= S((B_I \times C_I)(p_1, p_2)) \\ &= D_I(p_1, p_2), \end{aligned}$$

and

$$\begin{aligned} D_F(0, 0) &= (B_F \times C_F)(0, 0) = T(B_F(0), C_F(0)) \\ &\geq T(B_F(p_1), C_F(p_2)) \\ &= T((B_F \times C_F)(p_1, p_2)) \\ &= D_F(p_1, p_2). \end{aligned}$$

Also,

$$\begin{aligned} D_T(p_1, p_2) &= (B_T \times C_T)(p_1, p_2) = S(B_T(p_1), C_T(p_2)) \\ &\leq S\left(S(B_T(p_1 * q_1), B_T(q_1)), S(C_T(p_2 * q_2), C_T(q_2))\right) \\ &= S\left(S(B_T(p_1 * q_1), C_T(p_2 * q_2)), S(B_T(q_1), C_T(q_2))\right) \\ &= S\left((B_T \times C_T)(p_1 * q_1, p_2 * q_2), (B_T \times C_T)(q_1, q_2)\right) \\ &= S\left((B_T \times C_T)((p_1, p_2) * (q_1, q_2)), (B_T \times C_T)(q_1, q_2)\right) \\ &= S\left(D_T((p_1, p_2) * (q_1, q_2)), D_T(q_1, q_2)\right), \end{aligned}$$

$$\begin{aligned} D_I(p_1, p_2) &= (B_I \times C_I)(p_1, p_2) = S(B_I(p_1), C_I(p_2)) \\ &\leq S\left(S(B_I(p_1 * q_1), B_I(q_1)), S(C_I(p_2 * q_2), C_I(q_2))\right) \\ &= S\left(S(B_I(p_1 * q_1), C_I(p_2 * q_2)), S(B_I(q_1), C_I(q_2))\right) \\ &= S\left((B_I \times C_I)(p_1 * q_1, p_2 * q_2), (B_I \times C_I)(q_1, q_2)\right) \\ &= S\left((B_I \times C_I)((p_1, p_2) * (q_1, q_2)), (B_I \times C_I)(q_1, q_2)\right) \\ &= S\left(D_I((p_1, p_2) * (q_1, q_2)), D_I(q_1, q_2)\right), \end{aligned}$$

and

$$\begin{aligned}
 D_F(p_1, p_2) &= (B_F \times C_F)(p_1, p_2) = T(B_F(p_1), C_F(p_2)) \\
 &\geq T\left(T(B_F(p_1 * q_1), B_F(q_1)), T(C_F(p_2 * q_2), C_F(q_2))\right) \\
 &= T\left(T(B_F(p_1 * q_1), C_F(p_2 * q_2)), T(B_F(q_1), C_F(q_2))\right) \\
 &= T\left((B_F \times C_T)(p_1 * q_1, p_2 * q_2), (B_F \times C_F)(q_1, q_2)\right) \\
 &= T\left((B_F \times C_F)((p_1, p_2) * (q_1, q_2)), (B_F \times C_F)(q_1, q_2)\right) \\
 &= T\left(D_F((p_1, p_2) * (q_1, q_2)), D_F(q_1, q_2)\right).
 \end{aligned}$$

This completes the proof. \square

Definition 4.3. Let $B = (B_T, B_I, B_F)$ and $C = (C_T, C_I, C_F)$ be two neutrosophic sets of a BCK/BCI -algebra X . Then, (S, T) - product of B and C , written as $[B.C]_{(S,T)}$, are defined by

$$[B.C]_{(S,T)} = ([B_T.C_T]_S, [B_I.C_I]_S, [B_F.C_F]_T),$$

where

$$\begin{aligned}
 [B_T.C_T]_S(p) &= S(B_T(p), C_T(p)), \\
 [B_I.C_I]_S(p) &= S(B_I(p), C_I(p))
 \end{aligned}$$

and

$$[B_F.C_F]_T(p) = T(B_F(p), C_F(p))$$

for all $p \in X$.

Theorem 4.4. Let S and T be a t -conorm and s -norm, respectively. Let $B = (B_T, B_I, B_F)$ and $C = (C_T, C_I, C_F)$ be two (S, T) - normed doubt neutrosophic ideals of X . If S_1 is a t -conorm which dominates S , i.e.,

$$S_1(S(p, q), S(a, b)) \leq S(S_1(p, a), S_1(q, b))$$

and T_1 is a t -norm which dominates T , i.e.,

$$T_1(T(p, q), T(a, b)) \geq T(T_1(p, a), T_1(q, b))$$

for all $p, q, a, b \in [0, 1]$, then $([B_T.C_T]_{S_1}, [B_I.C_I]_{S_1}, [B_F.C_F]_{T_1})$ is an (S, T) -normed doubt neutrosophic ideal of X .

Proof. For any $p \in X$, we have

$$\begin{aligned} [B_T.C_T]_{S_1}(0) &= S_1(B_T(0), C_T(0)) \leq S_1(B_T(p), C_T(p)) = [B_T.C_T]_{S_1}(p), \\ [B_I.C_I]_{S_1}(0) &= S_1(B_I(0), C_I(0)) \leq S_1(B_I(p), C_I(p)) = [B_I.C_I]_{S_1}(p), \\ [B_F.C_F]_{T_1}(0) &= T_1(B_F(0), C_F(0)) \geq T_1(B_F(p), C_F(p)) = [B_F.C_F]_{T_1}(p). \end{aligned}$$

Also, for all $p, q \in X$, we have

$$\begin{aligned} [B_T.C_T]_{S_1}(p) &= S_1(B_T(p), C_T(p)) \\ &\leq S_1\left(S(B_T(p * q), B_T(q)), S(B_T(p * q), B_T(q))\right) \\ &\leq S\left(S_1(B_T(p * q), B_T(p * q)), S_1(B_T(q), B_T(q))\right) \\ &= S\left([B_T.C_T]_{S_1}(p * q), [B_T.C_T]_{S_1}(q)\right), \end{aligned}$$

$$\begin{aligned} [B_I.C_I]_{S_1}(p) &= S_1(B_I(p), C_I(p)) \\ &\leq S_1\left(S(B_I(p * q), B_I(q)), S(B_I(p * q), B_I(q))\right) \\ &\leq S\left(S_1(B_I(p * q), B_I(p * q)), S_1(B_I(q), B_I(q))\right) \\ &= S\left([B_I.C_I]_{S_1}(p * q), [B_I.C_I]_{S_1}(q)\right), \end{aligned}$$

and

$$\begin{aligned} [B_F.C_F]_{T_1}(p) &= T_1(B_F(p), C_F(p)) \\ &\geq T_1\left(T(B_F(p * q), B_F(q)), T(B_T(p * q), B_F(q))\right) \\ &\geq T\left(T_1(B_F(p * q), B_F(p * q)), T_1(B_F(q), B_F(q))\right) \\ &= T\left([B_F.C_F]_{T_1}(p * q), [B_F.C_F]_{T_1}(q)\right). \end{aligned}$$

This completes the proof. \square

5. Conclusions

In this paper, we have introduced the notions of (S, T) - normed doubt neutrosophic subalgebras and ideals of BCK/BCI -algebras and described the characteristic properties. Then, we have considered images and preimages of (S, T) - normed doubt neutrosophic ideals under homomorphism. Moreover, we have discussed the direct product and (S, T) - product of (S, T) -normed doubt neutrosophic ideals of BCK/BCI -algebras. We aim to extend our notions to

- (1) (S, T) - normed doubt generalized neutrosophic positive implicative ideals of BCK -algebras.
- (2) (S, T) - normed doubt generalized neutrosophic ideals of BCK/BCI -algebras.

(3) (S, T) - normed doubt cubic neutrosophic ideals of BCK/BCI -algebras.

Acknowledgment: We are indebted to Universiti Kebangsaan Malaysia for providing support and facilities for this research. Also, we are indebted to Zarqa University, since this paper is an extended paper of a short paper published in the 6th International Arab Conference on Mathematics and Computations (IACMC 2019), special session of Neutrosophic Set and Logic, organized by F. Smarandache and S. Alkhazaleh, Zarqa University, Jordan.

References

1. Y. Imai, and K. Iséki, On axiom systems of propositional calculi, Proc. Jpn. Acad. 42(1966), 19–21.
2. K. Iséki, An algebra related with a propositional calculus, Proc. Jpn. Acad. 42(1966), 26–29.
3. L.A. Zadeh, Fuzzy Sets, Information and Control, 8 (1965), 338–353.
4. O.G. Xi, Fuzzy BCK-algebras, Math. Japon. 24(1991), 935–942.
5. K. T. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems, 20 (1986), 87–96.
6. Y. B. Jun, and K. H. Kim, Intuitionistic fuzzy ideals of BCK-algebras, Internat. J. Math. & Math. Sci. 24(2000), 839849.
7. Y. B. Jun, Doubt fuzzy BCK/BCI-algebras, Soochow J. Math. 20(1994), 351–358.
8. T. Bej and M. Pal, Doubt intuitionistic fuzzy ideals in BCK/BCI-algebras, International Journal of Fuzzy Logic Systems, 5(1) (2015).
9. A. Al-Masarwah and A.G. Ahmad, Doubt bipolar fuzzy subalgebras and ideals in BCK/BCI-algebras, J. Math. Anal. 9(2018), 9–27.
10. A. Al-Masarwah and A.G. Ahmad, On some properties of doubt bipolar fuzzy H-ideals in BCK/BCI-algebras, Eur. J. Pure Appl. Math. 11(2018), 652–670.
11. A. Al-Masarwah and A.G. Ahmad, Novel concepts of doubt bipolar fuzzy H-ideals of BCK/BCI-algebras, Int. J. Innov. Comput., Inf. Control, 14(2018), 2025–2041.
12. A. Al-Masarwah, and A.G. Ahmad, m -polar fuzzy ideals of BCK/BCI-algebras, J. King Saud Univ.-Sci. 31(4)(2019), 1220-1226.
13. A. Al-Masarwah, and A.G. Ahmad, m -Polar (α, β) -fuzzy ideals in BCK/BCI-algebras, Symmetry, 11(2019), 44.
14. J. Zhan and Z. Tan, Doubt fuzzy BCI-algebras, Int. J. Math. Math. Sci. 30(2002), 49–56.
15. B. Schweizer and A. Sklar, Statistical metric spaces, Pacific J. Math. 10(1960), 314–334.
16. E.P. Klement, R. Mesiar and E. Pap, Triangular norms, Kluwer Acad. Publishers, Dordrecht, (2000).
17. T. Senapati, M. Bhowmik and M. Pal, Triangular norm based fuzzy BG-algebras, Afr. Mat. 27(2016), 187-199.
18. K. H. Kim, intuitionistic (T, S) -normed fuzzy subalgebras of BCK-algebras, J. of the chungcheong mathematical society, 20(3) (2007), 279–286.
19. S. Kutukcu and A. Tuna, Anti implicative IF-ideals in BCK/BCI-algebras, J. Computational Analysis and Applications, 25(2)(2018), 270–282.
20. F. Smarandache, Unifying field in logics: neutrosophic logic. Neutrosophy, Neutrosophic set, neutrosophic probability; American Reserch Press: Rehoboth, NM, USA, (1999).
21. F. Smarandache, Neutrosophic set-a generalization of the intuitionistic fuzzy set. Int. J. Pure Appl. Math. 24(2005), 287-297.
22. M. Abu Qamar and N. Hassan, An approach toward a Q-neutrosophic soft set and its application in decision making, Symmetry, 11(2019), 139.

23. M. Abu Qamar and N. Hassan, Entropy, measures of distance and similarity of Q-neutrosophic soft sets and some applications, *Entropy*, 20(2018), 672.
24. M. Abu Qamar and N. Hassan, Q-neutrosophic soft relation and its application in decision making, *Entropy*, 20(2018), 172.
25. H. Garg and Nancy, Some new biparametric distance measures on single-valued neutrosophic sets with applications to pattern recognition and medical diagnosis, *Information*, 8(2017), 126.
26. M. Abu Qamar and N. Hassan, Characterizations of group theory under Q-neutrosophic soft environment, *Neutrosophic Sets Syst.* 27(2019), 114–130.
27. M. Abu Qamar, A.G. Ahmad and N. Hassan, On Q-neutrosophic soft fields, *Neutrosophic Sets Syst.* 32(2020), 80–93.
28. Abdel-Basset, Mohamed, Mumtaz Ali, and Asmaa Atef. Uncertainty assessments of linear time-cost trade-offs using neutrosophic set. *Computers and Industrial Engineering* 141 (2020): 106286.
29. M. Abdel-Basset and R. Mohamed, A novel plithogenic TOPSIS-CRITIC model for sustainable supply chain risk management, *Journal of Cleaner Production*, 247(2020), 119586.
30. M. Abdel-Basset, M. Ali and A. Atef, Resource levelling problem in construction projects under neutrosophic environment, *The Journal of Supercomputing*, (2019), 1-25.
31. M. Abdel-Basset, M. Mohamed, M. Elhoseny, F. Chiclana, A.E.N.H. Zaied, Cosine similarity measures of bipolar neutrosophic set for diagnosis of bipolar disorder diseases, *Artificial Intelligence in Medicine*, 101(2019), 101735.
32. R.A. Borzooei, X.H. Zhang, F. Smarandache and Y.B. Jun, Commutative generalized neutrosophic ideals in BCK-algebras, *Symmetry* 2018, 10, 350; doi:10.3390/sym10080350.
33. Y.B. Jun, Neutrosophic subalgebras of several types in BCK/BCI-algebras, *Ann. Fuzzy Math. Inform.* 14(1) (2017), 7586.
34. Y.B. Jun, S.J. Kim and F. Smarandache, Interval neutrosophic sets with applications in BCK/BCI- algebra, *Axioms* 2018, 7, 23.
35. Y.B. Jun, F. Smarandache and H. Bordbar, Neutrosophic N-structures applied to BCK/BCI-algebras, *Information* 2017, 8, 128.
36. Y.B. Jun, F. Smarandache, S.Z. Song and M. Khan, Neutrosophic positive implicative N-ideals in BCK/BCI-algebras, *Axioms* 2018, 7, 3.
37. Y.B. Jun, F. Smarandache and M.A. Ozturk, Commutative falling neutrosophic ideals in BCK-algebras, *Neutrosophic Sets Syst.* 20(2018), 44–53.
38. S.J. Kim, S.Z. Song and Y.B. Jun, Generalizations of neutrosophic subalgebras in BCK/BCI-algebras based on neutrosophic points, *Neutrosophic Sets Syst.* 20(2018), 26–35.
39. R. Lowen, *Fuzzy set theory*, Dordrecht, Kluwer Acad. Publishers, (1996).
40. D.C. Chen, *Fuzzy rough set theory and method*; Science Press: Beijing, China, (2013).
41. J. Martina Jency and I. Arockiarani, Single Valued Neutrosophic Ideals of BCK-Algebras, *International Journal of Mathematics Trends and Technology*, 39(1)(2016), 8–17.

Received: 08, January, 2020 / Accepted: 02, May, 2020