

University of New Mexico

## UNM Digital Repository

---

Branch Mathematics and Statistics Faculty and  
Staff Publications

Branch Academic Departments

---

2014

### Neutrosophic Ideal Theory Neutrosophic Local Function and Generated Neutrosophic Topology

A. A. Salama

Florentin Smarandache

Follow this and additional works at: [https://digitalrepository.unm.edu/math\\_fsp](https://digitalrepository.unm.edu/math_fsp)



Part of the [Applied Mathematics Commons](#), [Logic and Foundations Commons](#), [Other Mathematics Commons](#), and the [Set Theory Commons](#)

---

# Neutrosophic Ideal Theory

## Neutrosophic Local Function and Generated Neutrosophic Topology

A. A. Salama & Florentin Smarandache

### ABSTRACT

**Abstract** In this paper we introduce the notion of ideals on neutrosophic set which is considered as a generalization of fuzzy and fuzzy intuitionistic ideals studies in [9,11], the important neutrosophic ideals has been given in [4]. The concept of neutrosophic local function is also introduced for a neutrosophic topological space. These concepts are discussed with a view to find new neutrosophic topology from the original one in [8]. The basic structure, especially a basis for such generated neutrosophic topologies and several relations between different neutrosophic ideals and neutrosophic topologies are also studied here. Possible application to GIS topology rules are touched upon.

**KEYWORDS:** Neutrosophic Set, Intuitionistic Fuzzy Ideal, Fuzzy Ideal, Neutrosophic Ideal, Neutrosophic Topology.

### 1-INTRODUCTION

The neutrosophic set concept was introduced by Smarandache [12, 13]. In 2012 neutrosophic sets have been investigated by Hanafy and Salama at el [4, 5, 6, 7, 8, 9, 10]. The fuzzy set was introduced by Zadeh [14] in 1965, where each element had a degree of membership. In 1983 the intuitionistic fuzzy set was introduced by K. Atanassov [1, 2, 3] as a generalization of fuzzy set, where besides the degree of membership and the degree of non-membership of each element. Salama at el [9] defined intuitionistic fuzzy ideal for a set and generalized the concept of fuzzy ideal concepts, first initiated by Sarker [10]. Neutrosophy has laid the foundation for a whole family of new mathematical theories generalizing both their classical and fuzzy counterparts. In this paper we will introduce the definitions of normal neutrosophic set, convex set, the concept of  $\alpha$ -cut and neutrosophic ideals, which can be discussed as generalization of fuzzy and fuzzy intuitionistic studies.

### 2-TERMINOLOGIES

We recollect some relevant basic preliminaries, and in particular, the work of Smarandache in [12, 13], and Salama at el. [4, 5, 6, 7, 8, 9, 10].

### 3- NEUTROSOPHIC IDEALS [4].

#### Definition.3.1

Let  $X$  is non-empty set and  $L$  a non-empty family of NSs. We will call  $L$  is a neutrosophic ideal (NL for short) on  $X$  if

- $A \in L$  and  $B \subseteq A \Rightarrow B \in L$  [heredity],

- $A \in L$  and  $B \in L \Rightarrow A \vee B \in L$  [Finite additivity].

A neutrosophic ideal  $L$  is called a  $\sigma$ -neutrosophic ideal if  $\bigvee_{j \in N} A_j \leq L$ , implies  $\bigvee_{j \in J} A_j \in L$  (countable additivity).

The smallest and largest neutrosophic ideals on a non-empty set  $X$  are  $O_N$  and NSs on  $X$ . Also,  $N.L_f$ ,  $N.L_c$  are denoting the neutrosophic ideals (NL for short) of neutrosophic subsets having finite and countable support of  $X$  respectively. Moreover, if  $A$  is a nonempty NS in  $X$ , then  $B \in NS : B \subseteq A$  is an NL on  $X$ . This is called the principal NL of all NSs of denoted by  $NL \langle A \rangle$ .

**Remark 3.1**

- If  $1_N \notin L$ , then  $L$  is called neutrosophic proper ideal.
- If  $1_N \in L$ , then  $L$  is called neutrosophic improper ideal.
- $O_N \in L$ .

**Example.3.1**

Any Intuitionistic fuzzy ideal  $\ell$  on  $X$  in the sense of Salama is obviously and NL in the form  $L = A : A = \langle x, \mu_A, \sigma_A, \nu_A \rangle \in \ell$ .

**Example.3.2**

Let  $X = a, b, c$ ,  $A = \langle x, 0.2, 0.5, 0.6 \rangle$ ,  $B = \langle x, 0.5, 0.7, 0.8 \rangle$ , and  $D = \langle x, 0.5, 0.6, 0.8 \rangle$ , then the family  $L = O_N, A, B, D$  of NSs is an NL on  $X$ .

**Example.3.3**

Let  $X = a, b, c, d, e$  and  $A = \langle x, \mu_A, \sigma_A, \nu_A \rangle$  given by:

$X$	$\mu_A$	$\sigma_A$	$\nu_A$
$a$	0.6	0.4	0.3
$b$	0.5	0.3	0.3
$c$	0.4	0.6	0.4
$d$	0.3	0.8	0.5
$e$	0.3	0.7	0.6

Then the family  $L = O_N, A$  is an NL on  $X$ .

**Definition.3.3**

Let  $L_1$  and  $L_2$  be two NL on  $X$ . Then  $L_2$  is said to be finer than  $L_1$  or  $L_1$  is coarser than  $L_2$  if  $L_1 \leq L_2$ . If also  $L_1 \neq L_2$ . Then  $L_2$  is said to be strictly finer than  $L_1$  or  $L_1$  is strictly coarser than  $L_2$ .

Two NL said to be comparable, if one is finer than the other. The set of all NL on  $X$  is ordered by the relation  $L_1$  is coarser than  $L_2$  this relation is induced the inclusion in NSs.

The next Proposition is considered as one of the useful result in this sequel, whose proof is clear.

**Proposition.3.1**

Let  $\{L_j : j \in J\}$  be any non - empty family of neutrosophic ideals on a set X. Then  $\bigcap_{j \in J} L_j$  and  $\bigcup_{j \in J} L_j$  are neutrosophic ideal on X,

In fact L is the smallest upper bound of the set of the  $L_j$  in the ordered set of all neutrosophic ideals on X.

**Remark.3.2**

The neutrosophic ideal by the single neutrosophic set  $O_N$  is the smallest element of the ordered set of all neutrosophic ideals on X.

**Proposition.3.3**

A neutrosophic set A in neutrosophic ideal L on X is a base of L iff every member of L contained in A.

**Proof**

(Necessity) Suppose A is a base of L. Then clearly every member of L contained in A.

(Sufficiency) Suppose the necessary condition holds. Then the set of neutrosophic subset in X contained in A coincides with L by the Definition 4.3.

**Proposition.3.4**

For a neutrosophic ideal  $L_1$  with base A, is finer than a fuzzy ideal  $L_2$  with base B iff every member of B contained in A.

**Proof**

Immediate consequence of Definitions

**Corollary.3.1**

Two neutrosophic ideals bases A, B, on X are equivalent iff every member of A, contained in B and via versa.

**Theorem.3.1**

Let  $\eta = \langle \mu_j, \sigma_j, \gamma_j \rangle : j \in J$  be a non empty collection of neutrosophic subsets of X. Then there exists a neutrosophic ideal  $L(\eta) = \{A \in NSs: A \subseteq \bigvee A_j\}$  on X for some finite collection  $\{A_j: j = 1,2, \dots, n \subseteq \eta\}$ .

**Proof**

Clear.

**Remark.3.3**

ii) The neutrosophic ideal  $L(\eta)$  defined above is said to be generated by  $\eta$  and  $\eta$  is called sub base of  $L(\eta)$ .

**Corollary.3.2**

Let  $L_1$  be an neutrosophic ideal on X and  $A \in NSs$ , then there is a neutrosophic ideal  $L_2$  which is finer than  $L_1$

and such that  $A \in L_2$  iff

$$A \vee B \in L_2 \text{ for each } B \in L_1.$$

**Corollary.3.3**

Let  $A = \langle x, \mu_A, \sigma_A, \nu_A \rangle \in L_1$  and  $B = \langle x, \mu_B, \sigma_B, \nu_B \rangle \in L_2$ , where  $L_1$  and  $L_2$  are neutrosophic ideals on the set  $X$ . then the neutrosophic set  $A^*B = \langle \mu_{A^*B}, \sigma_{A^*B}(x), \nu_{A^*B} \rangle \in L_1 \vee L_2$  on  $X$  where  $\mu_{A^*B} = \mu_A \wedge \mu_B, \sigma_{A^*B}(x) = \sigma_A(x) \vee \sigma_B(x)$  or  $\sigma_{A^*B}(x) = \sigma_A(x) \wedge \sigma_B(x)$  and  $\nu_{A^*B} = \nu_A \vee \nu_B$  or  $\nu_{A^*B} = \nu_A \wedge \nu_B, x \in X$ .

**4. Neutrosophic local Functions**

**Definition.4.1.** Let  $(X, \tau)$  be a neutrosophic topological spaces (NTS for short) and  $L$  be neutrosophic ideal (NL, for short) on  $X$ . Let  $A$  be any NS of  $X$ . Then the neutrosophic local function  $NA^*(L, \tau)$  of  $A$  is the union of all neutrosophic points (NP, for short)  $C(\alpha, \beta, \gamma)$  such that if  $U \in N(C(\alpha, \beta, \gamma))$  and  $NA^*(L, \tau) = \{C(\alpha, \beta, \gamma) \in X : A \wedge U \notin L \text{ for every } U \text{ nbd of } C(\alpha, \beta, \gamma)\}$ ,  $NA^*(L, \tau)$  is called a neutrosophic local function of  $A$  with respect to  $\tau$  and  $L$  which it will be denoted by  $NA^*(L, \tau)$ , or simply  $NA^*$ .

**Example .4.1.** One may easily verify that.

If  $L = \{0_N\}$ , then  $NA^*(L, \tau) = Ncl(A)$ , for any neutrosophic set  $A \in NSs$  on  $X$ .

If  $L =$  all NSs on  $X$  then  $NA^*(L, \tau) = 0_N$ , for any  $A \in NSs$  on  $X$ .

**Theorem.4.1.** Let  $(X, \tau)$  be a NTS and  $L_1, L_2$  be two neutrosophic ideals on  $X$ . Then for any neutrosophic sets  $A, B$  of  $X$ . then the following statements are verified

- i)  $A \subseteq B \Rightarrow NA^*(L, \tau) \subseteq NB^*(L, \tau)$ ,
- ii)  $L_1 \subseteq L_2 \Rightarrow NA^*(L_2, \tau) \subseteq NA^*(L_1, \tau)$ .
- iii)  $NA^* = Ncl(A^*) \subseteq Ncl(A)$ .
- iv)  $NA^{**} \subseteq NA^*$ .
- v)  $N(A \vee B)^* = NA^* \vee NB^*$ .
- vi)  $N(A \wedge B)^*(L) \leq NA^*(L) \wedge NB^*(L)$ .
- vii)  $\ell \in L \Rightarrow N(A \vee \ell)^* = NA^*$ .
- viii)  $NA^*(L, \tau)$  is neutrosophic closed set.

**Proof.**

- i) Since  $A \subseteq B$ , let  $p = C(\alpha, \beta, \gamma) \in NA^*(L_1)$  then  $A \wedge U \notin L$  for every  $U \in N(p)$ . By hypothesis we get  $B \wedge U \notin L$ , then  $p = C(\alpha, \beta, \gamma) \in NB^*(L_1)$ .
- ii) Clearly.  $L_1 \subseteq L_2$  implies  $NA^*(L_2, \tau) \subseteq NA^*(L_1, \tau)$  as there may be other IFSs which belong to  $L_2$  so that for GIFFP  $p = C(\alpha, \beta, \gamma) \in NA^*$  but  $C(\alpha, \beta, \gamma)$  may not be contained in  $NA^*(L_2)$ .
- iii) Since  $0_N \subseteq L$  for any NL on  $X$ , therefore by (ii) and Example 3.1,  $NA^*(L) \subseteq NA^*(0_N) = Ncl(A)$  for any NS  $A$  on  $X$ . Suppose  $p_1 = C_1(\alpha, \beta, \gamma) \in Ncl(NA^*(L_1))$ . So for every  $U \in N(p_1)$ ,  $NA^* \wedge U \neq 0_N$ , there exists  $p_2 = C_2(\alpha, \beta, \gamma) \in A^*(L_1 \wedge U)$  such that for every  $V$  nbd of  $p_2 \in N(p_2)$ ,  $A \wedge U \notin L$ . Since  $U \wedge V \in N(p_2)$  then  $A \wedge U \wedge V \notin L$  which leads to  $A \wedge U \notin L$ , for every  $U \in N(C(\alpha, \beta, \gamma))$  therefore  $p_1 = C(\alpha, \beta, \gamma) \in (A^*(L_1))$ .

and so  $Ncl \mathfrak{A}^* \leq NA^*$  While, the other inclusion follows directly. Hence  $NA^* = Ncl(NA^*)$ . But the inequality  $NA^* \leq Ncl(NA^*)$ .

iv) The inclusion  $NA^* \vee NB^* \leq N \mathfrak{A} \vee B^*$  follows directly by (i). To show the other implication, let  $p = C(\alpha, \beta, \gamma) \in N \mathfrak{A} \vee B^*$  then for every  $U \in N(p)$ ,  $\mathfrak{A} \vee B^* \wedge U \notin L$ , i.e.,  $\mathfrak{A} \wedge U \notin L$  or  $B^* \wedge U \notin L$ . then, we have two cases  $A \wedge U \notin L$  and  $B \wedge U \in L$  or the converse, this means that exist  $U_1, U_2 \in N \mathfrak{C}(\alpha, \beta, \gamma)$  such that  $A \wedge U_1 \notin L$ ,  $B \wedge U_1 \in L$ ,  $A \wedge U_2 \in L$  and  $B \wedge U_2 \notin L$ . Then  $A \wedge U_1 \wedge U_2 \in L$  and  $B \wedge U_1 \wedge U_2 \in L$  this gives  $\mathfrak{A} \vee B^* \wedge U_1 \wedge U_2 \in L$ ,  $U_1 \wedge U_2 \in N \mathfrak{C}(\alpha, \beta, \gamma)$  which contradicts the hypothesis. Hence the equality holds in various cases.

vi) By (iii), we have  $NA^{**} = Ncl(NA^*)^* \leq Ncl(NA^*) = NA^*$

Let  $\mathfrak{X}, \tau$  be a GIFTS and L be GIFL on X. Let us define the neutrosophic closure operator  $cl^*(A) = A \cup A^*$  for any GIFS A of X. Clearly, let  $Ncl^*(A)$  is a neutrosophic operator. Let  $N\tau^*(L)$  be NT generated by  $Ncl^*$

i.e  $N\tau^* \mathfrak{A} = A : Ncl^*(A^c) = A^c$ . Now  $L = O_N \Rightarrow Ncl^* \mathfrak{A} = A \cup NA^* = A \cup Ncl \mathfrak{A}$  for every neutrosophic set A. So,  $N\tau^*(O_N) = \tau$ . Again  $L = \text{all NSs on X} \Rightarrow Ncl^* \mathfrak{A} = A$ , because  $NA^* = O_N$ , for every neutrosophic set A so  $N\tau^* \mathfrak{A}$  is the neutrosophic discrete topology on X. So we can conclude by Theorem 4.1.(ii).  $N\tau^*(O_N) = N\tau^* \mathfrak{A}$  i.e.  $N\tau \subseteq N\tau^*$ , for any neutrosophic ideal  $L_1$  on X. In particular, we have for two neutrosophic ideals  $L_1$ , and  $L_2$  on X,  $L_1 \subseteq L_2 \Rightarrow N\tau^* \mathfrak{A}_1 \subseteq N\tau^* \mathfrak{A}_2$ .

**Theorem.4.2.** Let  $\tau_1, \tau_2$  be two neutrosophic topologies on X. Then for any neutrosophic ideal L on X,  $\tau_1 \leq \tau_2$  implies  $NA^*(L, \tau_2) \subseteq NA^*(L, \tau_1)$ , for every  $A \in L$  then  $N\tau^*_1 \subseteq N\tau^*_2$

**Proof.** Clear.

A basis  $N\beta \mathfrak{A}, \tau$  for  $N\tau^*(L)$  can be described as follows:

$N\beta \mathfrak{A}, \tau = A - B : A \in \tau, B \in L$  Then we have the following theorem

**Theorem 4.3.**  $N\beta \mathfrak{A}, \tau = A - B : A \in \tau, B \in L$  Forms a basis for the generated NT of the NT  $\mathfrak{X}, \tau$  with neutrosophic ideal L on X.

**Proof.** Straight forward.

The relationship between  $\tau$  and  $N\tau^*(L)$  established throughout the following result which have an immediately proof

**Theorem 4.4.** Let  $\tau_1, \tau_2$  be two neutrosophic topologies on X. Then for any neutrosophic ideal L on X,  $\tau_1 \subseteq \tau_2$  implies  $N\tau^*_1 \subseteq N\tau^*_2$ .

**Theorem 4.5 :** Let  $\mathfrak{X}, \tau$  be a NTS and  $L_1, L_2$  be two neutrosophic ideals on X. Then for any neutrosophic set A in X, we have

i)  $NA^* \mathfrak{A}_1 \vee L_2, \tau \supseteq NA^* \mathfrak{A}_1, N\tau^*(L_1) \wedge NA^* \mathfrak{A}_2, N\tau^*(L_2)$  ;ii)

$N\tau^*(L_1 \vee L_2) = \mathfrak{A} \tau^*(L_1) \wedge N \mathfrak{A}^*(L_2) \wedge (L_1)$

**Proof** Let  $p = C(\alpha, \beta) \notin \mathfrak{A}_1 \vee L_2, \tau$ , this means that there exists  $U_p \in N \mathfrak{P}$  such that  $A \wedge U_p \in \mathfrak{A}_1 \vee L_2$  i.e. There exists  $\ell_1 \in L_1$  and  $\ell_2 \in L_2$  such that  $A \wedge U_p \in \mathfrak{A}_1 \vee \ell_2$  because of the heredity of  $L_1$ , and assuming

$\ell_1 \wedge \ell_2 = O_N$ . Thus we have  $\mathfrak{A} \wedge U_p - \ell_1 = \ell_2$  and  $\mathfrak{A} \wedge U_p - \ell_2 = \ell_1$  therefore  $U_p - \ell_1 \wedge A = \ell_2 \in L_2$

and  $U_p - \ell_2 \wedge A = \ell_1 \in L_1$ . Hence  $p = C(\alpha, \beta, \gamma) \notin NA^* \mathfrak{A}_2, N\tau^* \mathfrak{A}_1$  or  $p = C(\alpha, \beta, \gamma) \notin NA^* \mathfrak{A}_1, N\tau^* \mathfrak{A}_2$  because

$p$  must belong to either  $\ell_1$  or  $\ell_2$  but not to both. This gives  $NA^* \mathfrak{A}_1 \vee L_2, \tau \supseteq NA^* \mathfrak{A}_1, N\tau^*(L_1) \wedge NA^* \mathfrak{A}_2, N\tau^*(L_2)$ .

To show the second inclusion, let us assume  $p = C(\alpha, \beta, \gamma) \notin NA^* \mathfrak{A}_1, N\tau^* \mathfrak{A}_2$ . This implies that there exist  $U_p \in N \mathfrak{P}$

and  $\ell_2 \in L_2$  such that  $U_p - \ell_2 \wedge A \in L_1$ . By the heredity of  $L_2$ , if we assume that  $\ell_2 \leq A$  and define

$\ell_1 = U_p - \ell_2 \wedge A$ . Then we have  $A \wedge U_p \in \mathfrak{A}_1 \vee \ell_2 \in L_1 \vee L_2$ . Thus,

$NA^* \mathbb{A}_1 \vee L_2, \tau \leq NA^* \mathbb{A}_1, \tau^*(L_1) \wedge NA^* \mathbb{A}_2, N\tau^*(L_2)$  and similarly, we can get  $A^* \mathbb{A}_1 \vee L_2, \tau \leq A^* \mathbb{A}_2, \tau^*(L_1)$ . This gives the other inclusion, which complete the proof.

**Corollary 4.1.** Let  $(X, \tau)$  be a NTS with neutrosophic ideal L on X. Then

- i)  $NA^*(L, \tau) = NA^*(L, \tau^*)$  and  $N\tau^*(L) = N(N\tau^*(L))^*(L)$ .
- ii)  $N\tau^*(L_1 \vee L_2) = N\tau^*(L_1) \vee N\tau^*(L_2)$

**Proof.** Follows by applying the previous statement.

**REFERENCES**

1. K. Atanassov, intuitionistic fuzzy sets, in V.Sgurev, ed., Vii ITKRS Session, Sofia (June 1983 central Sci. and Techn. Library, Bulg. Academy of Sciences (1984).
2. K. Atanassov, intuitionistic fuzzy sets, Fuzzy Sets and Systems 20, 87-96,(1986).
3. K. Atanassov, Review and new result on intuitionistic fuzzy sets, preprint IM-MFAIS-1-88, Sofia, (1988).
4. S. A. Albowi, A. A. Salama & Mohmed Eisa, New Concepts of Neutrosophic Sets, International Journal of Mathematics and Computer Applications Research (IJMCAR),Vol. 3, Issue 4, Oct 2013, 95-102.(2013).
5. I. Hanafy, A.A. Salama and K. Mahfouz, Correlation of neutrosophic Data, International Refereed Journal of Engineering and Science (IRJES) , Vol.(1), Issue 2 PP.39-43.(2012)
6. I.M. Hanafy, A.A. Salama and K.M. Mahfouz,," Neutrosophic Classical Events and Its Probability" International Journal of Mathematics and Computer Applications Research(IJMCAR) Vol.(3),Issue 1,Mar 2013, pp171-178.(2013)
7. A.A. Salama and S.A. Alblowi, "Generalized Neutrosophic Set and Generalized Neutrosophic Topological Spaces ", Journal computer Sci. Engineering, Vol. (2) No. (7) (2012).
8. A.A.Salama and S.A.Albolwi, Neutrosophic set and neutrosophic topological space, ISORJ. Mathematics,Vol.(3), Issue(4), pp-31-35.( 2012)
9. A.A. Salama and S.A. Albalwi, Intuitionistic Fuzzy Ideals Topological Spaces, Advances in Fuzzy Mathematics , Vol.(7), Number 1, pp. 51- 60, (2012).
10. A.A.Salama, and H.Elagamy, "Neutrosophic Filters" International Journal of Computer Science Engineering and Information Technology Research (IJCSEITR), Vol.3, Issue(1),Mar 2013, pp 307-312.(2013)
11. Debasis Sarker, Fuzzy ideal theory, Fuzzy local function and generated fuzzy topology, Fuzzy Sets and Systems 87, 117 – 123. (1997)
12. Florentin Smarandache, Neutrosophy and Neutrosophic Logic , First International Conference on Neutrosophy , Neutrosophic Logic , Set, Probability, and Statistics University of New Mexico, Gallup, NM 87301, USA(2002).
13. F. Smarandache. A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability. American Research Press, Rehoboth, NM, (1999).
14. L.A. Zadeh, Fuzzy Sets, Inform and Control 8, 338-353,(1965).