

7-15-2020

Generalized neutrosophic b-open sets in neutrosophic topological space

Suman Das

Surapati Pramanik

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Das, Suman and Surapati Pramanik. "Generalized neutrosophic b-open sets in neutrosophic topological space." *Neutrosophic Sets and Systems* 35, 1 (2020). https://digitalrepository.unm.edu/nss_journal/vol35/iss1/30

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu, lsloane@salud.unm.edu, sarahrk@unm.edu.



Generalized neutrosophic b -open sets in neutrosophic topological space

Suman Das¹, and Surapati Pramanik^{2,*}

¹Department of Mathematics, Tripura University, Agartala, 799022, Tripura, India.
Email: suman.mathematics@tripurauniv.in

²Department of Mathematics, Nandalal Ghosh B. T. College, Narayanpur, 743126, West Bengal, India.
Email: sura_pati@yahoo.co.in

* Correspondence: sura_pati@yahoo.co.in Tel.: (+91-9477035544)

Abstract: The purpose of the study is to introduce the notion of generalized neutrosophic b -open set in neutrosophic topological space. We define generalized neutrosophic b -open set, generalized neutrosophic b -interior, generalized neutrosophic b -closure and investigate some of their properties. By defining generalized neutrosophic b -open set, we prove some theorems on neutrosophic topological spaces. We also furnish some suitable examples.

Keywords: Neutrosophic set; neutrosophic b -open set; generalized neutrosophic b -open set; generalized neutrosophic b -interior; generalized neutrosophic b -closure

1. Introduction

Smarandache (1998) grounded the Neutrosophic Set (NS) in 1998. From then it became very popular and attracted many researchers' attention for theoretical and practical researches (Broumi et al., 2018; Khalid, 2020; Peng & Dai, 2018; Pramanik, 2013; 2016a; 2016b; 2020; Pramanik & Mallick, 2018; 2019; Pramanik & Mondal, 2016; Pramanik & Roy, 2014; Smarandache & Pramanik, 2016; 2018, Biswas, Pramanik & Giri, 2014; 2016a; 2016b; Dalapati et al., 2017; Dey, Pramanik, & Giri, 2016a; 2016b; Pramanik, Mallick, & Dasgupta, 2018; Mondal & Pramanik, 2015; Pramanik & Dalapati, 2018, Pramanik, Dey, & Smarandache, 2018; Pramanik, Mondal, & Smarandache, 2016a; 2016b).

Salama and Alblowi (2012a) grounded the "Neutrosophic Topological Space" (NTS). Salama and Alblowi (2012b) also presented generalized NS and generalized NTSs. Salama, Smarandache, & Alblowi (2014) studied the concept of neutrosophic crisp topological space. Arokiarani, Dhavaseelan, Jafari, and Parimala (2017) defined neutrosophic semi-open functions and established relation between them. Iswaraya and Bageerathi (2016) studied neutrosophic semi-closed set and neutrosophic semi-open set. Rao and Srinivasa (2017) introduced neutrosophic pre-open set and pre-closed set. Dhavaseelan and Jafari (2018) studied generalized neutrosophic closed sets. Pushpalatha and Nandhini (2019) defined the neutrosophic generalized closed sets in NTSs. Shanthi, Chandrasekar, Safina, and Begam (2018) presented the neutrosophic generalized semi closed sets in

NTSs. Ebenanjar, Immaculate, and Wilfred (2018) studied neutrosophic b -open sets in NTSs. Maheswari, Sathyabama, and Chandrasekar (2018) studied the neutrosophic generalized b -closed sets in NTSs.

Research gap: No investigation on neutrosophic generalized b -open set has been reported in the recent literature.

Motivation: In order to fill the research gap, we introduce neutrosophic generalized b -open set.

Remaining of the paper is designed as follows:

Section 2 recalls of NTS, neutrosophic b -closed sets and a theorem. Section 3 introduces neutrosophic generalized b -open set and proofs of some theorems on neutrosophic b -open sets. Section 4 presents concluding remarks.

2. Preliminaries and some properties

Definition 2.1 Assume that (W, τ) is an NTS. Then χ , an NS over W is said to be a Neutrosophic b -Open (N - b -open) set (Ebenanjar, Immaculate, & Wilfred, 2018) if and only if (iff) $\chi \subseteq N_{int}(N_{cl}(\chi)) \cup N_{cl}(N_{int}(\chi))$.

Definition 2.2 In an NTS (W, τ) , an NS χ is said to be a Neutrosophic b -Closed (N - b -closed) set (Ebenanjar, Immaculate, & Wilfred, 2018) iff $\chi \supseteq N_{int}(N_{cl}(\chi)) \cap N_{cl}(N_{int}(\chi))$.

Remark 2.1 An NS χ over W is said to be an N - b -closed set (Ebenanjar, Immaculate, & Wilfred, 2018) in (W, τ) iff χ^c is a N - b -open set in (W, τ) .

In 2018, Ebenanjar, Immaculate, and Wilfred (2018) studied the concept of N - b -open set in NTS but they did not check whether the union or intersection of two N - b -open sets (N - b -closed sets) is again an N - b -open set (N - b -closed set) or not. In this paper we show some results on the intersection and union of neutrosophic b -closed sets.

Theorem 2.1 The intersection of any two N - b -closed sets is again an N - b -closed set.

Proof. Assume that E, F be any two N - b -closed sets in an NTS (W, τ) . Then we have

$$E \supseteq N_{int}(N_{cl}(E)) \cap N_{cl}(N_{int}(E)) \tag{1}$$

$$\text{and } F \supseteq N_{int}(N_{cl}(F)) \cap N_{cl}(N_{int}(F)) \tag{2}$$

For any two NSs E and F We know that $E \cap F \subseteq E$ and $E \cap F \subseteq F$.

$$\text{Now } E \cap F \subseteq E \Rightarrow N_{int}(E \cap F) \subseteq N_{int}(E) \Rightarrow N_{cl}(N_{int}(E \cap F)) \subseteq N_{cl}(N_{int}(E)) \tag{3}$$

$$E \cap F \subseteq E \Rightarrow N_{cl}(E \cap F) \subseteq N_{cl}(E) \Rightarrow N_{int}(N_{cl}(E \cap F)) \subseteq N_{int}(N_{cl}(E)) \tag{4}$$

$$E \cap F \subseteq F \Rightarrow N_{int}(E \cap F) \subseteq N_{int}(F) \Rightarrow N_{cl}(N_{int}(E \cap F)) \subseteq N_{cl}(N_{int}(F)) \tag{5}$$

$$E \cap F \subseteq F \Rightarrow N_{cl}(E \cap F) \subseteq N_{cl}(F) \Rightarrow N_{int}(N_{cl}(E \cap F)) \subseteq N_{int}(N_{cl}(F)) \tag{6}$$

From (1) and (2) we have,

$$\begin{aligned} E \cap F &\supseteq N_{int}(N_{cl}(E)) \cap N_{cl}(N_{int}(E)) \cap N_{int}(N_{cl}(F)) \cap N_{cl}(N_{int}(F)) \\ &\supseteq N_{int}(N_{cl}(E \cap F)) \cap N_{cl}(N_{int}(E \cap F)) \cap N_{int}(N_{cl}(E \cap F)) \cap N_{cl}(N_{int}(E \cap F)) \\ &\hspace{15em} [\text{ by eqs (3), (4), (5) \& (6)}] \\ &= N_{int}(N_{cl}(E \cap F)) \cap N_{cl}(N_{int}(E \cap F)) \end{aligned}$$

$$\Rightarrow E \cap F \supseteq N_{cl}(N_{int}(E \cap F)) \cap N_{int}(N_{cl}(E \cap F)).$$

Therefore $E \cap F$ is an N - b -closed set.

Hence the intersection of any two N-*b*-closed sets is again an N-*b*-closed set.

Remark 2.2: The union of any two N-*b*-closed sets may not be an N-*b*-closed set. This is proved as follows:

Example 2.1: Assume that $W = \{p_1, p_2\}$ and $\tau = \{0_N, 1_N, \{(p_1, 0.5, 0.2, 0.4), (p_2, 0.6, 0.1, 0.3)\}, \{(p_1, 0.3, 0.5, 0.6), (p_2, 0.4, 0.4, 0.5)\}\}$ be the family of some NSs over W . Then τ is an NT on W . Now it can be verified that $E = \{(a, 0.6, 0.5, 0.6), (b, 0.5, 0.6, 0.7)\}$, $F = \{(a, 1, 0, 1), (b, 0.9, 0.1, 0.1)\}$ are two N-*b*-closed sets in (W, τ) . But their union $E \cup F = \{(a, 1, 0, 0.6), (b, 0.9, 0.1, 0.1)\}$ is not an N-*b*-closed set.

Definition 2.3 Assume that (W, τ) is an NTS and χ is an NS over W . Then the Neutrosophic *b*-Closure (N_{bcl}) and Neutrosophic *b*-Interior (N_{bint}) (Ebenanjar, Immaculate & Wilfred, 2018) of χ are defined by

$$N_{bcl}(\chi) = \cap \{ \psi : \psi \text{ is an N-}b\text{-closed set in } (W, \tau) \text{ and } \chi \subseteq \psi \};$$

$$N_{bint}(\chi) = \cup \{ \xi : \xi \text{ is an N-}b\text{-open set in } (W, \tau) \text{ and } \xi \subseteq \chi \}.$$

Remark 2.3 Clearly $N_{bint}(\chi)$ is the largest N-*b*-open set (Ebenanjar, Immaculate, & Wilfred, 2018) in (W, τ) which is contained in χ and $N_{bcl}(\chi)$ is the smallest N-*b*-closed set in (W, τ) which contains χ .

Definition 2.4 Assume that (W, τ) is an NTS. A neutrosophic subset E of (W, τ) is said to be a Neutrosophic Generalized Closed Set (NGCS) (Dhavaseelan & Jafari, 2018) if $N_{cl}(E) \subseteq F$ whenever $E \subseteq F$ and F is an NOS. A subset K of (W, τ) is called Neutrosophic Generalized Open Set (NGOS) iff K^c is an NGCS in (W, τ) .

3. Generalized neutrosophic *b*-open set

Definition 3.1 Assume that (W, τ) is an NTS. An NS G over W is called a Generalized Neutrosophic *b*-Open (g-N-*b*-open) set if \exists an N-*b*-closed set H (except 1_N) with $G \subseteq H$ such that $G \subseteq N_{int}(H)$. A neutrosophic subset K in (W, τ) is called a Generalized Neutrosophic *b*-Closed (g-N-*b*-closed) set iff K^c is a g-N-*b*-open set in (W, τ) .

Example 3.1 Assume that $W = \{p_1, p_2\}$ and $\tau = \{0_N, 1_N, \{(p_1, 0.5, 0.6, 0.7), (p_2, 0.6, 0.7, 0.8)\}, \{(p_1, 0.6, 0.5, 0.6), (p_2, 0.7, 0.6, 0.7)\}\}$ are the collection of some NSs over W . Then (W, τ) is clearly an NTS. Here $K = \{(p_1, 0.6, 0.7, 0.8), (p_2, 0.5, 0.8, 0.8)\}$ is a g-N-*b*-open set, because there exists an N-*b*-closed set $G = \{(p_1, 0.7, 0.3, 0.4), (p_2, 0.8, 0.3, 0.4)\}$ in (W, τ) with $K \subseteq G$ such that $K \subseteq N_{int}(G)$.

Proposition 3.1 In an NTS (W, τ) , 0_N is a g-N-*b*-open set but 1_N is not a g-N-*b*-open set.

Proof. Assume that (W, τ) is an NTS. Since a Neutrosophic Open Set (NOS) is an N-*b*-open set, so 1_N is an N-*b*-open set. Therefore, 0_N is an N-*b*-closed set (since it is the complement of N-*b*-open set 1_N). Now $0_N \subseteq 0_N$ and $0_N \subseteq N_{int}(0_N) = 0_N$.

Thus there exist an N-*b*-closed set 0_N (except 1_N) with $0_N \subseteq 0_N$ such that $0_N \subseteq N_{int}(0_N)$. Hence 0_N is a g-N-*b*-open set in (W, τ) .

But in case of NS 1_N , we cannot find any neutrosophic *b*-closed set H (except 1_N) with $1_N \subseteq H$ such that $1_N \subseteq N_{int}(H)$. Hence 1_N is not a g-N-*b*-open set in (W, τ) .

Proposition 3.2 Assume that ψ is a g-N-*b*-open set in an NTS (W, τ) . Then, every NS contained in ψ is a g-N-*b*-open set.

Proof. Assume that ψ be a g-N- b -open set in an NTS (W, τ) and ξ be any arbitrary NS over W which is contained in ψ . Since ψ is a g-N- b -open set, so there exists an N- b -closed set η (except 1_N) with $\psi \subseteq \eta$ such that $\psi \subseteq N_{int}(\eta)$.

Now ξ is contained in A , so

$$\begin{aligned} &\xi \subseteq \psi \\ \Rightarrow &\xi \subseteq \psi \subseteq \eta \ \& \ \xi \subseteq \psi \subseteq N_{int}(\eta). \end{aligned}$$

Therefore there exists an N- b -closed set η (except 1_N) with $\xi \subseteq \eta$ such that $\xi \subseteq N_{int}(\eta)$. Hence ξ is a g-N- b -open set. Thus each NS contained in ψ is again a g-N- b -open set in (W, τ) .

Definition 3.2 Assume that (W, τ) is an NTS and ψ be an NS over W . Then the Generalized Neutrosophic b -Interior ($g-N_{bint}$) and Generalized Neutrosophic b - Closure ($g-N_{bcl}$) of ψ are defined by

$$\begin{aligned} g-N_{bint}(\psi) &= \cup \{ \xi : \xi \text{ is a g-N-}b\text{-open set and } \xi \subseteq \psi \}; \\ g-N_{bcl}(\psi) &= \cap \{ \eta : \eta \text{ is a g-N-}b\text{-closed set and } \psi \subseteq \eta \}. \end{aligned}$$

Theorem 3.1 Assume that (W, τ) is an NTS. Then each neutrosophic open subset of (W, τ) is a g-N- b -open set.

Proof. Assume that ψ be an arbitrary NOS in an NTS (W, τ) . So $\psi = N_{int}(\psi)$. Since each neutrosophic closed set is an N- b -closed set so $N_{cl}(\psi)$ is an N- b -closed set. Also we know that $\psi \subseteq N_{cl}(\psi)$.

$$\begin{aligned} \text{Now } &\psi \subseteq N_{cl}(\psi) \\ \Rightarrow &N_{int}(\psi) \subseteq N_{int}(N_{cl}(\psi)) \\ \Rightarrow &\psi = N_{int}(\psi) \subseteq N_{int}(N_{cl}(\psi)) \\ \Rightarrow &\psi \subseteq N_{int}(N_{cl}(\psi)) \end{aligned}$$

Therefore there exists an N- b -closed set $N_{cl}(\psi)$ with $\psi \subseteq N_{cl}(\psi)$ such that $\psi \subseteq N_{int}(N_{cl}(\psi))$. Hence ψ is a g-N- b -open set in (W, τ) . Thus each neutrosophic open subset of (W, τ) is again a g-N- b -open set.

Remark 3.1 The converse of the theorem 3.1 is not true. This can be shown by the example 3.2.

Example 3.2 In example 3.1, it can be easily seen that $K = \{(a, 0.6, 0.7, 0.8), (b, 0.5, 0.8, 0.8)\}$ is a g-N- b -open set in (W, τ) but it is not an NOS.

Theorem 3.2 Assume that (W, τ) is an NTS. Then each Neutrosophic Pre-Open Set (NPOS) in (W, τ) is a g-N- b -open set.

Proof. Assume that (W, τ) is an NTS and ψ is an NPOS. Then $\psi \subseteq N_{int}(N_{cl}(\psi))$. Since for any NS ψ , $N_{cl}(\psi)$ is an N- b -closed set and $\psi \subseteq N_{cl}(\psi)$. Therefore there exists an N- b -closed set $N_{cl}(\psi)$ with $\psi \subseteq N_{cl}(\psi)$ such that $\psi \subseteq N_{int}(N_{cl}(\psi))$. Hence ψ is a g-N- b -open set in (W, τ) . Thus each NPOS in (W, τ) is again a g-N- b -open set.

Theorem 3.3 If ψ is both NOS and Neutrosophic Semi-Open Set (NSOS) in an NTS (W, τ) then it is a g-N- b -open set.

Proof. Assume that (W, τ) is an NTS and ψ is both NSOS and NOS. Since ψ is an NOS, so $\psi = N_{int}(\psi)$. Again since ψ is an NSOS, so $\psi \subseteq N_{cl}(N_{int}(\psi))$. It can be verified that $N_{cl}(N_{int}(\psi))$ is an N- b -closed set (since it is an NCS).

$$\begin{aligned} \text{Now } \psi &\subseteq N_{cl}(N_{int}(\psi)) \\ \Rightarrow N_{int}(\psi) &\subseteq N_{int}(N_{cl}(N_{int}(\psi))) \quad [\text{since } \psi \subseteq \delta \Rightarrow N_{int}(\psi) \subseteq N_{int}(\delta)] \\ \Rightarrow \psi = N_{int}(\psi) &\subseteq N_{int}(N_{cl}(N_{int}(\psi))) \quad [\text{since } \psi = N_{int}(\psi)] \\ \Rightarrow \psi &\subseteq N_{int}(N_{cl}(N_{int}(\psi))) \end{aligned}$$

Therefore there exists an N- b -closed set $N_{cl}(N_{int}(\psi))$ with $\psi \subseteq N_{cl}(N_{int}(\psi))$ in (W, τ) such that $\psi \subseteq N_{int}(N_{cl}(N_{int}(\psi)))$. Hence ψ is a g-N- b -open set.

Theorem 3.4 Assume that (W, τ) is an NTS and ψ is both neutrosophic α -open and neutrosophic open set. Then ψ is again a g-N- b -open set.

Proof. Assume that ψ is an arbitrary NS which is both neutrosophic α -open set and NOS. Since ψ is an NOS so $\psi = N_{int}(\psi)$. Again since ψ is a neutrosophic α -open set, so $\psi \subseteq N_{int}(N_{cl}(N_{int}(\psi)))$. Hence, it is clear that $N_{cl}(N_{int}(\psi))$ is an N- b -closed set (since it is an NCS) in (W, τ) .

$$\begin{aligned} \text{Now } \psi &= N_{int}(\psi) \\ \Rightarrow \psi = N_{int}(\psi) &\subseteq N_{cl}(N_{int}(\psi)) \\ \Rightarrow \psi &\subseteq N_{cl}(N_{int}(\psi)) \end{aligned}$$

Therefore there exists an N- b -closed set $N_{cl}(N_{int}(\psi))$ with $\psi \subseteq N_{cl}(N_{int}(\psi))$ such that $\psi \subseteq N_{int}(N_{cl}(N_{int}(\psi)))$. Hence ψ is a generalized N- b -open set in (W, τ) .

Theorem 3.5 The intersection of any two g-N- b -open sets in an NTS (W, τ) is again a g-N- b -open set.

Proof. Let ψ and ξ be any two g-N- b -open sets in an NTS (W, τ) . Then there exist two N- b -closed sets K, L with $\psi \subseteq K$, $\xi \subseteq L$ such that $\psi \subseteq N_{int}(K)$ and $\xi \subseteq N_{int}(L)$.

$$\text{Here } \psi \cap \xi \subseteq K \cap L.$$

We know that the intersection of two N- b -closed sets is again an N- b -closed set. So $K \cap L$ is an N- b -closed set in (W, τ) .

$$\begin{aligned} \text{Now } \psi \cap \xi &\subseteq N_{int}(K) \cap N_{int}(L) \quad [\text{since } \psi \subseteq N_{int}(K), \xi \subseteq N_{int}(L)] \\ &= N_{int}(K \cap L) \\ \Rightarrow \psi \cap \xi &\subseteq N_{int}(K \cap L). \end{aligned}$$

Therefore there exists an N- b -closed set $K \cap L$ with $\psi \cap \xi \subseteq K \cap L$ such that $\psi \cap \xi \subseteq N_{int}(K \cap L)$. Hence $\psi \cap \xi$ is a g-N- b -open set in (W, τ) . Thus the intersection of any two g-N- b -open sets in (W, τ) is again a g-N- b -open set.

Theorem 3.6 The union of two g-N- b -open sets is a g-N- b -open set if one is contained in the other.

Proof. Let ψ, ξ are any two g-N- b -open sets in (W, τ) such that $\psi \subseteq \xi$. Since ψ and ξ are g-N- b -open sets, so there exist two N- b -closed sets G_1, G_2 with $\psi \subseteq G_1$ and $\xi \subseteq G_2$ such that $\psi \subseteq N_{int}(G_1)$ and $\xi \subseteq N_{int}(G_2)$.

$$\begin{aligned} \text{Now } \psi \cup \xi &\subseteq \xi \quad [\text{since } \psi \subseteq \xi] \\ &\subseteq G_2 \\ \Rightarrow \psi \cup \xi &\subseteq G_2 \end{aligned}$$

Again $\psi \cup \xi \subseteq \xi \subseteq N_{int}(G_2)$, where G_2 is an N- b -closed set in (X, τ) .

Therefore there exists an N-*b*-closed set G_2 with $\psi \cup \xi \subseteq G_2$ in (X, τ) such that $\psi \cup \xi \subseteq N_{\text{int}}(G_2)$.

Hence the union of two g-N-*b*-open sets is again a g-N-open set if one is contained in the other.

Definition 3.3 An NS χ is called a g-N-*b*-open set relative to an NS ψ if there exists an N-*b*-closed set ξ with $\chi \subseteq \psi \cap \xi$ such that $\chi \subseteq N_{\text{int}}(\psi \cap \xi)$.

Theorem 3.7 Assume that (W, τ) is an NTS. If ξ is a g-N-*b*-open set relative to ψ and ψ is a g-N-*b*-open set relative to χ then ξ is a g-N-*b*-open set relative to χ .

Proof. Since ξ is a g-N-*b*-open set relative to ψ so there exists an N-*b*-closed set K with $\xi \subseteq \psi \cap K$ such that $\xi \subseteq N_{\text{int}}(\psi \cap K)$. Similarly, since ψ is a g-N-*b*-open set relative to χ then there exists an N-*b*-closed set L with $\psi \subseteq \chi \cap L$ such that $\psi \subseteq N_{\text{int}}(\chi \cap L)$.

We know that the intersection of two N-*b*-closed sets is again an N-*b*-closed set. So $K \cap L$ is an N-*b*-closed set.

$$\begin{aligned} \text{Now } \xi &\subseteq \psi \cap K \subseteq \chi \cap L \cap K \\ &= \chi \cap (L \cap K) \\ &= \chi \cap G, \text{ where } G = K \cap L \text{ is an N-}b\text{-closed set.} \end{aligned}$$

$$\begin{aligned} \text{Again } \xi &\subseteq N_{\text{int}}(\psi \cap K) \\ &\subseteq N_{\text{int}}(\chi \cap G). \end{aligned}$$

Therefore there exists an N-*b*-closed set G with $\xi \subseteq \chi \cap G$ such that $\xi \subseteq N_{\text{int}}(\chi \cap G)$

Hence ξ is a g-N-*b*-open relative to χ .

4. Conclusion

In this article, we introduce generalized neutrosophic *b*-open set, generalized neutrosophic *b*-interior, generalized neutrosophic *b*-closure and investigate some of their properties. By defining generalized neutrosophic *b*-open set, we prove some theorems on NTSs and few illustrative examples are provided. In the future, we hope that based on these notions in NTSs, many new investigations can be carried out.

References

- Arokiarani, I., Dhavaseelan, R., Jafari, S., & Parimala, M. (2017). On some new notations and functions in neutrosophic topological spaces. *Neutrosophic Sets and Systems*, 16,16-19. doi.org/10.5281/zenodo.831915
- Biswas, P, Pramanik, S. & Giri, B. C. (2014). A new methodology for neutrosophic multi-attribute decision-making with unknown weight information. *Neutrosophic Sets and Systems*, 3, 42-50. doi.org/10.5281/zenodo.571212
- Biswas, P, Pramanik, S. & Giri, B. C. (2016a). Aggregation of triangular fuzzy neutrosophic set information and its application to multi-attribute decision making. *Neutrosophic Sets and Systems*, 12, 20-40. doi.org/10.5281/zenodo.571125
- Biswas, P, Pramanik, S. & Giri, B. C. (2016b). Value and ambiguity index based ranking method of single-valued trapezoidal neutrosophic numbers and its application to multi-attribute decision making. *Neutrosophic Sets and Systems*, 12, 127-138. doi.org/10.5281/zenodo.571154

- Broumi, S., Bakali, A., Talea, M., Smarandache, F., Uluçay, V., Sahin, S, ..., & Pramanik, S. (2018). *Neutrosophic sets: An overview*. In F. Smarandache, & S. Pramanik (Eds., vol.2), *New trends in neutrosophic theory and applications* (pp. 403-434). Brussels: Pons Editions.
- Dalapati, S., Pramanik, S., Alam, S., Smarandache, S., & Roy, T. K. (2017). IN-cross entropy based MAGDM strategy under interval neutrosophic set environment. *Neutrosophic Sets and Systems*, 18, 43-57. <http://doi.org/10.5281/zenodo.1175162>
- Dey, P. P., Pramanik, S. & Giri, B. C. (2016a). An extended grey relational analysis based multiple attribute decision making in interval neutrosophic uncertain linguistic setting. *Neutrosophic Sets and Systems*, 11, 21-30. doi.org/10.5281/zenodo.571228
- Dey, P. P., Pramanik, S. & Giri, B. C. (2016b). Neutrosophic soft multi-attribute decision making based on grey relational projection method. *Neutrosophic Sets and Systems*, 11, 98-106. doi.org/10.5281/zenodo.571576
- Dhavaseelan, R., & Jafari, S. (2018). Generalized neutrosophic closed sets. In F. Smarandache, & S. Pramanik (Eds., vol.2), *New trends in neutrosophic theory and applications* (pp. 261-273). Brussels: Pons Editions.
- Ebenanjar, E., Immaculate, J., & Wilfred, C. B. (2018). On neutrosophic b -open sets in neutrosophic topological space. *Journal of Physics Conference Series*, 1139(1), 012062. [doi: 10.1088/1742-6596/1139/1/012062](http://doi.org/10.1088/1742-6596/1139/1/012062)
- Iswarya, P., & Bageerathi, K. (2016). On neutrosophic semi-open sets in neutrosophic topological spaces. *International Journal of Mathematical Trends and Technology*, 37(3), 214-223.
- Khalid, H. E. (2020). Neutrosophic geometric programming (NGP) with (max-product) operator, an innovative model. *Neutrosophic Sets and Systems*, 32, 269-281. [doi: 10.5281/zenodo.3723803](http://doi.org/10.5281/zenodo.3723803)
- Maheswari, C., Sathyabama, M., & Chandrasekar, S. (2018). Neutrosophic generalized b-closed sets in neutrosophic topological spaces. *Journal of Physics Conference Series*, 1139(1),012065. [doi: 10.1088/1742-6596/1139/1/012065](http://doi.org/10.1088/1742-6596/1139/1/012065)
- Mondal, K., & Pramanik, S. (2015). Neutrosophic decision making model for clay-brick selection in construction field based on grey relational analysis. *Neutrosophic Sets and Systems*, 9, 64-71. doi.org/10.5281/zenodo.34864
- Mondal, K., Pramanik, S. & Smarandache, F. (2016a). Rough neutrosophic TOPSIS for multi-attribute group decision making. *Neutrosophic Sets and Systems*, 13, 105-117. doi.org/10.5281/zenodo.570845
- Mondal, K., Pramanik, S. & Smarandache, F. (2016b). Multi-attribute decision making based on rough neutrosophic variational coefficient similarity measure. *Neutrosophic Sets and Systems*, 13, 3-17. doi.org/10.5281/zenodo.570854
- Peng, X., & Dai, J. (2018). A bibliometric analysis of neutrosophic set: two decades review from 1998 to 2017. *Artificial Intelligence Review*. <https://doi.org/10.1007/s10462-018-9652-0>
- Pramanik, S. (2013). A critical review of Vivekanada's educational thoughts for women education based on neutrosophic logic, *MS Academic*, 3(1), 191-198.
- Pramanik, S. (2016a). Neutrosophic linear goal programming. *Global Journal of Engineering Science and Research Management*, 3(7), 01-11. [doi:10.5281/zenodo.57367](http://doi.org/10.5281/zenodo.57367)

- Pramanik, S. (2016b). Neutrosophic multi-objective linear programming. *Global Journal of Engineering Science and Research Management*, 3(8), 36-46. doi:10.5281/zenodo.59949
- Pramanik, S. (2020). Rough neutrosophic set: an overview. In F. Smarandache, & Broumi, S. (Eds.), *Neutrosophic theories in communication, management and information technology* (pp.275-311). NewYork. Nova Science Publishers.
- Pramanik, S., & Dalapati, S. (2018). A revisit to NC-VIKOR based MAGDM strategy in neutrosophic cubic set environment. *Neutrosophic Sets and Systems*, 21, 131-141. <https://doi.org/10.5281/zenodo.1408665>
- Pramanik, S., Dey, P.P., & Smarandache, F. (2018). Correlation coefficient measures of interval bipolar neutrosophic sets for solving multi-attribute decision making problems. *Neutrosophic Sets and Systems*, 19,70-79. <http://doi.org/10.5281/zenodo.1235151>
- Pramanik, S., & Mallick, R. (2018). VIKOR based MAGDM strategy with trapezoidal neutrosophic numbers. *Neutrosophic Sets and Systems*, 22, 118-130. 10.5281/zenodo.2160840
- Pramanik, S., & Mallick, R. (2019). TODIM strategy for multi-attribute group decision making in trapezoidal neutrosophic number environment. *Complex & Intelligent Systems*, 5 (4), 379–389 <https://doi.org/10.1007/s40747-019-0110>.
- Pramanik, S., Mallick, R., & Dasgupta, A. (2018). Contributions of selected Indian researchers to multi-attribute decision making in neutrosophic environment. *Neutrosophic Sets and Systems*, 20, 108-131. <http://doi.org/10.5281/zenodo.1284870>
- Pramanik, S. & Mondal, K. (2016). Rough bipolar neutrosophic set. *Global Journal of Engineering Science and Research Management*, 3(6), 71-81.
- Pramanik, S., & Roy, T.K. (2014). Neutrosophic game theoretic approach to Indo-Pak conflict over Jammu-Kashmir. *Neutrosophic Sets and Systems*, 2, 82-101.
- Pushpalatha, A., & Nandhini, T. (2019). Generalized closed sets via neutrosophic topological spaces. *Malaya Journal of Matematik*,7(1), 50-54. Doi.org/10.26637/MJM0701/0010
- Rao, V. V., & Srinivasa, R. (2017). Neutrosophic pre-open sets and pre-closed sets in neutrosophic topology. *International Journal of Chem Tech Research*, 10(10), 449-458.
- Salama, A. A., & Alblowi, S. A. (2012a). Neutrosophic set and neutrosophic topological space. *ISOR Journal of Mathematics*, 3(4), 31-35.
- Salama, A. A., & Alblowi, S. A. (2012b). Generalized neutrosophic set and generalized neutrosophic topological space. *Computer Science and Engineering*, 2 (7), 129-132.
- Salama, A. A., Smarandache, F., & Alblowi, S. A. (2014). New neutrosophic crisp topological concepts. *Neutrosophic Sets and System*, 4, 50-54. doi.org/10.5281/zenodo.571462
- Shanthi, V. K., Chandrasekar, S., Safina, & Begam, K. (2018). Neutrosophic generalized semi closed sets in neutrosophic topological spaces. *International Journal of Research in Advent Technology*, 67, 1739-1743.
- Smarandache, F. (1998). *A unifying field in logics, neutrosophy: neutrosophic probability, set and logic*. Rehoboth: American Research Press.
- Smarandache, F. & Pramanik, S. (Eds.). (2016). *New trends in neutrosophic theory and applications*. Brussels: Pons Editions.

Smarandache, F. & Pramanik, S. (Eds.). (2018). *New trends in neutrosophic theory and applications*, Vol.2. Brussels: Pons Editions.

Received: Apr 20, 2020. Accepted: July 15 2020