

9-22-2020

## Fuzzy Neutrosophic Strongly Alpha Generalized Closed Sets in Fuzzy Neutrosophic Topological spaces

Shaymaa F. Matar

Fatimah M. Mohammed

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Matar, Shaymaa F. and Fatimah M. Mohammed. "Fuzzy Neutrosophic Strongly Alpha Generalized Closed Sets in Fuzzy Neutrosophic Topological spaces." *Neutrosophic Sets and Systems* 36, 1 (2020). [https://digitalrepository.unm.edu/nss\\_journal/vol36/iss1/13](https://digitalrepository.unm.edu/nss_journal/vol36/iss1/13)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact [amywinter@unm.edu](mailto:amywinter@unm.edu), [lsloane@salud.unm.edu](mailto:lsloane@salud.unm.edu), [sarahrk@unm.edu](mailto:sarahrk@unm.edu).



# Fuzzy Neutrosophic Strongly Alpha Generalized Closed Sets in Fuzzy Neutrosophic Topological spaces

Shaymaa F. Matar<sup>1</sup> and Fatimah M. Mohammed<sup>2,\*</sup>

<sup>1</sup> Salah Al-din general directorate for education, Tikrit, IRAQ;

shaimaa.1988@yahoo.com

<sup>2</sup>Department of Mathematics, College of Education for Pure Sciences, Tikrit University, Tikrit, IRAQ;

\*<sup>2</sup> Correspondence: e-mail dr.fatimahmahmood@tu.edu.iq

**Abstract:** In this paper, we will define a new set called fuzzy neutrosophic strongly alpha generalized closed set, so we will prove some theorems related to this concept. After that, we will give some interesting properties were investigated and referred to some results related to the new definitions by theorems, propositions to get some relationships among fuzzy neutrosophic strongly alpha generalized closed sets, fuzzy neutrosophic closed sets, fuzzy neutrosophic regular closed sets, fuzzy neutrosophic alpha closed sets, fuzzy neutrosophic alpha generalized closed sets and fuzzy neutrosophic pre closed sets which are compared with necessary examples based of fuzzy neutrosophic topological spaces.

**Keywords:** Fuzzy neutrosophic set, fuzzy neutrosophic topological space, fuzzy neutrosophic strongly alpha generalized closed set.

---

## 1. Introduction

The concept of fuzzy set "FS" was introduced by Lotfi Zadeh in 1965 [1], then Chang depended the fuzzy set to introduce the concept of fuzzy topological space "FTS" in 1968 [7]. After that the concept of fuzzy set was developed into the concept of intuitionistic fuzzy set "IFS" by Atanassov in 1983 [4-6], the intuitionistic fuzzy set gives a degree of membership and a degree of non-membership functions. Coker in 1997 [7] relied on intuitionistic fuzzy set to introduced the concept of intuitionistic fuzzy topological space."IFTS". In 2005 Smaradache [23] study the concept of neutrosophic set. "NS". After that and as developed the term of neutrosophic set, Salama has studied neutrosophic topological space "NTS" and many of its applications [18-21]. In 2013 Arockiarani Sumathi and Martina Jency [2] introduced the concept of fuzzy neutrosophic set as generalizes the concept of fuzzy set and intuitionistic fuzzy set. where each element had three associated defining functions on the universe of discourse X, namely the membership function (T), indeterminacy function (I), the non-membership function (F) that is added an indeterminacy degree between the

degree of membership and the degree of non- membership. In 2012 Salama and Alblowi defined fuzzy neutrosophic topological space [18].

In the present work, we will generalized the concept of strongly alpha generalized closed set in fuzzy neutrosophic topological spaces which was studied by Santhi and Sakthivel in 2011 [22] via intuitionistic topological spaces and generalizing our works in 2018 [ 9,10 ], the new set will called fuzzy neutrosophic strongly alpha generalized closed set in fuzzy neutrosophic topological spaces.

Finally, there are many application of neutrosophic sets in many fields so we can enhance our work, we will try in the future to applied this work in different fields such as many authors applications see [11] and [13-17].

## 2. Preliminaries:

In this section, we will define some basic definitions and some operations which are useful in our present study.

**Definition 2.1 [18]:** Let  $X$  be a non-empty fixed set. The fuzzy neutrosophic set (FNS, for short),  $\eta_N$  is an object having the form  $\eta_N = \{ \langle x, \mu_{\eta_N}(x), \sigma_{\eta_N}(x), \nu_{\eta_N}(x) \rangle : x \in X \}$  where the functions  $\mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N}: X \rightarrow [0, 1]$  denote the degree of membership function (namely  $\mu_{\eta_N}(x)$ ), the degree of indeterminacy function (namely  $\sigma_{\eta_N}(x)$ ) and the degree of non-membership (namely  $\nu_{\eta_N}(x)$ ) respectively of each element  $x \in X$  to the set  $\eta_N$  and  $0 \leq \mu_{\eta_N}(x) + \sigma_{\eta_N}(x) + \nu_{\eta_N}(x) \leq 3$ , for each  $x \in X$ .

**Remark 2.2 [18]:** FNS  $\eta_N = \{ \langle x, \mu_{\eta_N}(x), \sigma_{\eta_N}(x), \nu_{\eta_N}(x) \rangle : x \in X \}$  can be identified to an ordered triple  $\langle \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  in  $[0, 1]$  on  $X$ .

**Definition 2.3 [18]:** Let  $X$  be a non-empty set and the FNSs  $\eta_N$  and  $\gamma_N$  be in the form:

$\eta_N = \{ \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle : x \in X \}$  and  $\gamma_N = \{ \langle x, \mu_{\gamma_N}, \sigma_{\gamma_N}, \nu_{\gamma_N} \rangle : x \in X \}$  on  $X$  then:

- i.  $\eta_N \subseteq \gamma_N$  iff  $\mu_{\eta_N} \leq \mu_{\gamma_N}, \sigma_{\eta_N} \leq \sigma_{\gamma_N}$  and  $\nu_{\eta_N} \geq \nu_{\gamma_N}$ .
- ii.  $\eta_N = \gamma_N$  iff  $\eta_N \subseteq \gamma_N$  and  $\gamma_N \subseteq \eta_N$ ,
- iii.  $1_N^- \eta_N = \{ \langle x, \nu_{\eta_N}, 1 - \sigma_{\eta_N}, \mu_{\eta_N} \rangle : x \in X \}$ ,
- iv.  $\eta_N \cup \gamma_N = \{ \langle x, \text{Max}(\mu_{\eta_N}, \mu_{\gamma_N}), \text{Max}(\sigma_{\eta_N}, \sigma_{\gamma_N}), \text{Min}(\nu_{\eta_N}, \nu_{\gamma_N}) \rangle : x \in X \}$ ,
- v.  $\eta_N \cap \gamma_N = \{ \langle x, \text{Min}(\mu_{\eta_N}, \mu_{\gamma_N}), \text{Min}(\sigma_{\eta_N}, \sigma_{\gamma_N}), \text{Max}(\nu_{\eta_N}, \nu_{\gamma_N}) \rangle : x \in X \}$ ,
- vi.  $0_N = \langle x, 0, 0, 1 \rangle$  and  $1_N = \langle x, 1, 1, 0 \rangle$ .

**Definition 2.4 [18]:** "Fuzzy neutrosophic topology (FNT, for short) on a non-empty set  $X$  is a family  $\tau_N$  of fuzzy neutrosophic subsets in  $X$  satisfying the following axioms.

- i.  $0_N, 1_N \in \tau_N$ ,
- ii.  $\eta_{N1} \cap \eta_{N2} \in \tau_N$  for any  $\eta_{N1}, \eta_{N2} \in \tau_N$ ,
- iii.  $\cup \eta_{Ni} \in \tau_N, \forall \{ \eta_{Ni} : i \in J \} \subseteq \tau_N$ ."

In this case the pair  $(X, \tau_N)$  is called fuzzy neutrosophic topological space (FNTS, for short). The elements of  $\tau_N$  are called fuzzy neutrosophic open set (FNOS, for short). The complement of FNOS in the FNTS  $(X, \tau_N)$  is called fuzzy neutrosophic closed set (FNCS, for short).

**Definition 2.5 [18]:** Let  $(X, \tau_N)$  be FNTS and  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be FNS in  $X$ . Then the fuzzy neutrosophic closure of  $\eta_N$  (FNCL, for short) and fuzzy neutrosophic interior of  $\eta_N$  (FNIn, for short) are defined by:

$$\text{FNCL}(\eta_N) = \bigcap \{C_N: C_N \text{ is FNCS in } X \text{ and } \eta_N \subseteq C_N \},$$

$$\text{FNIn}(\eta_N) = \bigcup \{O_N: O_N \text{ is FNOS in } X \text{ and } O_N \subseteq \eta_N \}.$$

We know, FNCL( $\eta_N$ ) is FNCS and FNIn( $\eta_N$ ) is FNOS in  $X$ . Further,

- i.  $\eta_N$  is FNCS in  $X$  iff FNCL( $\eta_N$ ) =  $\eta_N$ ,
- ii.  $\eta_N$  is FNOS in  $X$  iff FNIn( $\eta_N$ ) =  $\eta_N$ .

**Proposition 2.6 [25]:** Let  $(X, \tau_N)$  is FNTS and  $\eta_N, \gamma_N$  are FNSs in  $X$ . Then the following properties hold:

- i. FNIn( $\eta_N$ )  $\subseteq$   $\eta_N$  and  $\eta_N \subseteq$  FNCL( $\eta_N$ ),
- ii.  $\eta_N \subseteq \gamma_N \implies$  FNIn( $\eta_N$ )  $\subseteq$  FNIn( $\gamma_N$ ) and  $\eta_N \subseteq \gamma_N \implies$  FNCL( $\eta_N$ )  $\subseteq$  FNCL( $\gamma_N$ ),
- iii. FNIn(FNIn( $\eta_N$ )) = FNIn( $\eta_N$ ) and FNCL(FNCL( $\eta_N$ )) = FNCL( $\eta_N$ ),
- iv. FNIn( $\eta_N \cap \gamma_N$ ) = FNIn( $\eta_N$ )  $\cap$  FNIn( $\gamma_N$ ) and FNCL( $\eta_N \cup \gamma_N$ ) = FNCL( $\eta_N$ )  $\cup$  FNCL( $\gamma_N$ ),
- v. FNIn( $1_N$ ) =  $1_N$  and FNCL( $1_N$ ) =  $1_N$ ,
- vi. FNIn( $0_N$ ) =  $0_N$  and FNCL( $0_N$ ) =  $0_N$ .

**Definition 2.7 [9]:** FNS  $\eta_N$  in FNTS  $(X, \tau_N)$  is called:

- i. Fuzzy neutrosophic regular closed set (FNRCFS, for short) if  $\eta_N = \text{FNCL}(\text{FNIn}(\eta_N))$ .
- ii. Fuzzy neutrosophic pre closed set (FNPCFS, for short) if  $\text{FNCL}(\text{FNIn}(\eta_N)) \subseteq \eta_N$ .
- iii. Fuzzy neutrosophic  $\alpha$  closed set (FN $\alpha$ CS, for short) if  $\text{FNCL}(\text{FNIn}(\text{FNCL}(\eta_N))) \subseteq \eta_N$ .

**Definition 2.8 [10]:** Let  $(X, \tau_N)$  be FNTS and  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be FNS in  $X$ . Then the fuzzy neutrosophic alpha closure of  $\eta_N$  (FN $\alpha$ CL, for short) and fuzzy neutrosophic alpha interior of  $\eta_N$  (FN $\alpha$ In,

for short) are defined by:

$$\text{FN}\alpha\text{CL}(\eta_N) = \bigcap \{C_N: C_N \text{ is FN}\alpha\text{CS in } X \text{ and } \eta_N \subseteq C_N \},$$

$$\text{FN}\alpha\text{In}(\eta_N) = \bigcup \{O_N: O_N \text{ is FN}\alpha\text{OS in } X \text{ and } O_N \subseteq \eta_N \}.$$

We know, FN $\alpha$ CL( $\eta_N$ ) is FN $\alpha$ CLS and FN $\alpha$ In( $\eta_N$ ) is FN $\alpha$ OS in  $X$ . Further,

- i.  $\eta_N$  is FN $\alpha$ CS in  $X$  iff FN $\alpha$ CL( $\eta_N$ ) =  $\eta_N$ ,
- ii.  $\eta_N$  is FN $\alpha$ OS in  $X$  iff FN $\alpha$ In( $\eta_N$ ) =  $\eta_N$ .

**Definition 2.9 [9,10]:** Fuzzy neutrosophic sub set  $\eta_N$  of FNTS  $(X, \tau_N)$  is called:

- i. fuzzy neutrosophic generalized closed set (FNGCS, for short ) if FNCL( $\eta_N$ )  $\subseteq$   $U_N$  wherever,  $\eta_N \subseteq U_N$  and  $U_N$  is FNOS in  $X$ . And  $\eta_N$  is said to be fuzzy neutrosophic generalized open set (FNGOS, for short) if the complement  $1_N - \eta_N$  is FNGCS set in  $(X, \tau_N)$ .
- ii. fuzzy neutrosophic alpha generalized closed set (FN $\alpha$ GCS, for short ) if FN $\alpha$ CL( $\eta_N$ )  $\subseteq$   $U_N$  wherever,  $\eta_N \subseteq U_N$  and  $U_N$  is FNOS in  $X$ . And  $\eta_N$  is said to be fuzzy neutrosophic

alpha generalized open set (FN $\alpha$ GOS, for short) if the complement  $1_N - \eta_N$  is FN $\alpha$ GCS set in  $(X, \tau_N)$ .

### 3. Fuzzy Neutrosophic Strongly Alpha Generalized Closed Sets in Fuzzy Neutrosophic Topological Spaces.

Now, we will introduce the concept of fuzzy neutrosophic strongly alpha generalized closed set in fuzzy neutrosophic topological spaces.

**Definition 3.1:** Fuzzy neutrosophic subset  $\eta_N$  of FNTS  $(X, \tau_N)$  is called fuzzy neutrosophic strongly alpha generalized closed set (FNS $\alpha$ GCS, for short) if FN $\alpha$ CL( $\eta_N$ )  $\subseteq$   $U_N$  wherever,  $\eta_N \subseteq U_N$  and  $U_N$  is FNGOS in  $X$ .

**Example 3.2:** Let  $X = \{a, b\}$  define FNS  $\eta_N$  in  $X$  as follows:

$\eta_N = \langle x, (0.2_{(a)}, 0.3_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.8_{(a)}, 0.7_{(b)}) \rangle$ , where the family  $\tau_N = \{0_N, 1_N, \eta_N\}$ .

If we take,  $\psi_N = \langle x, (0.8_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.1_{(a)}, 0_{(b)}) \rangle$ .

And,  $U_N = 1_N$  where  $U_N$  is FNGOS such that,  $\psi_N \subseteq U_N$ . Then, FN $\alpha$ CL( $\psi_N$ ) =  $1_N$ . So, FN $\alpha$ CL( $\psi_N$ )  $\subseteq$   $U_N$ .

Hence,  $\psi_N$  is FNS $\alpha$ GCS.

**Theorem 3.3:** For any FNSs, the following statements are true in general:

- i. Every FNOS is FNGOS.
- ii. Every FNCS is FN $\alpha$ CS.
- iii. Every FNCS is FNS $\alpha$ GCS.
- iv. Every FNRCS is FNS $\alpha$ GCS.
- v. Every FN $\alpha$ CS is FNS $\alpha$ GCS.
- vi. Every FN $\alpha$ GCS is FNS $\alpha$ GCS.
- vii. Every FNRCS is FNCS.
- viii. Every FN $\alpha$ CS is FN $\alpha$ GCS.

**Proof:**

i. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be FNOS in the FNTS  $(X, \tau_N)$ .

Then by **Definition 2.5 ii** we get, FNIn( $\eta_N$ ) =  $\eta_N$ .

Now, let  $U_N$  is FNCS such that,  $U_N \subseteq \eta_N$ . Therefore, FNIn( $\eta_N$ ) =  $\eta_N \supseteq U_N$ .

Hence,  $\eta_N$  is FNGOS in  $(X, \tau_N)$ .

ii. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be FNCLS in the FNTS  $(X, \tau_N)$ .

Then by **Definition 2.5 (i)** we get, FNCL( $\eta_N$ ) =  $\eta_N \dots \dots (1)$ .

And by **Proposition 2.6 i** we get, FNIn( $\eta_N$ )  $\subseteq$   $\eta_N$ .

So we get,  $\text{FNIn}(\text{FNCL}(\eta_N)) \subseteq \eta_N$

This implies  $\text{FNCL}(\text{FNIn}(\text{FNCL}(\eta_N))) \subseteq \text{FNCL}(\eta_N)$ .

So by **(1)** we get,  $\text{FNCL}(\text{FNIn}(\text{FNCL}(\eta_N))) \subseteq \eta_N$ .

Hence,  $\eta_N$  is  $\text{FN}\alpha\text{CS}$  in  $(X, \tau_N)$ .

iii. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be  $\text{FNCS}$  in  $\text{FNTS}(X, \tau_N)$ .

Then by **Definition 2.5 (i)** we get,  $\text{FNCL}(\eta_N) = \eta_N$ . Now, let  $U_N$  be  $\text{FNGOS}$  such that,  $\eta_N \subseteq U_N$ .

Since,  $\text{FN}\alpha\text{CL}(\eta_N) \subseteq \text{FNCL}(\eta_N)$  by **Definition 2.5 and Definition 2.8**.

So we get,  $\text{FN}\alpha\text{CL}(\eta_N) \subseteq \text{FNCL}(\eta_N) = \eta_N \subseteq U_N$ .

Hence,  $\eta_N$  is  $\text{FNS}\alpha\text{GCS}$  in  $(X, \tau_N)$ .

iv. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be  $\text{FNRCS}$  in the  $\text{FNTS}(X, \tau_N)$ .

Then,  $\text{FNCL}(\text{FNIn}(\eta_N)) = \eta_N \dots \dots (1)$ .

This implies,  $\text{FNCL}(\text{FNIn}(\eta_N)) = \text{FNCL}(\eta_N) \dots \dots (2)$ .

Now, let  $U_N$  be  $\text{FNGOS}$  such that,  $\eta_N \subseteq U_N$ .

From **(1) and (2)** we get,  $\text{FNCL}(\eta_N) = \eta_N$ .

That  $\eta_N$  is  $\text{FNCS}$  in  $X$ .

So by **iii** we get,  $\text{FN}\alpha\text{CL}(\eta_N) \subseteq \text{FNCL}(\eta_N) = \eta_N \subseteq U_N$ .

Hence,  $\eta_N$  is  $\text{FNS}\alpha\text{GCS}$  in  $(X, \tau_N)$ .

v. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be  $\text{FN}\alpha\text{CLOS}$  in the  $\text{FNTS}(X, \tau_N)$ .

Then by **Definition 2.8 i** we get,  $\text{FN}\alpha\text{CL}(\eta_N) = \eta_N$ .

Now, let  $U_N$  be  $\text{FNGOS}$  such that,  $\eta_N \subseteq U_N$ . So,  $\text{FN}\alpha\text{CL}(\eta_N) = \eta_N \subseteq U_N$ .

Hence,  $\eta_N$  is  $\text{FNS}\alpha\text{GCS}$  in  $(X, \tau_N)$ .

vi. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be  $\text{FN}\alpha\text{GCS}$  in the  $\text{FNTS}(X, \tau_N)$ .

Then,  $\text{FN}\alpha\text{CL}(\eta_N) \subseteq U_N$ ,  $\eta_N \subseteq U_N$  and  $U_N$  be  $\text{FNOS}$ , so by **i** we get,  $\text{FNOS}$  be  $\text{FNGOS}$  in  $(X, \tau_N)$ .

Therefore,  $\text{FN}\alpha\text{CL}(\eta_N) \subseteq U_N$ ,  $\eta_N \subseteq U_N$  and  $U_N$  be  $\text{FNGOS}$ . Hence,  $\eta_N$  is  $\text{FNS}\alpha\text{GCS}$  in  $(X, \tau_N)$ .

vii. Let  $\eta_N = \langle x, \mu_{\eta_N}, \sigma_{\eta_N}, \nu_{\eta_N} \rangle$  be  $\text{FN}\alpha\text{CS}$  in the  $\text{FNTS}(X, \tau_N)$ . Then,  $\text{FN}\alpha\text{CL}(\eta_N) = \eta_N$ .

Now, let  $U_N$  be  $\text{FNOS}$  such that,  $\eta_N \subseteq U_N$ , so,  $\text{FN}\alpha\text{CL}(\eta_N) = \eta_N \subseteq U_N$ .

Hence,  $\eta_N$  is  $\text{FN}\alpha\text{GCS}$  in  $(X, \tau_N)$ .

**Remark 3.4:** The convers of **Theorem 3.3** is not true and this can be clarified in the following examples.

**Example 3.5:**

- i.** Let  $X = \{a, b\}$  define FNS  $\eta_N$  in  $X$  as follows:  
 $\eta_N = \langle x, (0.5_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.5_{(a)}, 0.2_{(b)}) \rangle$ .  
 The family  $\tau_N = \{0_N, 1_N, \eta_N\}$  be FNT.  
 If we take,  $\psi_N = \langle x, (0.1_{(a)}, 0.6_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.9_{(a)}, 0.3_{(b)}) \rangle$ .  
 And let,  $U_N = 0_N$ , where  $U_N$  be FNCS such that,  $U_N \subseteq \psi_N$ .  
 Then,  $\text{FNIn}(\psi_N) = \langle x, (0_{(a)}, 0_{(b)}), (0_{(a)}, 0_{(b)}), (1_{(a)}, 1_{(b)}) \rangle \subseteq \langle x, (0.1_{(a)}, 0.6_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.9_{(a)}, 0.3_{(b)}) \rangle$  such that,  $(0_{(a)}, 0_{(b)}) \leq (0.1_{(a)}, 0.6_{(b)})$ ,  $(0_{(a)}, 0_{(b)}) \leq (0.5_{(a)}, 0.5_{(b)})$  and  $(1_{(a)}, 1_{(b)}) \geq (0.9_{(a)}, 0.3_{(b)}) = 0_N$ . So,  $\text{FNIn}(\psi_N) \supseteq U_N$ . Hence,  $\psi_N$  is FNGOS but, not FNOS.  
 Since  $\psi_N \notin \tau_N$ .
- ii.** Let  $X = \{a\}$  define the FNSs  $\eta_N$  and  $\gamma_N$  in  $X$  as follows:  
 $\eta_N = \langle x, (0.5_{(a)}), (0.4_{(a)}), (0.7_{(a)}) \rangle$ ,  $\gamma_N = \langle x, (0.4_{(a)}), (0.1_{(a)}), (0.8_{(a)}) \rangle$ .  
 The family  $\tau_N = \{0_N, 1_N, \eta_N, \gamma_N\}$  be FNT.  
 If we take,  $\psi_N = \langle x, (0.8_{(a)}), (0.6_{(a)}), (0.5_{(a)}) \rangle$ .  
 Then,  $\text{FNCL}(\psi_N) = \langle x, (0.8_{(a)}), (0.9_{(a)}), (0.4_{(a)}) \rangle$ . And,  $\text{FNIn}(\text{FNCL}(\psi_N)) = \langle x, (0.5_{(a)}), (0.4_{(a)}), (0.7_{(a)}) \rangle$ . So,  $\text{FNCL}(\text{FNIn}(\text{FNCL}(\psi_N))) = \langle x, (0.7_{(a)}), (0.6_{(a)}), (0.5_{(a)}) \rangle$ .  
 Therefore,  $\langle x, (0.7_{(a)}), (0.6_{(a)}), (0.5_{(a)}) \rangle \supseteq \psi_N$ .  
 Hence,  $\psi_N$  is  $\text{FN}\alpha\text{CS}$  but not FNCS. Since  $\psi_N \notin 1_N - \tau_N$ .
- iii.** Take **Example 3.2**. Then,  $\psi_N$  is  $\text{FNS}\alpha\text{GCS}$  but, not FNCS.  
 Since,  $\psi_N \notin 1_N - \tau_N$ .
- iv.** Take **Example 3.2**. Then  $\psi_N$  is  $\text{FNS}\alpha\text{GCS}$  but, not FNRCS.  
 Since,  $\text{FNIn}(\psi_N) = \langle x, (0.2_{(a)}, 0.3_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.8_{(a)}, 0.7_{(b)}) \rangle$  and  
 $\text{FNCL}(\text{FNIn}(\psi_N)) = \langle x, (0.8_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.2_{(a)}, 0.3_{(b)}) \rangle \neq \psi_N$ .
- v.** Let  $X = \{a, b\}$  define the FNSs  $\eta_N$  and  $\gamma_N$  in  $X$  as follows:  
 $\eta_N = \langle x, (0.4_{(a)}, 0.2_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.6_{(a)}, 0.7_{(b)}) \rangle$ ,  
 $\gamma_N = \langle x, (0.8_{(a)}, 0.8_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.2_{(a)}, 0.2_{(b)}) \rangle$ .  
 The family  $\tau_N = \{0_N, 1_N, \eta_N, \gamma_N\}$  be FNT.  
 Now if,  $\psi_N = \langle x, (0.6_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.4_{(a)}, 0.3_{(b)}) \rangle$ .  
 By **Theorem 3.3 i**. If  $U_N$  is FNOS then is FNGOS.  
 So,  $U_N = \gamma_N$  where,  $U_N$  be FNGOS such that,  $\psi_N \subseteq U_N$ .  
 By **Theorem 3.3 ii**. Every FNCS is  $\text{FN}\alpha\text{CS}$ .

Then,  $\text{FN}\alpha\text{CL}(\psi_N) = 1_N - \eta_N$ . Therefore  $\text{FN}\alpha\text{CL}(\psi_N) \subseteq U_N$ .

Hence,  $\psi_N$  is  $\text{FNS}\alpha\text{GCS}$  but, not  $\text{FN}\alpha\text{CS}$ .

Since,  $\text{FNCL}(\psi_N) = 1_N - \eta_N$ ,  $\text{FNIn}(\text{FNCL}(\psi_N)) = \eta_N$  and

$\text{FNCL}(\text{FNIn}(\text{FNCL}(\psi_N))) = 1_N - \eta_N \not\subseteq \psi_N$ .

**vi.** Let  $X = \{a\}$  define the FNSs  $\eta_N$  and  $\gamma_N$  in  $X$  as follows:

$\eta_N = \langle x, (0.5_{(a)}), (0.5_{(a)}), (0.5_{(a)}) \rangle$ ,  $\gamma_N = \langle x, (0.5_{(a)}), (0_{(a)}), (1_{(a)}) \rangle$ .

The family  $\tau_N = \{0_N, 1_N, \eta_N, \gamma_N\}$  be FNT.

Now if,  $\psi_N = \langle x, (0.6_{(a)}), (0.6_{(a)}), (0.6_{(a)}) \rangle$ .

Let  $U_N = \langle x, (1_{(a)}), (1_{(a)}), (0.4_{(a)}) \rangle$  be FNGOS such that,  $\psi_N \subseteq U_N$ .

Then,  $\text{FN}\alpha\text{CL}(\psi_N) = 1_N - \gamma_N$ . So  $\text{FN}\alpha\text{CL}(\psi_N) \subseteq U_N$ .

Hence,  $\psi_N$  is  $\text{FNS}\alpha\text{GCS}$  but, not  $\text{FN}\alpha\text{GCS}$ .

Since,  $U_N$  is FNGOS but not FNOS.

**vii.** Let  $X = \{a\}$  define the FNSs  $\eta_N$  and  $\gamma_N$  in  $X$  as follows:

$\eta_N = \langle x, (0.5_{(a)}), (0.5_{(a)}), (0.7_{(a)}) \rangle$ ,  $\gamma_N = \langle x, (0.4_{(a)}), (0_{(a)}), (1_{(a)}) \rangle$ .

The family  $\tau_N = \{0_N, 1_N, \eta_N, \gamma_N\}$  be FNT.

Now if,  $\psi_N = \langle x, (1_{(a)}), (1_{(a)}), (0.4_{(a)}) \rangle$ .

Then,  $\psi_N$  is FNCS. Since  $\psi_N \in 1_N - \tau_N$  but, not FNRCs.

Since  $\text{FNIn}(\psi_N) = \langle x, (0.5_{(a)}), (0.5_{(a)}), (0.7_{(a)}) \rangle$  and

$\text{FNCL}(\text{FNIn}(\psi_N)) = \langle x, (0.7_{(a)}), (0.5_{(a)}), (0.5_{(a)}) \rangle \neq \psi_N$ .

**viii.** Let  $X = \{a\}$  define the FNSs  $\eta_N$  and  $\gamma_N$  in  $X$  as follows:

$\eta_N = \langle x, (0.5_{(a)}), (0.5_{(a)}), (0.6_{(a)}) \rangle$ ,  $\gamma_N = \langle x, (0.5_{(a)}), (0_{(a)}), (1_{(a)}) \rangle$ .

The family  $\tau_N = \{0_N, 1_N, \eta_N, \gamma_N\}$  be FNT.

Now if,  $\psi_N = \langle x, (0.6_{(a)}), (0.6_{(a)}), (0.6_{(a)}) \rangle$ .

Let,  $U_N = 1_N$  be FNOS such that,  $\psi_N \subseteq U_N$ .

Then,  $\text{FNCL}(\psi_N) = \langle x, (1_{(a)}), (1_{(a)}), (0.5_{(a)}) \rangle$  and  $\text{FNCL}(\psi_N) \subseteq U_N$ .

Hence,  $\psi_N$  is  $\text{FN}\alpha\text{GCS}$  but, not  $\text{FN}\alpha\text{CS}$ .

Since,  $\text{FNCL}(\psi_N) = \langle x, (1_{(a)}), (1_{(a)}), (0.5_{(a)}) \rangle$ ,  $\text{FNIn}(\text{FNCL}(\psi_N)) = \langle x, (0.5_{(a)}), (0.5_{(a)}), (0.6_{(a)}) \rangle$  and

$\text{FNCL}(\text{FNIn}(\text{FNCL}(\psi_N))) = \langle x, (0.6_{(a)}), (0.5_{(a)}), (0.5_{(a)}) \rangle \not\subseteq \psi_N$ .

**Remark 3.6: i.** The relation between FNPCS and  $\text{FNS}\alpha\text{GCS}$  is independent and this can be clarified in the next example.



- ii. The intersection of two FNS $\alpha$ GCS is not FNS $\alpha$ GCS in general and we explained it in the next example.

**Example 3.7:**

i. (1) Let  $X = \{a, b\}$  define FNS  $\eta_N$  in  $X$  as follows:

$$\eta_N = \langle x, (0.5_{(a)}, 0.5_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.4_{(a)}, 0.5_{(b)}) \rangle.$$

The family  $\tau_N = \{0_N, 1_N, \eta_N\}$  be FNT.

$$\text{Now if, } \psi_N = \langle x, (0.5_{(a)}, 0.4_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.6_{(a)}, 0.5_{(b)}) \rangle.$$

Then,  $\text{FNIn}(\psi_N) = 0_N$  and  $\text{FNCL}(\text{FNIn}(\psi_N)) = 0_N$ . So,  $\text{FNCL}(\text{FNIn}(\psi_N)) \subseteq \psi_N$ .

Hence,  $\psi_N$  is FNPCS but, not FNS $\alpha$ GCS. Since

Let,  $U_N = \eta_N$ , where  $U_N$  be FNGOS such that,  $\psi_N \subseteq U_N$ . Then,  $\text{FN}\alpha\text{CL}(\psi_N) = 1_N$ . So  $\text{FN}\alpha\text{CL}(\psi_N) \not\subseteq U_N$ .

(2) Let  $X = \{a, b\}$  define the FNSs  $\eta_N$  and  $\gamma_N$  in  $X$  as follows:

$$\eta_N = \langle x, (0.5_{(a)}, 0.2_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.5_{(a)}, 0.7_{(b)}) \rangle,$$

$$\gamma_N = \langle x, (0.8_{(a)}, 0.8_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.2_{(a)}, 0.2_{(b)}) \rangle.$$

The family  $\tau_N = \{0_N, 1_N, \eta_N, \gamma_N\}$  be FNT.

$$\text{Now if, } \psi_N = \langle x, (0.5_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.5_{(a)}, 0.3_{(b)}) \rangle.$$

Let,  $U_N = \gamma_N$ , where  $U_N$  be FNGOS such that,  $\psi_N \subseteq U_N$ .

$$\text{Then, } \text{FN}\alpha\text{CL}(\psi_N) = \langle x, (0.5_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.5_{(a)}, 0.2_{(b)}) \rangle \subseteq U_N.$$

Hence,  $\psi_N$  is FNS $\alpha$ GCS but, not FNPCS.

$$\text{Since, } \text{FNIn}(\psi_N) = \eta_N \text{ and } \text{FNCL}(\text{FNIn}(\psi_N)) = \langle x, (0.5_{(a)}, 0.7_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.5_{(a)}, 0.2_{(b)}) \rangle.$$

So,  $\text{FNCL}(\text{FNIn}(\psi_N)) \not\subseteq \psi_N$ .

ii. Let  $X = \{a, b\}$  define FNS  $\eta_N$  in  $X$  as follows:  $\eta_N = \langle x, (0.5_{(a)}, 0_{(b)}), (0.5_{(a)}, 0.5_{(b)}), (0.1_{(a)}, 1_{(b)}) \rangle$ .


The family  $\tau_N = \{0_N, 1_N, \eta_N\}$  be FNT.

Now if,  $\psi_{N1} = \langle x, (0.2_{(a)}, 1_{(b)}), (1_{(a)}, 1_{(b)}), (0.7_{(a)}, 0_{(b)}) \rangle$  and  $\psi_{N2} = \langle x, (0.6_{(a)}, 0_{(b)}), (1_{(a)}, 1_{(b)}), (0.3_{(a)}, 1_{(b)}) \rangle$  are FNS $\alpha$ GCS. But,  $\psi_{N1} \cap \psi_{N2} = \langle x, (0.2_{(a)}, 0_{(b)}), (1_{(a)}, 1_{(b)}), (0.7_{(a)}, 1_{(b)}) \rangle$ .

Now let,  $U_N = \eta_N$ , where  $U_N$  be FNGOS such that,  $\psi_{N1} \cap \psi_{N2} \subseteq U_N$ . Then,  $\text{FN}\alpha\text{CL}(\psi_{N1} \cap \psi_{N2}) = 1_N \not\subseteq U_N$ .

Hence,  $\psi_{N1} \cap \psi_{N2}$  is not FNS $\alpha$ GCS.

**Remark 3.8:** The next diagram explains the relationships among different sets in the FNTS and the convers is not true in general.



**Diagram 3.1**

### 5. Conclusions

In this present paper, we have defined new class of neutrosophic generalized closed sets called, fuzzy neutrosophic strongly alpha generalized closed set in fuzzy neutrosophic topological spaces. Many results have been discussed with some properties. Further, we giving some theorems, propositions and provided some useful examples where such properties failed to be preserved in order to get the relations between fuzzy neutrosophic strongly alpha generalized closed set and existing fuzzy neutrosophic closed sets in fuzzy neutrosophic topological spaces . We think, our studied class of sets belongs to the new class of fuzzy neutrosophic sets which is useful not only in the deepening of our understanding of some special features of the well-known notions of fuzzy neutrosophic topology but also useful in neutrosophic control theory.

### Acknowledgments:

In this section the authors would like to thank the referees for their valuable suggestions to improve the paper.

### References

1. L A.. Zadeh. (1965). Fuzzy Sets, Inform. and Control, Vol. 8, 338- 353.
2. I. Arockiarani, I.R.Sumathi & J.Martina Jency. (2013). Fuzzy Neutrosophic Soft Topological Spaces, IJMA, Vol. 4, 225-238.
3. I. Arockiarani & J. Martina Jency. (2014). More on Fuzzy Neutrosophic Sets and Fuzzy Neutrosophic Topological Spaces, IJIRS, 3(5), 642-652.

4. K. Atanassov & S. Stoeva. (1983). Intuitionistic Fuzzy Sets, in : Polish Syrup. On Interval and Fuzzy Mathematics, Poznan, 23-26.
5. K. Atanassov. (1986). Intuitionistic Fuzzy Sets, Fuzzy Sets and Systems, Vol. 20, 87-96.
6. K. Atanassov. (1988). Review and New Results on Intuitionistic Fuzzy Sets, Preprint IM- MFAIS, Sofia, 1-88.
7. C.L. Chang. (1968). Fuzzy Topological Space, J. Math. Anal. Appl., Vol. 24, 182-190.
8. D. Coker. (1997). An Introduction to Intuitionistic Fuzzy Topological Spaces, Fuzzy Sets and System, Vol. 88, 81-89.
9. F. M. Mohammed & Shaymaa F. Matar. (2018). Fuzzy Neutrosophic  $\alpha^m$ - closed set in Fuzzy Neutrosophic Topological Spaces, Neutrosophic set and systems, Vol. 21, 56-65.
10. F. M. Mohammed, Anas A. Hijab & Shaymaa F. Matar . (2018). Fuzzy Neutrosophic Weakly-Generalized closed set in Fuzzy Neutrosophic Topological Spaces, University of Anbar for Pure Science, Vol. 12, 63-73.
11. F. M. Mohammed & Sarah W. Raheem. (2020). Generalized b Closed Sets and Generalized b Open Sets in Fuzzy Neutrosophic bi-Topological Spaces, Neutrosophic set and systems, Vol.35, 188-197.
12. D. Jayanthi. (2018).  $\alpha$  Generalized closed set in Neutrosophic Topological Spaces, IJMTT, ISSN: 2231-5373, 88-91.
13. Abdel-Basset, M., Mohamed, R., Elhoseny, M., & Chang, V. (2020). Evaluation framework for smart disaster response systems in uncertainty environment. Mechanical Systems and Signal Processing, 145, 106-941.
14. Abdel-Basset, M., Ali, M., & Atef, A. (2020). Uncertainty assessments of linear time-cost tradeoffs using neutrosophic set. Computers & Industrial Engineering, 141, 106-286
15. Abdel-Basset, M., Ali, M., & Atef, A. (2020). Resource levelling problem in construction projects under neutrosophic environment. The Journal of Supercomputing, 76(2), 964-988.
16. Abdel-Basset, M., Gamal, A., Son, L. H., & Smarandache, F. (2020). A Bipolar Neutrosophic Multi Criteria Decision Making Framework for Professional Selection. Applied Sciences, 10(4), 12-20.
17. Abdel-Basset, M., Mohamed, R., Zaied, A. E. N. H., Gamal, A., & Smarandache, F. (2020). Solving the supply chain problem using the best-worst method based on a novel Plithogenic model. In Optimization Theory Based on Neutrosophic and Plithogenic Sets (pp. 1-19). Academic Press.
18. A. A. Salama & S. A. Alblowi. (2012). Neutrosophic Set and Neutrosophic Topological Spaces, IOSR Journal of Mathematics, 3(4), 31-35.
19. A. A. Salama, Florentin Smarandache & S. A. Alblowi. (2014). Characteristic Function of Neutrosophic Set, Neutrosophic Sets and Systems, Vol. 3, 14-17.
20. A. A. Salama, Florentin Smarandache & Valeri Kromov. (2014). Neutrosophic Closed Set and Neutrosophic Continuous Functions, Neutrosophic Sets and Systems, Vol. 4, 4-8.
21. A. A. Salama. (2015). Basic Structure of Some Classes of Neutrosophic Crisp Nearly Open Sets & Possible Application to GIS topology, Neutrosophic Sets and Systems, Vol. 7, 18-22.
22. R. Santhi & K. Sakthivel. (2011). Strongly  $\alpha$  Generalized closed set in Intuitionistic Topological Spaces, International Journal Pure Applied Sciences and Technology, Vol. 3, 51-58.
23. F. Smaradache. (2005). Neutrosophic Set: A Generalization of the Intuitionistic Fuzzy Sets, Inter. J. Pure Appl. Math., Vol. 24, 287-297.

24. F. Smaradache. (2010). Neutrosophic Set: A Generalization of Intuitionistic Fuzzy Set, Journal of Defense Resources Management, Vol. 1, 1-10.
25. Y. Veereswari . (2017). An Introduction To Fuzzy Neutrosophic Topological Spaces, IJMA, 8(3), 145-149.

Received: May 2, 2020. Accepted: September 22, 2020