

University of New Mexico

UNM Digital Repository

Branch Mathematics and Statistics Faculty and
Staff Publications

Branch Academic Departments

2022

Introduction to Plithogenic Sociogram with preference representations by Plithogenic Number

Nivetha Martin

Florentin Smarandache

University of New Mexico, smarand@unm.edu

R. Priya

Follow this and additional works at: https://digitalrepository.unm.edu/math_fsp




Part of the [Mathematics Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Martin, Nivetha; Florentin Smarandache; and R. Priya. "Introduction to Plithogenic Sociogram with preference representations by Plithogenic Number." *Journal of Fuzzy Extension & Applications* (2022). https://digitalrepository.unm.edu/math_fsp/594

This Article is brought to you for free and open access by the Branch Academic Departments at UNM Digital Repository. It has been accepted for inclusion in Branch Mathematics and Statistics Faculty and Staff Publications by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.



 Paper Type: Research Paper

Introduction to Plithogenic Sociogram with preference representations by Plithogenic Number

Nivetha Martin ^{*1}, Florentin Smarandache ², R.Priya ³

¹Department of Mathematics, Arul Anandar College (Autonomous), Karumathur, India

²Department of Mathematics, University of New Mexico, Gallup, NM 87301, USA.

³Department of Mathematics, PKN Arts College, Madurai, India

PAPER INFO	ABSTRACT
Chronicle: Received: Reviewed: Revised: Accepted:	The theory of Plithogeny is gaining momentum in recent times as it generalizes the concepts of fuzzy, intuitionistic fuzzy, neutrosophy and other extended representations of fuzzy sets. The relativity of the comprehensive and accommodative nature of plithogenic sets in dealing with attributes shall be applied to handle the decision-making problems in the field of sociology. This paper introduces the concepts of Plithogenic Sociogram (PS) and Plithogenic Number (PN) where the former is the integration of plithogeny to the sociometric technique of sociogram and the latter is the generalization of fuzzy, intuitionistic fuzzy and neutrosophic numbers that shall be used in representations of preferences in group dynamics. This research work outlines the conceptual development of these two newly proposed concepts and discusses the merits of the existing theory of similar kind with suitable substantiation. The plithogenic sociogram model encompassing the attributive preferences with plithogenic number representation is also developed to explicate how it can be materialized in the real social field. A conjectural illustration is put forth to analyze the efficiency and the feasibility of the proposed plithogenic sociogram model and its function in decision-making. This paper also throws light on generalized plithogenic number, dominant attribute constrained plithogenic number and combined dominant attribute constrained plithogenic number together with its operations and suitable illustrations.
Keywords: Plithogeny. Plithogenic Sociogram. Attributes. Preferences generalized plithogenic number. dominant attribute constrained plithogenic number. combined dominant attribute constrained plithogenic number.	

*Corresponding author
E-mail address: nivetha.martin710@gmail.com
DOI:

1. Introduction

The social relationship is the resultant of the social interaction between persons and the longevity of their relationship depends on the likeness in thoughts, behaviour and sometimes the influence of one's attribute over another. The formation of social groups for carrying out group activities is sometimes deliberate but quite natural in any social setting ranging from small schools, organizations to mammoth industries. Should we concern about the strength of the interrelationship between the members of the group? Will making the bond strong between the members benefit the group? The answer is certainly yes, because the extent of functioning as a group with common objectives and the success in goal attainment depends on the coordination and cooperativeness of the members of the group. Hence, the study of interpersonal relationships in a group, preferably a social group has greater significance in group dynamics. Sociogram developed by Jacob Levy Moreno [1] is one of the sociometric techniques that is widely used in the quantitative study on interpersonal relationship. This technique is used to determine the structure of interrelationship in a group setting by determining the order of preferences of the members of the group to work with through a questionnaire. The preferential positions of the members determine the most influential and isolated people of the group and as the result, the decision-makers or the group coordinators can work on enhancing interpersonal relationship and make other alternatives for improving the group efficiency.

Conventional sociogram characterized by crisp preferential ordering, matrix and graphical representations finds several applications in a various social setting. The uncertainty in the order of preferences led to the development of fuzzy sociogram with fuzzy matrix and fuzzy graphical representations and it has made the researchers explore its applicability in determining the interrelationship between the members [2,3]. The decision-making environment is characterized not only by uncertainty but also indeterminacy, to handle such circumstances, Smarandache [4,5] introduced neutrosophic sets which consist of truth values, indeterminacy values and falsity values. Neutrosophic sets are used in decision-making on green supply chain management [6], decision support systems and in many other. Gustavo et al [7] extended fuzzy sociogram to neutrosophic sociogram to incorporate the notion of the existence of indeterminacy in relationships. The preferential ordering is certainly influenced by the indeterminacy that occurs when the members are not sure of certain attributes of others and also they may not sure of their compatibility or suitability to perform a particular task. A hypothetical example was used to illustrate the applicability of the neutrosophic sociogram model to group analysis. On profound analysis over the transition from conventional or the classical sociogram to neutrosophic sociogram, the order of preferences or the preferential ordering is influenced by certainty in the case of classical, uncertainty in the case of fuzzy and indeterminacy in the case of neutrosophic. This fact has led the authors to investigate the factors that influence preferential ordering as it is the deciding factor of the nature of the sociogram. This is the origin of the Plithogenic sociogram which encompasses the attributive preferential ordering, i.e order of preference based on the attributives of the members. Before making the order of preferences, in the sociograms of earlier kinds, the activities (such as quiz program, team-based tasks) that require group work are stated first and the members express their preference for working with others, but in the realm, the choice of choosing or giving preference to the members to get involved in activity also depends on the attributes possessed by the members that are essential to make partnership to take part in any particular activity and many times these attributes may be an essential requisite to take part in the activity or the activities may itself demand the same. In such circumstances, the preferential ordering will be characterized not just by stating the members preferred alone but it also carries the

additional information on why the members are being preferred and naturally it brings the attributes of the members and the extent to which the members possess in the perception of the choice-maker, i.e. the person who makes the preference. The making of choice in preferring a person depending on the attributes has led to the development of Plithogenic sociogram and on exploring will certainly yield better results.

Plithogeny is the recently evolved philosophy that deals with the evolution of entities and their attributes. Smarandache [8] introduced Plithogenic sets that are widely applied in decision making on sustainability [9], medical decision system model [10] and supply chain management [11]. Plithogenic sets are used in decision making as it is highly embedded with wide-ranging generalization approaches in accommodating crisp, fuzzy, intuitionistic fuzzy, neutrosophic sets and the other kinds of extended sets. The preferential ordering assumes either crisp, fuzzy or neutrosophic values, but if the preferential ordering presumes linguistic representation then the linguistic variable requires to be quantified using either fuzzy, intuitionistic fuzzy or neutrosophic numbers. To make such kind of representations more comprehensive, the notion of plithogenic number shall be used. This research work intends to investigate and unveil the Plithogenic sociogram with plithogenic number representing the preferential ordering.

The paper is structured into the following sections, section 2 introduces plithogenic number and discusses their nature; section 3 describes plithogenic sociogram and its utility in decision making and the last section concludes the work.

2. Plithogenic Number

Zadeh [12] introduced Fuzzy numbers and their arithmetic operations to characterize uncertainty. A fuzzy number is a fuzzy set if it is a normal fuzzy set with bounded support and alpha cut being a closed interval for every alpha belonging to $[0,1]$. The fuzzy numbers are the special kind of fuzzy sets used to quantify linguistic variables and it is applied to represent quantities that are uncertain in nature, for instance, the costs parameters, demand are represented as fuzzy numbers. Luciano et al [13,14] discussed fuzzy numbers, fuzzy arithmetic. Mike et al [15] presented the different kinds of fuzzy numbers and their properties. Saed et al [16] described special kinds of fuzzy numbers. Przemyslaw et al [17] illustrated the applications of fuzzy numbers. Thus, fuzzy numbers are the simple form of representing uncertainty and are extended to intuitionistic fuzzy fuzzy numbers which are the next higher or extended form that are extensively applied in decision-making models. Atanassov [18] introduced the concept of intuitionistic fuzzy sets. Intuitionistic fuzzy fuzzy numbers are characterized by membership and non-membership values. Mahapatra et al [19] briefed the applications of an intuitionistic fuzzy fuzzy number. Mijanu et al [20] presented the various kinds of intuitionistic fuzzy fuzzy numbers. Researchers have discussed the different ordering techniques of IFN [21-23]. Smarandache extended Intuitionistic fuzzy sets to neutrosophic sets and discussed the arithmetic operations of neutrosophic numbers. Neutrosophic numbers are the extended or the higher forms of representing uncertainty. Sapna et al [24] described single-valued neutrosophic number, Sun et al [25] elaborated interval-valued neutrosophic number, Faruk [26] explored Gaussian neutrosophic number, Avishek et al [27] discussed the applications of Cylindrical neutrosophic single-valued number in networking, decision making. Researchers like Rajesh et al [28], Nancy et al [29] stated the applications of neutrosophic number in various fields of decision making [30]. Neutrosophic numbers are the extended forms of intuitionistic fuzzy and fuzzy numbers and neutrosophic numbers can be stated as higher forms or super forms of fuzzy numbers. The defuzzification techniques of the extended higher/super forms of fuzzy numbers to its next sub forms

of fuzzy numbers are also discussed by Dragan et al [31], Radhika et al [32], Ali Mert [33], Luca et al [34], and many others. The above discussed forms of fuzzy numbers ranging from simple to higher versions shall be generalized into plithogenic number.

Classical Plithogenic set is characterized by (P, a, V, d, c) , where P is a set, a is the attribute, V is the set of attribute values, d is the degree of appurtenance stating the extent of elements belonging to P satisfying the attribute values and c is the contradiction degree. In this work, the plithogenic set is newly characterized as $(P, A, V A, d, c)$, where A is a system of attributes and $V A$ is the set of all possible attribute values corresponding to each attribute a in A . The classical characterization is with respect to a single attribute and this newly proposed pertains to the system of attributes. To define plithogenic number, the attributes should also be considered and the plithogenic number can also be differentiated into Plithogenic fuzzy number, plithogenic intuitionistic fuzzy number, plithogenic neutrosophic number based on the degree of appurtenance

Let U be a universe of discourse, and a non-empty set M included in U .

Let x be a generic element from M .

Let's consider the attributes A_1, A_2, \dots, A_n , for $n \geq 1$.

The attribute A_1 has the attribute values $A_{11}, A_{12}, \dots, A_{1m_1}$, where $m_1 \geq 1$.

The attribute A_2 has the attribute values $A_{21}, A_{22}, \dots, A_{2m_2}$, where $m_2 \geq 1$.

.....

.....

The attribute A_n has the attribute values $A_{n1}, A_{n2}, \dots, A_{nm}$ where $m, n \geq 1$.

The plithogenic fuzzy number will be of the form

$$M = \{x(A_{11}(t_{11}), A_{12}(t_{12}), \dots, A_{1m_1}(t_{1m_1}); \quad A_{21}(t_{21}), A_{22}(t_{22}), \dots, A_{2m_2}(t_{2m_2}); \quad \dots \quad A_{n1}(t_{n1}), A_{n2}(t_{n2}), A_{nm}(t_{nm}); \text{ with } x \text{ in } U \} ,$$

where

t_{11} is the degree of appurtenance of element x to the set M with respect to the attribute value A_{11} ;

t_{12} is the degree of appurtenance of element x to the set M with respect to the attribute value A_{12} ; etc.

The plithogenic intuitionistic fuzzy number will be of the form

$$M = \{x(A_{11}(t_{11}, f_{11}), A_{12}(t_{12}, f_{12}), \dots, A_{1m_1}(t_{1m_1}, f_{1m_1}); \quad A_{21}(t_{21}, f_{21}), A_{22}(t_{22}, f_{22}), \dots, A_{2m_2}(t_{2m_2}, f_{2m_2}); \quad \dots \quad A_{n1}(t_{n1}, f_{n1}), A_{n2}(t_{n2}, f_{n2}), A_{nm}(t_{nm}, f_{nm}); \text{ with } x \text{ in } U \} ,$$

Where

t_{11} is the degree of appurtenance of element x to the set M with respect to the attribute value A_{11} and f_{11} is the degree of non-appurtenance of element x to the set M with respect to the attribute value A_{11} ;

t_{12} is the degree of appurtenance of element x to the set M with respect to the attribute value A_{12} and f_{12} is the degree of non-appurtenance of element x to the set M with respect to the attribute value A_{12} ; etc.

The neutrosophic plithogenic set:

$$M = \{x(A_{11}(t_{11}, i_{11}, f_{11}), A_{12}(t_{12}, i_{12}, f_{12}), \dots, A_{1m_1}(t_{1m_1}, i_{1m_1}, f_{1m_1}); A_{21}(t_{21}, i_{21}, f_{21}), A_{22}(t_{22}, i_{22}, f_{22}), \dots, A_{2m_2}(t_{2m_2}, i_{2m_2}, f_{2m_2}); \dots A_{n1}(t_{n1}, i_{n1}, f_{n1}), A_{n2}(t_{n2}, i_{n2}, f_{n2}), A_{nm}(t_{nm}, i_{nm}, f_{nm}); \text{ with } x \text{ in } U \} ,$$

where

t_{11} is the degree of appurtenance of element x to the set M with respect to the attribute value A_{11} , i_{11} is the degree of indeterminacy of element x to the set M with respect to the attribute value A_{11} and f_{11} is the degree of non-appurtenance of element x to the set M with respect to the attribute value A_{11} ;

t_{12} is the degree of appurtenance of element x to the set M with respect to the attribute value A_{12} , i_{12} is the degree of indeterminacy of element x to the set M with respect to the attribute value A_{12} and f_{12} is the degree of non-appurtenance of element x to the set M with respect to the attribute value A_{12} ; etc.

Example

Let $U = \{ a, b, c, d, e, f \}$, $M = \{ b, c, e \}$, $A = \{ a_1, a_2, a_3 \}$, $V_{a1} = \{ A_{11}, A_{12}, A_{13} \}$, $V_{a2} = \{ A_{21}, A_{22} \}$, $V_{a3} = \{ A_{31}, A_{32}, A_{33}, A_{34} \}$.

Contradiction Degree	0	1/3	2/3	0	1/2	0	1/4	2/4	3/4	
Attribute Values	A_{11}	A_{12}	A_{13}	A_{21}	A_{22}	A_{31}	A_{32}	A_{33}	A_{34}	
Fuzzy Degree of Appurtenance	b	0.2	0.5	0.6	0.7	0.6	0.5	0.4	0.8	0.9
	c	0.3	0.5	0.6	0.5	0.8	0.9	0.7	0.5	0.6
	e	0.5	0.6	0.4	0.7	0.5	0.6	0.8	0.4	0.5
Intuitionistic fuzzy Degree of Appurtenance	b	(0.7,0.2)	(0.8,0.1)	(0.4,0.6)	(0.7,0.3)	(0.6,0.2)	(0.7,0.1)	(0.8,0.2)	(0.6,0.3)	(0.5,0.3)
	c	(0.8,0.1)	(0.7,0.3)	(0.5,0.3)	(0.6,0.2)	(0.7,0.3)	(0.6,0.3)	(0.7,0.2)	(0.8,0.1)	(0.6,0.2)
	e	(0.5,0.3)	(0.8,0.1)	(0.7,0.2)	(0.8,0.2)	(0.8,0.1)	(0.6,0.2)	(0.7,0.2)	(0.5,0.3)	(0.6,0.3)
Neutrosophic Degree of Appurtenance	b	(0.7,0.2,0.3)	(0.8,0.1,0.3)	(0.6,0.4,0.5)	(0.7,0.2,0.3)	(0.6,0.2,0.3)	(0.5,0.2,0.4)	(0.5,0.1,0.3)	(0.7,0.2,0.3)	(0.7,0.2,0.2)
	c	(0.5,0.1,0.3)	(0.7,0.2,0.3)	(0.5,0.1,0.3)	(0.5,0.2,0.4)	(0.7,0.2,0.3)	(0.6,0.4,0.2)	(0.4,0.1,0.3)	(0.7,0.2,0.2)	(0.8,0.1,0.3)
	e	(0.8,0.1,0.3)	(0.5,0.1,0.3)	(0.6,0.4,0.5)	(0.7,0.2,0.3)	(0.5,0.1,0.3)	(0.7,0.2,0.3)	(0.6,0.4,0.5)	(0.5,0.1,0.3)	(0.7,0.2,0.2)

The plithogenic number with fuzzy degree of appurtenance to all the attribute values will be of the form $P = \{b(A_{11}(0.2), A_{12}(0.5), A_{13}(0.6), A_{21}(0.7), A_{22}(0.6), A_{31}(0.5), A_{32}(0.4), A_{33}(0.8), A_{34}(0.9)), c(A_{11}(0.3), A_{12}(0.5), A_{13}(0.6), A_{21}(0.5), A_{22}(0.8), A_{31}(0.9), A_{32}(0.7), A_{33}(0.5), A_{34}(0.6))\}$ This plithogenic number may be termed as generalized plithogenic fuzzy number as it encompasses all the attribute values. From the values of intuitionistic fuzzy and neutrosophic degrees of appurtenance to all the attribute values the generalized plithogenic intuitionistic fuzzy and generalized plithogenic neutrosophic numbers can be defined.

2.1 Dominant Attribute Constrained Plithogenic Number

This section also proposes the concept of dominant attribute constrained Plithogenic number and it shall be defined by considering only the dominant attribute values.

Let $U = \{ a, b, c, d, e, f \}$, $M = \{ b, c, e \}$, $A = \{ a_1, a_2, a_3 \}$, $V_{a_1} = \{ A_{11}, A_{12}, A_{13} \}$, $V_{a_2} = \{ A_{21}, A_{22} \}$

$V_{a_3} = \{ A_{31}, A_{32}, A_{33}, A_{34} \}$.

In this example, the attribute values A_{11}, A_{21}, A_{31} are considered to be dominant and the plithogenic number considering the values of degree of appurtenance corresponding only to the dominant attribute values are called as **Dominant Attribute Constrained Plithogenic Number**.

Let $P_1 = \{b (A_{11}(0.5), A_{21}(0.7), A_{31}(0.8)) , c (A_{11}(0.4), A_{21}(0.5), A_{31}(0.6)), e (A_{11}(0.4), A_{21}(0.6), A_{31}(0.7))\}$ and $P_2 = \{b (A_{11}(0.6), A_{21}(0.5), A_{31}(0.3)) , c (A_{11}(0.5), A_{21}(0.2), A_{31}(0.5)), e (A_{11}(0.5), A_{21}(0.6), A_{31}(0.8))\}$ where P_1 and P_2 are the Dominant Attribute Constrained plithogenic fuzzy numbers with fuzzy degree of appurtenance with respect to the dominant attribute values.

The union of two Dominant Attribute Constrained plithogenic fuzzy numbers is $P_1 \cup_F P_2$ is defined as $\max \{a_1(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), a_2 (A_\alpha(t_\alpha), A_\beta (t_\beta), \dots, A_\lambda (t_\lambda)), \dots, a_m(A_\alpha(t_\alpha), A_\beta (t_\beta), \dots, A_\lambda (t_\lambda))\}$

Where $A_\alpha, A_\beta, \dots, A_\lambda$ are the dominant attribute values and $t_\alpha, t_\beta, \dots, t_\lambda$ are the respective fuzzy degree of appurtenance with respect to each elements of M .

$P_1 \cup_F P_2 = \{b (A_{11}(0.6), A_{21}(0.7), A_{31}(0.8)) , c (A_{11}(0.5), A_{21}(0.5), A_{31}(0.6)), e (A_{11}(0.5), A_{21}(0.6), A_{31}(0.8))\}$

The intersection of two Dominant Attribute Constrained plithogenic fuzzy numbers is $P_1 \cap_F P_2$ is defined as $\min \{a_1(A_\alpha(t_\alpha), A_\beta (t_\beta), \dots, A_\lambda (t_\lambda)), a_2 (A_\alpha(t_\alpha), A_\beta (t_\beta), \dots, A_\lambda (t_\lambda)), \dots, a_m(A_\alpha(t_\alpha), A_\beta (t_\beta), \dots, A_\lambda (t_\lambda))\}$

$P_1 \cap_F P_2 = \{b (A_{11}(0.5), A_{21}(0.5), A_{31}(0.3)) , c (A_{11}(0.4), A_{21}(0.2), A_{31}(0.5)), e (A_{11}(0.4), A_{21}(0.6), A_{31}(0.7))\}$

Let $P_1 = \{b (A_{11}(0.7,0.2), A_{21}(0.8,0.1), A_{31}(0.7,0.1)) , c (A_{11}(0.7,0.3), A_{21}(0.4,0.6), A_{31}(0.5,0.3)), e (A_{11}(0.6,0.2), A_{21}(0.5,0.3), A_{31}(0.7,0.2))\}$ and

$P_2 = \{ b (A_{11}(0.6,0.3), A_{21} (0.5,0.3), A_{31}(0.6,0.3)) , c (A_{11}(0.7,0.1), A_{21}(0.5,0.3), A_{31}(0.7,0.3)), e (A_{11}(0.5,0.3), A_{21}(0.6,0.3), A_{31} (0.8,0.2))\}$ where P_1 and P_2 are the Dominant Attribute Constrained

plithogenic intuitionistic fuzzy fuzzy numbers with intuitionistic fuzzy degree of appurtenance with respect to the dominant attribute values.

The union of two Dominant Attribute Constrained plithogenic intuitionistic fuzzy fuzzy numbers is $P1 \cup_{IF} P2$ is defined as $(\max \{a_1(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), a_2(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \dots, am(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \min \{a_1(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), a_2(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), \dots, am(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda))\}$

$$P1 \cup_{IF} P2 = \{b(A_{11}(0.7, 0.2), A_{21}(0.8, 0.1), A_{31}(0.7, 0.1)), c(A_{11}(0.7, 0.1), A_{21}(0.5, 0.3), A_{31}(0.7, 0.3)), e(A_{11}(0.6, 0.2), A_{21}(0.6, 0.3), A_{31}(0.8, 0.2))\}$$

The intersection of two Dominant Attribute Constrained plithogenic intuitionistic fuzzy fuzzy numbers is $P1 \cap_{IF} P2$ is defined as $(\min \{a_1(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), a_2(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \dots, am(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \max \{a_1(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), a_2(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), \dots, am(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda))\}$

$$P1 \cap_{IF} P2 = \{b(A_{11}(0.6, 0.3), A_{21}(0.5, 0.3), A_{31}(0.6, 0.3)), c(A_{11}(0.7, 0.3), A_{21}(0.4, 0.6), A_{31}(0.5, 0.3)), e(A_{11}(0.5, 0.3), A_{21}(0.5, 0.3), A_{31}(0.7, 0.2))\}$$

Let $P1 = \{b(A_{11}(0.7, 0.2, 0.3), A_{21}(0.8, 0.1, 0.3), A_{31}(0.6, 0.4, 0.5)), c(A_{11}(0.6, 0.4, 0.2), A_{21}(0.5, 0.1, 0.3), A_{31}(0.7, 0.2, 0.2)), e(A_{11}(0.6, 0.2, 0.1), A_{21}(0.5, 0.1, 0.3), A_{31}(0.7, 0.2, 0.3))\}$ and

$P2 = \{b(A_{11}(0.6, 0.2, 0.3), A_{21}(0.5, 0.2, 0.4), A_{31}(0.6, 0.4, 0.2)), c(A_{11}(0.7, 0.2, 0.3), A_{21}(0.5, 0.2, 0.4), A_{31}(0.7, 0.2, 0.3)), e(A_{11}(0.6, 0.4, 0.2), A_{21}(0.7, 0.2, 0.3), A_{31}(0.8, 0.1, 0.3))\}$ where $P1$ and $P2$ are the Dominant Attribute Constrained plithogenic neutrosophic numbers with neutrosophic degree of appurtenance with respect to the dominant attribute values.

The union of two Dominant Attribute Constrained plithogenic neutrosophic numbers is $P1 \cup_N P2$ is defined as $(\max \{a_1(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), a_2(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \dots, am(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \max \{a_1(A_\alpha(I_\alpha), A_\beta(I_\beta), \dots, A_\lambda(I_\lambda)), a_2(A_\alpha(I_\alpha), A_\beta(I_\beta), \dots, A_\lambda(I_\lambda)), \dots, am(A_\alpha(I_\alpha), A_\beta(I_\beta), \dots, A_\lambda(I_\lambda)), \min \{a_1(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), a_2(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), \dots, am(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda))\}$

$$P1 \cup_N P2 = \{b(A_{11}(0.7, 0.2, 0.3), A_{21}(0.8, 0.2, 0.3), A_{31}(0.6, 0.4, 0.2)), c(A_{11}(0.7, 0.4, 0.2), A_{21}(0.5, 0.2, 0.3), A_{31}(0.7, 0.2, 0.2)), e(A_{11}(0.6, 0.4, 0.1), A_{21}(0.7, 0.2, 0.3), A_{31}(0.8, 0.2, 0.3))\}$$

The intersection of two Dominant Attribute Constrained plithogenic neutrosophic numbers is $P1 \cap_N P2$ is defined as $(\min \{a_1(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), a_2(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \dots, am(A_\alpha(t_\alpha), A_\beta(t_\beta), \dots, A_\lambda(t_\lambda)), \max \{a_1(A_\alpha(I_\alpha), A_\beta(I_\beta), \dots, A_\lambda(I_\lambda)), a_2(A_\alpha(I_\alpha), A_\beta(I_\beta), \dots, A_\lambda(I_\lambda)), \dots, am(A_\alpha(I_\alpha), A_\beta(I_\beta), \dots, A_\lambda(I_\lambda)), \max \{a_1(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), a_2(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda)), \dots, am(A_\alpha(f_\alpha), A_\beta(f_\beta), \dots, A_\lambda(f_\lambda))\}$

$$P1 \cap_N P2 = \{b(A_{11}(0.6, 0.2, 0.3), A_{21}(0.5, 0.1, 0.4), A_{31}(0.6, 0.4, 0.5)), c(A_{11}(0.6, 0.4, 0.3), A_{21}(0.5, 0.2, 0.4), A_{31}(0.7, 0.2, 0.3)), e(A_{11}(0.6, 0.4, 0.2), A_{21}(0.5, 0.2, 0.3), A_{31}(0.7, 0.2, 0.3))\}$$

2.2 Combined Dominant Attribute Constrained Plithogenic Number

In **Combined Dominant Attribute Constrained Plithogenic Number**, the attribute values possess combined degree of appurtenance of the attribute values. For instance

$P_1 = \{b (A_{11}(0.7,0.2), A_{21}(0.8,0.1), A_{31} (0.7,0.1)) , c (A_{11}(0.5), A_{21} (0.5), A_{31} (0.3)), e (A_{11}(0.6,0.4,0.2), A_{21}(0.5,0.2,0.3), A_{31}(0.7,0.2,0.3))\}$ In this plithogenic representation, the element b has intuitionistic fuzzy degree of appurtenance with respect to the attribute values, the element c has fuzzy degree of appurtenance with respect to the attribute values and the element e has neutrosophic degree of appurtenance with respect to the attribute values.

On other hand the combined plithogenic number can also be represented as

$P_1 = \{b (A_{11}(0.7,0.2), A_{21} (0.8), A_{31}(0.7,0.1,0.1)) , c (A_{11} (0.5), A_{21}(0.7,0.2) , A_{31} (0.3)), e (A_{11}(0.6,0.4,0.2), A_{21}(0.5,0.2), A_{31} (0.7))\}$ in which the element b has the combination of intuitionistic fuzzy, fuzzy and neutrosophic degree of appurtenance with respect to the dominant attribute values and the other elements c and e also have a combination of degree of appurtenance.

The union and intersection of combined Plithogenic numbers shall be computed after converting the combined degrees of appurtenance into a same degree of appurtenance using 2.1, 2.2 or 2.3

2.1 Method I (Imprecision membership): Any neutrosophic fuzzy set $N_A = (T_A, I_A, F_A)$ including neutrosophic fuzzy values are transformed into intuitionistic fuzzy values or vague values as $\eta(A) = (T_A, f_A)$ where f_A is estimated the formula stated below which is called as Imprecision membership method. [35]

$$f_A = \begin{cases} F_A + \frac{[1-F_A-I_A][1-F_A]}{[F_A+I_A]} & \text{if } F_A = 0 \\ F_A + \frac{[1-F_A-I_A][F_A]}{[F_A+I_A]} & \text{if } 0 < F_A \leq 0.5 \\ F_A + [1 - F_A - I_A] \left[0.5 + \frac{F_A-0.5}{F_A+I_A}\right] & \text{if } 0.5 < F_A \leq 1 \end{cases}$$

2.2 Method II (Defuzzification): After Method I (Median membership), intuitionistic fuzzy (vague)

fuzzy values of the form $\eta(A) = (T_A, f_A)$ are transformed into fuzzy set including fuzzy values

$$\text{as } \langle \Delta(A) \rangle = \left\langle \frac{T_A}{[T_A+f_A]} \right\rangle. \quad [35]$$

2.3 The score function of the intuitionistic fuzzy set of the form (μ_A, ϑ_A) is $\mu_A - \vartheta_A$. [35]

Let $P_1 = \{b (A_{11}(0.7,0.2), A_{21} (0.8), A_{31}(0.7,0.1,0.1)) , c (A_{11} (0.5), A_{21}(0.7,0.2) , A_{31} (0.3)), e (A_{11}(0.6,0.4,0.2), A_{21}(0.5,0.2), A_{31} (0.7))\}$ and $P_2 = \{b (A_{11}(0.7), A_{21}(0.5,0.2), A_{31} (0.6)) , c (A_{11}(0.5,0.2), A_{21} (0.8), A_{31} (0.2)), e (A_{11}(0.6,0.4), A_{21}(0.5,0.2,0.1), A_{31} (0.5))\}$ be two combined plithogenic number with different degrees of appurtenance and it can be converted to plithogenic number with same degree of appurtenance using the above methods I and II. The modified plithogenic numbers are

$P'_1 = \{b (A_{11}(0.5), A_{21} (0.8), A_{31} (0.58)), c (A_{11}(0.5), A_{21} (0.5), A_{31} (0.3)), e (A_{11}(0.64), A_{21} (0.3), A_{31} (0.7))\}$ and $P'_2 = \{b (A_{11}(0.7), A_{21} (0.3), A_{31} (0.6)), c (A_{11}(0.3), A_{21} (0.8) , A_{31} (0.2)), e (A_{11} (0.2), A_{21} (0.6), A_{31} (0.5))\}$

$$P'_1 \cup P'_2 = \{b (A_{11} (0.7), A_{21} (0.8), A_{31} (0.6)), c (A_{11} (0.5), A_{21} (0.8), A_{31} (0.3)), e (A_{11} (0.64), A_{21} (0.6), A_{31} (0.7))\}$$

$$P'_1 \cap P'_2 = \{b (A_{11} (0.5), A_{21} (0.3), A_{31} (0.58)), c (A_{11} (0.3), A_{21} (0.5), A_{31} (0.2)), e (A_{11} (0.2), A_{21} (0.6), A_{31} (0.5))\}$$

3. Plithogenic Sociogram

In this section, the concept of Plithogenic sociogram is discussed with a simple illustration based on the conceptualization of Neutrosophic sociogram developed by Smarandache. A group of members are given a questionnaire to give their choices of preference in partaking as a team with other members based on certain attributes.

Let $S = \{s_1, s_2, s_3, s_4, s_5\}$ be the members interviewed with the following questions. The members are asked to give their preferential choices of teaming with respect to the attributes.

Write your friends with whom you want to work as a team with respect to their

Q₁: Degree of compatibility

Q₂: Optimistic approaches

Q₃: Disciplinary Knowledge

These questions are focusing on the attributive preferential choice making.

The attributes are the degree of compatibility, optimistic approach and disciplinary knowledge. The attribute values of the attributes are as follows

Degree of compatibility = {low (Q₁₁), moderate (Q₁₂), **high (Q₁₃)**}

Optimistic Approach = {**Dispositional (Q₂₁)**, Unrealistic (Q₂₂), comparative (Q₂₃)}

Disciplinary Knowledge = {**Excellent (Q₃₁)**, good (Q₃₂), average (Q₃₃)}

The preferential choice making of the members with respect to the dominant attributive values say high (Q₁₃), Dispositional (Q₂₁), Excellent (Q₃₁) are presented in the form of Dominant attribute constrained plithogenic number in Table 3.1

Table 3.1 Attributive Preferential Choice-making of the Members

Members	Attributive Preferential Choice-making
S ₁	{S ₂ (Q ₁₃ (0.5), Q ₂₁ (0.6), Q ₃₁ (0.8)), S ₄ (Q ₁₃ (0.6), Q ₂₁ (0.7), Q ₃₁ (0.8))}
S ₂	{S ₁ (Q ₁₃ (0.4), Q ₂₁ (0.7), Q ₃₁ (0.6)), S ₃ (Q ₁₃ (0.5), Q ₂₁ (0.6), Q ₃₁ (0.9)), S ₅ (Q ₁₃ (0.3), Q ₂₁ (0.4), Q ₃₁ (0.6))}
S ₃	{S ₂ (Q ₁₃ (0.5), Q ₂₁ (0.6), Q ₃₁ (0.7)), S ₄ (Q ₁₃ (0.4), Q ₂₁ (0.2), Q ₃₁ (0.5))}
S ₄	{S ₁ (Q ₁₃ (0.7), Q ₂₁ (0.8), Q ₃₁ (0.6)), S ₃ (Q ₁₃ (0.7), Q ₂₁ (0.5), Q ₃₁ (0.3))}
S ₅	{S ₂ (Q ₁₃ (0.5), Q ₂₁ (0.6), Q ₃₁ (0.7)), S ₄ (Q ₁₃ (0.5), Q ₂₁ (0.6), Q ₃₁ (0.6))}

S₁ prefers S₂ with the Plithogenic fuzzy degree of appurtenance of 0.5 to high degree of compatibility, 0.6 to dispositional optimistic approach and 0.8 to excellent disciplinary knowledge and similarly the

preference to S_4 can also be comprehended with the help of fuzzy degree of appurtenance. The approach of Plithogenic sociogram is based on the methodology of neutrosophic sociogram.

The evaluation matrix $M_k = (m_{gh})$, where m_{gh} assumes the degree of appurtenance (in this case, it is fuzzy) of the member s_g selecting s_h with respect to the dominant attribute values and when $g=h$ $m_{gh} = 0$. In neutrosophic sociogram the elements of the evaluation matrix assumes either 0 or 1 based on the number of times a member selects another.

The evaluation matrix M_1 for the dominant attribute value Q_{13} is

	s_1	s_2	s_3	s_4	s_5
s_1	0	0.5	0	0.6	0
s_2	0.4	0	0.5	0	0.3
s_3	0	0.5	0	0.4	0
s_4	0.7	0	0.7	0	0
s_5	0	0.5	0	0.5	0

The evaluation matrix M_2 for the dominant attribute value Q_{21} is

	s_1	s_2	s_3	s_4	s_5
s_1	0	0.6	0	0.7	0
s_2	0.7	0	0.6	0	0.4
s_3	0	0.6	0	0.2	0
s_4	0.8	0	0.5	0	0
s_5	0	0.6	0	0.6	0

The evaluation matrix M_3 for the dominant attribute value Q_{31} is

	s_1	s_2	s_3	s_4	s_5
s_1	0	0.8	0	0.8	0
s_2	0.6	0	0.9	0	0.6
s_3	0	0.7	0	0.5	0
s_4	0.6	0	0.3	0	0
s_5	0	0.7	0	0.6	0

In neutrosophic sociogram each question was given weightage but here in Plithogenic sociogram the dominant attributes are given weightage. By considering the weights of the dominant attributes

values, the final weighted evaluation matrix is determined by assigning the weights as 0.5, 0.25 and 0.25 to the dominant attribute values high (Q₁₃), Dispositional (Q₂₁) and Excellent (Q₃₁) respectively.

	S ₁	S ₂	S ₃	S ₄	S ₅
S ₁	0	0.6	0	0.675	0
S ₂	0.525	0	0.625	0	0.4
S ₃	0	0.575	0	0.375	0
S ₄	0.7	0	0.55	0	0
S ₅	0	0.575	0	0.55	0

The fuzzy amicable degree t_{gh} is calculated by using the formula

$$\frac{2}{t_{gh}} = \frac{1}{f_{gh}} + \frac{1}{f_{hg}}, \text{ where } f_{gh} \text{ represents the compatibility existing between the members } g \text{ and } h$$

h prefers f_{hg}

The $s_g(i)$ is

	S ₁	S ₂	S ₃	S ₄	S ₅
S ₁	0	0.56	0	0.69	0
S ₂	0.56	0	0.6	0	0.47
S ₃	0	0.6	0	0.45	0
S ₄	0.69	0	0.45	0	0
S ₅	0	0.47	0	0	0

which means the member g h and it is vice-versa for

final scores of the members 1,2,..5) of the group, $F(s_g)$ determined by $\frac{\sum_h t_{gh}}{\sum_g \sum_h t_{gh}}$

Table 3.2 Preferential scores of the members

S ₁	0.225632
S ₂	0.294224
S ₃	0.189531
S ₄	0.205776
S ₅	0.084838

Based on the scores as in Table 3.2, it is very vivid that the member s_2 has the maximum score and it represents the significance of the member s_2 in the group and his influencing attributes have made s_2 more preferable, on other hand, the member s_5 has the least score and it shows that the member is not much preferred as the attributes of s_5 may not seems to be influential. This preferential ranking is based on considering plithogenic fuzzy degree of appurtenance. Plithogenic intuitionistic fuzzy fuzzy,

Plithogenic neutrosophic degrees of appurtenance and the concept of combined plithogenic shall also be used to represent the attributive preferential choice making.

3.1 Plithogenic Sociogram in Decision-making

The approach of Plithogenic sociogram shall also be used in decision-making on the alternatives that satisfy the criteria. Let A be the set of alternative methods of food processing say

$A = \{A_1, A_2, A_3, A_4, A_5\}$ and C be the set of criteria or the attributives with attributive values.

$C = \{C_1, C_2, C_3\}$

$C = \{ \text{cost efficiency, energy efficiency, quality conservation} \}$

The attribute values are

Cost efficiency = { **highly economic** (C_{11}), moderately economic (C_{12}), lowly economic (C_{13})}

Energy efficiency = { **above 90%** (C_{21}), above 70% (C_{22}), above 50% (C_{23})}

Quality conservation = { **very good** (C_{31}), good (C_{32}), average (C_{33})}

The comparative attributive preferential choice making over compatibility of the alternatives from expert's point of view with respect to the dominant attribute values highly economic (C_{11}), above 90% (C_{21}) and very good (C_{31}) is presented in the Table 3.3

Table 3.3 Alternatives and its compatibility comparison

Alternatives	Comparative Attributive Preferential Choice-making over compatibility	
	Expert-I	Expert-II
A ₁	{ A ₃ (C ₁₁ (0.4), C ₂₁ (0.6), C ₃₁ (0.8), A ₄ (C ₁₁ (0.6), C ₂₁ (0.6), C ₃₁ (0.7)) }	{ A ₂ (C ₁₁ (0.6), C ₂₁ (0.6), C ₃₁ (0.8)), A ₄ (C ₁₁ (0.7), C ₂₁ (0.8), C ₃₁ (0.7)) }
A ₂	{ A ₁ (C ₁₁ (0.5), C ₂₁ (0.8), C ₃₁ (0.7)), A ₃ (C ₁₁ (0.7), C ₂₁ (0.5), C ₃₁ (0.8)), A ₄ (C ₁₁ (0.8), C ₂₁ (0.6), C ₃₁ (0.7)) }	{ A ₁ (C ₁₁ (0.6), C ₂₁ (0.6), C ₃₁ (0.7)), A ₃ (C ₁₁ (0.8), C ₂₁ (0.6), C ₃₁ (0.8)), A ₅ (C ₁₁ (0.9), C ₂₁ (0.6), C ₃₁ (0.7)) }
A ₃	{ A ₄ (C ₁₁ (0.5), C ₂₁ (0.7), C ₃₁ (0.9)), A ₅ (C ₁₁ (0.6), C ₂₁ (0.7), C ₃₁ (0.8)) }	{ A ₁ (C ₁₁ (0.6), C ₂₁ (0.7), C ₃₁ (0.8)), A ₂ (C ₁₁ (0.7), C ₂₁ (0.5), C ₃₁ (0.8)) }
A ₄	{ A ₂ (C ₁₁ (0.6), C ₂₁ (0.8), C ₃₁ (0.8)), A ₃ (C ₁₁ (0.6), C ₂₁ (0.5), C ₃₁ (0.7)) }	{ A ₁ (C ₁₁ (0.6), C ₂₁ (0.7), C ₃₁ (0.7)), A ₃ (C ₁₁ (0.5), C ₂₁ (0.6), C ₃₁ (0.8)) }
A ₅	{ A ₃ (C ₁₁ (0.7), C ₂₁ (0.6), C ₃₁ (0.7)), A ₄ (C ₁₁ (0.5), C ₂₁ (0.6), C ₃₁ (0.6)) }	{ A ₂ (C ₁₁ (0.8), C ₂₁ (0.6), C ₃₁ (0.5)), A ₄ (C ₁₁ (0.5), C ₂₁ (0.7), C ₃₁ (0.8)) }

With respect to the dominant attribute values, the alternative A₁ is compatible in comparison with the alternatives A₃ and A₄ according to the viewpoint of Expert I and compatible in comparison with the alternatives A₂ and A₄ according to the viewpoint of Expert II

The weights of the dominant attributes values are considered and the final weighted evaluation matrix is determined by assigning the weights as 0.5, 0.25 and 0.25 to the dominant attribute values highly economic (C_{11}), above 90% (C_{21}) and very good (C_{31}) respectively.

	A ₁	A ₂	A ₃	A ₄	A ₅
A ₁	0	0.325	0.275	0.675	0
A ₂	0.625	0	0.7125	0.3625	0.3875
A ₃	0.3375	0.3375	0	0.325	0.3
A ₄	0.325	0.35	0.6	0	0
A ₅	0	0.3375	0.3375	0.5875	0

The amicable degree is presented as in the below

	A ₁	A ₂	A ₃	A ₄	A ₅
A ₁	0	0.428	0.3031	0.439	0
A ₂	0.428	0	0.458	0.356	0.361
A ₃	0.3031	0.458	0	0.422	0.3176
A ₄	0.439	0.356	0.422	0	0
A ₅	0	0.361	0.3176	0	0

The score values of the alternatives are presented in Table 3.4

Table 3.4 Score values of Alternatives

A ₁	0.189662
A ₂	0.259831
A ₃	0.243249
A ₄	0.197264
A ₅	0.109994

The alternative A₂ is the most preferred method of food processing based on the satisfaction of the dominant attribute values and in comparison with other alternatives. This plithogenic sociogram is used to determine the most influential member in the group based on the attributes and the most preferred alternative in decision-making.

Conclusion

This paper introduces the concept of generalized plithogenic number, dominant attribute constrained plithogenic number, combined dominant attribute constrained plithogenic number and its utility in plithogenic sociogram. On comparing the proposed Plithogenic Sociogram with neutrosophic sociogram the former approach is more comprehensive in nature. In neutrosophic sociogram, the questions were deterministic and indeterminate in nature, in the sense, the members are asked to make

the selection of their choice with whom they are very sure to take part in a quiz or study and also they are not sure of teaming up for the group activities. The calculation was done separately by considering members of deterministic teaming and later together with the deterministic and indeterminate teaming. Finally, based on the neutrosophic amicable degree, the opportunity of enhancing the relationship between the members, leadership index and potential leadership index was discussed. But in the neutrosophic sociogram, the reasons for preferring and hesitance were not much explored which are very significant to enhance the relationship in future. The calculation of the numerical ranges representing the extent of the relationship shall become more meaningful if the attributes are considered. This is the origin of the Plithogenic sociogram in which the choice of the members are based on the attributes and the degree of appurtenance states the nature of their preference. The qualitative nature of the members plays a vital role in decision making on the choice of the members preferred. The score values of the members indicate their preference and significance in the group. The members with the least score can be subjected to counselling and made exposed to other kinds of training programs to enhance their attributes of group dynamics. Thus in the Plithogenic sociogram with dominant attribute constrained plithogenic number representing the degree of appurtenance, the attributive preferential choice-making appears to be more realistic and pragmatic in nature. This works on the principle of identifying the attribute deficiency of the members and finds the possibilities of enhancing it to improve the efficiency of teamwork. On enriching the attributes of the members then all the members of the group shall team up with each other without any constraints. The proposed concept shall be extended and employed in decision-making and the illustrations of plithogenic sociogram and plithogenic sociogram in decision making shall be discussed under intuitionistic fuzzy or neutrosophic degrees of appurtenance.

References

- [1] Bautista, E., E. Casas, I. Pineda, J.M. Bezanilla, L. Renero, and Y. Silva, (2009). Usefulness of the sociogram as tool for the analysis of group interactions (Utilidad del sociogramacomoherramienta para el análisis de lasinteraccionesgrupales)(In Spanish). *Psicologia para América Latina*, 2009(18),1-6.
- [2] Yanai, M., Yanai.A., Tsuda.E., Okuda.Y., Yamashita.H., and Inaida.J., (1996)Fuzzy Sociogram Analysis Applying Shapley Value, *Biomedical Fuzzy and Human Sciences*, 2(1), 63-68.
- [3] Uesu, H., Yamashita.H., Yanai.M., and Tomita.M.,(2001). Sociometry analysis applying fuzzy node fuzzy graph, In *Proceedings Joint 9th IFSA World Congress and 20th NAFIPS International Conference, IEEE*.
- [4] Smarandache, F. (1999). A Unifying Field in Logic Neutrosophy, Neutrosophic Probability Set and Logic, Rehoboth, American Research PressAbdel-Basset, M., Nabeeh, N. A., El-Ghareeb, H. A., &Aboelfetouh, A. (2019). Utilising neutrosophic theory to solve transition difficulties of IoT-based enterprises. *Enterprise Information Systems*, 1-21.
- [5] Smarandache, F., (2002). Neutrosophy, A New Branch of Philosophy, *Multiple Valued Logic*, 8(3), 297-384.
- [6] Abdel-Baset, M., Chang,V., &Gamal,A., (2019). Evaluation of the green supply chain management practices: A novel neutrosophic approach. *Computers in Industry*, 108, 210-220.
- [7] Gustavo Alvarez Gómez., Jorge Fernando GoyesGarcía., Sharon DinarzaÁlvarez Gómez., Florentin Smarandache.,(2020). Neutrosophic Sociogram for Group Analysis, Neutrosophic Sets and {Special Issue: Impact of neutrosophy in solving the Latin American's social problems }, 37.
- [8] Smarandache, F., (2018). Plithogeny, Plithogenic Set, Logic, Probability, and Statistics. arXiv preprint arXiv:1808.03948.
- [9] Abdel-Basset, M., El-hoseny, M., Gamal, A., & Smarandache, F. (2019). A novel model for green supply chain management practices based on plithogenic sets. *Artificial intelligence in medicine*, 100, 101710.

- [10] Abdel-Basset, M., Manogaran, G., Gamal, A., & Chang, V. (2019). A Novel Intelligent Medical Decision Support Model Based on Soft Computing and IoT. *IEEE Internet of Things Journal*.
- [11] Abdel-Basset, M., Mohamed, R., Zaied, A. E. N. H., & Smarandache, F. (2019). A hybrid plithogenic decision-making approach with quality function deployment for selecting supply chain sustainability metrics. *Symmetry*, 11(7), 1-21.
- [12] Zadeh, L. A., (1965)., Fuzzy set, *Inform and Control* 8, 338-353.
- [13] Stefanini.L., Sorini.L., Guerra.M.L.,(2008). in Handbook of Granular Computing (John Wiley & Sons, Ltd, 2008), chap. 12, pp. 249–283, <https://onlinelibrary.wiley.com/doi/abs/10.1002/9780470724163>.
- [14] Stefanini.L.,(2010). New Tools in Fuzzy Arithmetic with Fuzzy Numbers, in Information Processing and Management of Uncertainty in Knowledge-Based Systems. Applications, edited by E. Hüllermeier, R. Kruse, F. Hoffmann (Springer Berlin Heidelberg, Berlin, Heidelberg, 2010), 471–480.
- [15] Pathinathan. T., Ponnivalavan. K. and Mike Dison. E. (2015). Different Types of Fuzzy Numbers and Certain Properties, *Journal of computer and mathematical sciences*,6(11),631-651.
- [16] Saed. F.,Mallak and Duha.M.,Bedo.,(2013)., Particular Fuzzy Numbers and a Fuzzy Comparison Method between them., *International Journal of Fuzzy Mathematics and Systems*,3,113-123.
- [17] PrzemyslawGrzegorzewski,LucianoStefanini,(2014) “Fuzzy Numbers and Their Applications”,*Fuzzy Sets and Systems*,257.
- [18] Atanassov, K. T. (1986). Intuitionistic fuzzy fuzzy set,*Fuzzy set and Systems*, 20(1), 87-96
- [19] MahapatraG.S., Roy,T.K.,(2013), Intuitionistic fuzzy Fuzzy Number and Its Arithmetic Operation with Application on System Failure, *Journal of Uncertain Systems*, 7(2),92-107.
- [20] MijanurRahamanSeikh., Prasun Kumar Nayak., Madhumangal Pal.,(2013), Notes on triangular intuitionistic fuzzy fuzzy numbers, *International Journal of Mathematics in Operational Research*,5(4),446-462.
- [21] Lakshmana GomathNayagam.V.,Jeevaraj.S,GeethaSivaraman., (2016),Complete Ranking of Intuitionistic fuzzy Fuzzy Numbers, *Fuzzy Information and Engineering*,8(2) , 237-254.
- [22] Pardha saradhi I.B.,Madhuri.M.V.,Ravi Shankar.N.,(2017), Ordering Of Intuitionistic fuzzy Fuzzy Numbers Using Centroid Of Centroids Of Intuitionistic fuzzy Fuzzy Number, *International Journal of Mathematics Trends and Technology (IJMTT)* , 52 (5),276-285.
- [23] LakshmanaGomathiNayagamVelu., JeevarajSelvaraj., and DhanasekaranPonnialagan.,(2017), A New Ranking Principle For Ordering Trapezoidal Intuitionistic fuzzy Fuzzy Numbers,*complexity*, 2017,1-24.
- [24] Sapna Gahlotand Saraswat.R.N, (2020),Single Valued Neutrosophic Numbers and Applications in Selection Criteria, *Journal of Physics: Conference Series* ,1849,1-6.
- [25] Sun, Hong-Xiaa., Yang, Hao-Xionga .,Wu, Jian-Zhangb ., Ouyang, Yaoc,(2015), Interval neutrosophic numbers Choquet integral operator for multi-criteria decision making, *Journal of Intelligent & Fuzzy Systems*, 28(6), 2443-2455.
- [26] Faruk .K.,(2018)., Gaussian single-valued neutrosophic numbers and its application in multi-attribute decision making, *Neutrosophic Sets and Systems*, 22,101-117.
- [27] Avishek Chakraborty., Sankar Prasad Mondal., SharifulAlam., AnimeshMahata.,(2020), *Transactions on Intelligence Technology*,5(2), 68-77.
- [28] Rajesh Kumar Saini., AtulSangal.,ManishaManisha.,(2020), Application of Single Valued Trapezoidal Neutrosophic Numbers in Transportation Problem,*Neutrosophic Sets and Systems*, 35,563-583.
- [29] [Nancy El-Hefenawy.](#), [Mohamed A Metwally.](#), [Zenat M Ahmed.](#), [Ibrahim Mahmoud El-henawy.](#), (2016),A Review on the Applications of Neutrosophic Sets,*Journal of Computational and Theoretical Nanoscience*,13,936-944.
- [30] SurapatiPramanik., Durga Banerjee.,(2018)., Neutrosophic number goal programming for multi-objective linear programming problem in neutrosophic number environment ,*MOJ Curr Res & Rev*,1(3),135-141
- [31] Dragan.Z.,SaleticiDusan.M.,VelasevicNikos.E.,Mastorakis., (2002).,Analysis of Basic DefuzzificationTechniques.,*Proceedings of the 6th WSES international multicongress on circuits, systems, communications and computers*.

- [32] Radhika.C., and Parvathi.R.,(2016).,Defuzzification of intuitionistic fuzzy fuzzy sets, *Notes on Intuitionistic fuzzy Fuzzy Sets*, 22(5), 19-26.
- [33] Mert.A,(2020), On the WABL Defuzzification Method for Intuitionistic fuzzy Fuzzy Numbers, *Intelligent and Fuzzy Techniques in Big Data Analytics and Decision Making* ,39-47.
- [34] Deli.I., Emel .K.O.,(2020), A defuzzification method on single-valued trapezoidal neutrosophic numbers and multiple attribute decision making, *Cumhuriyet Science Journal*, 41(1), 2.
- [35] Solairaju and Shajahan, (2018),Transforming Neutrosophic Fuzzy Set into Fuzzy Set by Imprecision Method, *Journal of Computer and Mathematical Sciences*, Vol.9(10),1392-1399.