

University of New Mexico

## UNM Digital Repository

---

Branch Mathematics and Statistics Faculty and  
Staff Publications

Branch Academic Departments

---

2024

### On stratified single-valued soft topogenous structures

Fahad Alsharari

Yaser Saber

Hanan Alohal

Mesfer H. Alqahtani

Mubarak Ebodey

*See next page for additional authors*

Follow this and additional works at: [https://digitalrepository.unm.edu/math\\_fsp](https://digitalrepository.unm.edu/math_fsp)




Part of the [Mathematics Commons](#)

---

---

**Authors**

Fahad Alsharari, Yaser Saber, Hanan Alohalı, Mesfer H. Alqahtani, Mubarak Ebodey, Tawfik Elmasry, Jafar Alsharif, Amal F. Soliman, Florentin Smarandache, and Fahad Sikander



## Review article

## On stratified single-valued soft topogenous structures

Fahad Alsharari<sup>a</sup>, Yaser Saber<sup>b,c,\*</sup>, Hanan Alohalı<sup>d</sup>, Mesfer H. Alqahtani<sup>e</sup>,  
Mubarak Ebodey<sup>f</sup>, Tawfik Elmasry<sup>f</sup>, Jafar Alsharif<sup>f</sup>, Amal F. Soliman<sup>g,h</sup>,  
Florentin Smarandache<sup>i</sup>, Fahad Sikander<sup>j</sup>

<sup>a</sup> Department of Mathematics, College of Science, Jouf University, Sakaka 72311, Saudi Arabia

<sup>b</sup> Department of Mathematics, College of Science Al-Zulfi, Majmaah University, P. O. Box 66, Al-Majmaah 11952, Saudi Arabia

<sup>c</sup> Department of Mathematics, Faculty of Science, Al-Azhar University, Assiut, 71524, Egypt

<sup>d</sup> Department of Mathematics, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

<sup>e</sup> Department of Mathematics, University College of Umluj, University of Tabuk, Tabuk 48322, Saudi Arabia

<sup>f</sup> Department of Business Administration, Faculty of Science and Humanities at Hotat Sudair, Majmaah University, 11952, Riyadh, Saudi Arabia

<sup>g</sup> Department of Mathematics, College of Science and Humanities in Alkharj, Prince Sattam Bin Abdulaziz University, Alkharj, Saudi Arabia

<sup>h</sup> Department of Basic Science, Benha Faculty of Engineering, Benha University, Banha, Egypt

<sup>i</sup> Department of Mathematics, University of New Mexico, Gallup, NM 87301, USA

<sup>j</sup> Department of Basics Sciences, College of Science and Theoretical studies, Saudi Electronic University, Jeddah 23442, Saudi Arabia

## ARTICLE INFO

## Keywords:

Stratified

Single-valued neutrosophic soft

quasi-uniformity

Single-valued neutrosophic soft topogenous

order

Single-valued neutrosophic soft

quasi-proximity

## ABSTRACT

This paper presents novel concepts including stratified single-valued neutrosophic soft topogenous (*stratified svns-topogenous*), stratified single-valued neutrosophic soft filter (*stratified svns-filter*), stratified single-valued neutrosophic soft quasi uniformity (*stratified svnsq-uniformity*) and stratified single-valued neutrosophic soft quasi proximity (*stratified svnsq-proximity*). Additionally, we present the idea of single-valued neutrosophic soft topogenous structures, formed by integrating svns-topogenous with svns-filter, and discuss their properties. Furthermore, we explore the connections between these single-valued neutrosophic soft topological structures and their corresponding stratifications.

## 1. Introduction and preliminaries

Various methodologies have been scrutinized for effectively handling uncertainties, encompassing fuzzy set theory [1], intuitionistic fuzzy set theory [2], vague set theory, interval mathematics [3,4], and rough set theory [5]. However, these approaches have encountered significant challenges. Soft set theory, introduced by Molodtsov [6], has emerged as a promising alternative and has been successfully applied in diverse fields such as function smoothness [7], game theory [8], Riemann integration [9], and probability theory [10]. Notably, recent advancements in soft set theory and its applications have been particularly noteworthy in certain domains.

Fuzzy sets: Handle uncertainty by assigning degrees of membership, Intuitionistic fuzzy sets: Allow for more uncertainty with an extra degree of non-membership, Vague sets: Handle uncertainty by having a boundary region of membership, Interval mathematics: Deals with uncertainty by working with ranges of numbers, Rough sets: Handle uncertainty by approximating vague concepts, Stochastic programming: Models uncertainty through probability distributions, Robust optimization: Provides solutions that are valid

\* Corresponding author.

E-mail addresses: [y.sber@mu.edu.sa](mailto:y.sber@mu.edu.sa) (Y. Saber), [t.almasry@mu.edu.sa](mailto:t.almasry@mu.edu.sa) (T. Elmasry), [smarand@unm.edu](mailto:smarand@unm.edu) (F. Smarandache).

<https://doi.org/10.1016/j.heliyon.2024.e27926>

Received 22 September 2023; Received in revised form 3 March 2024; Accepted 8 March 2024

Available online 16 March 2024

2405-8440/© 2024 The Author(s).

Published by Elsevier Ltd.

This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

under all scenarios, Simulation: Mimics real-world processes to evaluate different strategies and Decision analysis: Uses decision trees to evaluate uncertain outcomes.

Maji et al. [11,12] provided an application for decision-making issues along with a few new definitions of soft sets. Dey and colleagues have examined the gray relational projection approach, generalized neutrosophic soft set and multiattribute decision-making in [13,14]. The findings reported in [11], started the study linking fuzzy and soft sets were enhanced by Aktas and Çağman [15], Feng et al. [16], Chen et al. [17], Ali et al. [18] and Sun et al. [19]. Subsequent research on the fuzzy soft sets notion was conducted by Yang et al. [20], Kharal and Ahmed [21] and Ahmed and Khara [22]. Shabir and Naz [23] distinguished a variety of issues utilizing soft sets, subsuming separation axioms. Tanay and Kandemir [24] first developed concepts of fuzzy soft topology by utilizing Chang’s concept of fuzzy topology. They explored the fundamental concepts by adopting Chang [25]’s definitions, whereas Pazar Varol and Aygün [26] defined the fuzzy soft topology in Lowen’s sense. Aygünoğlu et al. [27] defined Šostak’s fuzzy soft topology. Saber et al. [28] framed single-valued neutrosophic soft topological spaces  $(\mathbb{Y}, \mathcal{T}^v, \mathcal{T}^\mu, \mathcal{T}^\omega)$  (*svnst-space*). In the fuzzy paradigm, there exist three alternative ways to uniformity: Kotzé’s [29] uniform covering approach, Hutton’s [30] uniform operator approach and Lowen’s [31] entourage approach.

Smarandache originally introduced the concept of a neutrosophic set [32], which subsequently led to research on both single-valued neutrosophic sets (*svns*) and neutrosophic sets (*ns*) by Wang et al. [33] and Salama et al. [34,35]. Various applications have been explored in the works of several researchers [36–40]. Saber et al. conducted detailed studies on single-valued neutrosophic regularity space (*svnr-space*), single-valued neutrosophic ideals (*svnis*), stratification of *svnst-space*, single-valued neutrosophic soft sets (*svns*), and stratified modeling in soft fuzzy topological structures in extensive works [41–47].

It is widely acknowledged that theories such as fuzzy sets, intuitionistic fuzzy sets, and rough sets are viewed as extensions of neutrosophic set theory, serving as essential mathematical tools for managing uncertainty. The concepts of stratified single-valued neutrosophic soft topogenous, which build upon the ideas introduced by Varol et al. [26], Aygünoğlu et al. [27] and Abbas et al. [48,49], constitute a significant contribution of this paper.

Building upon the insights gained from previous analyses, we introduce the concepts of “stratified svnsq-uniformity” (or “stratified svns-topogenous order” and “stratified svnsq-proximity”) derived from predefined “svnsq-uniformity” (or “svns-topogenous order” and “svnsq-proximity”). We investigate several properties associated with these newly formulated structures. Additionally, we explore the interrelations between these single-valued neutrosophic soft topological structures and their respective stratifications.

Throughout this study,  $(\mathbb{Y}, \mathcal{V})$  denotes the collection of all *svns* sets on  $\mathbb{Y}$ , where  $\mathcal{V}$  is the set of all parameters on  $\mathbb{Y}$  and  $\mathbb{Y}$  indicated to an initial universe.

The *svns* set  $\mathcal{L}_\zeta$  on  $\mathbb{Y}$  is said to be *ι* – absolute *svn-soft sets* and indicated by  $\tilde{\mathcal{V}}^\iota$ , if  $\mathcal{L}_\zeta = \bar{\iota}$  for each  $e \in \mathcal{V}$ ,  $\iota \in \zeta$ ,  $\bar{\iota}(x) = \iota$  for every  $x \in \mathbb{Y}$  (where,  $[\tilde{\mathcal{V}}^\iota]^c = \tilde{\mathcal{V}}^{\iota^c}$ ,  $\zeta = [0, 1]$ ) and  $\zeta_0 = (0, 1]$ .

**Definition 1.** [32]. Let  $\mathbb{Y} \neq \emptyset$ . A neutrosophic set (*n-set*) on  $\mathcal{X}$  defined as

$$\Theta = \{ \langle y, \nu_\Theta(y), \mu_\Theta(y), \omega_\Theta(y) \mid y \in \mathbb{Y}, \nu_\Theta(y), \mu_\Theta(y), \omega_\Theta(y) \in ]^{-0}, 1^+ [ \} ,$$

representing the degree of membership where  $(\nu_\Theta(y))$ , the degree of indeterminacy  $(\mu_\Theta(y))$ , and degree of nonmembership  $(\omega_\Theta(y))$ ;  $\forall y \in \zeta$  to the set  $\Theta$ .

**Definition 2.** [33]. Let  $\mathbb{Y} \neq \emptyset$ . Then *svn-set*  $\Theta$  on  $\mathbb{Y}$  is defined as

$$\Theta = \{ \langle y, \nu_\Theta(y), \mu_\Theta(y), \omega_\Theta(y) \mid y \in \mathbb{Y}, \nu_\Theta(y), \mu_\Theta(y), \omega_\Theta(y) \in \zeta \} ,$$

where  $\nu_\Theta, \mu_\Theta, \omega_\Theta : \mathbb{Y} \rightarrow \zeta$  and  $0 \leq \nu_\Theta(y) + \mu_\Theta(y) + \omega_\Theta(y) \leq 3$ .

**Definition 3.** [28].  $f_A$  is a *svns-set* on  $\mathbb{Y}$  where,  $f : \mathcal{V} \rightarrow \zeta^\mathbb{Y}$ ; i.e.,  $f_e \triangleq f(e)$  is a *svn-set* on  $\mathbb{Y}$ , for all  $e \in A$  and  $f(e) = \langle 0, 1, 1 \rangle$ , if  $e \notin A$ . The *svn-set*  $f(e)$  is termed as an element of the *svns-set*  $f_A$ . Thus, a *svns-set*  $f_\mathcal{V}$  on  $\mathbb{Y}$  can be defined as:

$$(f, \mathcal{V}) = \{ \langle e, f(e) \mid e \in \mathcal{V}, f(e) \in \zeta^\mathbb{Y} \} \\ = \left\{ e, \langle \nu_f(e), \mu_f(e), \omega_f(e) \rangle \mid e \in \mathbb{Y}, f(e) \in \zeta^\mathbb{Y} \right\} ,$$

where  $\nu_f : \mathcal{V} \rightarrow \zeta$  ( $\nu_f$  is termed as a membership function),  $\mu_f : \mathcal{V} \rightarrow \zeta$  ( $\mu_f$  is termed as indeterminacy function), and  $\omega_f : \mathcal{V} \rightarrow \zeta$  ( $\omega_f$  is termed as a nonmembership function) of *svns-set*.

A *svns-set*  $f_\mathcal{V}$  on  $\mathbb{Y}$  is termed as a null *svns-set* (for short,  $\Phi$ ), if  $\nu_f(e) = 0$ ,  $\mu_f(e) = 1$  and  $\omega_f(e) = 1$ , for any  $e \in \mathcal{V}$ .

A *svns-set*  $f_\mathcal{V}$  on  $\mathbb{Y}$  is termed as an absolute *svns-set* (for short,  $\tilde{\mathcal{V}}$ ), if  $\nu_f(e) = 1$ ,  $\mu_f(e) = 0$  and  $\omega_f(e) = 0$ , for any  $e \in \mathcal{V}$ .

**Definition 4.** [45]  $(\mathbb{Y}, \mathcal{T}^v, \mathcal{T}^\mu, \mathcal{T}^\omega)$  is a *svnst*, if  $\mathcal{T}^v, \mathcal{T}^\mu, \mathcal{T}^\omega : \mathcal{V} \rightarrow \zeta^{\mathbb{Y}, \mathcal{V}}$  it meets the next criteria: for every  $e \in \mathcal{V}$ :

- (T<sub>1</sub>)  $\mathcal{T}_e^v(\Phi) = \mathcal{T}_e^v(\tilde{\mathcal{V}}) = 1$  and  $\mathcal{T}_e^\mu(\Phi) = \mathcal{T}_e^\mu(\tilde{\mathcal{V}}) = \mathcal{T}_e^\omega(\Phi) = \mathcal{T}_e^\omega(\tilde{\mathcal{V}}) = 0$ ,
- (T<sub>2</sub>)  $\mathcal{T}_e^v(\mathcal{L}_\sigma \sqcap \mathcal{h}_\alpha) \geq \mathcal{T}_e^v(\mathcal{L}_\sigma) \wedge \mathcal{T}_e^v(\mathcal{h}_\alpha)$ ,  $\mathcal{T}_e^\mu(\mathcal{L}_\sigma \sqcap \mathcal{h}_\alpha) \leq \mathcal{T}_e^\mu(\mathcal{L}_\sigma) \vee \mathcal{T}_e^\mu(\mathcal{h}_\alpha)$ ,  
 $\mathcal{T}_e^\omega(\mathcal{L}_\sigma \sqcap \mathcal{h}_\alpha) \leq \mathcal{T}_e^\omega(\mathcal{L}_\sigma) \vee \mathcal{T}_e^\omega(\mathcal{h}_\alpha)$ ,  $\forall \mathcal{L}_\sigma, \mathcal{h}_\alpha \in \mathbb{Y}, \mathcal{V}$ ,
- (T<sub>3</sub>)  $\mathcal{T}_e^v(\bigsqcup_{j \in J} [\mathcal{L}_\sigma]_j) \geq \bigwedge_{j \in J} \mathcal{T}_e^v([\mathcal{L}_\sigma]_j)$ ,  $\mathcal{T}_e^\mu(\bigsqcup_{j \in J} [\mathcal{L}_\sigma]_j) \leq \bigvee_{j \in J} \mathcal{T}_e^\mu([\mathcal{L}_\sigma]_j)$ ,

$$\mathcal{T}_e^\omega(\bigsqcup_{j \in J} [\mathcal{L}_\sigma]_j) \leq \bigvee_{j \in J} \mathcal{T}_e^\omega([\mathcal{L}_\sigma]_j), \quad \forall \mathcal{L}_\sigma \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}.$$

The *svnst*  $(\mathcal{T}^\nu, \mathcal{T}^\mu, \mathcal{T}^\omega)$  is said to be stratified if it meets the next condition

$$(\mathcal{T}_3) \mathcal{T}_e^\nu(\tilde{\nu}^i) = 1 \text{ and } \mathcal{T}_e^\mu(\tilde{\nu}^i) = \mathcal{T}_e^\omega(\tilde{\nu}^i) = 0 \text{ for each } e \in \mathfrak{Y}, i \in \zeta.$$

The quadrilateral  $(\mathfrak{X}, \mathcal{T}^\nu, \mathcal{T}^\mu, \mathcal{T}^\omega)$  is named *stratified svnst-space*. Representing the degree of openness  $(\mathcal{T}_e^\nu(\mathcal{L}_\sigma))$ , the degree of indeterminacy  $(\mathcal{T}_e^\mu(\mathcal{L}_\sigma))$ , and the degree of non-openness  $(\mathcal{T}_e^\omega(\mathcal{L}_\sigma))$ ; of a *svns* set with respect to that parameter  $e \in \mathfrak{Y}$ .

Sometimes, we will write  $\mathcal{T}^{\nu\mu\omega}$  for  $(\mathcal{T}^\nu, \mathcal{T}^\mu, \mathcal{T}^\omega)$ .

Now we mind some concepts and nomenclature that will be applied in this paper.

Assume  $\Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  denotes the collection of all mappings  $z : (\mathfrak{X}, \mathfrak{Y}) \rightarrow \widetilde{(\mathfrak{X}, \mathfrak{Y})}$  with the next properties, for each  $\mathcal{L}_\sigma \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$

- (z<sub>1</sub>)  $\mathcal{L}_\sigma = z(f_\sigma)$ ,
- (z<sub>2</sub>)  $z(\bigsqcup_{j \in J} (\mathcal{L}_\sigma)_j) = \bigsqcup_{j \in J} z((\mathcal{L}_\sigma)_j)$ .

For  $z, t \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  we define that  $z \circ t$  and  $z \sqcap t$  by

$$(z \circ t)(\mathcal{L}_\sigma) = z(t(\mathcal{L}_\sigma)),$$

and

$$(z \sqcap t)(\mathcal{L}_\sigma) = \sqcap \{ z(\mathfrak{h}_\alpha) \sqcup t(\mathfrak{g}_\zeta) \mid \mathcal{L}_\sigma = \mathfrak{h}_\alpha \sqcup \mathfrak{g}_\zeta \}.$$

Let  $\psi_\varphi : (\mathfrak{X}, \mathfrak{Y}) \rightarrow (\mathcal{U}, \mathcal{R})$  be a mapping,  $z \in \Psi(\widetilde{(\mathcal{U}, \mathcal{R})})$ , then  $\psi_\varphi^{-1} \circ z \circ \psi_\varphi \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ , (equivalently,  $\psi_\varphi^{-1}(z(\psi_\varphi(\mathcal{L}_\sigma))) \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$  for any  $\mathcal{L}_\sigma \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ).

For each  $z, t, c, v \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ , the next characteristics hold:

- (1) If  $z \leq z_1$  and  $t \leq t_1$ , then  $z \sqcap t \sqsubseteq z_1 \sqcap t_1$ ,
- (2)  $z \sqcap t \sqsubseteq z$ ,  $z \sqcap t \sqsubseteq t$  and  $z \sqcap z = z$ ,
- (3)  $(z \sqcap t) \sqcap c = z \sqcap (t \sqcap c)$ ,
- (4)  $(z \sqcap t) \circ (c \sqcap v) = (z \circ c) \sqcap (t \circ v)$ .

$\forall \mathcal{L}_\sigma \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ,  $i \in \zeta$  define the mapping  $\hat{i} : \widetilde{(\mathfrak{X}, \mathfrak{Y})} \rightarrow \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ , by

$$\hat{i}(\mathcal{L}_\sigma)(x) = \begin{cases} \tilde{\nu}^{sup(\mathcal{L}_\sigma)}, & \text{if } sup(\mathcal{L}_\sigma) \leq i, \\ \tilde{\nu}, & \text{if otherwise,} \end{cases}$$

where  $sup(\mathcal{L}_\sigma) = \bigvee_{x \in \mathfrak{X}} \mathcal{L}_\sigma(x)$ ,  $\forall e \in \sigma$ . The map  $\hat{i}$  satisfy the properties (z<sub>1</sub>) and (z<sub>2</sub>), that is,  $\hat{i} \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ . Furthermore,  $\hat{i}$  fulfills the next properties:

**Lemma 1.** Let  $i, i_1, i_2 \in \zeta$ :

- (1) If  $i_1 \leq i_2$ , then  $\hat{i}_2 \sqsubseteq \hat{i}_1$ ,
- (2)  $\hat{i} \circ \hat{i} = \hat{i}$  for each  $i \in \zeta$ ,
- (3)  $z \sqsubseteq \hat{i}$  holds for all  $z \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ .

**Lemma 2.** Let  $z_{(\mathcal{L}_\sigma)} : \widetilde{(\mathfrak{X}, \mathfrak{Y})} \rightarrow \widetilde{(\mathfrak{X}, \mathfrak{Y})}$  be a mapping defined by

$$z_{(\mathcal{L}_\sigma)}(\mathfrak{h}_\alpha) = \begin{cases} \mathcal{L}_\sigma, & \text{if } \mathfrak{h}_\alpha \sqsubseteq \mathcal{L}_\sigma, \\ \tilde{\nu}, & \text{if otherwise.} \end{cases}$$

Then,  $z_{\mathcal{L}_\sigma} \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  and  $z_{(\mathcal{L}_\sigma)} \circ z_{(\mathcal{L}_\sigma)} = z_{(\mathcal{L}_\sigma)}$ .

**Theorem 1.** Let  $(\mathfrak{X}, \mathcal{T}^{\nu\mu\omega})$  be a *svnst-space*, for every  $e \in \mathfrak{Y}$ ,  $\mathcal{L}_\sigma \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  we define the map  $(\mathcal{T}_{st}^\nu)_e : \mathfrak{Y} \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \mathfrak{Y})}}$ ,  $(\mathcal{T}_{st}^\mu)_e : \mathfrak{Y} \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \mathfrak{Y})}}$  and  $(\mathcal{T}_{st}^\omega)_e : \mathfrak{Y} \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \mathfrak{Y})}}$  as follows

$$(\mathcal{T}_{st}^\nu)_e(\mathcal{L}_\sigma) = \bigvee \left\{ \bigwedge_{j \in J} \mathcal{T}_e^\nu((\mathcal{L}_\sigma)_j) \mid \bigsqcup_{j \in J} ((\mathcal{L}_\sigma)_j \sqcap \tilde{\nu}^{i_j}) = \mathcal{L}_\sigma \right\},$$

$$(\mathcal{T}_{st}^\mu)_e(\mathcal{L}_\sigma) = \bigwedge \left\{ \bigvee_{j \in J} \mathcal{T}_e^\mu((\mathcal{L}_\sigma)_j) \mid \bigsqcup_{j \in J} ((\mathcal{L}_\sigma)_j \sqcap \tilde{\nu}^{i_j}) = \mathcal{L}_\sigma \right\},$$

$$(\mathcal{T}_{st}^\omega)_e(\mathcal{E}_\sigma) = \bigwedge \left\{ \bigvee_{j \in J} \mathcal{T}_e^\omega((\mathcal{E}_\sigma)_j) \mid \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{Y}^j) = \mathcal{E}_\sigma \right\},$$

where  $\bigvee$  and  $\bigwedge$  are taken over all families  $\{(\mathcal{E}_\sigma)_j : j \in J\}$  with  $\mathcal{E}_\sigma = \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{Y}^j)$ . Then  $(\mathcal{T}_{st}^{v\mu\omega})_\gamma$  is the coarsest stratified svnsq on  $\mathbb{Y}$  which is finer than  $\mathcal{T}_\gamma^{v\mu\omega}$ . And  $(\mathcal{T}_{st}^{v\mu\omega})_\gamma$  is called the stratification of a svnsq  $\mathcal{T}_\gamma^{v\mu\omega}$  on  $\mathbb{Y}$ .

## 2. Stratified single-valued neutrosophic soft quasi-uniform spaces

**Definition 5.** Let  $\mathcal{K}^v, \mathcal{K}^\mu, \mathcal{K}^\omega : \gamma \rightarrow \zeta^{\Psi(\widetilde{\mathbb{Y}, \gamma})}$  and  $\tilde{e} \in \gamma$ . Then  $(\mathcal{K}^v, \mathcal{K}^\mu, \mathcal{K}^\omega)$  is called *svnsq-uniformity* on  $\mathbb{Y}$ , if it satisfies these properties:

- ( $\mathcal{K}_1$ ) There exists  $z \in \Psi(\widetilde{\mathbb{Y}, \gamma})$  such that  $\mathcal{K}_e^v(z) = 1, \mathcal{K}_e^\mu(z) = 0$  and  $\mathcal{K}_e^\omega(z) = 0$ .
  - ( $\mathcal{K}_2$ ) If  $z, t \in \Psi(\widetilde{\mathbb{Y}, \gamma})$  and  $z \sqsubseteq t$ , then  $\mathcal{K}_e^v(z) \leq \mathcal{K}_e^v(t), \mathcal{K}_e^\mu(z) \geq \mathcal{K}_e^\mu(t)$  and  $\mathcal{K}_e^\omega(z) \geq \mathcal{K}_e^\omega(t)$ .
  - ( $\mathcal{K}_3$ ) For  $z, t \in \Psi(\widetilde{\mathbb{Y}, \gamma})$ ,  $\mathcal{K}_e^v(z \cap t) \geq \mathcal{K}_e^v(z) \wedge \mathcal{K}_e^v(t), \mathcal{K}_e^\mu(z \cap t) \leq \mathcal{K}_e^\mu(z) \vee \mathcal{K}_e^\mu(t)$  and  $\mathcal{K}_e^\omega(z \cap t) \leq \mathcal{K}_e^\omega(z) \vee \mathcal{K}_e^\omega(t)$ .
  - ( $\mathcal{K}_4$ ) For  $z \in \Psi(\widetilde{\mathbb{Y}, \gamma})$ ,  $\bigvee \{\mathcal{K}_e^v(z_1) \mid z_1 \circ z_1 = z\} \geq \mathcal{K}_e^v(z), \bigwedge \{\mathcal{K}_e^\mu(z_1) \mid z_1 \circ z_1 = z\} \leq \mathcal{K}_e^\mu(z)$  and  $\bigwedge \{\mathcal{K}_e^\omega(z_1) \mid z_1 \circ z_1 = z\} \leq \mathcal{K}_e^\omega(z)$ .
- The pair  $(\mathbb{Y}, \mathcal{K}_\gamma^{v\mu\omega})$  is called a *svnsq-uniform space*.  
A *svnsq-uniformity*  $\mathcal{K}_\gamma^{v\mu\omega}$  is said to be stratified if it provides that

$$(\mathcal{K}_5) \mathcal{K}_e^v(\tilde{e}) = 1, \mathcal{K}_e^\mu(\tilde{e}) = 0 \text{ and } \mathcal{K}_e^\omega(\tilde{e}) = 0 \text{ for any } \tilde{e} \in \zeta.$$

So, the pair  $(\mathbb{Y}, \mathcal{K}_\gamma^{v\mu\omega})$  is called a *stratified svnsq-uniform space*.

Let  $(\mathcal{K}_\gamma^{v\mu\omega})_1$  and  $(\mathcal{K}_\gamma^{v\mu\omega})_2$  be *stratified svnsq-uniformities* on  $\mathbb{Y}$ . We say that  $(\mathcal{K}_\gamma^{v\mu\omega})_1$  is finer than  $(\mathcal{K}_\gamma^{v\mu\omega})_2$  [ $(\mathcal{K}_\gamma^{v\mu\omega})_2$  is coarser than  $(\mathcal{K}_\gamma^{v\mu\omega})_1$ ] denoted by  $(\mathcal{K}_\gamma^{v\mu\omega})_2 \sqsubseteq (\mathcal{K}_\gamma^{v\mu\omega})_1$  if

$$(\mathcal{K}_e^v)_2(z) \leq (\mathcal{K}_e^v)_1(z), (\mathcal{K}_e^\mu)_2(z) \geq (\mathcal{K}_e^\mu)_1(z), (\mathcal{K}_e^\omega)_2(z) \geq (\mathcal{K}_e^\omega)_1(z),$$

for any  $e \in \gamma, z \in \Psi(\widetilde{\mathbb{Y}, \gamma})$ .

Assume that  $(\mathbb{Y}, \mathcal{K}_\gamma^{v\mu\omega})$  and  $(\mathcal{U}, \mathcal{G}_\mathcal{R}^{v\mu\omega})$  are *svnsq-uniform spaces*. Then,  $\psi_\varphi : \widetilde{\mathbb{Y}, \gamma} \rightarrow \widetilde{\mathcal{U}, \mathcal{R}}$  is named a *svns-uniformly continuous* iff

$$\mathcal{G}_{\varphi(e)}^v(t) \leq \mathcal{K}_e^v(\psi_\varphi^{-1} \circ t \circ \psi_\varphi), \quad \mathcal{G}_{\varphi(e)}^\mu(t) \geq \mathcal{K}_e^\mu(\psi_\varphi^{-1} \circ t \circ \psi_\varphi),$$

$$\mathcal{G}_{\varphi(e)}^\omega(t) \geq \mathcal{K}_e^\omega(\psi_\varphi^{-1} \circ t \circ \psi_\varphi),$$

for every  $t \in \Psi(\widetilde{\mathcal{U}, \mathcal{R}}), e \in \gamma$ .

Sometimes in this paper we will use  $\mathcal{K}^{v\mu\omega}$  instead of  $(\mathcal{K}^v, \mathcal{K}^\mu, \mathcal{K}^\omega)$ .

**Theorem 2.** Assume that  $(\mathbb{Y}, \mathcal{K}_\gamma^{v\mu\omega})$  is *svnsq-uniform space* on  $\mathbb{Y}$ . Define

$$(\mathcal{K}_{st}^v)_e(z) = \bigvee \{\mathcal{K}_e^v(t) \mid t \cap \hat{t} \sqsubseteq z\} \text{ for any } e \in \gamma, t \in \zeta, z \in \Psi(\widetilde{\mathbb{Y}, \gamma}),$$

$$(\mathcal{K}_{st}^\mu)_e(z) = \bigwedge \{\mathcal{K}_e^\mu(t) \mid t \cap \hat{t} \sqsubseteq z\} \text{ for any } e \in \gamma, t \in \zeta, z \in \Psi(\widetilde{\mathbb{Y}, \gamma}),$$

$$(\mathcal{K}_{st}^\omega)_e(z) = \bigwedge \{\mathcal{K}_e^\omega(t) \mid t \cap \hat{t} \sqsubseteq z\} \text{ for any } e \in \gamma, t \in \zeta, z \in \Psi(\widetilde{\mathbb{Y}, \gamma}).$$

Then  $(\mathcal{K}_{st}^{v\mu\omega})_\gamma$  is the coarsest stratified svnsq-uniformity which is finer than  $\mathcal{K}_\gamma^{v\mu\omega}$ .

**Proof.** ( $\mathcal{K}_1$ ) There exists  $z \in \Psi(\widetilde{\mathbb{Y}, \gamma})$  such that  $\mathcal{K}_e^v(z) = 1, \mathcal{K}_e^\mu(z) = 0, \mathcal{K}_e^\omega(z) = 0$ . Since,  $z = z \cap \hat{0}, (\mathcal{K}_{st}^v)_e(z) = 1, (\mathcal{K}_{st}^\mu)_e(z) = 0, (\mathcal{K}_{st}^\omega)_e(z) = 0$ .

( $\mathcal{K}_2$ ) Obvious.

( $\mathcal{K}_3$ ) Assume that there exists  $z_1, z_2 \in \Psi(\widetilde{\mathbb{Y}, \gamma})$  such that

$$(\mathcal{K}_{st}^v)_e(z_1 \cap z_2) \not\geq (\mathcal{K}_{st}^v)_e(z_1) \wedge (\mathcal{K}_{st}^v)_e(z_2),$$

$$(\mathcal{K}_{st}^\mu)_e(z_1 \cap z_2) \not\leq (\mathcal{K}_{st}^\mu)_e(z_1) \vee (\mathcal{K}_{st}^\mu)_e(z_2),$$

$$(\mathcal{K}_{st}^\omega)_e(z_1 \cap z_2) \not\leq (\mathcal{K}_{st}^\omega)_e(z_1) \vee (\mathcal{K}_{st}^\omega)_e(z_2).$$

From the concept of  $(\mathcal{K}_{st}^{v\mu\omega})_\gamma$  there are  $t_1, t_2 \in \Psi(\widetilde{\mathbb{Y}, \gamma}), t_1, t_2 \in \zeta$  with  $z_1 \supseteq t_1 \cap \hat{t}_1$  and  $z_2 \supseteq t_2 \cap \hat{t}_2$  such that

$$(\mathcal{K}_{st}^v)_e(z_1 \cap z_2) \not\geq \mathcal{K}_e^v(t_1) \wedge \mathcal{K}_e^v(t_2),$$

$$(\mathcal{K}_{st}^\mu)_e(z_1 \cap z_2) \not\leq \mathcal{K}_e^\mu(t_1) \vee \mathcal{K}_e^\mu(t_2),$$

(1)

$$(\mathcal{K}_{st}^\omega)_e(z_1 \sqcap z_2) \not\leq \mathcal{K}_e^\omega(t_1) \vee \mathcal{K}_e^\omega(t_2).$$

On another side,  $(z_1 \sqcap z_2) \sqsupseteq (t_1 \sqcap t_2) \sqcap (\hat{t}_1 \sqcap \hat{t}_2)$ . By Lemma 1(3), we obtain  $(\hat{t}_1 \sqcap \hat{t}_2) = \hat{t}_1$  or  $\hat{t}_2$ , then

$$\begin{aligned} (\mathcal{K}_{st}^v)_e(z_1 \sqcap z_2) &\geq \mathcal{K}_e^v(t_1 \sqcap t_2) \geq \mathcal{K}_e^v(t_1) \wedge \mathcal{K}_e^v(t_2), \\ (\mathcal{K}_{st}^\mu)_e(z_1 \sqcap z_2) &\leq \mathcal{K}_e^\mu(t_1 \sqcap t_2) \leq \mathcal{K}_e^\mu(t_1) \vee \mathcal{K}_e^\mu(t_2), \\ (\mathcal{K}_{st}^\omega)_e(z_1 \sqcap z_2) &\leq \mathcal{K}_e^\omega(t_1 \sqcap t_2) \leq \mathcal{K}_e^\omega(t_1) \vee \mathcal{K}_e^\omega(t_2). \end{aligned}$$

It is a contradiction for equation (1). Hence  $(\mathcal{K}_3)$  holds.

$(\mathcal{K}_4)$  Let  $z \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ , be a given such that

$$\begin{aligned} (\mathcal{K}_{st}^v)_e(z) &\not\leq \bigvee \{(\mathcal{K}_{st}^v)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}, \\ (\mathcal{K}_{st}^\mu)_e(z) &\not\leq \bigwedge \{(\mathcal{K}_{st}^\mu)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}, \\ (\mathcal{K}_{st}^\omega)_e(z) &\not\leq \bigwedge \{(\mathcal{K}_{st}^\omega)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}. \end{aligned}$$

From the concept of  $[(\mathcal{K}_{st}^v)_e(z), (\mathcal{K}_{st}^\mu)_e(z), (\mathcal{K}_{st}^\omega)_e(z)]$  there exist  $t \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ ,  $t \in \zeta$  with  $z \sqsupseteq t \sqcap \hat{t}$  such that

$$\begin{aligned} \mathcal{K}_e^v(t) &\not\leq \{(\mathcal{K}_{st}^v)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}, \\ \mathcal{K}_e^\mu(t) &\not\leq \{(\mathcal{K}_{st}^\mu)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}, \\ \mathcal{K}_e^\omega(t) &\not\leq \{(\mathcal{K}_{st}^\omega)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}. \end{aligned}$$

Since  $\mathcal{K}_V^{v\mu\omega}$  is *svnsq-uniformity* on  $\mathfrak{X}$ ,

$$\begin{aligned} \bigvee \{\mathcal{K}_e^v(c) \mid t \sqsupseteq c \circ c\} &\geq \mathcal{K}_e^v(t), \\ \bigwedge \{\mathcal{K}_e^\mu(c) \mid t \sqsupseteq c \circ c\} &\leq \mathcal{K}_e^\mu(t), \\ \bigwedge \{\mathcal{K}_e^\omega(c) \mid t \sqsupseteq c \circ c\} &\leq \mathcal{K}_e^\omega(t). \end{aligned}$$

There exist  $c \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  with  $c \circ c \sqsubseteq t$  such that

$$\begin{aligned} \mathcal{K}_e^v(c) &\not\leq \bigvee \{(\mathcal{K}_{st}^v)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}, \\ \mathcal{K}_e^\mu(c) &\not\leq \bigwedge \{(\mathcal{K}_{st}^\mu)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}, \\ \mathcal{K}_e^\omega(c) &\not\leq \bigwedge \{(\mathcal{K}_{st}^\omega)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\}. \end{aligned} \tag{2}$$

On another side,

$$(c \sqcap \hat{t}) \circ (c \sqcap \hat{t}) \sqsubseteq (c \circ c) \sqcap \hat{t} \sqsubseteq t \sqcap \hat{t} \sqsubseteq z,$$

that is,  $c \sqcap \hat{t} = z_1$  with  $z = z_1 \circ z_1$ ,

$$\begin{aligned} \bigvee \{(\mathcal{K}_{st}^v)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\} &\geq (\mathcal{K}_{st}^v)_e(z_1) \geq \mathcal{K}_e^v(c), \\ \bigwedge \{(\mathcal{K}_{st}^\mu)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\} &\leq (\mathcal{K}_{st}^\mu)_e(z_1) \leq \mathcal{K}_e^\mu(c), \\ \bigwedge \{(\mathcal{K}_{st}^\omega)_e(z_1) \mid z \sqsupseteq z_1 \circ z_1\} &\leq (\mathcal{K}_{st}^\omega)_e(z_1) \leq \mathcal{K}_e^\omega(c). \end{aligned}$$

In this case, it is a contradiction with the hypothesis as we can see from Equations (2). Hence  $(\mathcal{K}_4)$  holds.

$(\mathcal{K}_5)$  By Lemma 1(3), we have  $z \sqsubseteq \hat{0}$  holds for every  $z \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ . Since  $\mathfrak{X}_V^{v\mu\omega}$  satisfies the conditions  $\mathcal{K}_1$  and  $\mathcal{K}_2$ , we obtain  $\mathcal{K}_e^v(\hat{0}) = 1$ ,  $\mathcal{K}_e^\mu(\hat{0}) = 0$  and  $\mathcal{K}_e^\omega(\hat{0}) = 0$ . So,  $\hat{0} \sqcap \hat{t} = \hat{t}$  for every  $t \in \zeta$ , then  $(\mathcal{K}_{st}^v)_e(\hat{t}) = 1$ ,  $(\mathcal{K}_{st}^\mu)_e(\hat{t}) = 0$ ,  $(\mathcal{K}_{st}^\omega)_e(\hat{t}) = 0$ . Hence,  $(\mathcal{K}_{st}^{v\mu\omega})_V$  is stratified.

For  $t = t \sqcap \hat{0}$  and  $t \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ . Then,  $(\mathcal{K}_{st}^v)_e(t) \geq \mathcal{K}_e^v(t)$ ,  $(\mathcal{K}_{st}^\mu)_e(t) \leq \mathcal{K}_e^\mu(t)$  and  $(\mathcal{K}_{st}^\omega)_e(t) \leq \mathcal{K}_e^\omega(t)$ . Hence,  $(\mathcal{K}_{st}^{v\mu\omega})_V$  is finer than  $\mathcal{K}_V^{v\mu\omega}$ .

Finally, let  $\mathcal{G}_V^{v\mu\omega}$  be stratified svnsq-uniformity finer than  $\mathcal{K}_V^{v\mu\omega}$ .

Presume that there exists  $z \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  s.t.,

$$\mathcal{G}_e^v(z) \not\leq (\mathcal{K}_{st}^v)_e(z), \mathcal{G}_e^\mu(z) \not\leq (\mathcal{K}_{st}^\mu)_e(z), \mathcal{G}_e^\omega(z) \not\leq (\mathcal{K}_{st}^\omega)_e(z).$$

From the concept of  $[(\mathcal{K}_{st}^v)_e(z), (\mathcal{K}_{st}^\mu)_e(z), (\mathcal{K}_{st}^\omega)_e(z)]$ , there exist  $t \in \Psi(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ ,  $t \in \zeta$  with  $z \sqsupseteq t \sqcap \hat{t}$  such that

$$\mathcal{G}_e^v(z) \not\leq \mathcal{K}_e^v(t), \mathcal{G}_e^\mu(z) \not\leq \mathcal{K}_e^\mu(t), \mathcal{G}_e^\omega(z) \not\leq \mathcal{K}_e^\omega(t). \tag{3}$$

Since  $\mathcal{G}_V^{v\mu\omega}$  be stratified svnsq-uniformity,

$$\begin{aligned} \mathcal{G}_e^v(z) &\geq \mathcal{G}_e^v(t \sqcap \hat{t}) \geq \mathcal{G}_e^v(t) \wedge \mathcal{G}_e^v(\hat{t}) \geq \mathcal{G}_e^v(t) \geq \mathcal{K}_e^v(t), \\ \mathcal{G}_e^{\mu}(z) &\leq \mathcal{G}_e^{\mu}(t \sqcap \hat{t}) \leq \mathcal{G}_e^{\mu}(t) \wedge \mathcal{G}_e^{\mu}(\hat{t}) \leq \mathcal{G}_e^{\mu}(t) \leq \mathcal{K}_e^{\mu}(t), \\ \mathcal{G}_e^{\omega}(z) &\leq \mathcal{G}_e^{\omega}(t \sqcap \hat{t}) \leq \mathcal{G}_e^{\omega}(t) \wedge \mathcal{G}_e^{\omega}(\hat{t}) \leq \mathcal{G}_e^{\omega}(t) \leq \mathcal{K}_e^{\omega}(t). \end{aligned}$$

In this case, the hypothesis is contradicted, as we see in Equations (3). Therefore,  $(\mathcal{K}_{st}^{v\mu\omega})_{\mathcal{V}}$  is the coarsest stratified svnsq-uniformity which is finer than  $\mathcal{K}_{\mathcal{V}}^{v\mu\omega}$ .  $\square$

**Theorem 3.** Let  $(\mathbb{Y}, \mathcal{K}_{\mathcal{V}}^{v\mu\omega})$  and  $(\mathcal{U}, \mathcal{G}_{\mathcal{R}}^{v\mu\omega})$  be two svnsq-uniform spaces. If the mapping  $\psi_{\varphi} : (\mathbb{Y}, \mathcal{K}_{\mathcal{V}}^{v\mu\omega}) \rightarrow (\mathcal{U}, \mathcal{G}_{\mathcal{R}}^{v\mu\omega})$  is a svns-uniformly continuous, then the mapping  $\psi_{\varphi} : (\mathbb{Y}, (\mathcal{K}_{st}^{v\mu\omega})_{\mathcal{V}}) \rightarrow (\mathcal{U}, (\mathcal{G}_{st}^{v\mu\omega})_{\mathcal{R}})$  is a svns-uniformly continuous, where  $(\mathcal{K}_{st}^{v\mu\omega})_{\mathcal{V}}$  and  $(\mathcal{G}_{st}^{v\mu\omega})_{\mathcal{R}}$  are the stratification of  $\mathcal{K}_{\mathcal{V}}^{v\mu\omega}$  and  $\mathcal{G}_{\mathcal{R}}^{v\mu\omega}$ , respectively.

**Proof.** We will prove that

$$\begin{aligned} (\mathcal{K}_{st}^v)_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\geq (\mathcal{G}_{st}^v)_{\varphi(e)}(t), \\ (\mathcal{K}_{st}^{\mu})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\leq (\mathcal{G}_{st}^{\mu})_{\varphi(e)}(t), \\ (\mathcal{K}_{st}^{\omega})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\leq (\mathcal{G}_{st}^{\omega})_{\varphi(e)}(t), \end{aligned}$$

for each  $t \in \Psi(\widetilde{(\mathcal{U}, \mathcal{R})})$ . Assume that

$$\begin{aligned} (\mathcal{K}_{st}^v)_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\not\geq (\mathcal{G}_{st}^v)_{\varphi(e)}(t), \quad (\mathcal{K}_{st}^{\mu})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) \not\leq (\mathcal{G}_{st}^{\mu})_{\varphi(e)}(t), \\ (\mathcal{K}_{st}^{\omega})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\not\leq (\mathcal{G}_{st}^{\omega})_{\varphi(e)}(t). \end{aligned}$$

From the concept of  $[(\mathcal{G}_{st}^v)_{\varphi(e)}(t), (\mathcal{G}_{st}^{\mu})_{\varphi(e)}(t), (\mathcal{G}_{st}^{\omega})_{\varphi(e)}(t)]$ , there exist  $c \in \Psi(\widetilde{(\mathcal{U}, \mathcal{R})})$ ,  $t \sqsupseteq c \sqcap \hat{t}$  such that

$$\begin{aligned} (\mathcal{K}_{st}^v)_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\not\geq \mathcal{G}_{\varphi(e)}^v(c), \\ (\mathcal{K}_{st}^{\mu})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\not\leq \mathcal{G}_{\varphi(e)}^{\mu}(c), \\ (\mathcal{K}_{st}^{\omega})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\not\leq \mathcal{G}_{\varphi(e)}^{\omega}(c). \end{aligned} \tag{4}$$

Since  $\psi_{\varphi} : (\mathbb{Y}, \mathcal{K}_{\mathcal{V}}^{v\mu\omega}) \rightarrow (\mathcal{U}, \mathcal{G}_{\mathcal{R}}^{v\mu\omega})$  is a svns-uniformly continuous,

$$\begin{aligned} \mathcal{K}_e^v(\psi_{\varphi}^{-1} \circ c \circ \psi_{\varphi}) &\geq \mathcal{G}_{\varphi(e)}^v(c), \quad \mathcal{K}_e^{\mu}(\psi_{\varphi}^{-1} \circ c \circ \psi_{\varphi}) \leq \mathcal{G}_{\varphi(e)}^{\mu}(c), \\ \mathcal{K}_e^{\omega}(\psi_{\varphi}^{-1} \circ c \circ \psi_{\varphi}) &\leq \mathcal{G}_{\varphi(e)}^{\omega}(c). \end{aligned}$$

From the concept of  $[(\mathcal{K}_{st}^v)_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}), (\mathcal{K}_{st}^{\mu})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}), (\mathcal{K}_{st}^{\omega})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi})]$  we obtain

$$\begin{aligned} (\mathcal{K}_{st}^v)_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\geq \mathcal{K}_e^v(\psi_{\varphi}^{-1} \circ c \circ \psi_{\varphi}) \geq \mathcal{G}_{\varphi(e)}^v(c), \\ (\mathcal{K}_{st}^{\mu})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\leq \mathcal{K}_e^{\mu}(\psi_{\varphi}^{-1} \circ c \circ \psi_{\varphi}) \leq \mathcal{G}_{\varphi(e)}^{\mu}(c), \\ (\mathcal{K}_{st}^{\omega})_e(\psi_{\varphi}^{-1} \circ t \circ \psi_{\varphi}) &\leq \mathcal{K}_e^{\omega}(\psi_{\varphi}^{-1} \circ c \circ \psi_{\varphi}) \leq \mathcal{G}_{\varphi(e)}^{\omega}(c). \end{aligned}$$

In this case, the hypothesis is contradicted as we can see from Equations (4).  $\square$

**Theorem 4.** Let  $(\mathbb{Y}, \mathcal{T}_{\mathcal{V}}^{v\mu\omega})$  be a stratified svnsts. Define,

$$\begin{aligned} \mathcal{K}_e^v(z) &= \bigvee \left\{ \bigwedge_{j \in J} \mathcal{T}_e^v((\mathcal{E}_{\sigma})_j) \mid \forall z \in \Psi(\widetilde{(\mathbb{Y}, \mathcal{Y})}), \sqcap_{j \in J} z(\mathcal{E}_{\sigma})_j \sqsubseteq z \right\}, \\ \mathcal{K}_e^{\mu}(z) &= \bigwedge \left\{ \bigvee_{j \in J} \mathcal{T}_e^{\mu}((\mathcal{E}_{\sigma})_j) \mid \forall z \in \Psi(\widetilde{(\mathbb{Y}, \mathcal{Y})}), \sqcap_{j \in J} z(\mathcal{E}_{\sigma})_j \sqsubseteq z \right\}, \\ \mathcal{K}_e^{\omega}(z) &= \bigwedge \left\{ \bigvee_{j \in J} \mathcal{T}_e^{\omega}((\mathcal{E}_{\sigma})_j) \mid \forall z \in \Psi(\widetilde{(\mathbb{Y}, \mathcal{Y})}), \sqcap_{j \in J} z(\mathcal{E}_{\sigma})_j \sqsubseteq z \right\}, \end{aligned}$$

where  $\bigvee$  and  $\bigwedge$  are possessed over all collections  $\{(\mathcal{E}_{\sigma})_j \mid \sqcap_{j \in J} z(\mathcal{E}_{\sigma})_j \sqsubseteq z\}$ . Then  $\mathcal{K}_{\mathcal{V}}^{v\mu\omega}$  is stratified svnsq-uniformity on  $\mathbb{Y}$ .

**Proof.**  $(\mathcal{K}_1)$  and  $(\mathcal{K}_2)$  straightforward.

$(\mathcal{K}_3)$  Assume there exists  $z, t \in \Psi(\widetilde{(\mathbb{Y}, \mathcal{Y})})$ , such that  $\mathcal{K}_e^v(z \sqcap t) \not\geq \mathcal{K}_e^v(z) \wedge \mathcal{K}_e^v(t)$ ,  $\mathcal{K}_e^{\mu}(z \sqcap t) \not\leq \mathcal{K}_e^{\mu}(z) \vee \mathcal{K}_e^{\mu}(t)$ , and  $\mathcal{K}_e^{\omega}(z \sqcap t) \not\leq \mathcal{K}_e^{\omega}(z) \vee \mathcal{K}_e^{\omega}(t)$ . Then, by the concept of  $\mathcal{K}_{\mathcal{V}}^{v\mu\omega}$ , there exist two finite collections  $\{(\mathcal{H}_{\alpha})_i \mid t \sqsupseteq \sqcap_{i \in \Gamma} (\mathcal{H}_{\alpha})_i\}$  and  $\{(\mathcal{E}_{\sigma})_j \mid z \sqsupseteq \sqcap_{j \in J} z(\mathcal{E}_{\sigma})_j\}$  such that



$$\begin{aligned}
 \mathcal{K}_e^v(z \sqcap t) &\not\leq \left[ \bigwedge_{j \in J} \mathcal{T}_e^v((\mathcal{E}_\sigma)_j) \right] \wedge \left[ \bigwedge_{i \in \Gamma} \mathcal{T}_e^v((\mathcal{H}_\alpha)_i) \right], \\
 \mathcal{K}_e^\mu(z \sqcap t) &\not\leq \left[ \bigvee_{j \in J} \mathcal{T}_e^\mu((\mathcal{E}_\sigma)_j) \right] \vee \left[ \bigvee_{i \in \Gamma} \mathcal{T}_e^\mu((\mathcal{H}_\alpha)_i) \right], \\
 \mathcal{K}_e^\omega(z \sqcap t) &\not\leq \left[ \bigvee_{j \in J} \mathcal{T}_e^\omega((\mathcal{E}_\sigma)_j) \right] \vee \left[ \bigvee_{i \in \Gamma} \mathcal{T}_e^\omega((\mathcal{H}_\alpha)_i) \right].
 \end{aligned} \tag{5}$$

On another side,

$$z \sqcap t \sqsupseteq (\bigcap_{j \in J} z_{(\mathcal{E}_\sigma)_j}) \wedge (\bigcap_{i \in \Gamma} t_{(\mathcal{H}_\alpha)_i}) \sqsupseteq \bigcap_{j \in J, i \in \Gamma} (z \sqcap t)_{(\mathcal{E}_\sigma)_j \sqcap (\mathcal{H}_\alpha)_i}.$$

Then by the definition of  $\mathcal{K}_v^{\nu\mu\omega}$ ,

$$\begin{aligned}
 \mathcal{K}_e^v(z \sqcap t) &\geq \bigwedge_{j \in J, i \in \Gamma} \mathcal{T}_e^v((\mathcal{E}_\sigma)_j \sqcap (\mathcal{H}_\alpha)_i) \geq \bigwedge_{j \in J, i \in \Gamma} (\mathcal{T}_e^v((\mathcal{E}_\sigma)_j) \sqcap \mathcal{T}_e^v((\mathcal{H}_\alpha)_i)) \\
 &\geq \left[ \bigwedge_{j \in J} \mathcal{T}_e^v((\mathcal{E}_\sigma)_j) \right] \wedge \left[ \bigwedge_{i \in \Gamma} \mathcal{T}_e^v((\mathcal{H}_\alpha)_i) \right], \\
 \mathcal{K}_e^\mu(z \sqcap t) &\leq \bigvee_{j \in J, i \in \Gamma} \mathcal{T}_e^\mu((\mathcal{E}_\sigma)_j \sqcap (\mathcal{H}_\alpha)_i) \leq \bigvee_{j \in J, i \in \Gamma} (\mathcal{T}_e^\mu((\mathcal{E}_\sigma)_j) \sqcup \mathcal{T}_e^\mu((\mathcal{H}_\alpha)_i)) \\
 &\leq \left[ \bigvee_{j \in J} \mathcal{T}_e^\mu((\mathcal{E}_\sigma)_j) \right] \vee \left[ \bigvee_{i \in \Gamma} \mathcal{T}_e^\mu((\mathcal{H}_\alpha)_i) \right], \\
 \mathcal{K}_e^\omega(z \sqcap t) &\leq \bigvee_{j \in J, i \in \Gamma} \mathcal{T}_e^\omega((\mathcal{E}_\sigma)_j \sqcap (\mathcal{H}_\alpha)_i) \leq \bigvee_{j \in J, i \in \Gamma} (\mathcal{T}_e^\omega((\mathcal{E}_\sigma)_j) \sqcap \mathcal{T}_e^\omega((\mathcal{H}_\alpha)_i)) \\
 &\leq \left[ \bigvee_{j \in J} \mathcal{T}_e^\omega((\mathcal{E}_\sigma)_j) \right] \vee \left[ \bigvee_{i \in \Gamma} \mathcal{T}_e^\omega((\mathcal{H}_\alpha)_i) \right],
 \end{aligned}$$

which is a contradiction for equations (5), and then  $(\mathcal{K}_3)$  holds.

$(\mathcal{K}_4)$  Since,  $z_{\mathcal{E}_\sigma} \circ z_{\mathcal{E}_\sigma} = z_{\mathcal{E}_\sigma}$ . From the Lemma 2, then,  $(\mathcal{K}_4)$  holds.

$(\mathcal{K}_1)$  and  $(\mathcal{K}_5)$  There exists  $z = z_{\tilde{\gamma}^l} = \hat{\gamma}$ , then  $\mathcal{K}_e^v(\hat{\gamma}) \geq \mathcal{T}_e^v(\tilde{\gamma}^l) = 1$ ,  $\mathcal{K}_e^\mu(\hat{\gamma}) \leq \mathcal{T}_e^\mu(\tilde{\gamma}^l) = 0$  and  $\mathcal{K}_e^\omega(\hat{\gamma}) \leq \mathcal{T}_e^\omega(\tilde{\gamma}^l) = 0$ . Therefore,  $\mathcal{K}_e^v(\hat{\gamma}) = 1$ ,  $\mathcal{K}_e^\mu(\hat{\gamma}) = 0$  and  $\mathcal{K}_e^\omega(\hat{\gamma}) = 0$  for each  $l \in \zeta$ . Hence,  $\mathcal{K}_v^{\nu\mu\omega}$  is stratified.  $\square$

### 3. Stratified single-valued neutrosophic soft topogenous order spaces

**Definition 6.** Maps  $\mathcal{H}^v : \mathcal{V} \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \mathcal{V})} \times \widetilde{(\mathfrak{X}, \mathcal{V})}}$ ,  $\mathcal{H}^\mu : \mathcal{V} \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \mathcal{V})} \times \widetilde{(\mathfrak{X}, \mathcal{V})}}$  and  $\mathcal{H}^\omega : \mathcal{V} \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \mathcal{V})} \times \widetilde{(\mathfrak{X}, \mathcal{V})}}$  are said to be *svns-topogenous order* on  $\mathfrak{X}$  if it fulfills the next properties:  $\forall e \in \mathcal{V}$  and  $\mathcal{E}_\sigma, \mathcal{H}_\alpha \in \widetilde{(\mathfrak{X}, \mathcal{V})}$ ,

$$(H_1) \mathcal{H}_e^\mu(\tilde{\gamma}, \tilde{\gamma}) = \mathcal{H}_e^\mu(\Phi, \Phi) = \mathcal{H}_e^\omega(\tilde{\gamma}, \tilde{\gamma}) = \mathcal{H}_e^\omega(\Phi, \Phi) = 0, \mathcal{H}_e^v(\tilde{\gamma}, \tilde{\gamma}) = \mathcal{H}_e^v(\Phi, \Phi) = 1,$$

$$(H_2) \text{ If } \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{H}_\alpha) \neq 0, \mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathcal{H}_\alpha) \neq 1 \text{ and } \mathcal{H}_e^\omega(\mathcal{E}_\sigma, \mathcal{H}_\alpha) \neq 1, \text{ then } \mathcal{E}_\sigma \sqsubseteq \mathcal{H}_\alpha.$$

$$(H_3) \text{ If } \mathcal{E}_\sigma \sqsubseteq (\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1 \sqsubseteq \mathcal{H}_\alpha, \text{ then } \mathcal{H}_e^v((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \leq \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{H}_\alpha),$$

$$\mathcal{H}_e^\mu((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \geq \mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathcal{H}_\alpha) \text{ and } \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \geq \mathcal{H}_e^\omega(\mathcal{E}_\sigma, \mathcal{H}_\alpha).$$

$(H_4)$  (i)

$$\mathcal{H}_e^v((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcup (\mathcal{H}_\alpha)_2) \geq \mathcal{H}_e^v((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \wedge \mathcal{H}_e^v((\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_2),$$

$$\mathcal{H}_e^\mu((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcup (\mathcal{H}_\alpha)_2) \leq \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \vee \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_2),$$

$$\mathcal{H}_e^\omega((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcup (\mathcal{H}_\alpha)_2) \leq \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \vee \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_2).$$

(ii)

$$\mathcal{H}_e^v((\mathcal{E}_\sigma)_1 \sqcap (\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcap (\mathcal{H}_\alpha)_2) \geq \mathcal{H}_e^v((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \wedge \mathcal{H}_e^v((\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_2),$$

$$\mathcal{H}_e^\mu((\mathcal{E}_\sigma)_1 \sqcap (\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcap (\mathcal{H}_\alpha)_2) \leq \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \vee \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_2).$$

$$\mathcal{H}_e^\omega((\mathcal{E}_\sigma)_1 \sqcap (\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcap (\mathcal{H}_\alpha)_2) \leq \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_1, (\mathcal{H}_\alpha)_1) \vee \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_2, (\mathcal{H}_\alpha)_2).$$

Therefore,  $(\mathfrak{X}, \mathcal{H}_v^{\nu\mu\omega})$  is termed to be a *svns-topogenous order space*. Also,  $\mathcal{H}_v^{\nu\mu\omega}$  is said to be

(1) Symmetrical iff  $\mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{H}_\alpha) = \mathcal{H}_e^v(\mathcal{H}_\alpha^c, \mathcal{E}_\sigma^c)$ ,  $\mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathcal{H}_\alpha) = \mathcal{H}_e^\mu(\mathcal{H}_\alpha^c, \mathcal{E}_\sigma^c)$  and  $\mathcal{H}_e^\omega(\mathcal{H}_\alpha, \mathcal{E}_\sigma) = \mathcal{H}_e^\omega(\mathcal{H}_\alpha^c, \mathcal{E}_\sigma^c)$ .

(2) Perfect iff

$$\begin{aligned} \mathcal{H}_e^v(\sqcup_{j \in J}(\mathcal{E}_\sigma)_j, \sqcup_{j \in J}(\mathfrak{h}_\alpha)_j) &\geq \bigwedge_{j \in J} \mathcal{H}_e^v((\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j), \\ \mathcal{H}_e^\mu(\sqcup_{j \in J}(\mathcal{E}_\sigma)_j, \sqcup_{j \in J}(\mathfrak{h}_\alpha)_j) &\leq \bigvee_{j \in J} \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j), \\ \mathcal{H}_e^\omega(\sqcup_{j \in J}(\mathcal{E}_\sigma)_j, \sqcup_{j \in J}(\mathfrak{h}_\alpha)_j) &\leq \bigvee_{j \in J} \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j). \end{aligned}$$

(3) Stratified iff  $\mathcal{H}_V^{\nu\mu\omega}$  satisfies the condition:

$$(H_S) \text{ For every } \iota \in \zeta, \mathcal{H}_e^v(\tilde{\nu}^\iota, \tilde{\nu}^\iota) = 1, \mathcal{H}_e^\mu(\tilde{\nu}^\iota, \tilde{\nu}^\iota) = 0, \mathcal{H}_e^\omega(\tilde{\nu}^\iota, \tilde{\nu}^\iota) = 0.$$

Suppose that  $(\mathcal{H}_V^{\nu\mu\omega})_1$  and  $(\mathcal{H}_V^{\nu\mu\omega})_2$  be *svns-topogenous order space* on  $\mathcal{V}$ . In our opinion  $(\mathcal{H}_V^{\nu\mu\omega})_1$  is finer than  $(\mathcal{H}_V^{\nu\mu\omega})_2$  [ $(\mathcal{H}_V^{\nu\mu\omega})_2$  is coarser than  $(\mathcal{H}_V^{\nu\mu\omega})_1$ ] indicated by  $(\mathcal{H}_V^{\nu\mu\omega})_1 \supseteq (\mathcal{H}_V^{\nu\mu\omega})_2$  if

$$\begin{aligned} (\mathcal{H}_e^v)_2(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\leq (\mathcal{H}_e^v)_1(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad (\mathcal{H}_e^\mu)_2(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq (\mathcal{H}_e^\mu)_1(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\ (\mathcal{H}_e^\omega)_2(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq (\mathcal{H}_e^\omega)_1(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad \forall \mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathcal{Y})}, e \in \mathcal{V}. \end{aligned}$$

Suppose  $(\mathfrak{X}, (\mathcal{H}_V^{\nu\mu\omega})_1)$  and  $(\mathcal{U}, (\mathcal{H}_R^{\nu\mu\omega})_2)$  be two *svns-topogenous order spaces*.

Then a map  $\psi_\varphi : \widetilde{(\mathfrak{X}, \mathcal{Y})} \rightarrow \widetilde{(\mathcal{U}, \mathcal{R})}$  is called *snv-soft topogenous continuous* iff

$$\begin{aligned} (\mathcal{H}_{\varphi(e)}^v)_2(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\leq (\mathcal{H}_e^v)_1(\psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathfrak{h}_\alpha)), \\ (\mathcal{H}_{\varphi(e)}^\mu)_2(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq (\mathcal{H}_e^\mu)_1(\psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathfrak{h}_\alpha)), \\ (\mathcal{H}_{\varphi(e)}^\omega)_2(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq (\mathcal{H}_e^\omega)_1(\psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathfrak{h}_\alpha)), \end{aligned}$$

for any  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathcal{U}, \mathcal{R})}$ ,  $e \in \mathcal{V}$ .

**Theorem 5.** Let  $(\mathfrak{X}, \mathcal{H}_V^{\nu\mu\omega})$  be *svns-topogenous order space*. Define

$$\begin{aligned} (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= \bigvee_{\{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\nu}^{lj} \mid j \in J\} \in \mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left\{ \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\nu}^{li}) \in \mathcal{A}} \mathcal{H}_e^v((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \right\}, \\ (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= \bigwedge_{\{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\nu}^{lj} \mid j \in J\} \in \mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left\{ \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\nu}^{li}) \in \mathcal{A}} \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \right\}, \\ (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= \bigwedge_{\{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\nu}^{lj} \mid j \in J\} \in \mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left\{ \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\nu}^{li}) \in \mathcal{A}} \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \right\}, \end{aligned}$$

where  $\mathcal{A} = \{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\nu}^{lj} \mid j \in J\}$  and  $\mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = \{\{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\nu}^{lj} \mid j \in J, J \text{ is finite}\} \mid \mathcal{E}_\sigma \sqsubseteq \sqcup_{j \in J}(\mathcal{E}_\sigma)_j \cap \tilde{\nu}^{lj} \text{ and } \mathfrak{h}_\alpha \sqsupseteq \sqcup_{j \in J}(\mathfrak{h}_\alpha)_j \cap \tilde{\nu}^{lj}, \iota \in \zeta\}$ . Then  $(\mathcal{H}_{st}^{\nu\mu\omega})_\mathcal{V}$  is the coarsest stratified *svns-topogenous order* on  $\mathfrak{X}$  which is finer than  $\mathcal{H}_V^{\nu\mu\omega}$ .

**Proof.**  $(H_1), (H_2), (H_3)$  straightforward.

$(H_4)$  (i) Assume that there exist,  $(\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1, (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2 \in \widetilde{(\mathfrak{X}, \mathcal{Y})}$  such that

$$\begin{aligned} (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) &\not\geq (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1) \wedge (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2), \\ (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) &\not\leq (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1) \vee (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2), \\ (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) &\not\leq (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1) \vee (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2), \end{aligned}$$

therefore, there exists  $r \in \zeta_0$  such that

$$\begin{aligned}
 & (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) < r \\
 & \leq (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1) \wedge (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2), \\
 & (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) \geq 1 - r > (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1) \vee (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2), \\
 & (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) \geq 1 - r \\
 & > (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1) \vee (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2).
 \end{aligned} \tag{6}$$

From the concept of  $(\mathcal{H}_{st}^{\nu\mu\omega})_\gamma$ , there are  $\forall, i \in J$  and  $k \in \Gamma$ ,  $\{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i)\} \in \mathcal{N}((\mathcal{E}_\sigma)_1, (\mathfrak{h}_\alpha)_1)$  and  $\{((\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_k, \tilde{\gamma}^k)\} \in \mathcal{N}((\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_2)$ , such that

$$\begin{aligned}
 & (\mathcal{H}_{st}^v)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) \geq \bigwedge_{i,k} \mathcal{H}_e^v((\mathcal{E}_\sigma)_i \sqcup (\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_i \sqcup (\mathfrak{h}_\alpha)_k) \\
 & \geq \bigwedge_{i,k} \mathcal{H}_e^v((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \wedge \mathcal{H}_e^v((\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_k) \geq r, \\
 & (\mathcal{H}_{st}^\mu)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) \leq \bigvee_{i,k} \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_i \sqcup (\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_i \sqcup (\mathfrak{h}_\alpha)_k) \\
 & \leq \bigvee_{i,k} \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \vee \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_k) < 1 - r, \\
 & (\mathcal{H}_{st}^\omega)_e((\mathcal{E}_\sigma)_1 \sqcup (\mathcal{E}_\sigma)_2, (\mathfrak{h}_\alpha)_1 \sqcup (\mathfrak{h}_\alpha)_2) \leq \bigvee_{i,k} \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_i \sqcup (\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_i \sqcup (\mathfrak{h}_\alpha)_k) \\
 & \leq \bigvee_{i,k} \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \vee \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_k, (\mathfrak{h}_\alpha)_k) < 1 - r.
 \end{aligned}$$

In this case, it is a contradiction with the hypothesis, as is clear from Equation No. (6). Hence,  $\mathcal{H}_4(i)$  holds.

(ii) In the same way as used to prove (i).

$(\mathcal{H}_5)$  Since  $\tilde{\gamma}' = \tilde{\gamma}' \sqcap \tilde{\gamma}$ ,

$$\begin{aligned}
 & (\mathcal{H}_{st}^v)_e(\tilde{\gamma}', \tilde{\gamma}') \geq \mathcal{H}_e^v(\tilde{\gamma}, \tilde{\gamma}) = 1, \quad (\mathcal{H}_{st}^\mu)_e(\tilde{\gamma}', \tilde{\gamma}') \leq \mathcal{H}_e^\mu(\tilde{\gamma}, \tilde{\gamma}) = 0, \\
 & (\mathcal{H}_{st}^\omega)_e(\tilde{\gamma}', \tilde{\gamma}') \leq \mathcal{H}_e^\omega(\tilde{\gamma}, \tilde{\gamma}) = 0
 \end{aligned}$$

Hence,  $(\mathcal{H}_{st}^v)_e(\tilde{\gamma}', \tilde{\gamma}') = 1$ ,  $(\mathcal{H}_{st}^\mu)_e(\tilde{\gamma}', \tilde{\gamma}') = 0$  and  $(\mathcal{H}_{st}^\omega)_e(\tilde{\gamma}', \tilde{\gamma}') = 0$ ,  $\forall i \in \zeta$ .

On another side, since  $\mathcal{E}_\sigma \sqsubseteq \mathcal{E}_\sigma \sqcap \tilde{\gamma}$  and  $\mathfrak{h}_\alpha \supseteq \mathfrak{h}_\alpha \sqcap \tilde{\gamma}$  we obtain

$$\begin{aligned}
 & (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq \mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 & (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq \mathcal{H}_e^\omega(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad \forall \mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}.
 \end{aligned}$$

Hence,  $(\mathcal{H}_{st}^{\nu\mu\omega})_\gamma$  is the stratified svns-topogenous order on  $\gamma$  which is finer than  $\mathcal{H}_\gamma^{\nu\mu\omega}$ .

Finally, suppose that  $(\mathcal{H}_\gamma^{\nu\mu\omega})^*$  be stratified svns-topogenous and finer than  $\mathcal{H}_\gamma^{\nu\mu\omega}$  then,  $\forall \mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ,  $i \in \zeta, e \in \gamma$ .

$$\begin{aligned}
 & (\mathcal{H}_e^v)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad (\mathcal{H}_e^\mu)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq \mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 & (\mathcal{H}_e^\omega)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq \mathcal{H}_e^\omega(\mathcal{E}_\sigma, \mathfrak{h}_\alpha).
 \end{aligned}$$

Now we will prove that

$$\begin{aligned}
 & (\mathcal{H}_e^v)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad (\mathcal{H}_e^\mu)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 & (\mathcal{H}_e^\omega)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha).
 \end{aligned}$$

Suppose there exist  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ,  $e \in \gamma$  such that

$$\begin{aligned}
 & (\mathcal{H}_e^v)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \not\geq (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \quad (\mathcal{H}_e^\mu)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \not\leq (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 & (\mathcal{H}_e^\omega)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \not\leq (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha),
 \end{aligned}$$

then there is  $r \in \zeta$  such that

$$\begin{aligned}
 & (\mathcal{H}_e^v)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) < r \leq (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 & (\mathcal{H}_e^\mu)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq 1 - r > (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 & (\mathcal{H}_e^\omega)^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq 1 - r > (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha).
 \end{aligned} \tag{7}$$

From the concept of  $(\mathcal{H}_{st}^{v\mu\omega})_{\vee}$ , there exists  $\mathcal{A} = \{((\mathcal{E}_{\sigma})_j, (\mathcal{h}_{\alpha})_j, \tilde{\gamma}^{lj})\} \in \mathcal{N}(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha})$ , for every  $j \in J$  such that

$$\begin{aligned}
 (\mathcal{H}_e^v)^*(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha}) &\geq (\mathcal{H}_e^v)^*(\sqcup_{j \in J}[(\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}], \sqcup_{j \in J}[(\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}]) \\
 &\geq \bigwedge_{j \in J} (\mathcal{H}_e^v)^*((\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}, (\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}) \\
 &\geq \bigwedge_{j \in J} [(\mathcal{H}_e^v)^*((\mathcal{E}_{\sigma})_j, (\mathcal{h}_{\alpha})_j) \wedge (\mathcal{H}_e^v)^*(\tilde{\gamma}^{lj}, \tilde{\gamma}^{lj})] \\
 &\geq \bigwedge_{((\mathcal{E}_{\sigma})_i, (\mathcal{h}_{\alpha})_i, \tilde{\gamma}^i) \in \mathcal{A}} \mathcal{H}_e^v((\mathcal{E}_{\sigma})_i, (\mathcal{h}_{\alpha})_i) \geq r, \\
 (\mathcal{H}_e^{\mu})^*(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha}) &\leq (\mathcal{H}_e^{\mu})^*(\sqcup_{j \in J}[(\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}], \sqcup_{j \in J}[(\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}]) \\
 &\leq \bigvee_{j \in J} (\mathcal{H}_e^{\mu})^*((\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}, (\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}) \\
 &\leq \bigvee_{j \in J} [(\mathcal{H}_e^{\mu})^*((\mathcal{E}_{\sigma})_j, (\mathcal{h}_{\alpha})_j) \vee (\mathcal{H}_e^{\mu})^*(\tilde{\gamma}^{lj}, \tilde{\gamma}^{lj})] \\
 &\leq \bigvee_{((\mathcal{E}_{\sigma})_i, (\mathcal{h}_{\alpha})_i, \tilde{\gamma}^i) \in \mathcal{A}} \mathcal{H}_e^{\mu}((\mathcal{E}_{\sigma})_i, (\mathcal{h}_{\alpha})_i) \leq 1 - r, \\
 (\mathcal{H}_e^{\omega})^*(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha}) &\leq (\mathcal{H}_e^{\omega})^*(\sqcup_{j \in J}[(\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}], \sqcup_{j \in J}[(\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}]) \\
 &\leq \bigvee_{j \in J} (\mathcal{H}_e^{\omega})^*((\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}, (\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}) \\
 &\leq \bigvee_{j \in J} [(\mathcal{H}_e^{\omega})^*((\mathcal{E}_{\sigma})_j, (\mathcal{h}_{\alpha})_j) \vee (\mathcal{H}_e^{\omega})^*(\tilde{\gamma}^{lj}, \tilde{\gamma}^{lj})] \\
 &\leq \bigvee_{((\mathcal{E}_{\sigma})_i, (\mathcal{h}_{\alpha})_i, \tilde{\gamma}^i) \in \mathcal{A}} \mathcal{H}_e^{\omega}((\mathcal{E}_{\sigma})_i, (\mathcal{h}_{\alpha})_i) \leq 1 - r.
 \end{aligned}$$

In this instance, the hypothesis is contradicted as is clear from Equations (7). Therefore, the coarsest stratified svns- topogenous order on  $\vee$  is  $(\mathcal{H}_{st}^{v\mu\omega})_{\vee}$  which is finer than  $\mathcal{H}_V^{v\mu\omega}$ .  $\square$

**Theorem 6.** Let  $(\mathbb{X}, \mathcal{H}_V^{v\mu\omega})$  and  $(\mathcal{U}, (\mathcal{H}_R^{v\mu\omega})^*)$  be two svns-topogenous order spaces, if  $\psi_{\varphi} : (\vee, \mathcal{H}_V^{v\mu\omega}) \rightarrow (\mathcal{U}, (\mathcal{H}_R^{v\mu\omega})^*)$  be a svns-topogenous continuous, then  $\psi_{\varphi} : (\mathbb{X}, (\mathcal{H}_V^{v\mu\omega})_{st}) \rightarrow (\mathcal{U}, (\mathcal{H}_R^{v\mu\omega})_{st}^*)$  is a svns-topogenous continuous.

**Proof.** Assume that  $\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha} \in \widetilde{(\mathcal{U}, \mathcal{R})}$  with  $\sqcup_{j \in J}((\mathcal{E}_{\sigma})_j \cap \tilde{\mathcal{R}}^{lj}) \supseteq \mathcal{E}_{\sigma}$  and  $\sqcup_{j \in J}((\mathcal{h}_{\alpha})_j \cap \tilde{\mathcal{R}}^{lj}) \subseteq \mathcal{h}_{\alpha}$ . Then  $\sqcup_{j \in J} \psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_j \cap \tilde{\gamma}^{lj}) \supseteq \psi_{\varphi}^{-1}(\mathcal{E}_{\sigma})$  and  $\sqcup_{j \in J} \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_j \cap \tilde{\gamma}^{lj}) \subseteq \psi_{\varphi}^{-1}(\mathcal{h}_{\alpha})$ . For all collections  $\mathcal{N}(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha}) = \{B = \{((\mathcal{E}_{\sigma})_j, (\mathcal{h}_{\alpha})_j, \tilde{\mathcal{R}}^{lj})\} \mid \mathcal{E}_{\sigma} \subseteq \sqcup_{j \in J}((\mathcal{E}_{\sigma})_j \cap \tilde{\mathcal{R}}^{lj}) \text{ and } \mathcal{h}_{\alpha} \supseteq \sqcup_{j \in J}((\mathcal{h}_{\alpha})_j \cap \tilde{\mathcal{R}}^{lj})\} \forall, \mid j \in J \text{ and } \iota \in \zeta$ , we have

$$\begin{aligned}
 &(\mathcal{H}_e^v)_{st}(\psi_{\varphi}^{-1}(\mathcal{E}_{\sigma}), \psi_{\varphi}^{-1}(\mathcal{h}_{\alpha})) \\
 &\geq \bigvee_{(\psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_{j_1}), \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_{j_1}), \tilde{\gamma}^{lj_1}) \in \{\psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_j), \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_j), \tilde{\gamma}^{lj} \mid j \in J\}} \\
 &\mathcal{H}_e^v(\psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_{j_1}), \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_{j_1})) \\
 &\geq \bigvee_{((\mathcal{E}_{\sigma})_{j_1}, (\mathcal{h}_{\alpha})_{j_1}, \tilde{\mathcal{R}}^{lj_1}) \in B} (\mathcal{H}_{\varphi(e)}^v)^*((\mathcal{E}_{\sigma})_{j_1}, (\mathcal{h}_{\alpha})_{j_1}) \\
 &\geq \bigwedge_{\mathcal{N}(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha})} \left( \bigvee_{((\mathcal{E}_{\sigma})_{j_1}, (\mathcal{h}_{\alpha})_{j_1}, \tilde{\mathcal{R}}^{lj_1}) \in B} (\mathcal{H}_{\varphi(e)}^v)^*((\mathcal{E}_{\sigma})_{j_1}, (\mathcal{h}_{\alpha})_{j_1}) \right) \\
 &= (\mathcal{H}_{\varphi(e)}^v)_{st}^*(\mathcal{E}_{\sigma}, \mathcal{h}_{\alpha}) \\
 &(\mathcal{H}_e^{\mu})_{st}(\psi_{\varphi}^{-1}(\mathcal{E}_{\sigma}), \psi_{\varphi}^{-1}(\mathcal{h}_{\alpha})) \\
 &\leq \bigwedge_{(\psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_{j_1}), \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_{j_1}), \tilde{\gamma}^{lj_1}) \in \{\psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_j), \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_j), \tilde{\gamma}^{lj} \mid j \in J\}} \\
 &\mathcal{H}_e^{\mu}(\psi_{\varphi}^{-1}((\mathcal{E}_{\sigma})_{j_1}), \psi_{\varphi}^{-1}((\mathcal{h}_{\alpha})_{j_1}))
 \end{aligned}$$

$$\begin{aligned}
 &\leq \bigwedge_{((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}, \tilde{\mathcal{R}}^{j_1}) \in \mathcal{B}} (\mathcal{H}_{\varphi(e)}^\mu)^\star((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}) \\
 &\leq \bigvee_{\mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left( \bigwedge_{((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}, \tilde{\mathcal{R}}^{j_1}) \in \mathcal{B}} (\mathcal{H}_{\varphi(e)}^\mu)^\star((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}) \right) \\
 &= (\mathcal{H}_{\varphi(e)}^\mu)_{st}^\star(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \\
 &(\mathcal{H}_e^\omega)_{st} \left( \psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathfrak{h}_\alpha) \right) \\
 &\leq \bigwedge_{\left( \psi_\varphi^{-1}((\mathcal{E}_\sigma)_{j_1}), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_{j_1}), \tilde{\mathcal{E}}^{j_1} \right) \in \left\{ \psi_\varphi^{-1}((\mathcal{E}_\sigma)_j), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_j), \tilde{\mathcal{V}}^{j_1} \mid j \in J \right\}} \\
 &\mathcal{H}_e^\omega \left( \psi_\varphi^{-1}((\mathcal{E}_\sigma)_{j_1}), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_{j_1}) \right) \\
 &\leq \bigwedge_{((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}, \tilde{\mathcal{R}}^{j_1}) \in \mathcal{B}} (\mathcal{H}_{\varphi(e)}^\omega)^\star((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}) \\
 &\leq \bigvee_{\mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left( \bigwedge_{((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}, \tilde{\mathcal{R}}^{j_1}) \in \mathcal{B}} (\mathcal{H}_{\varphi(e)}^\omega)^\star((\mathcal{E}_\sigma)_{j_1}, (\mathfrak{h}_\alpha)_{j_1}) \right) \\
 &= (\mathcal{H}_{\varphi(e)}^\omega)_{st}^\star(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \quad \square
 \end{aligned}$$

**Theorem 7.** (1) If  $\mathcal{H}_V^{\nu\mu\omega}$  is symmetrical svns-topogenous order, then  $(\mathcal{H}_{st}^{\nu\mu\omega})_V$  is also symmetrical svns-topogenous order.

(2) If  $\mathcal{H}_V^{\nu\mu\omega}$  is perfect svns-topogenous order, then

(i)  $(\mathcal{T}_H^\nu)_e(\mathcal{E}_\sigma) = \mathcal{H}_e^\nu(\mathcal{E}_\sigma, \mathcal{E}_\sigma)$ ,  $(\mathcal{T}_H^\mu)_e(\mathcal{E}_\sigma) = \mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathcal{E}_\sigma)$  and  $(\mathcal{T}_H^\omega)_e(\mathcal{E}_\sigma) =$

$\mathcal{H}_e^\omega(\mathcal{E}_\sigma, \mathcal{E}_\sigma)$  is svns-topology related by  $\mathcal{H}_V^{\nu\mu\omega}$

(ii)  $(\mathcal{T}_{Hst}^{\nu\mu\omega})_V = [(\mathcal{T}_H^{\nu\mu\omega})_{st}]_V$ .

(3) If  $\mathcal{T}_V^{\nu\mu\omega}$  is a svnst, then

(i)  $(\mathcal{H}_T^\nu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = \bigvee \{ \mathcal{T}_e^\nu(\mathfrak{g}_C) \mid \mathcal{E}_\sigma \sqsubseteq \mathfrak{g}_C \sqsubseteq \mathfrak{h}_\alpha \}$ ,  $(\mathcal{H}_T^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = \bigwedge \{ \mathcal{T}_e^\mu(\mathfrak{g}_C) \mid$

$\mathcal{E}_\sigma \sqsubseteq \mathfrak{g}_C \sqsubseteq \mathfrak{h}_\alpha \}$  and  $(\mathcal{H}_T^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = \bigwedge \{ \mathcal{T}_e^\omega(\mathfrak{g}_C) \mid \mathcal{E}_\sigma \sqsubseteq \mathfrak{g}_C \sqsubseteq \mathfrak{h}_\alpha \}$  is perfect svns-topogenous order related by  $\mathcal{T}_V^{\nu\mu\omega}$ .

(ii)  $(\mathcal{H}_{Tst}^{\nu\mu\omega})_V = [(\mathcal{H}_T^{\nu\mu\omega})_{st}]_V$  is a perfect svns-topogenous order.

**Proof.** (1) We will prove that for each  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ , then

$$(\mathcal{H}_{st}^\nu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = (\mathcal{H}_{st}^\nu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \quad (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = (\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c),$$

$$(\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = (\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c).$$

Suppose that

$$(\mathcal{H}_{st}^\nu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \not\leq (\mathcal{H}_{st}^\nu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \quad (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \not\leq (\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c)$$

$$(\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \not\leq (\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c).$$

From the concept of  $(\mathcal{H}_{st}^{\nu\mu\omega})_V$ , there are  $\mathcal{A} = \left\{ (\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\mathcal{V}}^{j_1} \mid j \in J \right\} \in \mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)$  such that

$$(\mathcal{H}_{st}^\nu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \leq r \leq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^{i_1}) \in \mathcal{A}} \mathcal{H}_e^\nu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)$$

$$(\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \geq 1 - r \geq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^{i_1}) \in \mathcal{A}} \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i) \tag{8}$$

$$(\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \geq 1 - r \geq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^{i_1}) \in \mathcal{A}} \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i).$$

On another side,

$$(\mathcal{H}_{st}^\nu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \geq (\mathcal{H}_{st}^\nu)_e \left[ \prod_{j \in J} \left( (\mathfrak{h}_\alpha^c)_j \sqcup \tilde{\mathcal{V}}^{1-j_1} \right), \prod_{j \in J} \left( (\mathcal{E}_\sigma^c)_j \sqcup \tilde{\mathcal{V}}^{1-j_1} \right) \right]$$

$$\begin{aligned}
 &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{H}_{st}^v)_e \left[ (\mathfrak{h}_\alpha^c)_i \sqcup \tilde{\mathcal{V}}^{1-i}, (\mathcal{E}_\sigma^c)_i \sqcup \tilde{\mathcal{V}}^{1-i} \right] \\
 &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{H}_{st}^v)_e \left( (\mathfrak{h}_\alpha^c)_i, (\mathcal{E}_\sigma^c)_i \right) \wedge (\mathcal{H}_{st}^v)_e \left( \tilde{\mathcal{V}}^{1-i}, \tilde{\mathcal{V}}^{1-i} \right) \\
 &= \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{H}_{st}^v)_e \left( (\mathfrak{h}_\alpha^c)_i, (\mathcal{E}_\sigma^c)_i \right) \\
 &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} \mathcal{H}_e^v \left( (\mathfrak{h}_\alpha^c)_i, (\mathcal{E}_\sigma^c)_i \right) \\
 &= \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} \mathcal{H}_e^v \left( (\mathcal{E}_\sigma)_i, \mathfrak{h}_\alpha \right) \geq r.
 \end{aligned}$$

Similarly, by using an analogous line of reasoning, we can show that  $(\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \leq 1 - r$  and  $(\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \leq r - 1$  are incompatible with equations (8). Hence,

$$\begin{aligned}
 (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\leq (\mathcal{H}_{st}^v)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \quad (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \geq (\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c) \\
 (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq (\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c).
 \end{aligned} \tag{9}$$

Similarly, we can establish through a parallel argument that

$$\begin{aligned}
 (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq (\mathcal{H}_{st}^v)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \quad (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \leq (\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \\
 (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\leq (\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c).
 \end{aligned} \tag{10}$$

Based on (9) and (10), we have

$$\begin{aligned}
 (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= (\mathcal{H}_{st}^v)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \quad (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = (\mathcal{H}_{st}^\mu)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c), \\
 (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= (\mathcal{H}_{st}^\omega)_e(\mathfrak{h}_\alpha^c, \mathcal{E}_\sigma^c).
 \end{aligned}$$

(2) (i) straightforward.

(ii) Since  $(\mathcal{T}_{H_{st}}^v)_e(\tilde{\mathcal{V}}^i) = (\mathcal{H}_{st}^v)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^i) = 1$ ,  $(\mathcal{T}_{H_{st}}^\mu)_e(\tilde{\mathcal{V}}^i) = (\mathcal{H}_{st}^\mu)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^i) = 0$  and  $(\mathcal{T}_{H_{st}}^\omega)_e(\tilde{\mathcal{V}}^i) = (\mathcal{H}_{st}^\omega)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^i) = 0$  for each  $i \in \zeta$  we have  $(\mathcal{T}_{H_{st}}^{v\mu\omega})_\nu$  is stratified which is finer than  $(\mathcal{T}_H^{v\mu\omega})_\nu$ . Thus,  $(\mathcal{T}_{H_{st}}^{v\mu\omega})_\nu \sqsupseteq [(\mathcal{T}_H^{v\mu\omega})_{st}]_\nu$ .

Conversely, suppose that  $\mathcal{E}_\sigma \in \widetilde{(\mathfrak{X}, \mathcal{V})}$  such that

$$\begin{aligned}
 (\mathcal{T}_{H_{st}}^v)_e(\mathcal{E}_\sigma) &= (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathcal{E}_\sigma) \not\leq [(\mathcal{T}_H^v)_{st}]_e(\mathcal{E}_\sigma), \\
 (\mathcal{T}_{H_{st}}^\mu)_e(\mathcal{E}_\sigma) &= (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathcal{E}_\sigma) \not\geq [(\mathcal{T}_H^\mu)_{st}]_e(\mathcal{E}_\sigma), \\
 (\mathcal{T}_{H_{st}}^\omega)_e(\mathcal{E}_\sigma) &= (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathcal{E}_\sigma) \not\geq [(\mathcal{T}_H^\omega)_{st}]_e(\mathcal{E}_\sigma).
 \end{aligned} \tag{11}$$

From the concept of  $(\mathcal{H}_{st}^{v\mu\omega})_\nu$ , there exists a collection

$$\mathcal{A} = \{((\mathcal{E}_\sigma)_j, (\mathcal{E}_\sigma)_j, \tilde{\mathcal{V}}^{ij}) \in \mathcal{N}(\mathcal{E}_\sigma, \mathcal{E}_\sigma), r \in \zeta \text{ and } j \in J\}$$

such that

$$\begin{aligned}
 [(\mathcal{T}_H^v)_{st}]_e(\mathcal{E}_\sigma) \leq r &\leq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} \mathcal{H}_e^v \left( (\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i \right) \\
 &= \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{T}_H^v)_e \left( (\mathcal{E}_\sigma)_i \right), \\
 [(\mathcal{T}_H^\mu)_{st}]_e(\mathcal{E}_\sigma) \geq 1 - r &\geq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} \mathcal{H}_e^\mu \left( (\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i \right) \\
 &= \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{T}_H^\mu)_e \left( (\mathcal{E}_\sigma)_i \right), \\
 [(\mathcal{T}_H^\omega)_{st}]_e(\mathcal{E}_\sigma) \geq 1 - r &\geq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} \mathcal{H}_e^\omega \left( (\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i \right) \\
 &= \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{T}_H^\omega)_e \left( (\mathcal{E}_\sigma)_i \right).
 \end{aligned}$$

On another side,

$$\begin{aligned}
 [(\mathcal{T}_H^v)_e]_{st}(\mathcal{E}_\sigma) &= [(\mathcal{T}_H^v)_{st}]_e \left( \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j) \right) \geq \bigwedge_{j \in J} [(\mathcal{T}_H^v)_e]_{st} \left( (\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j \right) \\
 &\geq \bigwedge_{j \in J} [(\mathcal{T}_H^v)_e]_{st} \left( (\mathcal{E}_\sigma)_j \right) \wedge [(\mathcal{T}_H^v)_{st}]_e \left( \tilde{\mathcal{V}}^j \right) \\
 &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{T}_H^v)_e \left( (\mathcal{E}_\sigma)_i \right) \geq r, \\
 [(\mathcal{T}_H^\mu)_e]_{st}(\mathcal{E}_\nu) &= [(\mathcal{T}_H^\mu)_{st}]_e \left( \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^{1-j}) \right) \leq \bigvee_{j \in J} [(\mathcal{T}_H^\mu)_e]_{st} \left( (\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j \right) \\
 &\leq \bigvee_{j \in J} [(\mathcal{T}_H^\mu)_{st}]_e \left( (\mathcal{E}_\sigma)_j \right) \vee [(\mathcal{T}_H^\mu)_{st}]_e \left( \tilde{\mathcal{V}}^j \right) \\
 &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{V}}^i) \in \mathcal{A}} (\mathcal{T}_H^\mu)_e \left( (\mathcal{E}_\sigma)_i \right) \leq 1 - r, \\
 [(\mathcal{T}_H^\omega)_e]_{st}(\mathcal{E}_\sigma) &= [(\mathcal{T}_H^\omega)_{st}]_e \left( \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^{1-j}) \right) \leq \bigvee_{j \in J} [(\mathcal{T}_H^\omega)_e]_{st} \left( (\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j \right) \\
 &\leq \bigvee_{j \in J} [(\mathcal{T}_H^\omega)_{st}]_e \left( (\mathcal{E}_\sigma)_j \right) \vee [(\mathcal{T}_H^\omega)_{st}]_e \left( \tilde{\mathcal{V}}^j \right) \\
 &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{E}_\sigma)_i, \tilde{\mathcal{E}}^i) \in \mathcal{A}} (\mathcal{T}_H^\omega)_e \left( (\mathcal{E}_\sigma)_i \right) \leq 1 - r.
 \end{aligned}$$

This contradicts the hypothesis in equation (11). Thus,  $(\mathcal{T}_{H_{st}}^{v\mu\omega})_\nu = [(\mathcal{T}_H^{v\mu\omega})_{st}]_\nu$ .

(3) (i) Straightforward.

(ii) Obvious.  $\square$

**Definition 7.** A mapping  $\mathcal{Z}^v, \mathcal{Z}^\mu, \mathcal{Z}^\omega : \mathcal{Y} \rightarrow \zeta^{\widetilde{(\mathcal{Y}, \mathcal{Y})}}$  is called single-valued neutrosophic soft filter (svns-filter) on  $\mathcal{Y}$ . If the following criteria are met,  $\forall e \in \mathcal{Y}$  and  $\mathcal{E}_\sigma, \mathcal{h}_\alpha \in \widetilde{(\mathcal{Y}, \mathcal{Y})}$ :

- (Z<sub>1</sub>)  $\mathcal{Z}_e^v(\Phi) = 0, \mathcal{Z}_e^\mu(\Phi) = 1, \mathcal{Z}_e^\omega(\Phi) = 1$  and  $\mathcal{Z}_e^v(\tilde{\mathcal{V}}) = 1, \mathcal{Z}_e^\mu(\tilde{\mathcal{V}}) = 0, \mathcal{Z}_e^\omega(\tilde{\mathcal{V}}) = 0,$
- (Z<sub>2</sub>)  $\mathcal{Z}_e^v(\mathcal{E}_\sigma \cap \mathcal{h}_\alpha) \geq \mathcal{Z}_e^v(\mathcal{E}_\sigma) \wedge \mathcal{Z}_e^v(\mathcal{h}_\alpha), \quad \mathcal{Z}_e^\mu(\mathcal{E}_\sigma \cap \mathcal{h}_\alpha) \leq \mathcal{Z}_e^\mu(\mathcal{E}_\sigma) \vee \mathcal{Z}_e^\mu(\mathcal{h}_\alpha),$   
 $\mathcal{Z}_e^\omega(\mathcal{E}_\sigma \cap \mathcal{h}_\alpha) \leq \mathcal{Z}_e^\omega(\mathcal{E}_\sigma) \vee \mathcal{Z}_e^\omega(\mathcal{h}_\alpha),$
- (Z<sub>3</sub>) If  $\mathcal{E}_\sigma \sqsubseteq \mathcal{h}_\beta,$  then  $\mathcal{Z}_e^v(\mathcal{E}_\sigma) \leq \mathcal{Z}_e^v(\mathcal{h}_\alpha), \mathcal{Z}_e^\mu(\mathcal{E}_\sigma) \geq \mathcal{Z}_e^\mu(\mathcal{h}_\alpha)$   
 and  $\mathcal{Z}_e^\omega(\mathcal{E}_\sigma) \geq \mathcal{Z}_e^\omega(\mathcal{h}_\alpha).$

The svns-filter  $\mathcal{Z}^{v\mu\omega}$  is called stratified iff the following condition is met.

(Z<sub>S</sub>) For every  $e \in \mathcal{Y}, i \in \zeta$  and  $\mathcal{E}_\sigma \in \widetilde{(\mathcal{Y}, \mathcal{Y})}, \mathcal{Z}_e^v(\mathcal{E}_\sigma \cap \tilde{\mathcal{V}}^i) \geq \mathcal{Z}_e^v(\mathcal{E}_\sigma) \wedge (i), \mathcal{Z}_e^\mu(\mathcal{E}_\sigma \cap \tilde{\mathcal{V}}^i) \leq \mathcal{Z}_e^\mu(\mathcal{E}_\sigma) \vee (i),$  and  $\mathcal{Z}_e^\omega(\mathcal{E}_\sigma \cap \tilde{\mathcal{V}}^i) \leq \mathcal{Z}_e^\omega(\mathcal{E}_\sigma) \vee (i).$

The pair  $(\mathcal{Y}, \mathcal{Z}_\nu^{v\mu\omega})$  is said to be stratified svns-filters space.

If  $\mathcal{Z}_\nu^{v\mu\omega}$  and  $\mathcal{Z}_\nu^{*v\mu\omega}$  are svns-filters on  $\mathcal{Y},$  then  $\mathcal{Z}_\nu^{\pi\alpha\sigma}$  is finer than  $\mathcal{Z}_\nu^{*v\mu\omega}$  or  $\mathcal{Z}_\nu^{*v\mu\omega}$  is coarser than  $\mathcal{Z}_\nu^{v\mu\omega}$  indicated by  $\mathcal{Z}_\nu^{v\mu\omega} \sqsupseteq \mathcal{Z}_\nu^{*v\mu\omega}$  if  $\mathcal{Z}_e^v(\mathcal{E}_\sigma) \geq \mathcal{Z}_e^{*v}(\mathcal{E}_\sigma), \mathcal{Z}_e^\mu(\mathcal{E}_\sigma) \leq \mathcal{Z}_e^{*\mu}(\mathcal{E}_\sigma)$  and  $\mathcal{Z}_e^\omega(\mathcal{E}_\sigma) \leq \mathcal{Z}_e^{*\omega}(\mathcal{E}_\sigma).$

**Theorem 8.** A consider that  $(\mathcal{Y}, \mathcal{Z}_\nu^{v\mu\omega})$  is svns-filters space. Define the mapping  $\mathcal{Z}_{st}^v : \mathcal{Y} \rightarrow \zeta^{\widetilde{(\mathcal{Y}, \mathcal{Y})}}, \mathcal{Z}_{st}^\mu : \mathcal{Y} \rightarrow \zeta^{\widetilde{(\mathcal{Y}, \mathcal{Y})}}, \mathcal{Z}_{st}^\omega : \mathcal{Y} \rightarrow \zeta^{\widetilde{(\mathcal{Y}, \mathcal{Y})}}$  as next:  
 $\forall \mathcal{E}_\sigma \in \widetilde{(\mathcal{Y}, \mathcal{Y})}, e \in \mathcal{Y}$

$$\begin{aligned}
 \bigvee \left[ \bigwedge_{j \in J} \mathcal{Z}_e^v((\mathcal{E}_\sigma)_j) \wedge I_j \mid \sqcup_{i \in J} ((\mathcal{E}_\sigma)_i \cap \tilde{\mathcal{V}}^i) \sqsubseteq \mathcal{E}_\sigma \right] &= (\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma), \\
 \bigwedge \left[ \bigvee_{j \in J} \mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j) \vee I_j \mid \sqcup_{i \in J} ((\mathcal{E}_\sigma)_i \cap \tilde{\mathcal{V}}^i) \sqsubseteq \mathcal{E}_\sigma \right] &= (\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma), \\
 \bigwedge \left[ \bigvee_{j \in J} \mathcal{Z}_e^\omega((\mathcal{E}_\sigma)_j) \vee I_j \mid \sqcup_{i \in J} ((\mathcal{E}_\sigma)_i \cap \tilde{\mathcal{V}}^i) \sqsubseteq \mathcal{E}_\sigma \right] &= (\mathcal{Z}_{st}^\omega)_e(\mathcal{E}_\sigma),
 \end{aligned}$$

where  $\bigvee$  and  $\bigwedge$  are taken over all collections  $\{((\mathcal{E}_\sigma)_j, \tilde{\mathcal{V}}^j)\}$  for each  $j \in J$  and  $J$  is finite) with  $\mathcal{E}_\sigma \sqsupseteq \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j).$  Then  $(\mathcal{Z}_{st}^{v\mu\omega})_\nu$  is the coarsest stratified svns-filter on  $\mathcal{Y}$  which is finer than  $\mathcal{Z}_\nu^{v\mu\omega}.$  Also,  $(\mathcal{Z}_{st}^{v\mu\omega})_\nu$  is the stratification of a svns-filter  $\mathcal{Z}_\nu^{v\mu\omega}$  on  $\mathcal{Y}.$

**Proof.** At initial, we will prove that  $(\mathcal{Z}_{st}^{v\mu\omega})_\nu$  is stratified svns-filter:

(Z<sub>1</sub>) For any  $\iota \in \zeta$ , there are collections  $\{\tilde{\gamma}\}$  and  $\{\Phi\}$  with  $\tilde{\gamma}^\iota = \tilde{\gamma}^\iota \cap \tilde{\gamma}$ . We have

$$\begin{aligned} (\mathcal{Z}_{st}^v)_e(\tilde{\gamma}) &\geq \mathcal{Z}_e^v(\tilde{\gamma}) \wedge 1 = 1, & (\mathcal{Z}_{st}^\mu)_e(\tilde{\gamma}) &\leq \mathcal{Z}_e^\mu(\tilde{\gamma}) \vee 0 = 0, \\ (\mathcal{Z}_v^\omega)_{st}(\tilde{\gamma}) &\leq \mathcal{Z}_v^\omega(\tilde{\gamma}) \vee 0 = 0, \\ (\mathcal{Z}_{st}^v)_e(\tilde{\Phi}) &\geq \mathcal{Z}_e^v(\tilde{\Phi}) \wedge 0 = 0, & (\mathcal{Z}_{st}^\mu)_e(\tilde{\Phi}) &\leq \mathcal{Z}_e^\mu(\tilde{\Phi}) \vee 1 = 1, \\ (\mathcal{Z}_\Phi^\omega)_{st}(\tilde{\Phi}) &\leq \mathcal{Z}_v^\omega(\tilde{\Phi}) \vee 1 = 1. \end{aligned}$$

Hence,  $(\mathcal{Z}_{st}^v)_e(\tilde{\gamma}) = 1$ ,  $(\mathcal{Z}_{st}^\mu)_e(\tilde{\gamma}) = 0$ ,  $(\mathcal{Z}_v^\omega)_{st}(\tilde{\gamma}) = 0$  and  $(\mathcal{Z}_{st}^v)_e(\Phi) = 0$ ,  $(\mathcal{Z}_{st}^\mu)_e(\Phi) = 1$ ,  $(\mathcal{Z}_v^\omega)_{st}(\Phi) = 1$ .

(Z<sub>2</sub>) Let  $\mathcal{D}(\mathcal{E}_\sigma) = \{ \{((\mathcal{E}_\sigma)_j, \tilde{\gamma}^j)\} \mid \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\gamma}^j) \sqsubseteq \mathcal{E}_\sigma \}$  and  $\mathcal{D}(\mathcal{h}_\alpha) = \{ \{((\mathcal{h}_\alpha)_i, \tilde{\gamma}^i)\} \mid \sqcup_{i \in \Gamma} ((\mathcal{h}_\alpha)_i \cap \tilde{\gamma}^i) \sqsubseteq \mathcal{h}_\alpha \}$  for all  $j \in J$  and  $i \in \Gamma$ . Then,

$$\begin{aligned} \mathcal{E}_\sigma \cap \mathcal{h}_\alpha &\sqsupseteq (\sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\gamma}^j)) \cap (\sqcup_{i \in \Gamma} ((\mathcal{h}_\alpha)_i \cap \tilde{\gamma}^i)) \\ &= \sqcup_{j \in J} \sqcup_{i \in \Gamma} (((\mathcal{E}_\sigma)_j \cap \tilde{\gamma}^j) \cap ((\mathcal{h}_\alpha)_i \cap \tilde{\gamma}^i)) \\ &= \sqcup_{j \in J} \sqcup_{i \in \Gamma} (((\mathcal{E}_\sigma)_j \cap (\mathcal{h}_\alpha)_i) \cap (\tilde{\gamma}^j \cap \tilde{\gamma}^i)) \\ &= \sqcup_{l \in J \cup \Gamma} ((\mathbf{g}_c)_l \cap \tilde{\gamma}^{w_l}), \end{aligned}$$

where  $\tilde{\gamma}^{w_l} = \tilde{\gamma}^j \cap \tilde{\gamma}^i$ ,  $w_l = l_j \cap l_i$  and  $(\mathbf{g}_c)_l = (\mathcal{E}_\sigma)_j \cap (\mathcal{h}_\alpha)_i$ , which implies

$$\begin{aligned} (\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma \cap \mathcal{h}_\alpha) &\geq \bigwedge_{j \in J} (\mathcal{Z}_e^v((\mathcal{E}_\sigma)_j \cap (\mathcal{h}_\alpha)_i)) \wedge w_l \\ &\geq \bigwedge_j (\mathcal{Z}_e^v((\mathcal{E}_\sigma)_j)) \wedge \bigwedge_i (\mathcal{Z}_e^v((\mathcal{h}_\alpha)_i)) \wedge (l_j \wedge l_i) \\ &\geq \left[ \bigwedge_j (\mathcal{Z}_e^v((\mathcal{E}_\sigma)_j) \wedge l_j) \right] \wedge \left[ \bigwedge_i (\mathcal{Z}_e^v((\mathcal{h}_\alpha)_i) \wedge l_i) \right] \\ &\geq (\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma) \wedge (\mathcal{Z}_{st}^v)_e(\mathcal{h}_\alpha), \\ (\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma \cap \mathcal{h}_\alpha) &\leq \bigvee_{j \in J} (\mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j \cap (\mathcal{h}_\alpha)_i)) \vee w_l \\ &\leq \bigvee_j (\mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j)) \vee \bigvee_i (\mathcal{Z}_e^\mu((\mathcal{h}_\alpha)_i)) \vee (l_j \wedge l_i) \\ &\leq \left[ \bigvee_j (\mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j) \vee l_j) \right] \vee \left[ \bigvee_i (\mathcal{Z}_e^\mu((\mathcal{h}_\alpha)_i) \vee l_i) \right] \\ &\geq (\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma) \vee (\mathcal{Z}_{st}^\mu)_e(\mathcal{h}_\alpha), \\ (\mathcal{Z}_{st}^\omega)_{st}(\mathcal{E}_\sigma \cap \mathcal{h}_\alpha) &\leq \bigvee_{j \in J} (\mathcal{Z}_e^{\omega}((\mathcal{E}_\sigma)_j \cap (\mathcal{h}_\alpha)_i)) \vee w_l \\ &\leq \left[ \bigvee_j (\mathcal{Z}_e^{\omega}((\mathcal{E}_\sigma)_j)) \right] \vee \left[ \bigvee_i (\mathcal{Z}_e^{\omega}((\mathcal{h}_\alpha)_i)) \vee (l_j \wedge l_i) \right] \\ &\leq \left[ \bigvee_j (\mathcal{Z}_e^{\omega}((\mathcal{E}_\sigma)_j) \vee l_j) \right] \vee \left[ \bigvee_i (\mathcal{Z}_e^{\omega}((\mathcal{h}_\alpha)_i) \vee l_i) \right] \\ &\geq (\mathcal{Z}_{st}^\omega)_{st}(\mathcal{E}_\sigma) \vee (\mathcal{Z}_{st}^\omega)_{st}(\mathcal{h}_\alpha). \end{aligned}$$

Thus, the proof of (Z<sub>2</sub>) is complete.

(Z<sub>3</sub>) Straightforward.

(Z<sub>5</sub>) Suppose

$$\begin{aligned} \mathcal{D}(\mathcal{E}_\sigma) &= \{ \{((\mathcal{E}_\sigma)_j, \tilde{\gamma}^j) \mid \mathcal{E}_\sigma \sqsupseteq \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\gamma}^j)\}, \text{ for all } j \in J, \\ (\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma \cap \tilde{\gamma}^\iota) &\geq \left[ \bigwedge_{j \in J} (\mathcal{Z}_e^v((\mathcal{E}_\sigma)_j) \wedge \tilde{\gamma}^\iota) \wedge l_j \right] \geq \left[ \bigwedge_{j \in J} (\mathcal{Z}_e^v((\mathcal{E}_\sigma)_j) \wedge \iota) \wedge l_j \right] \\ &\geq \left[ \bigwedge_{j \in J} \mathcal{Z}_e^v((\mathcal{E}_\sigma)_j) \wedge l_j \right] \wedge \iota \geq (\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma) \wedge \iota, \end{aligned}$$



$$\begin{aligned} (\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma \cap \tilde{\mathcal{V}}) &\leq \left[ \bigvee_j (\mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j) \vee \tilde{\mathcal{V}}^j) \vee \iota_j \right] \leq \left[ \bigvee_j (\mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j) \vee \iota) \vee \iota_j \right] \\ &\leq \left[ \bigvee_j \mathcal{Z}_e^\mu((\mathcal{E}_\sigma)_j) \vee \iota_j \right] \vee \iota \leq (\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma) \vee \iota. \end{aligned}$$

Similarly, by using an analogous line of reasoning, we can show that  $(\mathcal{Z}_{st}^\omega)_e(\mathcal{E}_\sigma \cap \tilde{\mathcal{V}}) \leq (\mathcal{Z}_{st}^\omega)_e(\mathcal{E}_\sigma) \vee \iota$ . Thus, the proof of  $(\mathcal{Z}_S)$  is complete.

Secondly, for every  $\mathcal{E}_\sigma \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ,  $e \in \mathfrak{V}$ , there exist collections  $\{\tilde{\mathcal{V}}\}$  with  $\mathcal{E}_\sigma = \mathcal{E}_A \cap \tilde{\mathcal{V}}$ . Then,  $(\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma) \geq \mathcal{Z}_e^v(\mathcal{E}_\sigma)$ ,  $(\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma) \leq \mathcal{Z}_e^\mu(\mathcal{E}_\sigma)$  and  $(\mathcal{Z}_{st}^\omega)_e(\mathcal{E}_\sigma) \leq \mathcal{Z}_e^\omega(\mathcal{E}_\sigma)$ . Thus,  $(\mathcal{Z}_{st}^{v\mu\omega})_\mathfrak{V}$  is finer than  $\mathcal{Z}_\mathfrak{V}^{v\mu\omega}$ .

Finally, let  $(\mathcal{Z}_{*st}^v, \mathcal{Z}_{*st}^\mu, \mathcal{Z}_{*st}^\omega)_\mathfrak{V}$  be stratified snvs-filter which is finer than  $(\mathcal{Z}^v, \mathcal{Z}^\mu, \mathcal{Z}^\omega)_\mathfrak{V}$  on  $\mathfrak{Y}$  and

$$\mathcal{D}(\mathcal{E}_\sigma) = \{ \{ (\mathcal{E}_\sigma)_j, \tilde{\mathcal{V}}^j \} \mid \mathcal{E}_\sigma \supseteq \bigcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j), \forall j \in J, \}$$

then we have,

$$\begin{aligned} (\mathcal{Z}_{*st}^v)_e(\mathcal{E}_\sigma) &\geq (\mathcal{Z}_{*st}^v)_e(\bigcup_{j \in J} ((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j)) \geq \bigwedge_{j \in J} (\mathcal{Z}_{*st}^v)_e((\mathcal{E}_\sigma)_j \cap \tilde{\mathcal{V}}^j) \\ &\geq \bigwedge_{j \in J} ((\mathcal{Z}_{*st}^v)_e((\mathcal{E}_\sigma)_j) \wedge \iota_j) \\ &\geq \bigwedge_{(\mathcal{E}_\sigma)_k, \tilde{\mathcal{V}}^k \in \{ (\mathcal{E}_\sigma)_j, \tilde{\mathcal{V}}^j \mid j \in J \}} \mathcal{Z}_e^v((\mathcal{E}_\sigma)_p \wedge \iota_k) \geq (\mathcal{Z}_{st}^v)_e(\mathcal{E}_\sigma). \end{aligned}$$

Likewise, using related reasoning, we can determine that  $(\mathcal{Z}_{*st}^\mu)_e(\mathcal{E}_\sigma) \leq (\mathcal{Z}_{st}^\mu)_e(\mathcal{E}_\sigma)$  and  $(\mathcal{Z}_{*st}^\omega)_e(\mathcal{E}_\sigma) \leq (\mathcal{Z}_{st}^\omega)_e(\mathcal{E}_\sigma)$ . Hence  $(\mathcal{Z}_{st}^{v\mu\omega})_\mathfrak{V}$  is the coarsest stratified snvs-filter finer than  $\mathcal{Z}_\mathfrak{V}^{v\mu\omega}$ .  $\square$

**Theorem 9.** Let  $\Theta(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  and  $\Omega(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  be collections of all snvs-filters and, snvs-topogenous correspondingly. Define,  $\forall, \mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ,  $e \in \mathfrak{V}$ ,

$$\begin{aligned} \mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) &= \bigvee_{\mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^v(\mathfrak{h}_\alpha, \mathcal{E}_\sigma) \wedge \mathcal{Z}_e^v(\mathcal{E}_\sigma) \} \\ \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) &= \bigwedge_{\mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^\mu(\mathfrak{h}_\alpha, \mathcal{E}_\sigma) \vee \mathcal{Z}_e^\mu(\mathcal{E}_\sigma) \} \\ \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) &= \bigwedge_{\mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^\omega(\mathfrak{h}_\alpha, \mathcal{E}_\sigma) \vee \mathcal{Z}_e^\omega(\mathcal{E}_\sigma) \} \end{aligned}$$

where  $\mathcal{H}_\mathfrak{V}^{v\mu\omega} \in \Omega(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$  and  $\mathcal{Z}_\mathfrak{V}^{v\mu\omega} \in \Theta(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ . Then:

- (1)  $\mathcal{O}_\mathfrak{V}^{v\mu\omega}(\mathcal{H}, \mathcal{Z}) \in \Theta(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ .
- (2)  $\mathcal{O}_\mathfrak{V}^{v\mu\omega}(\mathcal{H}, \mathcal{Z}) \sqsubseteq \mathcal{Z}_\mathfrak{V}^{v\mu\omega}$  for all  $\mathcal{Z}_\mathfrak{E}^{v\mu\omega} \in \Theta(\widetilde{(\mathfrak{X}, \mathfrak{Y})})$ .
- (3)  $\mathcal{O}_\mathfrak{V}^{v\mu\omega}(\mathcal{H}, \mathcal{H}_{\mathcal{E}_\sigma}) = (\mathcal{H}_{\mathcal{E}_\sigma}^{v\mu\omega})_\mathfrak{V}$ .
- (4)  $\mathcal{O}_\mathfrak{V}^{v\mu\omega}(\mathcal{H}_{st}, \mathcal{Z}_{st}) = [(\mathcal{O}_{st}^{v\mu\omega})_\mathfrak{V}](\mathcal{H}, \mathcal{Z})$ .

**Proof.**  $(\mathcal{Z}_1)$  Since  $\mathcal{Z}_e^v(\Phi) = 0, \mathcal{Z}_e^\mu(\Phi) = 1, \mathcal{Z}_e^\omega(\Phi) = 1$  and  $\mathcal{Z}_e^v(\tilde{\mathcal{V}}) = 1, \mathcal{Z}_e^\mu(\tilde{\mathcal{V}}) = 0, \mathcal{Z}_e^\omega(\tilde{\mathcal{V}}) = 0$ , we obtain;  $\forall e \in \mathfrak{V}$ ,

$$\begin{aligned} \mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\Phi) &= \bigvee_{\mathfrak{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^v(\mathfrak{g}_C, \Phi) \wedge \mathcal{Z}_e^v(\Phi) \}, \\ \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\Phi) &= \bigwedge_{\mathfrak{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^\mu(\mathfrak{g}_C, \Phi) \vee \mathcal{Z}_e^\mu(\Phi) \}, \\ \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\Phi) &= \bigwedge_{\mathfrak{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^\omega(\mathfrak{g}_C, \Phi) \vee \mathcal{Z}_e^\omega(\Phi) \}, \end{aligned}$$

thus,  $\mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\Phi) = 0, \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\Phi) = 1, \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\Phi) = 1$  and  $\mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\tilde{\mathcal{V}}) = 1, \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\tilde{\mathcal{V}}) = 0, \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\tilde{\mathcal{V}}) = 0$ .

$(\mathcal{Z}_2)$  Let  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ . Then we obtain

$$\mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma \cap \mathfrak{h}_\alpha) = \bigvee_{\mathfrak{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \{ \mathcal{H}_e^v(\mathfrak{g}_C, \mathcal{E}_\sigma \cap \mathfrak{h}_\alpha) \wedge \mathcal{Z}_e^v(\mathcal{E}_\sigma \cap \mathfrak{h}_\alpha) \}$$

$$\begin{aligned}
 &\geq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{(\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{E}_\sigma) \wedge \mathcal{H}_e^v(\mathbf{g}_C, \mathcal{h}_\alpha)) \wedge (\mathcal{Z}_e^v(\mathcal{E}_\sigma) \wedge \mathcal{Z}_e^v(\mathcal{h}_\alpha))\} \\
 &\geq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{(\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{E}_\sigma) \wedge \mathcal{Z}_e^v(\mathcal{E}_\sigma))\} \wedge \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{(\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{h}_\alpha) \wedge \mathcal{Z}_e^v(\mathcal{h}_\alpha))\} \\
 &\geq \mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) \wedge \mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\mathcal{h}_\alpha), \\
 \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma \sqcap \mathcal{h}_\alpha) &= \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{E}_\sigma \sqcap \mathcal{h}_\alpha) \vee \mathcal{Z}_e^\mu(\mathcal{E}_\sigma \sqcap \mathcal{h}_\alpha)\} \\
 &\leq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{(\mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{E}_\sigma) \vee \mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{h}_\alpha)) \vee (\mathcal{Z}_e^\mu(\mathcal{E}_\sigma) \vee \mathcal{Z}_e^\mu(\mathcal{h}_\alpha))\} \\
 &\leq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{(\mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{E}_\sigma) \vee \mathcal{Z}_e^\mu(\mathcal{E}_\sigma))\} \vee \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{(\mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{h}_\alpha) \vee \mathcal{Z}_e^\mu(\mathcal{h}_\alpha))\} \\
 &\leq \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) \vee \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\mathcal{h}_\alpha).
 \end{aligned}$$

Likewise, using related reasoning, we can determine that

$$\mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma \sqcap \mathcal{h}_\alpha) \leq \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) \vee \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\mathcal{h}_\alpha).$$

(Z<sub>3</sub>) If  $\mathcal{E}_\sigma \sqsubseteq \mathcal{h}_\alpha$ , then;

$$\begin{aligned}
 \mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) &= \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{E}_\sigma) \wedge \mathcal{Z}_e^v(\mathcal{E}_\sigma)\} \\
 &\leq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{h}_\alpha) \wedge \mathcal{Z}_e^v(\mathcal{h}_\alpha)\} = \mathcal{O}_e^v(\mathcal{H}, \mathcal{Z})(\mathcal{h}_\alpha), \\
 \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) &= \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{E}_\sigma) \vee \mathcal{Z}_e^\mu(\mathcal{E}_\sigma)\} \\
 &\geq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^\mu(\mathbf{g}_C, \mathcal{h}_\alpha) \vee \mathcal{Z}_e^\mu(\mathcal{h}_\alpha)\} = \mathcal{O}_e^\mu(\mathcal{H}, \mathcal{Z})(\mathcal{h}_\alpha), \\
 \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\mathcal{E}_\sigma) &= \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^\omega(\mathbf{g}_C, \mathcal{E}_\sigma) \vee \mathcal{Z}_e^\omega(\mathcal{E}_\sigma)\} \\
 &\geq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^\omega(\mathbf{g}_C, \mathcal{h}_\alpha) \vee \mathcal{Z}_e^\omega(\mathcal{h}_\alpha)\} = \mathcal{O}_e^\omega(\mathcal{H}, \mathcal{Z})(\mathcal{h}_\alpha).
 \end{aligned}$$

(2) It is obvious from the definition.

(3) From (2), we obtain  $\mathcal{O}_v^{\nu\mu\omega}(\mathcal{H}, \mathcal{H}_{\mathcal{E}_\sigma}) \sqsubseteq (\mathcal{H}_{\mathcal{E}_\sigma}^{\nu\mu\omega})_E$ . Now we just need to prove that  $\mathcal{O}_v^{\nu\mu\omega}(\mathcal{H}, \mathcal{H}_{\mathcal{E}_\sigma}) \sqsupseteq (\mathcal{H}_{\mathcal{E}_\sigma}^{\nu\mu\omega})_V$ . Let  $\Phi \neq \mathcal{h}_\alpha \in \widetilde{(\mathbb{Y}, \mathbb{Y})}$ .

Then we obtain

$$\begin{aligned}
 \mathcal{O}_v^v(\mathcal{H}, \mathcal{H}_{\mathcal{E}_\sigma})(\mathcal{h}_\alpha) &= \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{h}_\alpha) \wedge (\mathcal{H}_{\mathcal{E}_\sigma}^v)_e(\mathcal{h}_\alpha)\} \\
 &= \bigvee_{\mathbf{g}_C \in \widetilde{(\mathbb{Y}, \mathbb{V})}} \{\mathcal{H}_e^v(\mathbf{g}_C, \mathcal{h}_\alpha) \wedge \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{h}_\alpha)\} \\
 &\geq \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{h}_\alpha) \wedge \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{h}_\alpha) = \mathcal{H}_e^v(\mathcal{E}_\sigma, \mathcal{h}_\alpha) = (\mathcal{H}_{\mathcal{E}_\sigma}^v)_e(\mathcal{h}_\alpha).
 \end{aligned}$$

Similarly, we can establish through a parallel argument that

$$\mathcal{O}_v^\mu(\mathcal{H}, \mathcal{H}_{\mathcal{E}_\sigma})(\mathcal{h}_\alpha) \geq (\mathcal{H}_{\mathcal{E}_\sigma}^\mu)_e(\mathcal{h}_\alpha), \quad \mathcal{O}_v^\omega(\mathcal{H}, \mathcal{H}_{\mathcal{E}_\sigma})(\mathcal{h}_\alpha) \geq (\mathcal{H}_{\mathcal{E}_\sigma}^\omega)_e(\mathcal{h}_\alpha)$$

(4) Theorems 5 and 8 provide a clear and simple explanation.  $\square$

#### 4. Stratified single-valued neutrosophic soft quasi-proximity

**Definition 8.** A mapping  $\mathcal{Q}^v, \mathcal{Q}^\mu, \mathcal{Q}^\omega : \mathbb{V} \rightarrow \zeta^{\widetilde{(\mathbb{Y}, \mathbb{V})} \times \widetilde{(\mathbb{Y}, \mathbb{V})}}$  is said to be *svnsq-proximity* on  $\mathbb{Y}$  if the following criteria are met,  $\forall e \in \mathbb{V}$ ,  $\mathcal{E}_A, \mathcal{h}_B \in \widetilde{(\mathbb{Y}, \mathbb{V})}$ :

$$(\mathcal{Q}_1) \mathcal{Q}_e^v(\tilde{\mathbb{Y}}, \Phi) = \mathcal{Q}_e^v(\Phi, \tilde{\mathbb{Y}}) = 0, \quad \mathcal{Q}_e^\mu(\tilde{\mathbb{Y}}, \Phi) = \mathcal{Q}_e^\mu(\Phi, \tilde{\mathbb{Y}}) = 1, \quad \mathcal{Q}_e^\omega(\tilde{\mathbb{Y}}, \Phi) = \mathcal{Q}_e^\omega(\Phi, \tilde{\mathbb{Y}}) = 1.$$

- (Q<sub>2</sub>) If  $Q_e^v(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \neq 1$ ,  $Q_e^\mu(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \neq 0$  and  $Q_e^\omega(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \neq 0$ , then  $\mathcal{F}_\sigma \sqsubseteq \mathcal{H}_\alpha^c$ .
- (Q<sub>3</sub>) If  $\mathcal{F}_\sigma \sqsubseteq \mathcal{H}_\alpha$ , then  $Q_e^v(\mathcal{F}_\sigma, \mathbf{g}_C) \leq Q_e^v(\mathcal{H}_\alpha, \mathbf{g}_C)$ ,  $Q_e^\mu(\mathcal{F}_\sigma, \mathbf{g}_C) \geq Q_e^\mu(\mathcal{H}_\alpha, \mathbf{g}_C)$ ,  
 $Q_e^\omega(\mathcal{F}_\sigma, \mathbf{g}_C) \geq Q_e^\omega(\mathcal{H}_\alpha, \mathbf{g}_C)$  for any  $\mathcal{F}_\sigma, \mathcal{H}_\alpha, \mathbf{g}_C \in \overline{(\mathbb{X}, \mathcal{Y})}$ .
- (Q<sub>4</sub>)  $Q_e^v((\mathcal{F}_\sigma)_1 \sqcap (\mathcal{F}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcup (\mathcal{H}_\alpha)_2) \leq Q_e^v((\mathcal{F}_\sigma)_1 \sqcap (\mathcal{H}_\alpha)_1) \wedge Q_e^v((\mathcal{F}_\sigma)_2 \sqcup (\mathcal{H}_\alpha)_2)$ ,  
 $Q_e^\mu((\mathcal{F}_\sigma)_1 \sqcap (\mathcal{F}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcup (\mathcal{H}_\alpha)_2) \geq Q_e^\mu((\mathcal{F}_\sigma)_1 \sqcap (\mathcal{H}_\alpha)_1) \vee Q_e^\mu((\mathcal{F}_\sigma)_2 \sqcup (\mathcal{H}_\alpha)_2)$ ,  
 $Q_e^\omega((\mathcal{F}_\sigma)_1 \sqcap (\mathcal{F}_\sigma)_2, (\mathcal{H}_\alpha)_1 \sqcup (\mathcal{H}_\alpha)_2) \geq Q_e^\omega((\mathcal{F}_\sigma)_1 \sqcap (\mathcal{H}_\alpha)_1) \vee Q_e^\omega((\mathcal{F}_\sigma)_2 \sqcup (\mathcal{H}_\alpha)_2)$ .
- (Q<sub>5</sub>)  $Q_e^v(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \geq \bigwedge_{\mathbf{g}_C \in \overline{(\mathbb{X}, \mathcal{Y})}} [Q_e^v(\mathcal{F}_\sigma, \mathbf{g}_C) \vee Q_e^v(\mathbf{g}_C^c, \mathcal{H}_\alpha)]$ ,  
 $Q_e^\mu(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \leq \bigvee_{\mathbf{g}_C \in \overline{(\mathbb{X}, \mathcal{Y})}} [Q_e^\mu(\mathcal{F}_\sigma, \mathbf{g}_C) \wedge Q_e^\mu(\mathbf{g}_C^c, \mathcal{H}_\alpha)]$ ,  
 $Q_e^\omega(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \leq \bigvee_{\mathbf{g}_C \in \overline{(\mathbb{X}, \mathcal{Y})}} [Q_e^\omega(\mathcal{F}_\sigma, \mathbf{g}_C) \wedge Q_e^\omega(\mathbf{g}_C^c, \mathcal{H}_\alpha)]$ .

The *svnsq-proximity*  $Q_V^{v\mu\omega}$  is called stratified iff the following condition is met.

(Q<sub>S</sub>)  $Q_e^v(\tilde{\mathcal{V}}^l, \tilde{\mathcal{V}}^{1-l}) = 0$ ,  $Q_e^\mu(\tilde{\mathcal{V}}^l, \tilde{\mathcal{V}}^{1-l}) = Q_e^\omega(\tilde{\mathcal{V}}^l, \tilde{\mathcal{V}}^{1-l}) = 1$ ,  $\forall l \in \zeta$ .

In this case, a pair  $(\mathbb{X}, Q_V^{v\mu\omega})$  is said to be *stratified svnsq-proximity space*.

Let  $(Q_V^{v\mu\omega})_1$  and  $(Q_V^{v\mu\omega})_2$  be *svnsq-proximity* on  $\mathcal{Y}$ . We say  $(Q_V^{v\mu\omega})_1$  is finer than  $(Q_V^{v\mu\omega})_2$  [ $(Q_V^{v\mu\omega})_2$  is coarser than  $(Q_V^{v\mu\omega})_1$ ] if  $(Q_e^v)_2(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \geq (Q_e^v)_1(\mathcal{F}_\sigma, \mathcal{H}_\alpha)$ ,  $(Q_e^\mu)_2(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \leq (Q_e^\mu)_1(\mathcal{F}_\sigma, \mathcal{H}_\alpha)$ ,  $(Q_e^\omega)_2(\mathcal{F}_\sigma, \mathcal{H}_\alpha) \leq (Q_e^\omega)_1(\mathcal{F}_\sigma, \mathcal{H}_\alpha)$ .

**Definition 9.** Let  $(\mathbb{X}, (Q_V^{v\mu\omega})_1)$  and  $(\mathcal{U}, (Q_R^{v\mu\omega})_2)$  be *svnsq-proximity spaces*. A mapping  $\psi_\varphi : (\mathbb{X}, (Q_V^{v\mu\omega})_1) \rightarrow (\mathcal{U}, (Q_R^{v\mu\omega})_2)$  is called *svnsq-proximity continuous* iff

$$(Q_e^v)_1(\psi_\varphi^{-1}(\mathcal{H}_\alpha), \psi_\varphi^{-1}(\mathbf{g}_C)) \geq (Q_{\varphi(e)}^v)_2(\mathcal{H}_\alpha, \mathbf{g}_C),$$

$$(Q_e^\mu)_1(\psi_\varphi^{-1}(\mathcal{H}_\alpha), \psi_\varphi^{-1}(\mathbf{g}_C)) \leq (Q_{\varphi(e)}^\mu)_2(\mathcal{H}_\alpha, \mathbf{g}_C)$$

$$(Q_e^\omega)_1(\psi_\varphi^{-1}(\mathcal{H}_\alpha), \psi_\varphi^{-1}(\mathbf{g}_C)) \leq (Q_{\varphi(e)}^\omega)_2(\mathcal{H}_\alpha, \mathbf{g}_C),$$

for any  $\mathbf{g}_C, \mathcal{H}_\alpha \in \overline{(\mathcal{U}, \mathcal{R})}$ ,  $e \in \mathcal{Y}$ .

**Theorem 10.** Let  $(\mathbb{X}, Q_V^{v\mu\omega})$  be *svnsq-proximity space*. For each  $e \in \mathcal{Y}$ , define

$$(Q_{st}^v)_e(\mathcal{F}_\sigma, \mathcal{H}_\alpha) = \bigwedge_{\{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{V}}^{lj} \mid j \in J\} \in D(\mathcal{F}_\sigma, \mathcal{H}_\alpha)}$$

$$\left\{ \bigvee_{\{(\mathcal{F}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathcal{V}}^{li} \} \in \{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{V}}^{lj} \mid j \in J\}} Q_e^v((\mathcal{F}_\sigma)_i, (\mathcal{H}_\alpha)_i) \right\},$$

$$(Q_{st}^\mu)_e(\mathcal{F}_\sigma, \mathcal{H}_\alpha) = \bigvee_{\{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{V}}^{lj} \mid j \in J\} \in D(\mathcal{F}_\sigma, \mathcal{H}_\alpha)}$$

$$\left\{ \bigwedge_{\{(\mathcal{F}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathcal{V}}^{li} \} \in \{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{V}}^{lj} \mid j \in J\}} Q_e^\mu((\mathcal{F}_\sigma)_i, (\mathcal{H}_\alpha)_i) \right\},$$

$$(Q_{st}^\omega)_e(\mathcal{F}_\sigma, \mathcal{H}_\alpha) = \bigvee_{\{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{V}}^{lj} \mid j \in J\} \in D(\mathcal{F}_\sigma, \mathcal{H}_\alpha)}$$

$$\left\{ \bigwedge_{\{(\mathcal{F}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathcal{V}}^{li} \} \in \{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{V}}^{lj} \mid j \in J\}} Q_e^\omega((\mathcal{F}_\sigma)_i, (\mathcal{H}_\alpha)_i) \right\},$$

where  $D(\mathcal{F}_\sigma, \mathcal{H}_\alpha) = \{ \{(\mathcal{F}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{E}}^j\} \mid j \in J, J \text{ is finite} \} \mid \mathcal{F}_\sigma \sqsubseteq \bigcup_{j \in J} ((\mathcal{F}_\sigma)_j \sqcap \tilde{\mathcal{E}}^j) \text{ and } \mathcal{H}_\alpha \sqsubseteq \bigcap_{j \in J} ((\mathcal{H}_\alpha)_j \sqcup \tilde{\mathcal{E}}^{1-l_j}), l \in \zeta \}$ . Then  $(Q_{st}^{v\mu\omega})_V$  is the coarsest stratified *svnsq-proximity* on  $\mathbb{X}$  which is finer than  $Q_V^{v\mu\omega}$ .

**Proof.** We will prove (Q<sub>S</sub>) only; Conditions (Q<sub>1</sub>) to (Q<sub>4</sub>) are similar to proving Theorem 5.

(Q<sub>5</sub>) Presume there exists  $\mathcal{E}_\sigma, \mathcal{H}_\alpha \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}$ ,  $e \in \mathfrak{Y}$  such that

$$\begin{aligned} (Q_{st}^v)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &\not\leq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ (Q_{st}^v)_e(\mathcal{E}_\sigma, \mathbf{g}_C) \vee (Q_{st}^v)_e(\mathbf{g}_C^c, \mathcal{H}_\alpha) \right], \\ (Q_{st}^\mu)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &\not\leq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ (Q_{st}^\mu)_e(\mathcal{E}_\sigma, \mathbf{g}_C) \wedge (Q_{st}^\mu)_e(\mathbf{g}_C^c, \mathcal{H}_\alpha) \right], \\ (Q_{st}^\omega)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &\not\leq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ (Q_{st}^\omega)_e(\mathcal{E}_\sigma, \mathbf{g}_C) \wedge (Q_{st}^\omega)_e(\mathbf{g}_C^c, \mathcal{H}_\alpha) \right], \end{aligned}$$

then there exists  $r \in \zeta_0$  such that

$$\begin{aligned} (Q_{st}^v)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &< r \leq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ (Q_{st}^v)_e(\mathcal{E}_\sigma, \mathbf{g}_C) \vee (Q_{st}^v)_e(\mathbf{g}_C^c, \mathcal{H}_\alpha) \right], \\ (Q_{st}^\mu)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &> 1 - r \geq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ (Q_{st}^\mu)_e(\mathcal{E}_\sigma, \mathbf{g}_C) \wedge (Q_{st}^\mu)_e(\mathbf{g}_C^c, \mathcal{H}_\alpha) \right], \\ (Q_{st}^\omega)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &> 1 - r \geq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ (Q_{st}^\omega)_e(\mathcal{E}_\sigma, \mathbf{g}_C) \wedge (Q_{st}^\omega)_e(\mathbf{g}_C^c, \mathcal{H}_\alpha) \right]. \end{aligned} \tag{12}$$

From the concept of  $(Q_{st}^{\mu\omega})_{\mathfrak{Y}}$ , there exists a collection  $\mathcal{A} = \{(\mathcal{E}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathfrak{Y}}^{lj} \mid j \in J\} \in \mathcal{D}(\mathcal{E}_\sigma, \mathcal{H}_\alpha)$  such that

$$\begin{aligned} (Q_{st}^v)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &= \bigwedge \left[ \bigvee_{[(\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathfrak{Y}}^{li}] \in \mathcal{A}} Q_e^v((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i) \right] \\ &\geq \bigwedge_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ \bigvee_{[(\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathfrak{Y}}^{li}] \in \mathcal{A}} \left[ (Q_{st}^v)_e((\mathcal{E}_\sigma)_i, (\mathbf{g}_C)_i) \vee (Q_{st}^v)_e((\mathbf{g}_C^c)_i, (\mathcal{H}_\alpha)_i) \right] \right] \geq r, \\ (Q_{st}^\mu)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &= \bigvee \left[ \bigwedge_{[(\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathfrak{Y}}^{li}] \in \mathcal{A}} Q_e^\mu((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i) \right] \\ &\leq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ \bigwedge_{[(\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathfrak{Y}}^{li}] \in \mathcal{A}} \left[ (Q_{st}^\mu)_e((\mathcal{E}_\sigma)_i, (\mathbf{g}_C)_i) \wedge (Q_{st}^\mu)_e((\mathbf{g}_C^c)_i, (\mathcal{H}_\alpha)_i) \right] \right] \leq 1 - r, \\ (Q_{st}^\omega)_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &= \bigvee \left[ \bigwedge_{[(\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathfrak{Y}}^{li}] \in \mathcal{A}} Q_e^\omega((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i) \right] \\ &\leq \bigvee_{\mathbf{g}_C \in \widetilde{(\mathfrak{X}, \mathfrak{Y})}} \left[ \bigwedge_{[(\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\mathfrak{Y}}^{li}] \in \mathcal{A}} \left[ (Q_{st}^\omega)_e((\mathcal{E}_\sigma)_i, (\mathbf{g}_C)_i) \wedge (Q_{st}^\omega)_e((\mathbf{g}_C^c)_i, (\mathcal{H}_\alpha)_i) \right] \right] \leq 1 - r. \end{aligned}$$

A contradiction for equation (12).  $\square$

**Theorem 11.** Let  $(\mathfrak{X}, Q_V^{\nu\mu\omega})$  and  $(\mathcal{U}, (Q_R^{\nu\mu\omega})^*)$  be two svnsq-proximity spaces. If  $\psi_\varphi : (\mathfrak{X}, Q_V^{\nu\mu\omega}) \rightarrow (\mathcal{U}, (Q_R^{\nu\mu\omega})^*)$  be a svnsq-proximity continuous, then  $\psi_\varphi : (\mathfrak{X}, (Q_V^{\nu\mu\omega})_{st}) \rightarrow (\mathcal{U}, (Q_R^{\nu\mu\omega})_{st}^*)$  is a svnsq-proximity continuous

**Proof.** Let  $\mathcal{E}_\sigma, \mathcal{H}_\alpha \in \widetilde{(\mathcal{U}, \mathcal{R})}$  with  $\mathcal{E}_\sigma \sqsubseteq \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \sqcap \tilde{\mathcal{R}}^{lj})$  and  $\mathcal{H}_\alpha \sqsubseteq \sqcup_{j \in J} ((\mathcal{H}_\alpha)_j \sqcap \tilde{\mathcal{R}}^{1-lj})$ . Then  $\psi_\varphi^{-1}(\mathcal{E}_\sigma) \sqsubseteq \sqcup_{j \in J} \psi_\varphi^{-1}((\mathcal{E}_\sigma)_j) \sqcap \tilde{\mathfrak{Y}}^{lj}$  and  $\psi_\varphi^{-1}(\mathcal{H}_\alpha) \sqsubseteq \sqcup_{j \in J} \psi_\varphi^{-1}((\mathcal{H}_\alpha)_j) \sqcap \tilde{\mathfrak{Y}}^{1-lj}$ . For each collection  $\mathcal{D}(\mathcal{E}_\sigma, \mathcal{H}_\alpha) = \{ \{ ((\mathcal{E}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{R}}^{lj}) \mid j \in J, J \text{ is finite} \} \mid \mathcal{E}_\sigma \sqsubseteq \sqcup_{j \in J} ((\mathcal{E}_\sigma)_j \sqcap \tilde{\mathcal{R}}^{lj}) \text{ and } \mathcal{H}_\alpha \sqsubseteq \sqcup_{j \in J} ((\mathcal{H}_\alpha)_j \sqcap \tilde{\mathcal{R}}^{1-lj}), l \in \zeta \}$ , and put  $\mathcal{A} = \{ (\mathcal{E}_\sigma)_j, (\mathcal{H}_\alpha)_j, \tilde{\mathcal{R}}^{lj} \mid j \in J \}$ , we have

$$\begin{aligned} &(Q_{st}^v)_e(\psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathcal{H}_\alpha)) \\ &\geq \bigvee_{(\psi_\varphi^{-1}((\mathcal{E}_\sigma)_j), \psi_\varphi^{-1}((\mathcal{H}_\alpha)_l), \tilde{\mathfrak{Y}}^{lj}) \in \{ \psi_\varphi^{-1}((\mathcal{E}_\sigma)_j), \psi_\varphi^{-1}((\mathcal{H}_\alpha)_j), \tilde{\mathfrak{Y}}^{lj} \mid j \in J \}} \\ &Q_e^v(\psi_\varphi^{-1}((\mathcal{E}_\sigma)_j), \psi_\varphi^{-1}((\mathcal{H}_\alpha)_j)) \end{aligned}$$

$$\begin{aligned}
 &\geq \bigvee_{((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{R}}^{j_i} \in \mathcal{A}} (\mathcal{Q}_{\varphi(e)}^v)^*((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}) \\
 &\geq \bigwedge_{D(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left( \bigvee_{((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{R}}^{j_i} \in \mathcal{A}} (\mathcal{Q}_{\varphi(e)}^v)^*((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}) \right) \\
 &= (\mathcal{Q}_{\varphi(e)}^v)_{st}^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \\
 &(\mathcal{Q}_e^\mu)_{st} \left( \psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathfrak{h}_\alpha) \right) \\
 &\leq \bigwedge_{\left( \psi_\varphi^{-1}((\mathcal{E}_\sigma)_{j_i}), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{V}}^{j_i} \right) \in \left\{ \psi_\varphi^{-1}((\mathcal{E}_\sigma)_j), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_j), \tilde{\mathcal{V}}^{j_i} \mid j \in J \right\}} \\
 &\mathcal{Q}_e^\mu \left( \psi_\varphi^{-1}((\mathcal{E}_\sigma)_{j_i}), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_{j_i}) \right) \\
 &\leq \bigwedge_{((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{R}}^{j_i} \in \mathcal{A}} (\mathcal{Q}_{\varphi(e)}^\mu)^*((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}) \\
 &\leq \bigvee_{D(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left( \bigwedge_{((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{R}}^{j_i} \in \mathcal{A}} (\mathcal{Q}_{\varphi(e)}^\mu)^*((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}) \right) \\
 &= (\mathcal{Q}_{\varphi(e)}^\mu)_{st}^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) \\
 &(\mathcal{Q}_e^\omega)_{st} \left( \psi_\varphi^{-1}(\mathcal{E}_\sigma), \psi_\varphi^{-1}(\mathfrak{h}_\alpha) \right) \\
 &\leq \bigwedge_{\left( \psi_\varphi^{-1}((\mathcal{E}_\sigma)_{j_i}), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{V}}^{j_i} \right) \in \left\{ \psi_\varphi^{-1}((\mathcal{E}_\sigma)_j), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_j), \tilde{\mathcal{V}}^{j_i} \mid j \in J \right\}} \\
 &\mathcal{Q}_e^\omega \left( \psi_\varphi^{-1}((\mathcal{E}_\sigma)_{j_i}), \psi_\varphi^{-1}((\mathfrak{h}_\alpha)_{j_i}) \right) \\
 &\leq \bigwedge_{((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{R}}^{j_i} \in \mathcal{A}} (\mathcal{Q}_{\varphi(e)}^\omega)^*((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}) \\
 &\leq \bigvee_{D(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)} \left( \bigwedge_{((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}), \tilde{\mathcal{R}}^{j_i} \in \mathcal{A}} (\mathcal{Q}_{\varphi(e)}^\omega)^*((\mathcal{E}_\sigma)_{j_i}, (\mathfrak{h}_\alpha)_{j_i}) \right) \\
 &= (\mathcal{Q}_{\varphi(e)}^\omega)_{st}^*(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)
 \end{aligned}$$

By Definition 5, we have  $\psi_\varphi : (\mathbb{Y}, (\mathcal{Q}_V^{v\mu\omega})_{st}) \rightarrow (\mathcal{U}, (\widetilde{\mathcal{Q}}_{\mathcal{R}}^{v\mu\omega})_{st}^*)$  is a *svnsq-proximity continuous*.  $\square$

**Theorem 12.** Let  $(\mathbb{Y}, \mathcal{Q}_V^{v\mu\omega})$  be svnsq-proximity space. Then, for  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in (\widetilde{\mathbb{Y}}, \mathcal{Y}), e \in \mathcal{Y}$ .

- (i)  $\mathcal{H}_Q^v, \mathcal{H}_Q^\mu, \mathcal{H}_Q^\omega : \mathcal{Y} \rightarrow \zeta^{(\mathbb{Y}, \mathcal{Y}) \times (\mathbb{Y}, \mathcal{Y})}$  defined by,  $(\mathcal{H}_Q^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = [\mathcal{Q}_e^v(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)]^c$ ,  $(\mathcal{H}_Q^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = [\mathcal{Q}_e^\mu(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)]^c$  and  $(\mathcal{H}_Q^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = [\mathcal{Q}_e^\omega(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)]^c$  is symmetrical svn-topogenous order on  $\mathbb{Y}$ .
- (ii)  $(\mathcal{H}_{Q_{st}}^{v\mu\omega})_V = [(\mathcal{H}_Q^{v\mu\omega})_{st}]_V$ .

**Proof.** (i) Straightforward.

(ii) Since  $(\mathcal{H}_{Q_{st}}^v)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^i) = [(\mathcal{Q}_{st}^v)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^{1-i})]^c = 1$ ,  $(\mathcal{H}_{Q_{st}}^\mu)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^i) = [(\mathcal{Q}_{st}^\mu)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^{1-i})]^c = 0$  and  $(\mathcal{H}_{Q_{st}}^\omega)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^i) = [(\mathcal{Q}_{st}^\omega)_e(\tilde{\mathcal{V}}^i, \tilde{\mathcal{V}}^{1-i})]^c = 0$ , for every  $i \in \zeta$ , we obtain  $[(\mathcal{H}_{Q_{st}}^{v\mu\omega})_{st}]_V$  is stratified which is finer than  $(\mathcal{H}_Q^{v\mu\omega})_V$ . Hence,  $(\mathcal{H}_{Q_{st}}^{v\mu\omega})_V \supseteq [(\mathcal{H}_Q^{v\mu\omega})_{st}]_V$ .

Conversely, assume there exist  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in (\widetilde{\mathbb{Y}}, \mathcal{Y}), e \in \mathcal{Y}$  such that,

$$\begin{aligned}
 (\mathcal{H}_{Q_{st}}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= [(\mathcal{Q}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)]^c \not\leq [(\mathcal{H}_Q^v)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 (\mathcal{H}_{Q_{st}}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= [(\mathcal{Q}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)]^c \not\leq [(\mathcal{H}_Q^\mu)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 (\mathcal{H}_{Q_{st}}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= [(\mathcal{Q}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)]^c \not\leq [(\mathcal{H}_Q^\omega)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha).
 \end{aligned}$$

By the concept of  $(\mathcal{Q}_{st}^{v\mu\omega})_E$ , there exists a collection  $\mathcal{A} = \{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\mathcal{V}}^{j_i} \mid j \in J\} \in D(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)$  such that

$$\begin{aligned}
 [(\mathcal{H}_Q^v)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\not\leq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{A}} [\mathcal{Q}_e^v((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)]^c, \\
 [(\mathcal{H}_Q^\mu)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\not\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{A}} [\mathcal{Q}_e^\mu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)]^c, \\
 [(\mathcal{H}_Q^\omega)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\not\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{A}} [\mathcal{Q}_e^\omega((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)]^c.
 \end{aligned} \tag{13}$$

On another side,  $M = \{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha^c)_j, \tilde{\gamma}^{lj} \mid j \in J\} \in \mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)$  we have

$$\begin{aligned}
 [(\mathcal{H}_Q^v)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in M} (\mathcal{H}_Q^v)_e((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i) \\
 &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in M} [\mathcal{Q}_e^v((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)]^c, \\
 [(\mathcal{H}_Q^\mu)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in M} (\mathcal{H}_Q^\mu)_e((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i) \\
 &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in M} [\mathcal{Q}_e^\mu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)]^c, \\
 [(\mathcal{H}_Q^\omega)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in M} (\mathcal{H}_Q^\omega)_e((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i) \\
 &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in M} [\mathcal{Q}_e^\omega((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i)]^c,
 \end{aligned}$$

which is a contradiction of equations (13). Therefore,  $(\mathcal{H}_{Q_{st}}^{v\mu\omega})_\gamma = [(\mathcal{H}_Q^{v\mu\omega})_{st}]_\gamma$ .  $\square$

**Theorem 13.** Let  $\mathcal{H}_\gamma^{v\mu\omega}$  be symmetrical svn-topogenous order on  $\gamma$ . Then, for all  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \gamma)}$ ,  $e \in \gamma$ .

- (i)  $\mathcal{Q}_{\mathcal{H}}^v, \mathcal{Q}_{\mathcal{H}}^\mu, \mathcal{Q}_{\mathcal{H}}^\omega : \gamma \rightarrow \zeta^{\widetilde{(\mathfrak{X}, \gamma)} \times \widetilde{(\mathfrak{X}, \gamma)}}$  defined by,  $(\mathcal{Q}_{\mathcal{H}}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = [(\mathcal{H}_e^v(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c))]^c$ ,  $(\mathcal{Q}_{\mathcal{H}}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = [(\mathcal{H}_e^\mu(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c))]^c$  and  $(\mathcal{Q}_{\mathcal{H}}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) = [(\mathcal{H}_e^\omega(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c))]^c$  is svnq-proximity on  $\mathfrak{X}$ .
- (ii)  $(\mathcal{Q}_{\mathcal{H}_{st}}^{v\mu\omega})_\gamma = [(\mathcal{Q}_{\mathcal{H}}^{v\mu\omega})_{st}]_\gamma$ .

**Proof.** (i) Obvious.

(ii) Since  $(\mathcal{Q}_{\mathcal{H}_{st}}^v)_e(\tilde{\gamma}^l, \tilde{\gamma}^l) = 1 - (\mathcal{H}_{st}^v)_e(\tilde{\gamma}^l, \tilde{\gamma}^{1-l}) = 1 - 0 = 1$ ,  $(\mathcal{Q}_{\mathcal{H}_{st}}^\mu)_e(\tilde{\gamma}^l, \tilde{\gamma}^l) = 1 - (\mathcal{H}_{st}^\mu)_e(\tilde{\gamma}^l, \tilde{\gamma}^{1-l}) = 1 - 1 = 0$  and  $(\mathcal{Q}_{\mathcal{H}_{st}}^\omega)_e(\tilde{\gamma}^l, \tilde{\gamma}^l) = 1 - (\mathcal{H}_{st}^\omega)_e(\tilde{\gamma}^l, \tilde{\gamma}^{1-l}) = 1 - 1 = 0$ , for every  $l \in \zeta$ , we have  $[(\mathcal{Q}_{\mathcal{H}_{st}}^{v\mu\omega})_{st}]_\gamma$  is stratified which is finer than  $(\mathcal{Q}_{\mathcal{H}}^{v\mu\omega})_\gamma$ . Hence,  $(\mathcal{Q}_{\mathcal{H}_{st}}^{v\mu\omega})_\gamma \supseteq [(\mathcal{Q}_{\mathcal{H}}^{v\mu\omega})_{st}]_\gamma$ .

Conversely, suppose that there exist  $\mathcal{E}_\sigma, \mathfrak{h}_\alpha \in \widetilde{(\mathfrak{X}, \gamma)}$ ,  $e \in \gamma$  such that,

$$\begin{aligned}
 (\mathcal{Q}_{\mathcal{H}_{st}}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= 1 - (\mathcal{H}_{st}^v)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c) \not\leq [(\mathcal{Q}_{\mathcal{H}}^v)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 (\mathcal{Q}_{\mathcal{H}_{st}}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= 1 - (\mathcal{H}_{st}^\mu)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c) \not\leq [(\mathcal{Q}_{\mathcal{H}}^\mu)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha), \\
 (\mathcal{Q}_{\mathcal{H}_{st}}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &= 1 - (\mathcal{H}_{st}^\omega)_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c) \not\leq [(\mathcal{Q}_{\mathcal{H}}^\omega)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha).
 \end{aligned}$$

By the concept of  $(\mathcal{H}_{st}^{v\mu\omega})_E$ , there exists a collection  $\{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha)_j, \tilde{\gamma}^{lj} \mid j \in J\} = \mathcal{A} \in \mathcal{N}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha^c)$  such that

$$\begin{aligned}
 [(\mathcal{Q}_{\mathcal{H}}^v)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\not\leq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{A}} 1 - \mathcal{H}_e^v((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i), \\
 [(\mathcal{Q}_{\mathcal{H}}^\mu)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\not\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{A}} 1 - \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i), \\
 [(\mathcal{Q}_{\mathcal{H}}^\omega)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\not\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{A}} 1 - \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i).
 \end{aligned} \tag{14}$$

On the other hand,  $P = \{(\mathcal{E}_\sigma)_j, (\mathfrak{h}_\alpha^c)_j, \tilde{\gamma}^{lj} \mid j \in J\} \in \mathcal{D}(\mathcal{E}_\sigma, \mathfrak{h}_\alpha)$  we have

$$\begin{aligned}
 [(\mathcal{Q}_{\mathcal{H}}^v)_{st}]_e(\mathcal{E}_\sigma, \mathfrak{h}_\alpha) &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in P} (\mathcal{Q}_{\mathcal{H}}^v)_e((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i) \\
 &\geq \bigwedge_{((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha^c)_i, \tilde{\gamma}^i) \in P} 1 - \mathcal{H}_e^v((\mathcal{E}_\sigma)_i, (\mathfrak{h}_\alpha)_i),
 \end{aligned}$$

$$\begin{aligned}
 [(\mathcal{Q}_H^\mu)_{st}]_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{P}} (\mathcal{Q}_H^\mu)_e((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i) \\
 &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{P}} 1 - \mathcal{H}_e^\mu((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i), \\
 [(\mathcal{Q}_H^\omega)_{st}]_e(\mathcal{E}_\sigma, \mathcal{H}_\alpha) &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{P}} (\mathcal{Q}_H^\omega)_e((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i) \\
 &\leq \bigvee_{((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i, \tilde{\gamma}^i) \in \mathcal{P}} 1 - \mathcal{H}_e^\omega((\mathcal{E}_\sigma)_i, (\mathcal{H}_\alpha)_i),
 \end{aligned}$$

which is a contradiction of equations (14). Hence,  $(\mathcal{Q}_{H_{st}}^{\nu\mu\omega})_\gamma = [(\mathcal{Q}_H^{\nu\mu\omega})_{st}]_\gamma \cdot (\mathcal{H}_{Q_{st}}^{\nu\mu\omega})_\gamma = [(\mathcal{H}_Q^{\nu\mu\omega})_{st}]_\gamma$ .  $\square$

## 5. Conclusions

Theoretically, we have advanced a set of overarching concepts derived from the principles outlined in previous works [48,49]. These include the stratified single-valued neutrosophic soft quasi proximity, stratified single-valued neutrosophic soft topogenous order, stratified single-valued neutrosophic soft filter, and stratified single-valued neutrosophic soft quasi uniformity, along with an exploration of their respective characteristics. Additionally, we have studied the interconnectedness between these single-valued neutrosophic soft topological constructs and their stratifications.

Regarding further research, we intend to evaluate the correlations between our findings and advancements in neural networks [50–54], multidimensional systems and signal processing [55–57], as well as optimization algorithms [58] and global optimization [59]. Finally, we present a practical application of these concepts in solving decision-making problems. This inventive expansion has the potential to greatly expand existing theoretical frameworks for managing indeterminacy, while also opening up new paths for application and research.

### For forthcoming papers.

The theory can be extended in the next normal methods,

1-Basic concepts can be studied of neutrosophic metric topological spaces using the notion of single-valued neutrosophic soft quasi-uniform present in this article;

2-Examine the connected, separation axioms and soft closure spaces in the context of neutrosophic soft quasi-uniform.

### CRedit authorship contribution statement

**Fahad Alsharari:** Writing – review & editing, Writing – original draft, Funding acquisition, Conceptualization. **Yaser Saber:** Writing – review & editing. **Hanan Alohalı:** Writing – review & editing, Writing – original draft. **Mesfer H. Alqahtani:** Writing – original draft, Supervision, Resources, Funding acquisition. **Mubarak Ebodey:** Writing – review & editing. **Tawfik Elmasry:** Writing – review & editing. **Jafar Alsharif:** Writing – review & editing. **Amal F. Soliman:** Writing – review & editing, Data curation. **Florentin Smarandache:** Writing – review & editing. **Fahad Sikander:** Funding acquisition, Formal analysis.

### Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Yaser Saber reports financial support was provided by Majmaah University.

### Data availability statement

The article outlines the research that used no data.

### Acknowledgements

The author extends the appreciation to the Deanship of Postgraduate Studies and Scientific Research at Majmaah University for funding this research work through the project number (R-2024-1005). Also, this study is supported via funding from Prince Sattam bin Abdulaziz University with project number (PSAU/2024/R/1445).

### References

- [1] L.A. Zadeh, Fuzzy sets, Inf. Control 8 (3) (1965) 338–353, [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X).
- [2] K.T. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets Syst. 20 (1) (1986) 87–96, [https://doi.org/10.1016/S0165-0114\(86\)80034-3](https://doi.org/10.1016/S0165-0114(86)80034-3).

- [3] K.T. Atanassov, Operators over interval-valued intuitionistic fuzzy sets, *Fuzzy Sets Syst.* 64 (2) (1994) 159–174, [https://doi.org/10.1016/0165-0114\(94\)90331-X](https://doi.org/10.1016/0165-0114(94)90331-X).
- [4] M.B. Gorzalczy, A method of inference in approximate reasoning based on interval-valued fuzzy sets, *Fuzzy Sets Syst.* 21 (1) (1987) 1–17.
- [5] Z. Pawlak, Rough sets, *Int. J. Comput. Inf. Sci.* 11 (5) (1982) 341–356.
- [6] D.A. Molodtsov, Soft set theory—first results, *Comput. Math. Appl.* 37 (4–5) (1999) 19–31, [https://doi.org/10.1016/S0898-1221\(99\)00056-5](https://doi.org/10.1016/S0898-1221(99)00056-5).
- [7] N.A. Shirokov, *Analytic Functions Smooth up to the Boundary*, *Lecture Notes Math.*, vol. 1312, Springer, Berlin, 1988.
- [8] J.W. Weibull, *Evolutionary Game Theory*, MIT Press, Cambridge, 1997, <https://mitpress.mit.edu/9780262731218/evolutionary-game-theory/>.
- [9] S. Biswas, S. Mui, S.P. Sarkar, Neutrosophic Riemann integration and its properties, *Soft Comput.* 25 (2021) 13987–13999, <https://doi.org/10.1007/s00500-021-06200-7>.
- [10] K. Kasahara, R. Masayama, K. Okita, N. Matubayasi, Elucidating protein–ligand binding kinetics based on returning probability theory, *J. Chem. Phys.* 159 (2023) 134103, <https://doi.org/10.1063/5.0165692>.
- [11] P.K. Maji, R. Biswas, A.R. Roy, Soft set theory, *Comput. Math. Appl.* 45 (4–5) (2003) 555–562, [https://doi.org/10.1016/S0898-1221\(03\)00016-6](https://doi.org/10.1016/S0898-1221(03)00016-6).
- [12] P.K. Maji, A.R. Roy, R. Biswas, An application of soft sets in a decision-making problem, *Comput. Math. Appl.* 44 (8–9) (2002) 1077–1083, [https://doi.org/10.1016/S0898-1221\(02\)00216-X](https://doi.org/10.1016/S0898-1221(02)00216-X).
- [13] P.P. Dey, S. Pramanik, B.C. Gir, Generalized neutrosophic soft multi-attribute group decision making based on TOPSIS, *Crit. Rev.* 11 (2015) 41–55.
- [14] P.P. Dey, S. Pramanik, B.C. Gir, Neutrosophic soft multi-attribute decision making based on grey relational projection method, *Neutrosophic Sets Syst.* 11 (2016) 98–106.
- [15] H. Aktas, N. Çağman, Soft sets and soft groups, *Inf. Sci.* 177 (3) (2007) 2726–2735, <https://doi.org/10.1016/j.ins.2006.12.008>.
- [16] F. Feng, Y.B. Jun, X. Zhao, Soft semirings, *Comput. Math. Appl.* 56 (10) (2008) 2621–2628, <https://doi.org/10.1016/j.camwa.2008.05.011>.
- [17] D. Chen, E.C.C. Tsang, D.S. Yeung, E. Wang, The parameterization reduction of soft sets and its applications, *Comput. Math. Appl.* 49 (5–6) (2005) 757–763, <https://doi.org/10.1016/j.camwa.2004.10.036>.
- [18] M.I. Ali, F. Feng, X. Liu, W.K. Min, M. Shabir, On some new operation in soft set theory, *Comput. Math. Appl.* 57 (9) (2009) 1547–1553, <https://doi.org/10.1016/j.camwa.2008.11.009>.
- [19] Q.M. Sun, Z.L. Zhang, J. Liu, Soft sets and soft modules, *Lect. Notes Comput. Sci.* 5009 (2008) 403–409.
- [20] X. Yang, T.S. Lin, J. Yang, Y. Li, D. Yu, Combination of interval-valued fuzzy set and soft set, *Comput. Math. Appl.* 58 (3) (2011) 521–527, <https://doi.org/10.1016/j.camwa.2009.04.019>.
- [21] A. Kharal, B. Ahmed, Mappings on fuzzy soft classes, *Adv. Fuzzy Syst.* (2009) 407890, <https://doi.org/10.1155/2009/407890>.
- [22] B. Ahmed, A. Kharal, On fuzzy soft sets, *Adv. Fuzzy Syst.* (2009) 586507, <https://doi.org/10.1155/2009/586507>.
- [23] M. Shabir, M. Naz, On soft topological spaces, *Comput. Math. Appl.* 61 (7) (2011) 1786–1799, <https://doi.org/10.1016/j.camwa.2011.02.006>.
- [24] B. Tanay, M.B. Kandemir, Topological structure of fuzzy soft sets, *Comput. Math. Appl.* 61 (10) (2011) 2952–2957, <https://doi.org/10.1016/j.camwa.2011.03.056>.
- [25] C.L. Chang, Fuzzy topological spaces, *J. Math. Anal. Appl.* 24 (1) (1968) 182–190, [https://doi.org/10.1016/0022-247X\(68\)90057-7](https://doi.org/10.1016/0022-247X(68)90057-7).
- [26] B.P. Varol, H. Aygün, Fuzzy soft topology, *Hacet. J. Math. Stat.* 41 (3) (2012) 407–419.
- [27] A. Aygünöglü, V. Çetkin, H. Aygün, An introduction to fuzzy soft topological spaces, *Hacet. J. Math. Stat.* 43 (2) (2014) 197–208.
- [28] Y.M. Saber, F. Alsharari, F. Smarandache, An introduction to single-valued neutrosophic soft topological structure, *Soft Comput.* 26 (2022) 7107–7122, <https://doi.org/10.1007/s00500-022-07150-4>.
- [29] W. Kotzé, *Uniform Spaces Handbook of Fuzzy Sets Series*, Chapter 8, vol. 3, Kluwer Academic Publisher, Dordrecht, Boston, London, 1999, pp. 553–580.
- [30] B. Hutton, Uniformities in fuzzy topological spaces, *J. Math. Anal. Appl.* 58 (1977) 559–571.
- [31] R. Lowen, Fuzzy uniform spaces, *J. Math. Anal. Appl.* 82 (2) (1981) 370–385, [https://doi.org/10.1016/0022-247X\(81\)90202-X](https://doi.org/10.1016/0022-247X(81)90202-X).
- [32] F. Smarandache, *A Unifying Field in Logics: Neutrosophic Logic: Neutrosophy, Neutrosophic Set, Neutrosophic Probability and Statistics*, 6th ed., InfoLearnQuest, Ann Arbor, MI, USA, 2007, available online: <http://fs.gallup.unm.edu/eBook-neutrosophics6.pdf>. (Accessed 10 February 2019).
- [33] H. Wang, F. Smarandache, Y.Q. Zhang, R. Sunderraman, Single valued neutrosophic sets, *Multisp. Multistruct.* 4 (2010) 410–413.
- [34] A.A. Salama, S.A. Alblowi, Neutrosophic set and neutrosophic topological spaces, *IOSR J. Math.* 3 (2012) 31–35.
- [35] A.A. Salama, F. Smarandache, Neutrosophic Crisp Set Theory, The Educational Publisher Columbus, Columbus, OH, USA.
- [36] P.A. Rodrigo, S. Maheswari, Neutrosophic  $gs\alpha^*$  – Open and closed maps in neutrosophic topological spaces, *Neutrosophic Syst. Appl.* 8 (2023) 42–49, <https://doi.org/10.61356/j.nswa.2023.39>.
- [37] F. Smarandache, New types of topologies and neutrosophic topologies, *Neutrosophic Syst. Appl.* 1 (2023) 1–3, <https://doi.org/10.61356/j.nswa.2023.1>.
- [38] R. Dhar, Compactness and neutrosophic topological space via grills, *Neutrosophic Syst. Appl.* 2 (2023) 1–7, <https://doi.org/10.61356/j.nswa.2023.6>.
- [39] S. Dey, G.C. Ray, Separation axioms in neutrosophic topological spaces, *Neutrosophic Syst. Appl.* 2 (2023) 38–54, <https://doi.org/10.61356/j.nswa.2023.9>.
- [40] S. Broumi, S. Mohanaselvi, T. Witezak, M. Talea, A. Bakali, F. Smarandache, Complex fermatean neutrosophic graph and application to decision making, *Decis. Mak. Appl. Manag. Eng.* 6 (1) (2023) 474–501, <https://doi.org/10.31181/dmame24022023b>.
- [41] Y.M. Saber, M.A. Abdel-Sattar, Ideals on fuzzy topological spaces, *Appl. Math. Sci.* 8 (34) (2014) 1667–1691, <https://doi.org/10.12988/ams.2014.33194>.
- [42] Y.M. Saber, F. Alsharari,  $\zeta\mathcal{O}_v^{st}$ -fuzzy closure operator, *New Math. Nat. Comput.* 16 (1) (2020) 123–141, <https://doi.org/10.1142/S1793005720500088>.
- [43] Y.M. Saber, F. Alsharari, Generalized fuzzy ideal closed sets on fuzzy topological spaces in Šostak sense, *Int. J. Fuzzy Log. Intell. Syst.* 18 (3) (2018) 161–166, <https://doi.org/10.5391/IJFIS.2018.18.3.161>.
- [44] Y.M. Saber, F. Alsharari, F. Smarandache, On single-valued neutrosophic ideals in Šostak sense, *Symmetry* 12 (2) (2020) 193, <https://doi.org/10.3390/sym12020193>.
- [45] Y.M. Saber, F. Alsharari, F. Smarandache, A. Abdel-Sattar, Connectedness and stratification of single-valued neutrosophic topological spaces, *Symmetry* 12 (9) (2020) 1464, <https://doi.org/10.3390/sym12091464>.
- [46] Y.M. Saber, F. Alsharari, F. Smarandache, A. Abdel-Sattar, On single valued neutrosophic regularity spaces, *Comput. Model. Eng. Sci.* 130 (3) (2022) 1625–1648, <https://doi.org/10.32604/cmescs.2022.017782>.
- [47] F. Alsharari, F. Smarandache, Y.M. Saber, Compactness on single-valued neutrosophic ideal topological spaces, *Neutrosophic Sets Syst.* 41 (2021) 127–145.
- [48] S.E. Abbas, A.A. Abd-Allah, Stratified double L-topological structures, *Math. Comput. Model.* 54 (1–2) (2011) 423–439, <https://doi.org/10.1016/j.mcm.2011.02.032>.
- [49] S.E. Abbas, E. El-sanowsy, A. Atef, Stratified modeling in soft fuzzy topological structures, *Soft Comput.* 22 (2018) 1603–1613.
- [50] B. Gu, V.S. Sheng, A robust regularization path algorithm for  $v$  support vector classification, *IEEE Trans. Neural Netw. Learn. Syst.* 28 (5) (2017) 1241–1248.
- [51] B. Gu, X. Sun, V.S. Sheng, Structural minimax probability machine, *IEEE Trans. Neural Netw. Learn. Syst.* 28 (7) (2017) 1646–1656.
- [52] W. Deng, H.M. Zhao, X.H. Yang, J.X. Xiong, M. Sun, B. Li, Study on an improved adaptive PSO algorithm for solving multi-objective gate assignment, *Appl. Soft Comput.* 59 (2017) 288–302.
- [53] B. Gu, V.S. Sheng, K.Y. Tay, W. Romano, S. Li, Incremental support vector learning for ordinal regression, *IEEE Trans. Neural Netw. Learn. Syst.* 26 (7) (2015) 1403–1416.
- [54] B. Wang, X. Gu, L. Ma, S. Yan, Temperature error correction based on BP neural network in meteorological WSN, *Int. J. Sens. Netw.* 23 (4) (2017) 265–278.
- [55] J. Wang, S. Lian, Y.Q. Shi, Hybrid multiplicative multiwatermarking in DWT domain, *Multidimens. Syst. Signal Process.* 28 (2) (2017) 617–636.
- [56] L. Xiong, Z. Xu, Y.Q. Shi, An integer wavelet transform-based scheme for reversible data hiding in encrypted images, *Multidimens. Syst. Signal Process.* 29 (2018) 1191–1202.



- [57] Y. Sun, F. Gu, Compressive sensing of piezoelectric sensor response signal for phased array structural health monitoring, *Int. J. Sens. Netw.* 23 (4) (2017) 258–264.
- [58] W. Deng, H.M. Zhao, L. Zou, G.Y. Li, X.H. Yang, D.Q. Wu, A novel collaborative optimization algorithm in solving complex optimization problems, *Soft Comput.* 21 (15) (2017) 4387–4398.
- [59] Y. Xue, J. Jiang, B. Zaho, T. Ma, A self-adaptive artificial bee colony algorithm based on global best for global optimization, *Soft Comput.* 22 (2018) 2935–2952.