

A Theoretical and Analytical Approach for Fundamental Framework of Composite Mappings on Fuzzy Hypersoft Classes

Muhammad Ahsan

Muhammad Saeed

Atiqe Ur Rahman



A Theoretical and Analytical Approach for Fundamental Framework of Composite Mappings on Fuzzy Hypersoft Classes

Muhammad Ahsan¹, Muhammad Saeed^{1,*} and Atiqe Ur Rahman²

^{1,2} Department of Mathematics, University of Management and Technology Lahore, Pakistan.;

ahsan1826@gmail.com, muhammad.saeed@umt.edu.pk, aurkhh@gmail.com *Correspondence:

ahsan1826@gmail.com

Abstract. Mapping is a fundamental mathematical concept that is used in many elementary areas of science and mathematics and has numerous applications. The core purpose of this study is to provide a theoretical and analytical approach for carving out a basic structure of composite mappings on the classes of Fuzzy Hypersoft (FHS) sets. It is a comprehensive study of existing concepts regarding mappings on fuzzy soft, soft and hesitant fuzzy soft classes through characterizing of composite mappings on FHS classes. Moreover, certain generalized properties of mappings on FHS classes like FHS images and FHS inverse images, are established. Some related results are verified with the help of illustrative examples.

Keywords: Fuzzy soft, Soft classes, Hesitant fuzzy soft classes, Composite mappings, Fuzzy hypersoft set.

1. Introduction

In 1965, Zadeh introduced the theory of fuzzy sets [24]. It has been utilized in different decision making problem [20]- [21]- [22]. There are some theories, theory of likelihood, theory of intuitionistic fuzzy sets [2], [5], theory of vague sets [10], the theory of interval mathematics [2], [11], and theory of rough sets [13] which can be considered as scientific apparatuses for dealing with uncertainties and ambiguous. Despite that, every one of these speculations has their innate challenges as brought up in [12]. The main reason behind these troubles is potentially the inadequacy of the parametrization device of the hypothesis.

Therefore, Molodtsov [12] started the idea of a soft set (SS) as a numerical device for dealing with uncertainties which are liberated from the above challenges (We know about the SS characterized by Pawlak [14], which is an alternate idea and helpful to understand some other kind of issues).

Karaaslan [8] introduced soft class and its pertinent activities. Athar et al. [3], [4] introduced the concept of mappings on fuzzy soft classes and mappings on soft classes in 2009 and 2011 individually. They considered the properties of the soft image and soft inverse image. They also characterized the properties of fuzzy SS, fuzzy S-image, fuzzy S-inverse image of fuzzy S-sets and supported them with examples. Manash et al. [7] gave the idea of composite mappings on hesitant fuzzy soft classes in 2016 and discussed some interesting properties of this idea.

In a diversity of real-life applications, the attributes should be further sub-partitioned into attribute values for more clear understanding. Samarandache [9] fulfilled this need and developed the concept of the HSS as a generalization of the SS. He opened numerous fields in this way of thinking and generalized SS to the hyper-soft set by changing the planning F into a multi-contention function. At that point, he made the differentiation between the sorts of initial universes, crisp, fuzzy, intuitionistic fuzzy, neutrosophic, and plithogenic respectively. Thus, he also showed that a HS set can be crisp, fuzzy, intuitionistic fuzzy, neutrosophic and plithogenic respectively. Saeed et al. [19, 25] explained some basic concepts like HS subset, HS complement, not HS set, absolute set, union, intersection, AND, OR, restricted union, extended intersection, relevant complement, restricted difference, restricted symmetric difference, HS set relation, sub relation, complement relation, HS representation in matrices form, and different operations on matrices. Saeed et al. [23] characterized mapping under a hypersoft set environment, then some of its essential properties like HS images, HS inverse images were also discussed.

The core purpose of this study is to provide a theoretical and analytical approach for carving out a basic structure of composite mappings on the classes of FHS sets. It is an comprehensive study of existing concepts regarding mappings on fuzzy soft, soft classes and hesitant fuzzy soft classes through characterizing of composite mappings on FHS classes. Moreover, certain generalized properties of mappings on FHS classes like FHS images and FHS inverse images, are established. Some related results are proved with the help of illustrative examples. The ordering of the following portion is working out as follows.

In Section 2, some pivotal regarding fuzzy set, SS, fuzzy soft class, soft class, hypersoft set, and fuzzy hypersoft set (FHSS) are re-imagined. In Section 3, composite mappings on FHS classes, FHS image, FHS inverse image, and its relevant theorems with their essential properties are considered. In the last section, some concluding remarks are described.

1.1. Motivation

In a diversity of real-life applications, the attributes should be further sub-partitioned into attribute values for more clear understanding. Samarandache [9] fulfilled this need and developed the concept of the FHSS as a generalization of the fuzzy soft set. Now, It will be a question that how do we define composite mappings for FHSS classes? FHSS set is significant? To answers these questions and getting inspiration from the above writing, it is relevant to broaden the idea of mappings for those sets managing disjoint arrangements of attributed values, i.e FHSS. In this investigation, an extension is made in existing theories with respect to mappings on fuzzy soft, soft classes and hesitant fuzzy soft classes by characterizing composite mappings on FHS classes. The striking component of composite mappings on FHS classes is that it can mirror the interrelationship between the multi-input contentions. Moreover, certain generalized properties of mappings on FHS classes like FHS images and FHS inverse images, are established. Some related results are proved with the help of illustrative examples.

2. Preliminaries

Throughout the following, let $L = F_1 \times F_2 \times F_3 \times \dots \times F_n$, $M = F'_1 \times F'_2 \times F'_3 \times \dots \times F'_n$, $N = F''_1 \times F''_2 \times F''_3 \times \dots \times F''_n$, $O = F'''_1 \times F'''_2 \times F'''_3 \times \dots \times F'''_n$, $\alpha = (\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n)$ and $\beta = (\beta_1, \beta_2, \beta_3, \dots, \beta_n)$.

Definition 2.1. [21] The fuzzy set $X = \{(x, \xi_X(x)) | x \in X\}$ such that

$$\xi_X : X \rightarrow [0, 1]$$

where $\xi_X(x)$ describes the membership percentage of $x \in X$.

Definition 2.2. [12] A pair (F, A) is said to be soft set over X , where F is a mapping given as

$$F : A \rightarrow P(X)$$

On the other way, a SS is the parameterized family of subsets of the universe X . In other words, a soft set over X is a parameterized family of subsets of the universe. For $\epsilon \in A$. $F(\epsilon)$ may be considered as the set of ϵ approximate elements of the soft set (F, A) .

Definition 2.3. [6] Let X be an initial universe, indexed class of fuzzy sets $\{f_i : f_i : X \rightarrow [0, 1], i = 1, 2, \dots, n\}$ is called a fuzzy class.

Definition 2.4. [1] Let X be a universe and E a set of attributes. Then the pair (X, E) denotes the collection of all fuzzy soft sets on X with attributes from E and is called a fuzzy soft class.

Definition 2.5. [9] Let $a_1, a_2, a_3, \dots, a_n$ be the distinct attributes whose corresponding attribute values belongs to the sets $F_1, F_2, F_3, \dots, F_n$ respectively, where $F_i \cap F_j = \Phi$ for $i \neq j$. A pair (Υ, L) is called a hypersoft set over the universal set X , where Υ is the mapping given by $\Upsilon : L \longrightarrow P(X)$.

For more definition see [15–19].

Definition 2.6. [19] Suppose X and $F(X)$ be the universal set and all fuzzy subsets of X respectively. Let $a_1, a_2, a_3, \dots, a_n$ be the distinct attributes whose corresponding attribute values belongs to the sets $F_1, F_2, F_3, \dots, F_n$ respectively, where $F_i \cap F_j = \Phi$ for $i \neq j$ and $i, j \in \{1, 2, 3, \dots, n\}$. Then the FHSS is the pair (Σ_L, L) over X defined by a map $\Sigma_L : L \rightarrow F(X)$.

3. Main results

Definition 3.1. Suppose X and $F(X)$ be the universal set and all fuzzy subsets of X respectively, let $a_1, a_2, a_3, \dots, a_n$ be the distinct attributes whose corresponding attribute values belongs to the sets $F_1, F_2, F_3, \dots, F_n$ respectively, where $F_i \cap F_j = \Phi$ for $i \neq j$, let $F = \{\varsigma_i : i = 1, 2, \dots, n\}$ be a collection of decision makers. Indexed class of FHSS $\{\vartheta_{\varsigma_i} : \vartheta_{\varsigma_i} : L \rightarrow F(X), \varsigma_i \in F\}$, is said to be fuzzy hypersoft class and it can be symbolized in such a form ϑ_F . If for any $\varsigma_i \in F$, $\vartheta_{\varsigma_i} = \Phi$, the FHSS $\vartheta_{\varsigma_i} \notin \vartheta_F$.

Example 3.2. Let $X = \{a = \text{Holstein}, b = \text{Angus}, c = \text{Charolais}\}$ be the set of cow categories. Peter decide to purchase a cow for milk to get vitamin B12 and iodine of doctor instruction. They visit Bos tarus (a Europeon Cattle) to buy such cow which fulfills his requirements. Let $a_1 = \text{vision and hearing}$, $a_2 = \text{Cost}$, $a_3 = \text{Colour}$, distinct attributes whose attribute values belong to the sets F_1, F_2, F_3 . Let $F_1 = \{f_1 = \text{Excellent peripheral vision}, f_2 = \text{Low peripheral vision}\}$, $F_2 = \{f_3 = \text{High}, f_4 = \text{Low}\}$, $F_3 = \{f_5 = \text{White}\}$ and let $F = \{\varsigma_1, \varsigma_1, \varsigma_1\}$ be a set of decision makers. If we consider FHS sets given as

$$\begin{aligned}\vartheta_{\varsigma_1} &= \left\{ \begin{aligned} &((f_1, f_3, f_5), \{0.5/a, 0.7/b\}), ((f_1, f_4, f_5), \{0.1/c\}), \\ &((f_2, f_3, f_5), \{0.4/b, 0.2/c\}), ((f_2, f_4, f_5), \{0.02/b\}) \end{aligned} \right\} \\ \vartheta_{\varsigma_2} &= \left\{ \begin{aligned} &((f_1, f_3, f_5), \{0.05/b, 0.006/c\}), ((f_1, f_4, f_5), \{0.08/a\}), \\ &((f_2, f_3, f_5), \{0.55/a, 0.75/c\}), ((f_2, f_4, f_5), \{0.52/b\}) \end{aligned} \right\} \\ \vartheta_{\varsigma_3} &= \left\{ \begin{aligned} &(f_1, f_3, f_5)\{0.008/c, 0.25/a\}), ((f_1, f_4, f_5), \{0.12/b\}), \\ &((f_2, f_3, f_5), \{0.05/b, 0.64/c\}), ((f_2, f_4, f_5), \{0.28/c\}) \end{aligned} \right\} \\ \text{and} \\ g_{\varsigma_1} &= \left\{ \begin{aligned} &((f_1, f_3, f_5), \{0.87/a\}), ((f_1, f_4, f_5), \{0.23/a\}), \\ &((f_2, f_3, f_5), \{0.09/c, 0.54/a\}), ((f_2, f_4, f_5), \{0.53/a\}) \end{aligned} \right\} \\ g_{\varsigma_2} &= \left\{ \begin{aligned} &((f_1, f_3, f_5), \{0.05/c\}), ((f_1, f_4, f_5), \{0.34/b\}), \\ &((f_2, f_3, f_5), \{0.32/b, 0.27/c\}), ((f_2, f_4, f_5), \{0.08/c\}) \end{aligned} \right\} \end{aligned}$$

$$g_{s3} = \left\{ \begin{array}{l} ((f_1, f_3, f_5), \{0.27/b\}), ((f_1, f_4, f_5), \{0.52/b\}), \\ ((f_2, f_3, f_5), \{0.37/a, 0.38/c\}), ((f_2, f_4, f_5), \{0.001/a\}) \end{array} \right\}$$

Then FHS classes can be written as $\{\vartheta_{s1}, \vartheta_{s2}, \vartheta_{s3}\}, \{g_{s1}, g_{s2}, g_{s3}\}$.




FIGURE 1. Representation of Fuzzy Hypersoft Mapping

Definition 3.3. Let $\overline{(X, L)}$ and $\overline{(Y, M)}$ be classes of FHS sets over X and Y with attributes from L and M respectively. Let $\omega : X \rightarrow Y$ and $v : L \rightarrow M$ be mappings. Then a FHS mappings $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ is defined as follows, for a FHSS P_A in $\overline{(X, L)}$, $\chi(P_A)$ is a FHSS in $\overline{(Y, M)}$ obtained as follows, for $\beta \in v(L) \subseteq M$ and $y \in Y$, $\chi(P_A)(\beta)(y) = \cup_{\alpha \in v^{-1}(\beta) \cap A, s \in \omega^{-1}(y)} (\alpha) \mu_s \chi(P_A)$ is called a fuzzy hypersoft image of a FHSS P_A . Hence $(P_A, \chi(P_A)) \in \chi$, where $P_A \subseteq \overline{(X, L)}$, $\chi(P_A) \subseteq \overline{(Y, M)}$. For more detail see fig 1.

Definition 3.4. If $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ be a FHS mapping, then FHS class $\overline{(X, L)}$ is called the domain of χ and the FHS class $G_B \in \overline{(Y, M)} : G_B = \chi(H_A)$, for some $H_A \in \overline{(X, L)}$ is called the range of χ . The FHS class $\overline{(Y, M)}$ is called co-domain of χ .

Definition 3.5. Let $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ be a FHS mapping and G_B a FHSS in $\overline{(Y, M)}$, where $\omega : X \rightarrow Y$, $v : L \rightarrow M$ and $B \subseteq M$. Then $\chi^{-1}(G_B)$ is a FHSS in $\overline{(X, L)}$ defined as follows, for $\alpha \in v^{-1}(B) \subseteq L$ and $x \in X$, $\chi^{-1}(G_B)(\alpha)(x) = (v(\alpha))\mu_{p(x)}\chi^{-1}(G_B)$ is called a FHS inverse image of G_B . For more detail see fig 2




FIGURE 2. Representation of Fuzzy Hypersoft Inverse Mapping

Example 3.6. Let $X = \{a = \text{Holstein}, b = \text{Angus}, c = \text{Charolais}\}$ and $Y = \{x = \text{Murrah}, y = \text{Siamese}, z = \text{Surti}\}$ be the two universal sets of Cows and Buffalo categories respectively. Peter plans his dairy farm business and wants to understand the difference between a strategy and a tactic. For this purpose, he creates the relationship between these two types of cattle categories which is helpful regrading his dairy farm business. Let $a_1 = \text{vision and hearing}$, $a_2 = \text{Cost}$, $a_3 = \text{Colour}$, and $b_1 = \text{appearance}$, $b_2 = \text{colour}$, $b_3 = \text{price}$ be the two types of distinct attributes whose corresponding attribute values belong to the sets F_1, F_2, F_3 and F'_1, F'_2, F'_3 respectively. Let $F_1 = \{f_1 = \text{Excellent peripheral vision}, f_2 = \text{Low peripheral vision}\}$, $F_2 = \{f_3 = \text{High}\}$, $F_3 = \{f_4 = \text{White}, f_5 = \text{brown}\}$. Similarly, $F'_1 = \{f'_1 = \text{Good}, f'_2 = \text{massive}\}$, $F'_2 = \{f'_3 = \text{black}\}$, $F'_3 = \{f'_4 = \text{low price}\}$ and $\overline{(X, L)}, \overline{(Y, M)}$ be two

Muhammad Ahsan, Muhammad Saeed, Atiqe Ur Rahman, A Theoretical and Analytical Approach for Fundamental Framework of Composite mappings on Fuzzy Hypersoft Classes

classes of FHS sets, where $L = F_1 \times F_2 \times F_3$ and $M = F'_1 \times F'_2 \times F'_3$. Let $\omega : X \rightarrow Y$, $v : F_1 \times F_2 \times F_3 \rightarrow F'_1 \times F'_2 \times F'_3$ be mappings as follows

$\omega(a) = y, \omega(b) = x, \omega(c) = y$ and

$v(f_1, f_3, f_4) = (f'_2, f'_3, f'_4), v(f_1, f_3, f_5) = (f'_1, f'_3, f'_4),$

$v(f_2, f_3, f_4) = (f'_2, f'_3, f'_4), v(f_2, f_3, f_5) = (f'_1, f'_3, f'_4).$

Consider a FHSS P_A in $\overline{(X, L)}$ as

$$P_A = \left\{ \left\{ \begin{array}{l} (f_1, f_3, f_4) = \{ \langle a, \{0.5, 0.3\} \rangle, \\ \langle b, \{0.9, 0.3, 0.5\} \rangle, \langle c, \{0.3\} \rangle \end{array} \right\}, \left\{ \begin{array}{l} (f_1, f_3, f_5) = \{ \langle a, \{0.3, 0.9\} \rangle, \\ \langle b, \{0.5, 0.1\} \rangle, \langle c, \{0.1\} \rangle \end{array} \right\} \right\}$$

Then the FHS image of P_A under $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ is obtained as

$$\chi(P_A)(f_{1'}, f_{3'}, f_{4'})(x) = \cup_{\alpha \in v^{-1}(f_{1'}, f_{3'}, f_{4'}) \cap A, s \in \omega^{-1}(x)} (\alpha) \mu_s = \cup_{\alpha \in (f_1, f_3, f_5) \cap A, s \in b} (\alpha) \mu_s \\ = (f_1, f_3, f_5) \mu_b = \{0.5, 0.1\},$$

$$\chi(P_A)(f_{1'}, f_{3'}, f_{4'})(y) = \{0.3, 0.9\},$$

$$\chi(P_A)(f_{1'}, f_{3'}, f_{4'})(z) = \{0\}$$

$$\chi(P_A)(f_{2'}, f_{3'}, f_{4'})(x) = \{0.9, 0.3, 0.5\}$$

$$\chi(P_A)(f_{2'}, f_{3'}, f_{4'})(y) = \{0.3, 0.5\}$$

$$\chi(P_A)(f_{2'}, f_{3'}, f_{4'})(z) = \{0\}$$

$$\chi(P_A) = \left\{ \left\{ \begin{array}{l} (f'_{1'}, f'_{3'}, f'_{4'}) = \{ \langle x, \{0.5, 0.1\} \rangle, \\ \langle y, \{0.3, 0.9\} \rangle, \langle z, \{0\} \rangle \end{array} \right\}, \left\{ \begin{array}{l} (f'_{2'}, f'_{3'}, f'_{4'}) = \{ \langle x, \{0.9, 0.3, 0.5\} \rangle, \\ \langle y, \{0.3, 0.5\} \rangle, \langle z, \{0\} \rangle \end{array} \right\} \right\}$$

Again consider a FHSS P'_B in $\overline{(Y, M)}$ as

$$P'_B = \left\{ \left\{ \begin{array}{l} (f'_1, f'_3, f'_4) = \{ \langle x, \{0.3, 0.4\} \rangle, \\ \langle y, \{0.4, 0.5, 0.1\} \rangle, \langle z, \{0.9, 0.3\} \rangle \end{array} \right\}, \left\{ \begin{array}{l} (f'_2, f'_3, f'_4) = \{ \langle x, \{0.4, 0.5\} \rangle, \\ \langle y, \{0.9, 0.3\} \rangle, \langle z, \{0.5, 0.4\} \rangle \end{array} \right\} \right\}$$

Therefore,

$$\chi^{-1}(P'_B)(f_1, f_3, f_4)(a) = v((f_1, f_3, f_4)) \mu_{\omega(a)} \\ = (f'_2, f'_3, f'_4) \mu_{\omega(a)} = (f'_2, f'_3, f'_4) \mu_y = \{0.9, 0.3\}$$

$$\chi^{-1}(P'_B)(f_1, f_3, f_4)(b) = \{0.4, 0.5\}$$

$$\chi^{-1}(P'_B)(f_1, f_3, f_4)(c) = \{0.9, 0.3\}$$

$$\chi^{-1}(P'_B)(f_1, f_3, f_5)(a) = \{0.4, 0.5, 0.1\}$$

$$\chi^{-1}(P'_B)(f_1, f_3, f_5)(b) = \{0.3, 0.4\}$$

$$\chi^{-1}(P'_B)(f_1, f_3, f_5)(c) = \{0.4, 0.5, 0.1\}$$

$$\chi^{-1}(P'_B)(f_2, f_3, f_4)(a) = \{0.9, 0.3\}$$

$$\chi^{-1}(P'_B)(f_2, f_3, f_4)(b) = \{0.4, 0.5\}$$

$$\chi^{-1}(P'_B)(f_2, f_3, f_4)(c) = \{0.9, 0.3\}$$

$$\chi^{-1}(P'_B)(f_2, f_3, f_5)(a) = \{0.4, 0.5, 0.1\}$$

$$\chi^{-1}(P'_B)(f_2, f_3, f_5)(b) = \{0.3, 0.4\}$$

$$\chi^{-1}(P'_B)(f_2, f_3, f_5)(c) = \{0.4, 0.5, 0.1\}$$

Similarly,

$$\chi^{-1}(P'_B) = \left\{ \begin{array}{l} \left\{ \begin{array}{l} \{(f_1, f_3, f_4) = \{< a, \{0.9, 0.3\} >, \\ < b, \{0.4, 0.5\} >, < c, \{0.9, 0.3\} >\} \} \\ \{(f_1, f_3, f_5) = \{< a, \{0.4, 0.5, 0.1\} >, \\ < b, \{0.3, 0.4\} >, < c, \{0.4, 0.5, 0.1\} >\} \} \end{array} \right\}, \\ \left\{ \begin{array}{l} \{(f_2, f_3, f_4) = \{< a, \{0.9, 0.3\} >, \\ < b, \{0.4, 0.5\} >, < c, \{0.9, 0.3\} >\} \} \\ \{(f_2, f_3, f_5) = \{< a, \{0.4, 0.5, 0.1\} >, \\ < b, \{0.3, 0.4\} >, < c, \{0.4, 0.5, 0.1\} >\} \} \end{array} \right\} \end{array} \right\}.$$

In case of $\chi : (X, L) \rightarrow (Y, M)$, Peter should purchase those buffalo having attribute values (f_1 = Excellent peripheral vision, f_3 = High, f_4 = White) or can purchase those cow having this attributes values (f'_1 = Good, f'_3 = black, f'_4 = low price) because both of the categories are interrelated and fulfills the requirements individually. For more clarity see 3.




FIGURE 3. Representation of Fuzzy Hypersoft Mapping

Similarly, for inverse FHSS see fig 4.




FIGURE 4. Representation of Fuzzy Hypersoft Inverse Mapping

Definition 3.7. Let $\chi = (\omega, v)$ be a FHS mapping of a FHS class $\overline{(X, L)}$ into a FHS class $\overline{(Y, M)}$. Then

- (1) χ is said to be a one-one (or injection) FHS mapping if for both $\omega : X \rightarrow Y$ and $v : L \rightarrow M$ are one-one.
- (2) χ is said to be a onto (or surjection) FHS mapping if for both $\omega : X \rightarrow Y$ and $v : L \rightarrow M$ are onto.

If χ is both one-one and onto then χ is called a one-one onto (or bijective) correspondence of FHS mapping.

Theorem 3.8. Let $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(X, M)}$ and $\phi = (m, n) : \overline{(X, L)} \rightarrow \overline{(X, M)}$ are two FHS mappings. Then χ and ϕ are equal if and only if $\omega = m$ and $v = n$.

Proof. Obvious.

Theorem 3.9. Two FHS mappings χ and ϕ of a FHS class $\overline{(X, L)}$ into a FHS class $\overline{(Y, M)}$ are equal if and only if $\chi(P_A) = \phi(P_A)$, for all $P_A \in \overline{(X, L)}$.

Proof. Let $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(X, M)}$ and $\phi = (m, n) : \overline{(X, L)} \rightarrow \overline{(X, M)}$ are two FHS mappings. Since ω and v are equal, this implies $\omega = m$ and $v = n$, let $\beta \in v(L) \subseteq M$ and $y \in Y$, $\chi(P_A)(\beta)(y) = \cup_{\alpha \in v^{-1}(\beta) \cap A, s \in \omega^{-1}(y)} (\alpha) \mu_s = \cup_{\alpha \in n^{-1}(\beta) \cap A, s \in m^{-1}(y)} (\alpha) \mu_s$. Hence $\chi(P_A) = \phi(P_A)$.

Conversely,

let $\chi(P_A) = \phi(P_A)$, for all $P_A \in \overline{(X, L)}$, let $(P, Q) \in \chi$, where $P \in \overline{(X, L)}$ and $Q \in \overline{(Y, M)}$.

Therefore $Q = \chi(P) = \phi(P)$, this gives $(P, Q) \in \phi$. Therefore $\chi \subseteq \phi$. Similarly, it can be proved that $\phi \subseteq \chi$. Hence $\phi = \chi$.

Definition 3.10. If $\chi = (\omega, \nu) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ and $\phi = (m, n) : \overline{(Y, M)} \rightarrow \overline{(Z, N)}$ are two FHS mappings, then their composite $\phi \circ \chi$ is a FHS mapping of $\overline{(X, L)}$ into $\overline{(Z, N)}$ such that for every $P_A \in \overline{(X, L)}$, $(\phi \circ \chi)(P_A) = \phi(\chi(P_A))$. We defined as for $\beta \in n(M) \subseteq N$ and $y \in Z$, $\phi(\chi(P_A))(\beta)(y) = \cup_{\alpha \in n^{-1}(\beta) \cap \chi(A), s \in m^{-1}(y)} (\alpha) \mu_s$. For more detail see fig 5.




FIGURE 5. Representation of Composite Fuzzy Hypersoft Mapping

Example 3.11. From 3.6, consider the FHS mapping $\phi = (m, n) : \overline{(Y, M)} \rightarrow \overline{(Z, N)}$ in such a way $m(x) = h_2, m(y) = h_3, m(z) = h_2$, and

$$\begin{aligned} n(f'_1, f'_3, f'_4) &= (f''_1, f''_2, f''_3), \\ n(f'_2, f'_3, f'_4) &= (f''_1, f''_2, f''_4) \end{aligned}$$

where $Z = \{h_1, h_2, h_3\}$, $N = F''_1 \times F''_2 \times F''_3 = \{(f''_1, f''_2, f''_3), (f''_1, f''_2, f''_4)\}$. Therefore

$$\begin{aligned} \phi(\chi(P_A))(f''_1, f''_2, f''_3)(h_1) &= \cup_{\alpha \in n^{-1}(f''_1, f''_2, f''_3) \cap A, s \in m^{-1}(h_1)} (\alpha) \mu_s \\ &= \cup_{\alpha \in (f'_1, f'_3, f'_4) \cap \chi(A), s \in \Phi} (\alpha) \mu_s = (f'_1, f'_3, f'_4) \mu_\Phi = \{0\}, \\ \phi(\chi(P_A))(f''_1, f''_2, f''_3)(h_2) &= \{0.3, 0.4, 0.9\}, \end{aligned}$$

$$\phi(\chi(P_A))(f_1'', f_2'', f_3'')(h_3) = \{0.4, 0.5, 0.1\},$$

So,

$$\begin{aligned} \phi(\chi(P_A))(f_1'', f_2'', f_4'')(h_1) &= \{0\} \\ \phi(\chi(P_A))(f_1'', f_2'', f_4'')(h_2) &= \{0.4, 0.5\}, \\ \phi(\chi(P_A))(f_1'', f_2'', f_4'')(h_3) &= \{0.9, 0.3\} \\ (\phi \circ \chi)(P_A) = \phi(\chi(P_A)) &= \left\{ \left\{ \begin{array}{l} (f_1'', f_2'', f_4'') = \{ < h_1, \{0\} >, \\ < h_2, \{0.3, 0.4, 0.9\} >, < h_3, \{0.4, 0.5, 0.1\} > \end{array} \right\} \right\} \right\} \\ &\left\{ \left\{ \begin{array}{l} (f_1'', f_2'', f_4'') = \{ < h_1, \{0\} >, \\ < h_2, \{0.4, 0.5\} >, < h_3, \{0.9, 0.3\} > \end{array} \right\} \right\} \right\} \end{aligned}$$

Theorem 3.12. Let $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ and $\phi = (m, n) : \overline{(Y, M)} \rightarrow \overline{(Z, N)}$ are two FHS mappings. Then

- (1) if χ and ϕ are one-one then so is $\phi \circ \chi$.
- (2) if ϕ and χ are onto then so is $\chi \circ \phi$.
- (3) if χ and ϕ are both bijections then so is $\phi \circ \chi$.

Example 3.13. Fom 3.6, 3.11, let $\omega(a) = z, \omega(b) = x, \omega(c) = y,$

$$v(f_1, f_3, f_4) = (f_2', f_3', f_5'),$$

$$v(f_1, f_3, f_5) = (f_1', f_3', f_4')$$

and

$$m(x) = h_2, m(y) = h_3, m(z) = h_1$$

$$n(f_1', f_3', f_4') = (f_1'', f_2'', f_3''),$$

$$n(f_2', f_3', f_4') = (f_1'', f_2'', f_4'').$$

Also we consider a FHSSP_A in $\overline{(X, L)}$ as

$$P_A = \left\{ \begin{array}{l} \{(f_1, f_3, f_4) = \left\{ \begin{array}{l} < a, \{0.5, 0.3\} >, \\ < b, \{0.9, 0.3, 0.5\} >, \\ < c, \{0.3\} > \end{array} \right\} \} \\ \{(f_1, f_3, f_5) = \left\{ \begin{array}{l} < a, \{0.3, 0.9\} >, \\ < b, \{0.5, 0.1\} >, \\ < c, \{0.1\} > \end{array} \right\} \} \end{array} \right\}$$

Then the FHS image of P_A under $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ is obtained as

$$\chi(P_A)(f_1', f_3', f_4')(x) = \cup_{\alpha \in v^{-1}(f_1', f_3', f_4') \cap A, s \in \omega^{-1}(x)} (\alpha) \mu_s$$

$$= \cup_{\alpha \in (f_1, f_3, f_5) \cap A, s \in b(\alpha)} \mu_s = (f_1, f_3, f_5) \mu_b = \{0.5, 0.1\},$$

$$\chi(P_A)(f_1', f_3', f_4')(y) = \{0.1\},$$

$$\chi(P_A)(f_1', f_3', f_4')(z) = \{0.3, 0.9\},$$

$$\chi(P_A)(f'_2, f'_3, f'_4)(x) = \{0.9, 0.3, 0.5\},$$

$$\chi(P_A)(f'_2, f'_3, f'_4)(y) = \{0.5, 0.3\},$$

$$\chi(P_A)(f'_2, f'_3, f'_4)(z) = \{0.3\}$$

$$\chi(P_A) = \left\{ \left\{ (f'_1, f'_3, f'_4) = \left\{ \begin{array}{l} < x, \{0.5, 0.1\} >, \\ < y, \{0.1\} >, \\ < z, \{0.3, 0.9\} > \end{array} \right\} \right\}, \right. \\ \left. \left\{ (f'_2, f'_3, f'_5) = \left\{ \begin{array}{l} < x, \{0.9, 0.3, 0.5\} >, \\ < y, \{0.3, 0.5\} >, \\ < z, \{0.3\} > \end{array} \right\} \right\} \right\}$$

Again,

$$\phi(\chi(P_A))(f''_1, f''_2, f''_3)(h_1) = \cup_{\alpha \in n^{-1}(f''_1, f''_2, f''_3) \cap A, s \in m^{-1}(h_1)} (\alpha) \mu_s$$

$$= \cup_{\alpha \in (f'_1, f'_3, f'_4) \cap \chi(A), s \in z} (\alpha) \mu_s = (f'_1, f'_3, f'_4) \mu_z = \{0.9, 0.3\},$$

$$\phi(\chi(P_A))(f''_1, f''_2, f''_3)(h_2) = \{0.5, 0.1\},$$

$$\phi(\chi(P_A))(f''_1, f''_2, f''_3)(h_3) = \{0.1\},$$

$$\phi(\chi(P_A))(f''_1, f''_2, f''_4)(h_1) = \{0.3\},$$

$$\phi(\chi(P_A))(f''_1, f''_2, f''_4)(h_2) = \{0.9, 0.3, 0.5\},$$

$$\phi(\chi(P_A))(f''_1, f''_2, f''_4)(h_3) = \{0.3, 0.5\},$$

$$(\phi \circ \chi)(P_A) = \phi(\chi(P_A)) = \left\{ \left\{ \begin{array}{l} (f''_1, f''_2, f''_4) = \{ < h_1, \{0.9, 0.3\} >, \\ < h_2, \{0.5, 0.1\} >, < h_3, \{0.1\} > \end{array} \right\} \right\} \\ \left\{ \left\{ \begin{array}{l} (f''_1, f''_2, f''_4) = \{ < h_1, \{0.3\} >, \\ < h_2, \{0.9, 0.3, 0.5\} >, < h_3, \{0.3, 0.5\} > \end{array} \right\} \right\} \right\}$$

Therefore, from above example we see that, if χ and ϕ are one-one then so is $\phi \circ \chi$. Similarly, for onto as well as bijections.

Theorem 3.14. Let us consider three FHS mappings $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$, $\phi : \overline{(Y, M)} \rightarrow \overline{(Z, N)}$ and $\varphi : \overline{(Z, N)} \rightarrow \overline{(A, O)}$. Then $\varphi \circ (\phi \circ \chi) = (\varphi \circ \phi) \circ \chi$.

Proof. Let $P_A \in \overline{(X, L)}$, now from definition we have, $[\varphi \circ (\phi \circ \chi)](P_A) = (\varphi \circ \phi) \circ \chi(P_A) = \varphi[\phi(\chi(P_A))]$, also $[(\varphi \circ \phi) \circ \chi](P_A) = (\varphi \circ \phi)(\chi(P_A)) = \varphi[\phi(\chi(P_A))]$. Hence $\varphi \circ (\phi \circ \chi) = (\varphi \circ \phi) \circ \chi$.

Theorem 3.15. A FHS mapping $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ is said to be many one FHS mapping if two (or more than two) FHS sets in $\overline{(X, L)}$ have the same FHS image in $\overline{(Y, M)}$.

Example 3.16. From example 3.6, consider the FHSS $Q_A \in \overline{(X, L)}$,

$$Q_A = \left\{ \left\{ (f_1, f_3, f_4) = \begin{cases} < a, \{0.5, 0.1\} >, \\ < b, \{0.9, 0.3, 0.5\} >, \\ < c, \{0.3\} > \end{cases} \right\}, \right. \\ \left. \left\{ (f_1, f_3, f_5) = \begin{cases} < a, \{0.3, 0.9\} >, \\ < b, \{0.5, 0.1\} >, \\ < c, \{0.1\} > \end{cases} \right\} \right\}$$

Then the FHS image of P_A under $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ can be written as

$$\chi(Q_A) = \left\{ \left\{ (f'_1, f'_3, f'_4) = \begin{cases} \{ < x, \{0.5, 0.1\} >, \\ < y, \{0.3, 0.9\} >, \\ < z, \{0\} > \} \right\}, \right. \\ \left. \left\{ (f'_2, f'_3, f'_5) = \begin{cases} < x, \{0.9, 0.3, 0.5\} >, \\ < y, \{0.3, 0.5\} >, \\ < z, \{0\} > \end{cases} \right\} \right\}$$

Therefore $\chi(P_A) = \chi(Q_A)$. Hence χ is many one FHS mapping.

Definition 3.17. Let $i = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(X, L)}$ be a FHS mapping, where $\omega : X \rightarrow X$ and $v : L \rightarrow L$. Then χ is said to be a FHS identity mapping if both ω and v are identity mappings.

Remark 3.18. $i = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(X, L)}$ be a FHS identity mapping, then $i(P_A) = P_A$, where $P_A \in \overline{(X, L)}$.

Definition 3.19. Let $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ be a FHS mapping and let $i = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(X, L)}$ and $j = (r, t) : \overline{(Y, M)} \rightarrow \overline{(Y, M)}$ are FHS identity mappings then $\chi \circ i = \chi$ and $j \circ \chi = \chi$.

Example 3.20. Consider the following example, we consider P_A from example 3.6 and consider the FHS mappings $i = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(X, L)}$, where $\omega : X \rightarrow X$ and $v : L \rightarrow L$, such that $\omega(a) = a, \omega(b) = b, \omega(c) = c$, and

$$v(f_1, f_3, f_4) = (f_1, f_3, f_4), v(f_1, f_3, f_5) = (f_1, f_3, f_5)$$

$$v(f_2, f_3, f_4) = (f_2, f_3, f_4), v(f_2, f_3, f_5) = (f_2, f_3, f_4)$$

Therefore,

$$i(P_A)(f_1, f_3, f_4)(a) = \cup_{\alpha \in v^{-1}(f_1, f_3, f_4) \cap A, s \in \omega^{-1}(a)} (\alpha) \mu_s$$

$$\cup_{\alpha \in (f_1, f_3, f_4), s \in \{a\}} (\alpha) \mu_s = (f_1, f_3, f_4) \mu_a = \{0.5, 0.1\}$$

$$i(P_A)(f_1, f_3, f_4)(b) = \{0.9, 0.3, 0.5\},$$

$$i(P_A)(f_1, f_3, f_4)(c) = \{0.3\},$$

So,

$$i(P_A)(f_1, f_3, f_5)(a) = \{0.3, 0.9\},$$

$$i(P_A)(f_1, f_3, f_5)(b) = \{0.5, 0.1\},$$

$$i(P_A)(f_1, f_3, f_5)(c) = \{0.1\}$$

we get,

$$i(P_A) = \left\{ \left\{ (f_1, f_3, f_4) = \begin{cases} < a, \{0.5, 0.3\} >, \\ < b, \{0.9, 0.3, 0.5\} >, \\ < c, \{0.3\} > \end{cases} \right\} \right\}, \left\{ \left\{ (f_1, f_3, f_5) = \begin{cases} < a, \{0.3, 0.9\} >, \\ < b, \{0.5, 0.1\} >, \\ < c, \{0.1\} > \end{cases} \right\} \right\} \right\}$$

Hence $i(P_A) = P_A \Rightarrow \chi(i(P_A)) = \chi(P_A) \Rightarrow (\chi o i)(P_A) = \chi(P_A) \Rightarrow \chi o i = \chi$.

Similarly, we get $\chi(P_A) \in \overline{(X, L)}$ and $j(\chi(P_A)) = (\chi(P_A)) \Rightarrow (j o \chi)(P_A) = \chi(P_A)$.

Hence $j o \chi = \chi$.

Definition 3.21. A one-one onto FHS mapping $\chi = (\omega, v) : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ is called FHS invertible mapping. Its FHS inverse mapping is denoted by $\chi^{-1} = (\omega^{-1}, v^{-1}) : \overline{(Y, M)} \rightarrow \overline{(X, L)}$.

Remark 3.22. In a FHS invertible mapping $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$, for $P_A \in \overline{(X, L)}, P_B \in \overline{(Y, M)}, \chi^{-1}(P_B) = P_A$, whenever $\chi(P_A) = P_B$.

Example 3.23. We consider $\chi(L_A) = P_B$ (see 3.13). Therefore,

$$\chi^{-1}(P_B)(f_1, f_3, f_4)(a) = (v(f_1, f_3, f_4)\mu_{\omega(a)} = (f'_1, f'_3, f'_4)\mu_z = \{0.3, 0.9\}$$

$$\chi^{-1}(P_B)(f_1, f_3, f_4)(b) = \{0.5, 0.1\},$$

$$\chi^{-1}(P_B)(f_1, f_3, f_4)(c) = \{0.1\},$$

So,

$$\chi^{-1}(P_B)(f_1, f_3, f_5)(a) = \{0.3\},$$

$$\chi^{-1}(P_B)(f_1, f_3, f_5)(b) = \{0.9, 0.3, 0.5\},$$

$$\chi^{-1}(P_B)(f_1, f_3, f_5)(c) = \{0.3, 0.5\}$$

we get,

$$\chi^{-1}(P_B) = \left\{ \left\{ (f_1, f_3, f_4) = \begin{Bmatrix} < a, \{0.9, 0.3\} >, \\ < b, \{0.5, 0.1\} >, \\ < c, \{0.1\} > \end{Bmatrix} \right\}, \right. \\ \left. \left\{ (f_1, f_3, f_5) = \begin{Bmatrix} < a, \{0.3\} >, \\ < b, \{0.9, 0.3, 0.5\} >, \\ < c, \{0.3, 0.5, \} > \end{Bmatrix} \right\} \right\}$$

Hence, $\chi^{-1}(P_B) = L_A$.

Theorem 3.24. Let $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ be a FHS invertible mapping. Therefore its FHS inverse mapping is unique.

Proof. Let χ^{-1} and ϕ^{-1} are two FHS inverse mappings of χ . Therefore, $\chi^{-1}(P_B) = P_A$, whenever $\chi(P_A) = P_B, P_A \in \overline{(X, L)}, P_B \in \overline{(Y, M)}$, and $\phi^{-1}(P_B) = H_A$, whenever $\phi(H_A) = P_B, H_A \in \overline{(X, L)}, P_B \in \overline{(Y, M)}$. Thus $\chi(P_A) = \phi(H_A)$. Since χ is one-one, therefore $P_A = H_A$. Hence $\chi^{-1}(P_B) = \phi^{-1}(P_B)$ i.e $\chi^{-1} = \phi^{-1}$.

3.1. Theorem

Let $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ and $\phi : \overline{(Y, M)} \rightarrow \overline{(Z, N)}$ are two one-one onto FHS mappings. If $\chi^{-1} : \overline{(Y, M)} \rightarrow \overline{(X, L)}$ and $\phi^{-1} : \overline{(Z, N)} \rightarrow \overline{(Y, M)}$ are FHS inverse mappings of χ and ϕ , respectively, then the inverse of the mapping $\phi \circ \chi : \overline{(X, L)} \rightarrow \overline{(Z, N)}$ is the FHS mapping $\chi^{-1} \circ \phi^{-1} : \overline{(Z, N)} \rightarrow \overline{(X, L)}$. For more detail see fig 6

Proof. Obvious.

Theorem 3.25. A FHS mapping $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ is invertible if and only if there exists a FHS inverse mapping $\chi^{-1} : \overline{(Y, M)} \rightarrow \overline{(X, L)}$ such that $\chi^{-1} \circ \chi = i_{\overline{(X, L)}}$ and $\chi \circ \chi^{-1} = i_{\overline{(Y, M)}}$, where $i_{\overline{(X, L)}}$ and $i_{\overline{(Y, M)}}$ is FHS identity mapping on $\overline{(X, L)}$ and $\overline{(Y, M)}$, respectively.

Proof. Let $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ be a FHS invertible mapping. Therefore, by definition we have $\chi^{-1}(P_B) = P_A$, whenever $\chi(P_A) = P_B, P_A \in \overline{(X, L)}, P_B \in \overline{(Y, M)}$. Since $(\chi^{-1} \circ \chi)(P_A) = \chi^{-1}(\chi(P_A)) = \chi^{-1}(P_B) = P_A$. Therefore, $\chi^{-1} \circ \chi = i_{\overline{(X, L)}}$. Similarly, we prove that $\chi \circ \chi^{-1} = i_{\overline{(Y, M)}}$.

Theorem 3.26. If $\chi : \overline{(X, L)} \rightarrow \overline{(Y, M)}$ and $\phi : \overline{(Y, M)} \rightarrow \overline{(Z, N)}$ are two one-one onto FHS mapping then $(\phi \circ \chi)^{-1} = \chi^{-1} \circ \phi^{-1}$.

Proof. Since χ and ϕ are one-one onto FHS mapping, then there exists $\chi^{-1} : \overline{(Y, M)} \rightarrow \overline{(X, L)}$ and $\phi^{-1} : \overline{(Z, N)} \rightarrow \overline{(Y, M)}$ such that $\chi^{-1}(P_B) = P_A$, whenever $\chi(P_A) = P_B, P_A \in \overline{(X, L)}, P_B \in \overline{(Y, M)}$, and $\phi^{-1}(H_C) = P_B$, whenever $\phi(P_B) = H_C, H_C \in \overline{(Z, N)}, P_B \in \overline{(Y, M)}$. Therefore, $(\phi \circ \chi)(P_A) = \phi[\chi(P_A)] = \phi(P_B) = H_C$. As $\phi \circ \chi$ is one-one onto, therefore




FIGURE 6. Representation of Composite Fuzzy Hypersoft Inverse Mapping

$(\phi \circ \chi)^{-1}$ exists such that $(\phi \circ \chi)(P_A) = H_C \Rightarrow (\phi \circ \chi)^{-1}(H_C) = P_A$. Also $(\phi^{-1} \circ \phi^{-1})(H_C) = \chi^{-1}[\phi^{-1}(H_C)] = \chi^{-1}(P_B) = P_A$. Hence $(\phi \circ \chi)^{-1}(H_C) = (\chi^{-1} \circ \phi^{-1})(H_C) \Rightarrow (\phi \circ \chi)^{-1} = \chi^{-1} \circ \phi^{-1}$.

4. Conclusion

A basic structure of composite mapping of FHS classes is established along with generalization of certain properties and results. It is very helpful for solving problems involving uncertainty and vagueness. In the future, we will expand our exploration in the domain of Neutrosophic Hypersoft Set, Plithogenic Crisp Hypersoft Set, Plithogenic Fuzzy Hypersoft Set, Plithogenic Intuitionistic Fuzzy Hypersoft Set, Plithogenic Neutrosophic Hypersoft Set, complex Intuitionistic Fuzzy Hypersoft Set, complex Neutrosophic Hypersoft Set, and their hybrid structures. We will apply them in medical imaging issues, pattern recognition, recommender frameworks, social, the monetary framework estimated thinking, image processing, and game theory.

References

- [1] Kharal, A., and Ahmad, B. (2009). Mappings on fuzzy soft classes. *Advances in Fuzzy Systems*, 2009.
- [2] Atanassov, K. T. (1989). More on intuitionistic fuzzy sets. *Fuzzy sets and systems*, 33(1), 37-45.

Muhammad Ahsan, Muhammad Saeed, Atiqe Ur Rahman, A Theoretical and Analytical Approach for Fundamental Framework of Composite mappings on Fuzzy Hypersoft Classes

- [3] Kharal, A., and Ahmad, B. (2011). Mappings on soft classes. *New Mathematics and Natural Computation*, 7(03), 471-481.
- [4] Kharal, A., and Ahmad, B. (2009). Mappings on Fuzzy Soft Classes. *Advances in Fuzzy Systems*, 2009(1), 1-6. doi:10.1155/2009/40789.
- [5] Atanassov, K. T. (1994). Operators over interval valued intuitionistic fuzzy sets. *Fuzzy sets and systems*, 64(2), 159-174.
- [6] Behounek, L., and Cintula, P. (2005). *Fuzzy class theory*. *Fuzzy Sets and Systems*, 154(1), 34-55.
- [7] Borah, M. J., and Hazarika, B. (2016). Composite mapping on hesitant fuzzy Soft Classes. *arXiv preprint arXiv:1605.01304*.
- [8] Karaaslan, F. (2016). Soft classes and soft rough classes with applications in decision making. *Mathematical problems in engineering*, 2016(1), 1-12. doi:10.1155/2016/1584528.
- [9] Smarandache, F. (2018). Extension of soft set to hypersoft set, and then to plithogenic hypersoft set. *Neutrosophic Sets and Systems*, 22(1), 168-170.
- [10] Gau, W. L., and Buehrer, D. J. (1993). Vague sets. *IEEE transactions on systems, man, and cybernetics*, 23(2), 610-614.
- [11] Gorzalczany, M. B. (1987). *A method of inference in approximate reasoning based on interval-valued fuzzy sets*. *Fuzzy sets and systems*, 21(1), 1-17.
- [12] Molodtsov, D. (1999). Soft set theory first results. *Computers and Mathematics with Applications*, 37(4-5), 19-31.
- [13] Pawlak, Z. (1982). Rough sets. *International journal of computer and information sciences*, 11(5), 341-356.
- [14] Pawlak, Z. (1994). Hard set and soft sets. ICS Research Report.
- [15] Rana, S., Qayyum, M., Saeed, M., Smarandache, F., and Khan, B. A. (2019). Plithogenic fuzzy whole hypersoft set, construction of operators and their application in frequency matrix multi attribute decision making technique. *Neutrosophic Sets and Systems*, 28(1), 34-50.
- [16] Saqlain, M., Jafar, N., Moin, S., Saeed, M., and Broumi, S. (2020). Single and multi-valued neutrosophic hypersoft set and tangent similarity measure of single valued neutrosophic hypersoft sets. *Neutrosophic Sets and Systems*, 32(1), 317-329.
- [17] Saqlain, M., Moin, S., Jafar, N., Saeed, M., and Smarandache, F. (2020). Aggregate Operators of Neutrosophic Hypersoft Set, *Neutrosophic Sets and Systems*, 32(1), 294-306.
- [18] Rahman, A. U., Saeed, M., Smarandache, F., and Ahmad, M. R. (2020). Development of Hybrids of Hypersoft Set with Complex Fuzzy Set, Complex Intuitionistic Fuzzy set and Complex Neutrosophic Set, *Neutrosophic Sets and Systems*, 38(1), 335-354.
- [19] Saeed, M., Ahsan, M., Siddique, M. K., and Ahmad, M. R. (2020). A study of the fundamentals of hypersoft set theory. *International Journal of Scientific and Engineering Research*, 11(1), 2229-5518.
- [20] Yager, R. R. (1978). Fuzzy decision making including unequal objectives. *Fuzzy sets and systems*, 1(2), 87-95.
- [21] Zadeh, L. A. (1973). Outline of a new approach to the analysis of complex systems and decision processes. *IEEE Transactions on systems, Man, and Cybernetics*, (1), 28-44.
- [22] Zimmermann, H. J. (1987). Fuzzy sets, decision making, and expert systems, *Springer Science and Business Media*, 1, 71-124.
- [23] Saeed, M., Ahsan, M., and Rahman, A.U. (2020). A Novel Approach to Mappings on Hypersoft Classes with Application, *Neutrosophic Sets and Systems*, 38(1), 175-191.
- [24] Zadeh, L. A. (1965). Fuzzy sets. *Information and Control*, 8(3), 338-353. doi:https://doi.org/10.1016/S0019-9958(65)90241-X.

- [25] Saeed, M., Rahman, A. U., Ahsan, M., and Smarandache, F. (2021). *An Inclusive Study on Fundamentals of Hypersoft Set*. In F. Smarandache, M. Saeed, M. Abdel-Baset and M. Saqlain (Eds) *Theory and Application of Hypersoft Set*(pp. 1-23). Belgium, Brussels: Pons Publishing House. ISBN 978-1-59973-699-0.

Received: May 2, 2021. Accepted: August 3, 2021