

8-28-2021

## Types of System of the neutrosophic linear equations and Cramer's rule

Yaser Ahmad Alhasan

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Alhasan, Yaser Ahmad. "Types of System of the neutrosophic linear equations and Cramer's rule." *Neutrosophic Sets and Systems* 45, 1 (). [https://digitalrepository.unm.edu/nss\\_journal/vol45/iss1/26](https://digitalrepository.unm.edu/nss_journal/vol45/iss1/26)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).



# Types of System of the neutrosophic linear equations and Cramer's rule

Yaser Ahmad Alhasan

Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, KSA; y.alhasan@psau.edu.sa

**Abstract:** The purpose of this article is studying the types of system of neutrosophic linear equations, where the neutrosophic linear equation and the solution to the neutrosophic linear equation are defined. Also, finding a solution to a linear equation with two variables, the general situation of the solution. In addition to studying the n-variable neutrosophic linear equation and the system of neutrosophic homogeneous linear equations. The most important is the introduction of the concept of Cramer's rule to solve the system of neutrosophic linear equations. Provide enough examples for each case to enhance understanding.

**Keywords:** neutrosophic liner equation; Cramer's rule; solution of the neutrosophic linear equation.

## 1. Introduction

As an alternative to the existing logics, Smarandache proposed the Neutrosophic Logic to represent a mathematical model of uncertainty, vagueness, ambiguity, imprecision, undefined, unknown, incompleteness, inconsistency, redundancy, contradiction, where the concept of neutrosophy is a new branch of philosophy introduced by Smarandache [3-13]. He presented the definition of the standard form of neutrosophic real number and conditions for the division of two neutrosophic real numbers to exist, he defined the standard form of neutrosophic complex number, and found root index  $n \geq 2$  of a neutrosophic real and complex number [2-4], studying the concept of the Neutrosophic probability [3-5], the Neutrosophic statistics [4][6], and professor Smarandache entered the concept of preliminary calculus of the differential and integral calculus, where he introduced for the first time the notions of neutrosophic mereo-limit, mereo-continuity, mereoderivative, and mereo-integral [1-8]. Madeleine Al- Taha presented results on single valued neutrosophic (weak) polygroups [9].Edalatpanah proposed a new direct algorithm to solve the neutrosophic linear programming where the variables and right-hand side represented with triangular neutrosophic numbers [10]. Chakraborty used pentagonal neutrosophic number in networking problem, and Shortest Path Problem [11-12]. Y.Alhasan studied the concepts of neutrosophic complex numbers, the general exponential form of a neutrosophic complex, and the neutrosophic integrals and integration methods [7-14-17]. On the other hand, M.Abdel-Basset presented study in the science of neutrosophic about an approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic

number [15]. Also, neutrosophic sets played an important role in applied science such as health care, industry, and optimization [16]. Recently, Alhasan, Y., and Alfahal, A presented study in the neutrosophic differential equations that translate into linear [18].

Mathematical equations are used to solve real problems in our daily life, for example, mathematical equations are used in electronic chips used in all modern machines and devices, such as washing machines and dryers, cars, airplanes, ships, cell phones, computers, space programs and so on. One may be surprised when he learns that there are about 2 million algorithms and mathematical equations in mobile and computer devices.

The paper consists of 5 sections. In 1th section, provides an introduction, in which neutrosophic science review has given. In 2th section, some definitions and examples of neutrosophic real number. The 3th section frames studying types of system of the neutrosophic linear equations. The 4th section introduces the concept of Cramer's rule to solve the system of neutrosophic linear equations. In 5th section, a conclusion to the paper is given.

## 2. Preliminaries

### 2.1. Neutrosophic Real Number [4]

Suppose that  $w$  is a neutrosophic number, then it takes the following standard form:  $w = a + bI$  where  $a, b$  are real coefficients, and  $I$  represent indeterminacy, such  $0.I = 0$  and  $I^n = I$ , for all positive integers  $n$ .

### 2.2. Division of neutrosophic real numbers [4]

Suppose that  $w_1, w_2$  are two neutrosophic numbers, where

$$w_1 = a_1 + b_1I, \quad w_2 = a_2 + b_2I$$

To find  $(a_1 + b_1I) \div (a_2 + b_2I)$ , we can write:

$$\frac{a_1 + b_1I}{a_2 + b_2I} \equiv x + yI$$

where  $x$  and  $y$  are real unknowns.

$$a_1 + b_1I \equiv (a_2 + b_2I)(x + yI)$$

$$a_1 + b_1I \equiv a_2x + (b_2x + a_2y + b_2y)I$$

by identifying the coefficients, we get

$$a_1 = a_2x$$

$$b_1 = b_2x + (a_2 + b_2)y$$

We obtain unique one solution only, provided that:

$$\begin{vmatrix} a_2 & 0 \\ b_2 & a_2 + b_2 \end{vmatrix} \neq 0 \Rightarrow a_2(a_2 + b_2) \neq 0$$

Hence:  $a_2 \neq 0$  and  $a_2 \neq -b_2$  are the conditions for the division of two neutrosophic real numbers to exist.

Then:

$$\frac{a_1 + b_1I}{a_2 + b_2I} = \frac{a_1}{a_2} + \frac{a_2b_1 - a_1b_2}{a_2(a_2 + b_2)} \cdot I$$

## 3. The neutrosophic linear equation

**Defination3.1:**

The neutrosophic linear equation of  $n$  variables  $x_1, x_2, x_3, \dots, x_n$ , is each equation that takes the form:

$$(a_1 + b_1I)x_1 + (a_2 + b_2I)x_2 + (a_3 + b_3I)x_3 + \dots + (a_n + b_nI)x_n = c + dI$$

Where:

$a_1, a_2, a_3, \dots, a_n, b_1, b_2, b_3, \dots, b_n, c, d$  are real coefficients, and  $I$  represent indeterminacy.

We call  $(a_1 + b_1I), (a_2 + b_2I), (a_3 + b_3I), \dots, (a_n + b_nI)$  neutrosophic coefficients of the borders of the equation, and  $c + dI$  constant neutrosophic border of the equation.

**Remarks3.1:**

We call each equation of the form:

$$(a_1 + b_1I)x + (a_2 + b_2I)y = c + dI$$

the two-variable neutrosophic linear equation, where:

$a_1, a_2, a_3, \dots, a_n, b_1, b_2, b_3, \dots, b_n, c, d$  are real coefficients, and  $I$  represent indeterminacy.

**Remarks3.2:**

We call each equation of the form:

$$(a_1 + b_1I)x + (a_2 + b_2I)y + (a_3 + b_3I)z = c + dI$$

the three-variable neutrosophic linear equation, where:

$a_1, a_2, a_3, \dots, a_n, b_1, b_2, b_3, \dots, b_n, c, d$  are real coefficients, and  $I$  represent indeterminacy.

**Example3.1:**

- ✓  $(2 + 3I)x + (4 - 5I)y + (-1 + I)z = 5 + 4I$
- ✓  $(3 - I)x + (9 - 5I)y + (-1 + I)z + (3 - 2I)w = 7 - I$
- ✓  $(1 + I)x + (2 - 5I)y = 6 - 2I$

**Defination3.2:**

Solution of the neutrosophic linear equation:

$$(a_1 + b_1I)x_1 + (a_2 + b_2I)x_2 + (a_3 + b_3I)x_3 + \dots + (a_n + b_nI)x_n = c + dI$$

is finding the values of the variables  $x_1, x_2, x_3, \dots, x_n$  that make its first term equal to its second term.

where  $a_1, a_2, b_1, b_2, c, d$  are real coefficients.

**Example3.2:**

The equation:  $(1 + I)x + (2 - 5I)y = 6 - 2I$  accepts the solution

$$x = 1 - I, \quad y = \frac{5}{2} - \frac{23}{6}I$$

because fulfills the equation:

$$L_1 = (1 + I)(1 - I) + (2 - 5I)\left(\frac{5}{2} - \frac{23}{6}I\right) = 1 - I + 5 - \frac{46}{6}I - \frac{25}{2}I + \frac{115}{6}I = 6 - 2I = L_2$$

**3.1 Solution of the two-variables neutrosophic linear equation**

For the neutrosophic linear equation:

$$(a_1 + b_1I)x + (a_2 + b_2I)y = c + dI$$

unlimited number of solutions defined by form:

$$S = \left\{ (x, y) \in R^2 \cup \{I\}: y = \left( \frac{a_1}{a_2} + \frac{a_2b_1 - a_1b_2}{a_2(a_2 + b_2)} \cdot I \right) x + \frac{c}{a_2} + \frac{a_2d - cb_2}{a_2(a_2 + b_2)} \cdot I \right\}$$

where  $a_1, a_2, b_1, b_2, c, d$  are real coefficients,  $a_2 \neq 0$  and  $a_2 \neq -b_2$ , by given a value for one of the two variables, we obtain a value for the other variable.

**Example3.1.1:**

Find solution of the equation:

$$(1 + I)x + (2 - 5I)y = 6 - 2I$$

Solution:

$$y = \frac{-(1 + I)}{(2 - 5I)}x + \frac{6 - 2I}{(2 - 5I)}$$

$$y = \left(\frac{-1}{2} + \frac{7}{6}I\right)x + 3 + \frac{13}{3}I$$

Then the set of solutions is:

$$S = \left\{ (x, y) \in R^2 \cup \{I\} : y = \left(\frac{-1}{2} + \frac{7}{6}I\right)x + 3 + \frac{13}{3}I \right\}$$

By given any value for the variables  $x$ , we obtain a value of the variable  $y$ .

### 3.2 General situation: Solution of the n-variable neutrosophic linear equation

For the neutrosophic linear equation:

$$(a_1 + b_1I)x_1 + (a_2 + b_2I)x_2 + (a_3 + b_3I)x_3 + \dots + (a_n + b_nI)x_n = c + dI$$

unlimited number of solutions, where  $a_1, a_2, a_3, \dots, a_n, b_1, b_2, b_3, \dots, b_n, c, d$  are real coefficients.

### 4. System of the neutrosophic linear equations

It is a system of neutrosophic linear equations that given by the form:

$$\begin{cases} (a_{11} + b_{11}I)x_1 + (a_{12} + b_{12}I)x_2 + (a_{13} + b_{13}I)x_3 + \dots + (a_{1n} + b_{1n}I)x_n = c_1 + d_1I \\ (a_{21} + b_{21}I)x_1 + (a_{22} + b_{22}I)x_2 + (a_{23} + b_{23}I)x_3 + \dots + (a_{2n} + b_{2n}I)x_n = c_2 + d_2I \\ \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots = \dots \\ \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots = \dots \\ (a_{m1} + b_{m1}I)x_1 + (a_{m2} + b_{m2}I)x_2 + (a_{m3} + b_{m3}I)x_3 + \dots + (a_{mn} + b_{mn}I)x_n = c_m + d_mI \end{cases}$$

Where:

$a_{ij}, b_{ij}, c_j, d_j$  are real coefficients,  $i = 1, \dots, n, j = 1, \dots, m$ , and  $I$  represent indeterminacy.

#### 4.1 Solution of system of the neutrosophic linear equations

Solution of system of the neutrosophic linear equation:

$$\begin{cases} (a_{11} + b_{11}I)x_1 + (a_{12} + b_{12}I)x_2 + (a_{13} + b_{13}I)x_3 + \dots + (a_{1n} + b_{1n}I)x_n = c_1 + d_1I \\ (a_{21} + b_{21}I)x_1 + (a_{22} + b_{22}I)x_2 + (a_{23} + b_{23}I)x_3 + \dots + (a_{2n} + b_{2n}I)x_n = c_2 + d_2I \\ \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots = \dots \\ \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots = \dots \\ (a_{m1} + b_{m1}I)x_1 + (a_{m2} + b_{m2}I)x_2 + (a_{m3} + b_{m3}I)x_3 + \dots + (a_{mn} + b_{mn}I)x_n = c_m + d_mI \end{cases} \quad (*)$$

Where:

$a_{ij}, b_{ij}, c_j, d_j$  are real coefficients,  $i = 1, \dots, n, j = 1, \dots, m$ , and  $I$  represent indeterminacy.

is finding the values of the variables  $x_1, x_2, x_3, \dots, x_n$  that fulfill each of the system equations. We call this solution the system solution.

#### Remarks4.1.1:

We distinguish three cases to solve the system(\*):

1. It may have unique solution
2. It may be impossible to solve
3. It may have Unlimited number of solutions

#### 4.2 Cramer's rule to solve system of the neutrosophic linear equations

Let the system of the neutrosophic linear equations:



- If  $\det(x_i) = 0 + 0I$ ;  $i = 1, 2, \dots, n$ , then the system has unlimited number of solutions.

**Note:** To verify the solution, we substitute the value of  $x$  and  $y$  into one of the equations.

**Example4.2.1:**

Let:

$$\begin{aligned}(2 + I)x + 3y &= 5 + I \\ (3 - 2I)x + 2y &= I\end{aligned}$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 2 + I & 3 \\ 3 - 2I & 2 \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2 + I & 3 \\ 3 - 2I & 2 \end{vmatrix} = 4 + 2I - 9 + 6I = -5 + 8I \neq 0 + 0I$$

Then the system has one solution, is:

$$\det x = \begin{vmatrix} 5 + I & 3 \\ I & 2 \end{vmatrix} = 10 - I, \quad \det y = \begin{vmatrix} 2 + I & 5 + I \\ 3 - 2I & I \end{vmatrix} = -15 + 12I$$

$$x = \frac{\det x}{\det A} = \frac{10 - I}{-5 + 8I} = -2 + 5I \quad \text{and} \quad y = \frac{\det y}{\det A} = \frac{-15 + 12I}{-5 + 8I} = 3 - 4I$$

$$(x, y) = (-2 + 5I, 3 - 4I)$$

To verify the solution, we substitute the value of  $x$  and  $y$  into the first equation:

$$\begin{aligned}(2 + I)x + 3y &= (2 + I)(-2 + 5I) + 3(3 - 4I) \\ &= -4 - 2I + 10I + 5I + 9 - 12I = 5 + I\end{aligned}$$

**Example4.2.2:**

Let:

$$\begin{aligned}2Ix + 7y &= I \\ 3Ix + y &= 2I\end{aligned}$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 2I & 7 \\ 3I & 1 \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2I & 7 \\ 3I & 1 \end{vmatrix} = 2I - 21I = 0 - 19I$$

This does not fulfill the condition:  $\det A = a + bI$ ,  $a \neq 0$  and  $a \neq -b$

Because:

$$\det x = \begin{vmatrix} I & 7 \\ 2I & 1 \end{vmatrix} = I - 14I = 0 - 13I$$

Then:

$$x = \frac{\det x}{\det A} = \frac{0 - 13I}{0 - 19I} \text{ (undefined)}$$

So, the system is impossible to solve.

**Example4.2.3:**

Let:

$$\begin{aligned}(2 + I)x + 3y &= 5 + I \\ (3 - 2I)x + y &= I\end{aligned}$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 2 + I & 3 \\ 3 - 2I & 1 \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2 + I & 3 \\ 3 - 2I & 1 \end{vmatrix} = 2 + I - 9 + 6I = -7 + 7I$$

W This does not fulfill the condition:  $\det A = a + bI$ ,  $a \neq 0$  and  $a \neq -b$ , where: ( $a = -7, b = 7$  and  $a = -b$ ).

So, the system is impossible to solve.

#### Example4.2.4:

Let:

$$(1 + I)x + (3 - I)y = 2 - 3I$$

$$(2 + 2I)x + (6 - 2I)y = 4 - 6I$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 1 + I & 3 - I \\ 2 + 2I & 6 - 2I \end{bmatrix}$$

$$\det A = \begin{vmatrix} 1 + I & 3 - I \\ 2 + 2I & 6 - 2I \end{vmatrix} = 0 + 0I$$

$$\det x = \begin{vmatrix} 2 - 3I & 3 - I \\ 4 - 6I & 6 - 2I \end{vmatrix} = 0 + 0I, \quad \det y = \begin{vmatrix} 1 + I & 2 - 3I \\ 2 + 2I & 4 - 6I \end{vmatrix} = 0 + 0I$$

As:  $\det A = \det x = \det y = 0 + 0I$ , so the system has Unlimited number of solutions defined by form:

$$S = \left\{ (x, y) \in R^2 \cup \{I\}: y = \left( \frac{1}{3} + \frac{2}{3} \cdot I \right) x + \frac{2}{3} - \frac{7}{6} \cdot I \right\}$$

By given any value for the variable  $x$ , we obtain a value of the variable  $y$ .

#### 4.2.2 Solve the system of three linear equations by three variables

Let:

$$(a_{11} + b_{11}I)x + (a_{12} + b_{12}I)y + (a_{13} + b_{13}I)z = c_1 + d_1I \quad (1)$$

$$(a_{21} + b_{21}I)x + (a_{22} + b_{22}I)y + (a_{23} + b_{23}I)z = c_2 + d_2I \quad (2)$$

$$(a_{33} + b_{33}I)x + (a_{33} + b_{33}I)y + (a_{33} + b_{33}I)z = c_3 + d_3I \quad (3)$$

$$A = \begin{bmatrix} a_{11} + b_{11}I & a_{12} + b_{12}I & a_{13} + b_{13}I \\ a_{21} + b_{21}I & a_{22} + b_{22}I & a_{23} + b_{23}I \\ a_{33} + b_{33}I & a_{33} + b_{33}I & a_{33} + b_{33}I \end{bmatrix}$$

$$\det A = \begin{vmatrix} a_{11} + b_{11}I & a_{12} + b_{12}I & a_{13} + b_{13}I \\ a_{21} + b_{21}I & a_{22} + b_{22}I & a_{23} + b_{23}I \\ a_{33} + b_{33}I & a_{33} + b_{33}I & a_{33} + b_{33}I \end{vmatrix}$$

$$\det x = \begin{vmatrix} c_1 + d_1I & a_{12} + b_{12}I & a_{13} + b_{13}I \\ c_2 + d_2I & a_{22} + b_{22}I & a_{23} + b_{23}I \\ c_3 + d_3I & a_{33} + b_{33}I & a_{33} + b_{33}I \end{vmatrix}, \quad \det y = \begin{vmatrix} a_{11} + b_{11}I & c_1 + d_1I & a_{13} + b_{13}I \\ a_{21} + b_{21}I & c_2 + d_2I & a_{23} + b_{23}I \\ a_{33} + b_{33}I & c_3 + d_3I & a_{33} + b_{33}I \end{vmatrix}$$

$$\det z = \begin{vmatrix} a_{11} + b_{11}I & a_{12} + b_{12}I & c_1 + d_1I \\ a_{21} + b_{21}I & a_{22} + b_{22}I & c_2 + d_2I \\ a_{33} + b_{33}I & a_{33} + b_{33}I & c_3 + d_3I \end{vmatrix}$$

We will discuss the second case of (3):



If  $\det(A) = 0 + 0I$  and  $\det(x_i) = 0 + 0I$ ;  $i = 1, 2, \dots, n$ , then we are looking for the solutions of two of the system of equations, such as  $\{(1), (2)\}$ :

- ✓ If the system  $\{(1), (2)\}$  is impossible to solve, then the system  $\{(1), (2), (3)\}$  is impossible to solve.
- ✓ If the system  $\{(1), (2)\}$  has unlimited number of solutions from the form  $x = g(z), y = h(z)$ , (maybe  $g(z)$  or  $h(z)$  is constant), then we substitution in (3) to obtain on the equation of the form:

$$0 \cdot z = \beta; \beta \in R \cup \{I\}$$

This equation is impossible to solve or it have unlimited number of solutions (According to  $\beta$  value).

#### Example4.2.5:

Let:

$$(2 + 2I)x + (3 + 3I)y + (-5 - 5I)z = 1 \quad (1)$$

$$(1 + I)x + (-1 - I)y + (1 + I)z = 2 \quad (2)$$

$$(5 + 5I)x + (5 + 5I)y + (-9 - 9I)z = 4 \quad (3)$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 2 + 2I & 3 + 3I & -5 - 5I \\ 1 + I & -1 - I & 1 + I \\ 5 + 5I & 5 + 5I & -9 - 9I \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2 + 2I & 3 + 3I & -5 - 5I \\ 1 + I & -1 - I & 1 + I \\ 5 + 5I & 5 + 5I & -9 - 9I \end{vmatrix} = 0 + 0I$$

$$\det x = \begin{vmatrix} 1 & 3 + 3I & -5 - 5I \\ 2 & -1 - I & 1 + I \\ 4 & 5 + 5I & -9 - 9I \end{vmatrix} = 0 + 0I, \quad \det y = \begin{vmatrix} 2 + 2I & 1 & -5 - 5I \\ 1 + I & 2 & 1 + I \\ 5 + 5I & 4 & -9 - 9I \end{vmatrix} = 0 + 0I$$

$$\det z = \begin{vmatrix} 2 + 2I & 3 + 3I & 1 \\ 1 + I & -1 - I & 2 \\ 5 + 5I & 5 + 5I & 4 \end{vmatrix} = 0 + 0I$$

As:  $\det A = 0 + 0I = \det x = \det y = \det z = 0 + 0I$ , so we are looking for a system solution  $\{(1), (2)\}$ , then:

$$x = \frac{1}{5} \left( 2z + 7 - \frac{7}{2}I \right)$$

$$y = \frac{1}{5} \left( 7z - 3 + \frac{3}{2}I \right)$$

By substitution in (3), we get:

$$(0 + 0I)z = 0 + 0I$$

This equation has unlimited number of solutions, so the system  $\{(1), (2), (3)\}$  has unlimited number of solutions given by:

$$S = \left\{ (x, y, z) \in R^3 \cup \{I\}; x = \frac{1}{5} \left( 2z + 7 - \frac{7}{2}I \right), y = \frac{1}{5} \left( 7z - 3 + \frac{3}{2}I \right) \right\}$$

Or:

$$S = \left\{ \left( \frac{1}{5} \left( 2z + 7 - \frac{7}{2}I \right), \frac{1}{5} \left( 7z - 3 + \frac{3}{2}I \right), z \right); z \in R \cup \{I\} \right\}$$

#### Example4.2.6:

Let:

$$(2 + I)x + (1 + I)y + (3 - I)z = 2 + I$$

$$(-1 + I)x + (3 - 2I)y + (1 + 3I)z = 4 + 2I$$

$$(3 + 2I)x + (4 - I)y + (2 - 3I)z = 5 - I$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 2 + I & 1 + I & 3 - I \\ -1 + I & 3 - 2I & 1 + 3I \\ 3 + 2I & 4 - I & 2 - 3I \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2 + I & 1 + I & 3 - I \\ -1 + I & 3 - 2I & 1 + 3I \\ 3 + 2I & 4 - I & 2 - 3I \end{vmatrix} = -30 + 21I$$

$$\det x = \begin{vmatrix} 2 + I & 1 + I & 3 - I \\ 4 + 2I & 3 - 2I & 1 + 3I \\ 2 - 3I & 4 - I & 2 - 3I \end{vmatrix} = 4 + 29I, \quad \det y = \begin{vmatrix} 2 + I & 2 + I & 3 - I \\ -1 + I & 4 + 2I & 1 + 3I \\ 3 + 2I & 2 - 3I & 2 - 3I \end{vmatrix} = -35 - 31I$$

$$\det z = \begin{vmatrix} 2 + I & 1 + I & 2 + I \\ -1 + I & 3 - 2I & 4 + 2I \\ 3 + 2I & 4 - I & 2 - 3I \end{vmatrix} = -11 + 14I$$

Then:

$$x = \frac{\det x}{\det A} = \frac{4 + 29I}{-30 + 21I} = \frac{-2}{15} - \frac{477}{135}I$$

$$y = \frac{\det y}{\det A} = \frac{-35 - 31I}{-30 + 21I} = \frac{7}{6} + \frac{37}{6}I$$

$$z = \frac{\det z}{\det A} = \frac{-11 + 14I}{-30 + 21I} = \frac{11}{30} - \frac{7}{10}I$$

$$(x, y, z) = \left( \frac{-2}{15} - \frac{477}{135}I, \frac{7}{6} + \frac{37}{6}I, \frac{11}{30} - \frac{7}{10}I \right)$$

To verify the solution, we substitute the value of  $x$  and  $y$  into the first equation:

$$(2 + I)\left(\frac{-2}{15} - \frac{477}{135}I\right) + (1 + I)\left(\frac{7}{6} + \frac{37}{6}I\right) + (3 - I)\left(\frac{11}{30} - \frac{7}{10}I\right) = 2 + I$$

**Example 4.2.7:**

Let:

$$(2 + I)x + (1 + I)y + (3 - I)z = 1 + I \quad (1)$$

$$(4 + 2I)x + (2 + 2I)y + (6 - 2I)z = 1 + 2I \quad (2)$$

$$(6 + 3I)x + (3 + 3I)y + (9 - 3I)z = -1 + 3I \quad (3)$$

Find solution of the system.

Solution:

$$A = \begin{bmatrix} 2 + I & 1 + I & 3 - I \\ 4 + 2I & 2 + 2I & 6 - 2I \\ 6 + 3I & 3 + 3I & 9 - 3I \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2 + I & 1 + I & 3 - I \\ 4 + 2I & 2 + 2I & 6 - 2I \\ 6 + 3I & 3 + 3I & 9 - 3I \end{vmatrix} = 0 + 0I$$



## 5. Conclusions

This study was presented based on the importance of linear equations in our lives, where we introduced the concept of neutrosophic linear equation and its types. In addition to the introduction of Cramer's rule to solve the neutrosophic system of equations. This study can be generalized and applied in several fields of application in our current reality, traffic as an example.

**Acknowledgments:** This publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia.

## References

- [1] Smarandache, F., "Introduction to Neutrosophic Measure, Neutrosophic Integral, and Neutrosophic Probability", Sitech-Education Publisher, Craiova – Columbus, 2013.
- [2] Smarandache, F., "Finite Neutrosophic Complex Numbers, by W. B. Vasantha Kandasamy", Zip Publisher, Columbus, Ohio, USA, pp.1-16, 2011.
- [3] Smarandache, F., "Neutrosophy. / Neutrosophic Probability, Set, and Logic, American Research Press", Rehoboth, USA, 1998.
- [4] Smarandache, F., "Introduction to Neutrosophic statistics", Sitech-Education Publisher, pp.34-44, 2014.
- [5] Smarandache, F., "A Unifying Field in Logics: Neutrosophic Logic", Preface by Charles Le, American Research Press, Rehoboth, 1999, 2000. Second edition of the Proceedings of the First International Conference on Neutrosophy, Neutrosophic Logic, Neutrosophic Set, Neutrosophic Probability and Statistics, University of New Mexico, Gallup, 2001.
- [6] Smarandache, F., "Proceedings of the First International Conference on Neutrosophy", Neutrosophic Set, Neutrosophic Probability and Statistics, University of New Mexico, 2001.
- [7] Alhasan, Y., "Concepts of Neutrosophic Complex Numbers", International Journal of Neutrosophic Science, Volume 8, Issue 1, pp. 9-18, 2020.
- [8] Smarandache, F., "Neutrosophic Precalculus and Neutrosophic Calculus", book, 2015.
- [9] Al- Tahan, M., "Some Results on Single Valued Neutrosophic (Weak) Polygroups", International Journal of Neutrosophic Science, Volume 2, Issue 1, pp. 38-46, 2020.
- [10] Edalatpanah. S., "A Direct Model for Triangular Neutrosophic Linear Programming", International Journal of Neutrosophic Science, Volume 1, Issue 1, pp. 19-28, 2020.
- [11] Chakraborty, A., "A New Score Function of Pentagonal Neutrosophic Number and its Application in Networking Problem", International Journal of Neutrosophic Science, Volume 1, Issue 1, pp. 40-51, 2020.
- [12] Chakraborty, A., "Application of Pentagonal Neutrosophic Number in Shortest Path Problem", International Journal of Neutrosophic Science, Volume 3, Issue 1, pp. 21-28, 2020.
- [13] Smarandache, F., "Neutrosophy and Neutrosophic Logic, First International Conference on Neutrosophy", Neutrosophic Logic, Set, Probability, and Statistics, University of New Mexico, Gallup, NM 87301, USA 2002.
- [14] Alhasan, Y., "The General Exponential form of a Neutrosophic Complex Number", International Journal of Neutrosophic Science, Volume 11, Issue 2, pp. 100-107, 2020.

[15] Abdel-Basset, M., "An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number", *Applied Soft Computing*, pp.438-452, 2019.

[16] Abdel-Baset, M., Chang, V., Gamal, A., Smarandache, F., "An integrated neutrosophic ANP and VIKOR method for achieving sustainable supplier selection: A case study in importing field", *Comput. Ind.*, pp.94–110, 2019.

[17] Alhasan, Y., "The neutrosophic integrals and integration methods", *Neutrosophic Sets and Systems*, Volume 43, pp. 290-301, 2021.

[18] Alhasan, Y., and Alfahal, A "Neutrosophic Differential Equations That Translate Into Linear", *International Journal of Neutrosophic Science*, Volume 15, Issue 1, pp. 29-42, 2021.

Received: May 24, 2021. Accepted: August 28, 2021