

2-15-2022

Anti Neutrosophic multi fuzzy ideals of near ring

K. Hemabala

B. Srinivasa Kumar

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Hemabala, K. and B. Srinivasa Kumar. "Anti Neutrosophic multi fuzzy ideals of near ring." *Neutrosophic Sets and Systems* 48, 1 (2022). https://digitalrepository.unm.edu/nss_journal/vol48/iss1/6

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



Anti Neutrosophic multi fuzzy ideals of γ near ring

K. Hemabala ^{1,*} and B. Srinivasa Kumar ²

1. Research Scholar, Mathematics department, Koneru Lakshmaiah Education Foundation, Vaddeswaram, Guntur- 522502, Andhra Pradesh, India.

hemaram.magi@gmail.com

2. Assoc. Prof., Mathematics department, Koneru Lakshmaiah Education Foundation, Vaddeswaram, Guntur- 522502, Andhra Pradesh, India

sk_bhavirisetty@kluniversity.in

* Correspondence: sk_bhavirisetty@kluniversity.in

Abstract: The theory of anti neutrosophic multi fuzzy ideals of γ near ring is dispensed in this work and various algebraic properties such as intersection, union of anti neutrosophic multi fuzzy ideals of γ near ring are examined. Further we examined the direct anti product of anti neutrosophic multi fuzzy ideals of γ near ring and also we proved the homorphic images and pre images of anti neutrosophic multi fuzzy ideals of γ near ring.

Keywords: Anti neutrosophic fuzzy set, γ near ring, anti neutrosophic multi fuzzy set, anti neutrosophic multi fuzzy ideal of γ near ring, anti product of anti neutrosophic multi fuzzy ideals of γ near ring.

1. Introduction

In 1965, Zadeh[25] proposed the notion of fuzzy set. Later A.Rosenfeld[16] developed fuzzy groups. The numerous authors like Bh.Satyanarayana[3,4,5] proposed the concept of fuzzy γ near ring. The authors S. Ragamai, Y. Bhargavi, T. Eswarlal[19] developed theory of fuzzy and L fuzzy ideals of γ near ring. Later the properties of γ near ring in multi fuzzy sets were extended by K. Hemabala and Srinivasa kumar[13]. After the theory of fuzzy sets, Florentin Smarandache[7,8] established as a new field of philosophy which is a neutrosophic theory, in 1995. The main base of neutrosophic logic is neutrosophy that includes indeterminacy. It is an agumentation of fuzzy set and intuitionistic fuzzy set. In neutrosophic logic each proposition is estimated by three components T,I,F. The neutrosophic set theory have seen great triumph in several fields such as image processing ,medical diagnosis, robotic, decision making problem and so on. I. Arockiarani[3] extended the theory of neutrosophic fuzzy set. A.Solairaju and S.Thiruvani[2] verified the algebraic properties of fuzzy neutrosophic set in near rings. In fuzzy neutrosophic set, the three components T,I,F can take single values between 0 and 1. There is some ambiguity irrespective of the distance to the element is. The neutrosophic fuzzy set theory on its own is not sufficient to study real world problems. F. Smarandache[9] developed notion of neutrosophic multi sets, an extension of neutrosophic set, in 2016. Authors like Vakkas Ulucay and Memet sahin[23] verified the concepts of neutrosophic multi fuzzy set in groups and verified the group properties. We carry the neutrosophic multi fuzzy notion in γ near ring and hence some properties of algebra are verified.

2. Preliminaries:

Basic definitions of fuzzy set, multi fuzzy set, neutrosophic set and neutrosophic multi set, γ near ring are presenting in this section. Fuzzy set can take a single value between [0,1]

2.1 Definition:

Let E be a non empty set and \tilde{A} be a fuzzy set over E is defined by[25]

$$\tilde{A} = \{\tilde{A}(a) / a \in E\} \text{ where } \tilde{A}: E \rightarrow [0,1].$$

2.2 Definition:

Let E be a non empty set and M_f be a multi fuzzy set over E is defined as[20,21]

$$M_f = \{ \langle a, M_f^1(a), M_f^2(a), M_f^3(a), \dots, M_f^i(a) \rangle : a \in E \} \text{ where } M_f^n: E \rightarrow [0,1] \text{ for all } n \in \{1,2,\dots,i\} \text{ and}$$

$$a \in E$$

2.3 Definition:

Let E be a non empty set then neutrosophic fuzzy set \tilde{N} [7] in E is defined as

$$\tilde{N} = \{ \langle a, \check{t}_{\tilde{N}}(a), \check{i}_{\tilde{N}}(a), \check{f}_{\tilde{N}}(a) \rangle : a \in E \text{ and } \check{t}_{\tilde{N}}(a), \check{i}_{\tilde{N}}(a), \check{f}_{\tilde{N}}(a) \in [0,1] \}$$

Where $\check{t}_{\tilde{N}}(a)$ is the truth membership function, $\check{i}_{\tilde{N}}(a)$ is the indeterminacy membership function

and $\check{f}_{\tilde{N}}$ falsity membership function and $0 \leq \check{t}_{\tilde{N}}(x) + \check{i}_{\tilde{N}}(x) + \check{f}_{\tilde{N}} \leq 1$.

2.4 Definition:

Let E be a non empty set. A neutrosophic multi fuzzy set \tilde{N} on E can be defined as follows

$$\tilde{N} = \{ \langle a, (\check{t}_{\tilde{N}}^1(a), \check{t}_{\tilde{N}}^2(a), \dots, \check{t}_{\tilde{N}}^i(a)), (\check{i}_{\tilde{N}}^1(a), \check{i}_{\tilde{N}}^2(a), \dots, \check{i}_{\tilde{N}}^i(a)), (\check{f}_{\tilde{N}}^1(a), \check{f}_{\tilde{N}}^2(a), \dots, \check{f}_{\tilde{N}}^i(a)) \rangle : a \in E \}$$

Where $\check{t}_{\tilde{N}}^1(a), \check{t}_{\tilde{N}}^2(a), \dots, \check{t}_{\tilde{N}}^i(a): E \rightarrow [0,1]$

$$\check{i}_{\tilde{N}}^1(a), \check{i}_{\tilde{N}}^2(a), \dots, \check{i}_{\tilde{N}}^i(a): E \rightarrow [0,1]$$

$$\check{f}_{\tilde{N}}^1(a), \check{f}_{\tilde{N}}^2(a), \dots, \check{f}_{\tilde{N}}^i(a) : E \rightarrow [0,1]$$

$$0 \leq \sup \check{t}_{\tilde{N}}^n(a) + \sup \check{i}_{\tilde{N}}^n(a) + \sup \check{f}_{\tilde{N}}^n(a) \leq 1 \quad \text{for } n=1 \text{ to } i$$

$(\check{t}_{\tilde{N}}^1(a), \check{t}_{\tilde{N}}^2(a), \dots, \check{t}_{\tilde{N}}^i(a))$, $(\check{i}_{\tilde{N}}^1(a), \check{i}_{\tilde{N}}^2(a), \dots, \check{i}_{\tilde{N}}^i(a))$, $(\check{f}_{\tilde{N}}^1(a), \check{f}_{\tilde{N}}^2(a), \dots, \check{f}_{\tilde{N}}^i(a))$ are the sequences of truth membership values, indeterminacy membership values and falsity membership values. In addition i is called the dimension of neutrosophic multi fuzzy set \tilde{N} denoted by $d(\tilde{N})$.

The sequence of truth membership values are arranged in decreasing order, but the corresponding indeterminacy membership and falsity membership values may not be in any order.

2.5 Definition:

Let \mathcal{L} and \mathcal{R} neutrosophic multi fuzzy sets

where $\mathcal{L} = \{(\check{t}_{\mathcal{L}}^1(a), \check{t}_{\mathcal{L}}^2(a), \dots, \check{t}_{\mathcal{L}}^i(a)), (\check{i}_{\mathcal{L}}^1(a), \check{i}_{\mathcal{L}}^2(a), \dots, \check{i}_{\mathcal{L}}^i(a)), (\check{f}_{\mathcal{L}}^1(a), \check{f}_{\mathcal{L}}^2(a), \dots, \check{f}_{\mathcal{L}}^i(a))\}$ and

$\mathcal{R} = \{(\check{t}_{\mathcal{R}}^1(a), \check{t}_{\mathcal{R}}^2(a), \dots, \check{t}_{\mathcal{R}}^i(a)), (\check{i}_{\mathcal{R}}^1(a), \check{i}_{\mathcal{R}}^2(a), \dots, \check{i}_{\mathcal{R}}^i(a)), (\check{f}_{\mathcal{R}}^1(a), \check{f}_{\mathcal{R}}^2(a), \dots, \check{f}_{\mathcal{R}}^i(a))\}$ then we have

the following relations and operations

- 1) $\mathcal{L} \subseteq \mathcal{R}$ iff $\check{t}_{\mathcal{L}}^n(a) \leq \check{t}_{\mathcal{R}}^n(a)$, $\check{i}_{\mathcal{L}}^n(a) \geq \check{i}_{\mathcal{R}}^n(a)$, $\check{f}_{\mathcal{L}}^n(a) \geq \check{f}_{\mathcal{R}}^n(a)$, $a \in E$ and $n= 1$ to i .
- 2) $\mathcal{L} = \mathcal{R}$ iff $\check{t}_{\mathcal{L}}^n(a) = \check{t}_{\mathcal{R}}^n(a)$, $\check{i}_{\mathcal{L}}^n(a) = \check{i}_{\mathcal{R}}^n(a)$, $\check{f}_{\mathcal{L}}^n(a) = \check{f}_{\mathcal{R}}^n(a)$, $a \in E$ and $n= 1$ to i .
- 3) $\mathcal{L} \cup \mathcal{R} = \{a, \max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)), \min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)), \min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a))\}$, $a \in E$ and $n= 1$ to i .
- 4) $\mathcal{L} \cap \mathcal{R} = \{a, \min(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)), \max(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)), \max(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a))\}$, $a \in E$ and $n= 1$ to i .

2.6 Definition:

A non empty set E with the binary operations '+'(addition) and '.'(multiplication) is called a near ring[3] if the following conditions hold:

- 1) $(E, +)$ is a group
- 2) (E, \cdot) is a semigroup
- 3) $(e_1 + e_2) \cdot e_3 = e_1 \cdot e_3 + e_2 \cdot e_3$ for all $e_1, e_2, e_3 \in E$

To be precise, it is called right near ring .Since it satisfies the right distributive law. But the word near ring is intended to mean right near ring. We use gh instead of $g \cdot h$

A γ near ring M is a triple $(M, +, \gamma)$ where

- 1) $(M, +)$ is a group
- 2) γ is a non empty set of binary operations on M such that $\tau \in \gamma$, $(M, +, \tau)$ is a near ring.
- 3) $e_1 \tau (e_2 \sigma e_3) = (e_1 \tau e_2) \sigma e_3$ for all $e_1, e_2, e_3 \in M$ and $\tau, \sigma \in \gamma$.

3. Anti Neutrosophic multi fuzzy set of Υ near ring

In this section, we introduce the definition of anti neutrosophic multi fuzzy sets of Υ near ring. We proved that union of two anti neutrosophic multi fuzzy ideals of \mathcal{L} and \mathcal{R} is an anti neutrosophic multi fuzzy ideal. We also prove that the intersection of two anti neutrosophic multi fuzzy ideals of \mathcal{L} and \mathcal{R} is also an anti neutrosophic multi fuzzy ideal.

3.1 Definition:

A neutrosophic multi fuzzy set

$$\mathcal{L} = \{(\check{t}_{\mathcal{L}}^1(a), \check{t}_{\mathcal{L}}^2(a), \dots, \check{t}_{\mathcal{L}}^i(a)), (\check{i}_{\mathcal{L}}^1(a), \check{i}_{\mathcal{L}}^2(a), \dots, \check{i}_{\mathcal{L}}^i(a)), (\check{f}_{\mathcal{L}}^1(a), \check{f}_{\mathcal{L}}^2(a), \dots, \check{f}_{\mathcal{L}}^i(a))\}$$
 in a Υ near ring

M is called anti neutrosophic multi fuzzy sub Υ near ring of M if

- 1) $\check{t}_{\mathcal{L}}^n(a - b) \leq \max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)),$
 $\check{i}_{\mathcal{L}}^n(a - b) \geq \min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)),$
 $\check{f}_{\mathcal{L}}^n(a - b) \geq \min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b)), n=1$ to $i.$
- 2) $\check{t}_{\mathcal{L}}^n(a\tau b) \leq \max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)),$
 $\check{i}_{\mathcal{L}}^n(a\tau b) \geq \min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)),$
 $\check{f}_{\mathcal{L}}^n(a\tau b) \geq \min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b)), n=1$ to $i.$

3.2 Definition:

Let M be a Υ near ring. An anti neutrosophic multi fuzzy set \mathcal{L} in a Υ near ring M is called anti neutrosophic multi fuzzy left (resp. right) ideal of M if for all $a, b, \rho_1, \rho_2 \in M, \tau \in \Upsilon,$
 $n=1,2,\dots,i$

- 1) $\check{t}_{\mathcal{L}}^n(a - b) \leq \max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)),$
 $\check{i}_{\mathcal{L}}^n(a - b) \geq \min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)),$

$$\check{f}_{\mathcal{L}}^n(a - b) \geq \min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b))$$

2) $\check{t}_{\mathcal{L}}^n(b + a - b) \leq \check{t}_{\mathcal{L}}^n(a),$

$$\check{i}_{\mathcal{L}}^n(b + a - b) \geq \check{i}_{\mathcal{L}}^n(a),$$

$$\check{f}_{\mathcal{L}}^n(b + a - b) \geq \check{f}_{\mathcal{L}}^n(a),$$

3) $\check{t}_{\mathcal{L}}^n(\rho_1\tau(a + \rho_2) - \rho_1\tau\rho_2) \leq \check{t}_{\mathcal{L}}^n(a),$

$$\check{i}_{\mathcal{L}}^n(\rho_1\tau(a + \rho_2) - \rho_1\tau\rho_2) \geq \check{i}_{\mathcal{L}}^n(a),$$

$$\check{f}_{\mathcal{L}}^n(\rho_1\tau(a + \rho_2) - \rho_1\tau\rho_2) \geq \check{f}_{\mathcal{L}}^n(a),$$

[resp. right

$$\check{t}_{\mathcal{L}}^n(a\tau\rho_1) \leq \check{t}_{\mathcal{L}}^n(a)$$

$$\check{i}_{\mathcal{L}}^n(a\tau\rho_1) \geq \check{i}_{\mathcal{L}}^n(a),$$

$$\check{f}_{\mathcal{L}}^n(a\tau\rho_1) \geq \check{f}_{\mathcal{L}}^n(a)]$$

\mathcal{L} is called a anti neutrosophic multi fuzzy ideal of M if \mathcal{L} both left and right anti neutrosophic multi fuzzy ideal of M .

3.1 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy left ideal of M . Then $\mathcal{L} \cup \mathcal{R}$ is a anti neutrosophic multi fuzzy left ideal of M .

Proof:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy left ideal of M .

Let $a, b, \rho_1, \rho_2 \in M, \tau \in \gamma$

1) $\check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a - b) = \max\{\check{t}_{\mathcal{L}}^n(a - b), \check{t}_{\mathcal{R}}^n(a - b)\}$

$$\begin{aligned} &\leq \max\{\{\max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)), \max(\check{t}_{\mathcal{R}}^n(a), \check{t}_{\mathcal{R}}^n(b))\}\} \\ &\leq \max\{\{\max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)), \max(\check{t}_{\mathcal{L}}^n(b), \check{t}_{\mathcal{R}}^n(b))\}\} \\ &\leq \max(\check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a), \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(b)) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a - b) &= \min\{\check{i}_{\mathcal{L}}^n(a - b), \check{i}_{\mathcal{R}}^n(a - b)\} \\ &\geq \min\{\{\min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)), \min(\check{i}_{\mathcal{R}}^n(a), \check{i}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min\{\{\min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)), \min(\check{i}_{\mathcal{L}}^n(b), \check{i}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min(\check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a), \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(b)) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(a - b) &= \min\{\check{f}_{\mathcal{L}}^n(a - b), \check{f}_{\mathcal{R}}^n(a - b)\} \\ &\geq \min\{\{\min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b)), \min(\check{f}_{\mathcal{R}}^n(a), \check{f}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min\{\{\min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a)), \min(\check{f}_{\mathcal{L}}^n(b), \check{f}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min(\check{f}_{\mathcal{L} \cup \mathcal{R}}^n(a), \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(b)) \end{aligned}$$

$$\begin{aligned} 2) \quad \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(b + a - b) &= \max\{\check{t}_{\mathcal{L}}^n(b + a - b), \check{t}_{\mathcal{R}}^n(b + a - b)\} \\ &\leq \max\{\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)\} \\ &\leq \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(b + a - b) &= \min\{\check{i}_{\mathcal{L}}^n(b + a - b), \check{i}_{\mathcal{R}}^n(b + a - b)\} \\ &\geq \min\{\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)\} \\ &\geq \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a) \end{aligned}$$

$$\check{f}_{\mathcal{L} \cup \mathcal{R}}^n(b + a - b) = \min\{\check{f}_{\mathcal{L}}^n(b + a - b), \check{f}_{\mathcal{R}}^n(b + a - b)\}$$

$$\geq \min\{\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a)\}$$

$$\geq \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(a)$$

$$\begin{aligned} 3. \quad & \check{t}_{\mathcal{L} \cup \mathcal{R}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2)) \\ &= \max\{\check{t}_{\mathcal{L}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2)), \check{t}_{\mathcal{R}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2))\} \\ &\leq \max\{\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)\} \\ &\leq \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a) \end{aligned}$$

$$\begin{aligned} & \check{i}_{\mathcal{L} \cup \mathcal{R}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2)) \\ &= \min\{\check{i}_{\mathcal{L}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2)), \check{i}_{\mathcal{R}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2))\} \\ &\geq \min\{\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)\} \\ &\geq \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a) \end{aligned}$$

$$\begin{aligned} & \check{f}_{\mathcal{L} \cup \mathcal{R}}^n((\theta_1 \tau(x + \theta_2) - \theta_1 \tau \theta_2)) \\ &= \min\{\check{f}_{\mathcal{L}}^n((\theta_1 \tau(x + \theta_2) - \theta_1 \tau \theta_2)), \check{f}_{\mathcal{R}}^n((\theta_1 \tau(x + \theta_2) - \theta_1 \tau \theta_2))\} \\ &\geq \min\{\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a)\} \\ &\geq \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(a) \end{aligned}$$

$\therefore \mathcal{L} \cup \mathcal{R}$ is a anti neutrosophic multi fuzzy left ideal of M .

3.2 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy right ideal of M then $\mathcal{L} \cup \mathcal{R}$ is a anti neutrosophic multi fuzzy right ideal of M .

Proof:

Let \mathcal{L} and \mathcal{R} neutrosophic multi fuzzy right ideal of M .

Let $a, b, \rho_1, \rho_2 \in M, \tau \in \gamma$

$$\begin{aligned}
 1. \quad \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a - b) &= \max\{\check{t}_{\mathcal{L}}^n(a - b), \check{t}_{\mathcal{R}}^n(a - b)\} \\
 &\leq \max\{\{\max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)), \max(\check{t}_{\mathcal{R}}^n(a), \check{t}_{\mathcal{R}}^n(b))\}\} \\
 &\leq \max\{\{\max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)), \max(\check{t}_{\mathcal{L}}^n(b), \check{t}_{\mathcal{R}}^n(b))\}\} \\
 &\leq \max\{\check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a), \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(b)\}
 \end{aligned}$$

$$\begin{aligned}
 \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a - b) &= \min\{\check{i}_{\mathcal{L}}^n(a - b), \check{i}_{\mathcal{R}}^n(a - b)\} \\
 &\geq \min\{\{\min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)), \min(\check{i}_{\mathcal{R}}^n(a), \check{i}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min\{\{\min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)), \min(\check{i}_{\mathcal{L}}^n(b), \check{i}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min\{\check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a), \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(b)\}
 \end{aligned}$$

$$\begin{aligned}
 \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(a - b) &= \min\{\check{f}_{\mathcal{L}}^n(a - b), \check{f}_{\mathcal{R}}^n(a - b)\} \\
 &\geq \min\{\{\min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b)), \min(\check{f}_{\mathcal{R}}^n(a), \check{f}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min\{\{\min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a)), \min(\check{f}_{\mathcal{L}}^n(b), \check{f}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min\{\check{f}_{\mathcal{L} \cup \mathcal{R}}^n(a), \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(b)\}
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(b + a - b) &= \max\{\check{t}_{\mathcal{L}}^n(b + a - b), \check{t}_{\mathcal{R}}^n(b + a - b)\} \\
 &\leq \max\{\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)\} \\
 &\leq \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(a)
 \end{aligned}$$

$$\begin{aligned}
 \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(b + a - b) &= \min\{\check{i}_{\mathcal{L}}^n(b + a - b), \check{i}_{\mathcal{R}}^n(b + a - b)\} \\
 &\geq \min\{\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)\} \\
 &\geq \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(a)
 \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b}) &= \min\{\check{f}_{\mathcal{L}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b}), \check{f}_{\mathcal{R}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b})\} \\ &\geq \min\{\check{f}_{\mathcal{L}}^n(\mathbf{a}), \check{f}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} 3. \quad \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a} \tau \rho_1) &= \max\{\check{t}_{\mathcal{L}}^n(\mathbf{a} \tau \rho_1), \check{t}_{\mathcal{R}}^n(\mathbf{a} \tau \rho_1)\} \\ &\leq \max\{\check{t}_{\mathcal{L}}^n(\mathbf{a}), \check{t}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\leq \check{t}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a} \tau \rho_1) &= \min\{\check{i}_{\mathcal{L}}^n(\mathbf{a} \tau \rho_1), \check{i}_{\mathcal{R}}^n(\mathbf{a} \tau \rho_1)\} \\ &\geq \min\{\check{i}_{\mathcal{L}}^n(\mathbf{a}), \check{i}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{i}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a} \tau \rho_1) &= \min\{\check{f}_{\mathcal{L}}^n(\mathbf{a} \tau \rho_1), \check{f}_{\mathcal{R}}^n(\mathbf{a} \tau \rho_1)\} \\ &\geq \min\{\check{f}_{\mathcal{L}}^n(\mathbf{a}), \check{f}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{f}_{\mathcal{L} \cup \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$\therefore \mathcal{L} \cup \mathcal{R}$ is a anti neutrosophic multi fuzzy right ideal of M .

3.3 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy ideal of M then $\mathcal{L} \cup \mathcal{R}$ is a anti neutrosophic multi fuzzy ideal of M .

Proof: It is clear.

3.4 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy left ideal of M and then $\mathcal{L} \cap \mathcal{R}$ is a anti neutrosophic multi fuzzy left ideal of M .

Proof:

Let \mathcal{L} and \mathcal{R} neutrosophic multi fuzzy left ideal of ξ .

$$\text{Let } a, b, \rho_1, \rho_2 \in M, \tau \in \gamma$$

$$\begin{aligned} 1. \quad \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(a - b) &= \min\{\check{t}_{\mathcal{L}}^n(a - b), \check{t}_{\mathcal{R}}^n(a - b)\} \\ &\leq \min\{\{\max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)), \max(\check{t}_{\mathcal{R}}^n(a), \check{t}_{\mathcal{R}}^n(b))\}\} \\ &\leq \max\{\{\min(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)), \min(\check{t}_{\mathcal{L}}^n(b), \check{t}_{\mathcal{R}}^n(b))\}\} \\ &\leq \max(\check{t}_{\mathcal{L} \cap \mathcal{R}}^n(a), \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(b)) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(a - b) &= \max\{\check{i}_{\mathcal{L}}^n(a - b), \check{i}_{\mathcal{R}}^n(a - b)\} \\ &\geq \max\{\{\min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)), \min(\check{i}_{\mathcal{R}}^n(a), \check{i}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min\{\{\max(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)), \max(\check{i}_{\mathcal{L}}^n(b), \check{i}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min(\check{i}_{\mathcal{L} \cap \mathcal{R}}^n(a), \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(b)) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(a - b) &= \max\{\check{f}_{\mathcal{L}}^n(a - b), \check{f}_{\mathcal{R}}^n(a - b)\} \\ &\geq \max\{\{\min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b)), \min(\check{f}_{\mathcal{R}}^n(a), \check{f}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min\{\{\max(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a)), \max(\check{f}_{\mathcal{L}}^n(b), \check{f}_{\mathcal{R}}^n(b))\}\} \\ &\geq \min(\check{f}_{\mathcal{L} \cap \mathcal{R}}^n(a), \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(b)) \end{aligned}$$

$$\begin{aligned} 2. \quad \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(b + a - b) &= \min\{\check{t}_{\mathcal{L}}^n(b + a - b), \check{t}_{\mathcal{R}}^n(b + a - b)\} \\ &\leq \min\{\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)\} \\ &\leq \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(a) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(b + a - b) &= \max\{\check{i}_{\mathcal{L}}^n(b + a - b), \check{i}_{\mathcal{R}}^n(b + a - b)\} \\ &\geq \max\{\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)\} \end{aligned}$$

$$\geq \ddot{i}_{\mathcal{L} \cap \mathcal{R}}^n(a)$$

$$\ddot{f}_{\mathcal{L} \cap \mathcal{R}}^n(b + a - b) = \max\{\ddot{f}_{\mathcal{L}}^n(b + a - b), \ddot{f}_{\mathcal{R}}^n(b + a - b)\}$$

$$\geq \max\{\ddot{f}_{\mathcal{L}}^n(a), \ddot{f}_{\mathcal{R}}^n(a)\}$$

$$\geq \ddot{f}_{\mathcal{L} \cap \mathcal{R}}^n(a)$$

3. $\ddot{t}_{\mathcal{L} \cap \mathcal{R}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2))$

$$= \min\{\ddot{t}_{\mathcal{L}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2)), \ddot{t}_{\mathcal{R}}^n((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2))\}$$

$$\leq \min\{\ddot{t}_{\mathcal{L}}^n(a), \ddot{t}_{\mathcal{R}}^n(a)\}$$

$$\leq \ddot{t}_{\mathcal{L} \cap \mathcal{R}}^n(a)$$

$$\ddot{i}_{\mathcal{L} \cap \mathcal{R}}^m((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2))$$

$$= \max\{\ddot{i}_{\mathcal{L}}^m((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2)), \ddot{i}_{\mathcal{R}}^m((\rho_1 \tau(a + \rho_2) - \rho_1 \tau \rho_2))\}$$

$$\geq \max\{\ddot{i}_{\mathcal{L}}^m(a), \ddot{i}_{\mathcal{R}}^m(a)\}$$

$$\geq \ddot{i}_{\mathcal{L} \cap \mathcal{R}}^m(a)$$

$$\ddot{f}_{\mathcal{L} \cap \mathcal{R}}^n((\theta_1 \tau(x + \theta_2) - \theta_1 \tau \theta_2))$$

$$= \max\{\ddot{f}_{\mathcal{L}}^n((\theta_1 \tau(x + \theta_2) - \theta_1 \tau \theta_2)), \ddot{f}_{\mathcal{R}}^n((\theta_1 \tau(x + \theta_2) - \theta_1 \tau \theta_2))\}$$

$$\geq \max\{\ddot{f}_{\mathcal{L}}^n(a), \ddot{f}_{\mathcal{R}}^n(a)\}$$

$$\geq \ddot{f}_{\mathcal{L} \cap \mathcal{R}}^n(a)$$

∴ $\mathcal{L} \cap \mathcal{R}$ is a anti neutrosophic multi fuzzy left ideal of M .

3.5 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy right ideal of M then $\mathcal{L} \cap \mathcal{R}$ is a anti neutrosophic multi fuzzy right ideal of M .

Proof:

Let \mathcal{L} and \mathcal{R} neutrosophic multi fuzzy right ideal of M .

Let $a, b, \rho_1, \rho_2 \in M, \tau \in \gamma$

$$\begin{aligned}
 1. \quad \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(a - b) &= \min\{\check{t}_{\mathcal{L}}^n(a - b), \check{t}_{\mathcal{R}}^n(a - b)\} \\
 &\leq \min\{\{\max(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{L}}^n(b)), \max(\check{t}_{\mathcal{R}}^n(a), \check{t}_{\mathcal{R}}^n(b))\}\} \\
 &\leq \max\{\{\min(\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)), \min(\check{t}_{\mathcal{L}}^n(b), \check{t}_{\mathcal{R}}^n(b))\}\} \\
 &\leq \max(\check{t}_{\mathcal{L} \cap \mathcal{R}}^n(a), \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(b))
 \end{aligned}$$

$$\begin{aligned}
 \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(a - b) &= \max\{\check{i}_{\mathcal{L}}^n(a - b), \check{i}_{\mathcal{R}}^n(a - b)\} \\
 &\geq \max\{\{\min(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{L}}^n(b)), \min(\check{i}_{\mathcal{R}}^n(a), \check{i}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min\{\{\max(\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(a)), \max(\check{i}_{\mathcal{L}}^n(b), \check{i}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min(\check{i}_{\mathcal{L} \cap \mathcal{R}}^n(a), \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(b))
 \end{aligned}$$

$$\begin{aligned}
 \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(a - b) &= \max\{\check{f}_{\mathcal{L}}^n(a - b), \check{f}_{\mathcal{R}}^n(a - b)\} \\
 &\geq \max\{\{\min(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{L}}^n(b)), \min(\check{f}_{\mathcal{R}}^n(a), \check{f}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min\{\{\max(\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(a)), \max(\check{f}_{\mathcal{L}}^n(b), \check{f}_{\mathcal{R}}^n(b))\}\} \\
 &\geq \min(\check{f}_{\mathcal{L} \cap \mathcal{R}}^n(a), \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(b))
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(b + a - b) &= \min\{\check{t}_{\mathcal{L}}^n(b + a - b), \check{t}_{\mathcal{R}}^n(b + a - b)\} \\
 &\leq \min\{\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(a)\} \\
 &\leq \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(a)
 \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b}) &= \max\{\check{i}_{\mathcal{L}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b}) , \check{i}_{\mathcal{R}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b})\} \\ &\geq \max\{\check{i}_{\mathcal{L}}^n(\mathbf{a}), \check{i}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b}) &= \max\{\check{f}_{\mathcal{L}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b}) , \check{f}_{\mathcal{R}}^n(\mathbf{b} + \mathbf{a} - \mathbf{b})\} \\ &\geq \max\{\check{f}_{\mathcal{L}}^n(\mathbf{a}), \check{f}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} 3. \quad \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a} \tau \rho_1) &= \min\{\check{t}_{\mathcal{L}}^n(\mathbf{a} \tau \rho_1) , \check{t}_{\mathcal{R}}^n(\mathbf{a} \tau \rho_1)\} \\ &\leq \min\{\check{t}_{\mathcal{L}}^n(\mathbf{a}), \check{t}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\leq \check{t}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a} \tau \rho_1) &= \max\{\check{i}_{\mathcal{L}}^n(\mathbf{a} \tau \rho_1) , \check{i}_{\mathcal{R}}^n(\mathbf{a} \tau \rho_1)\} \\ &\geq \max\{\check{i}_{\mathcal{L}}^n(\mathbf{a}), \check{i}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{i}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a} \tau \rho_1) &= \max\{\check{f}_{\mathcal{L}}^n(\mathbf{a} \tau \rho_1) , \check{f}_{\mathcal{R}}^n(\mathbf{a} \tau \rho_1)\} \\ &\geq \max\{\check{f}_{\mathcal{L}}^n(\mathbf{a}), \check{f}_{\mathcal{R}}^n(\mathbf{a})\} \\ &\geq \check{f}_{\mathcal{L} \cap \mathcal{R}}^n(\mathbf{a}) \end{aligned}$$

∴ $\mathcal{L} \cap \mathcal{R}$ is a anti neutrosophic multi fuzzy right ideal of M .

3.6 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy ideal of M then $\mathcal{L} \cap \mathcal{R}$ is also a anti neutrosophic multi fuzzy ideal of M .

Proof: It is clear.

4. Anti Product of anti neutrosophic multi fuzzy ideals

In this section we define anti product of anti neutrosophic multi fuzzy Υ near ring M . We proved that anti product of anti neutrosophic multi fuzzy ideals of M is a anti neutrosophic multi fuzzy ideal of M .

4.1 Definition:

Let \mathcal{L} and \mathcal{R} are two anti neutrosophic multi fuzzy ideals of Υ near rings M_1 and M_2 resp. Then the anti product of anti neutrosophic multi fuzzy subset of Υ near ring is defined by

$\mathcal{L} \times \mathcal{R}: M_1 \times M_2 \rightarrow [0,1]$ such that

$$\mathcal{L} \times \mathcal{R} = \{ \langle (a, b), \check{t}_{\mathcal{L} \times \mathcal{R}}^n(a, b), \check{i}_{\mathcal{L} \times \mathcal{R}}^n(a, b), \check{f}_{\mathcal{L} \times \mathcal{R}}^n(a, b) \rangle : a \in M_1, b \in M_2 \}$$

Where $\check{t}_{\mathcal{L} \times \mathcal{R}}^n(a, b) = \max\{\check{t}_{\mathcal{L}}^n(a), \check{t}_{\mathcal{R}}^n(b)\}$

$$\check{i}_{\mathcal{L} \times \mathcal{R}}^n(a, b) = \min\{\check{i}_{\mathcal{L}}^n(a), \check{i}_{\mathcal{R}}^n(b)\}$$

$$\check{f}_{\mathcal{L} \times \mathcal{R}}^n(a, b) = \min\{\check{f}_{\mathcal{L}}^n(a), \check{f}_{\mathcal{R}}^n(b)\}$$

4.2 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy left ideal of Υ near rings M_1 and M_2 then $\mathcal{L} \times \mathcal{R}$ is also a anti neutrosophic multi fuzzy left ideal of $M_1 \times M_2$.

Proof:

Let \mathcal{L} and \mathcal{R} be anti neutrosophic fuzzy left ideals of $M_1 \times M_2$ respectively

Let $(a_1, a_2), (b_1, b_2), (\rho_1, \rho_2) \in M_1 \times M_2$

$$\begin{aligned} 1. \quad \check{t}_{\mathcal{L} \times \mathcal{R}}^n((a_1, a_2) - (b_1, b_2)) &= \check{t}_{\mathcal{L} \times \mathcal{R}}^n(a_1 - b_1, a_2 - b_2) \\ &= \max(\check{t}_{\mathcal{L}}^n(a_1 - b_1), \check{t}_{\mathcal{R}}^n(a_2 - b_2)) \end{aligned}$$

$$\begin{aligned} &\leq \max \{ \max (\check{t}_{\mathcal{L}}^n (a_1), \check{t}_{\mathcal{L}}^n (b_1)), \max [\check{t}_{\mathcal{R}}^n (a_2), \check{t}_{\mathcal{R}}^n (b_2)] \} \\ &\leq \max \{ \max [\check{t}_{\mathcal{L}}^n (a_1), \check{t}_{\mathcal{R}}^n (a_2)], \max [\check{t}_{\mathcal{L}}^n (b_1), \check{t}_{\mathcal{R}}^n (b_2)] \} \\ &\leq \max (\check{t}_{\mathcal{L}\times\mathcal{R}}^n (a_1, a_2), \check{t}_{\mathcal{L}\times\mathcal{R}}^n (b_1, b_2)) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L}\times\mathcal{R}}^n ((a_1, a_2) - (b_1, b_2)) &= \check{i}_{\mathcal{L}\times\mathcal{R}}^n (a_1 - b_1, a_2 - b_2) \\ &= \min (\check{i}_{\mathcal{L}}^n (a_1 - b_1), \check{i}_{\mathcal{R}}^n (a_2 - b_2)) \\ &\geq \min \{ \min (\check{i}_{\mathcal{L}}^n (a_1), \check{i}_{\mathcal{L}}^n (b_1)), \min [\check{i}_{\mathcal{R}}^n (a_2), \check{i}_{\mathcal{R}}^n (b_2)] \} \\ &\geq \min \{ \min [\check{i}_{\mathcal{L}}^n (a_1), \check{i}_{\mathcal{R}}^n (a_2)], \min [\check{i}_{\mathcal{L}}^n (b_1), \check{i}_{\mathcal{R}}^n (b_2)] \} \\ &\geq \min (\check{i}_{\mathcal{L}\times\mathcal{R}}^n (a_1, a_2), \check{i}_{\mathcal{L}\times\mathcal{R}}^n (b_1, b_2)) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L}\times\mathcal{R}}^n ((a_1, a_2) - (b_1, b_2)) &= \check{f}_{\mathcal{L}\times\mathcal{R}}^n (a_1 - b_1, a_2 - b_2) \\ &= \min (\check{f}_{\mathcal{L}}^n (a_1 - b_1), \check{f}_{\mathcal{R}}^n (a_2 - b_2)) \\ &\geq \min \{ \min (\check{f}_{\mathcal{L}}^n (a_1), \check{f}_{\mathcal{L}}^n (b_1)), \min [\check{f}_{\mathcal{R}}^n (a_2), \check{f}_{\mathcal{R}}^n (b_2)] \} \\ &\geq \min \{ \min [\check{f}_{\mathcal{L}}^n (a_1), \check{f}_{\mathcal{R}}^n (a_2)], \min [\check{f}_{\mathcal{L}}^n (b_1), \check{f}_{\mathcal{R}}^n (b_2)] \} \\ &\geq \min (\check{f}_{\mathcal{L}\times\mathcal{R}}^n (a_1, a_2), \check{f}_{\mathcal{L}\times\mathcal{R}}^n (b_1, b_2)) \end{aligned}$$

$$\begin{aligned} 2. \quad \check{t}_{\mathcal{L}\times\mathcal{R}}^n ((b_1, b_2) + (a_1, a_2) - (b_1, b_2)) &= \check{t}_{\mathcal{L}\times\mathcal{R}}^n (b_1 + a_1 - b_1, b_2 + a_2 - b_2) \\ &= \max (\check{t}_{\mathcal{L}}^n (b_1 + a_1 - b_1), \check{t}_{\mathcal{R}}^n (b_2 + a_2 - b_2)) \\ &\leq \max \{ \check{t}_{\mathcal{L}}^n (a_1), \check{t}_{\mathcal{R}}^n (a_2) \} \\ &\leq \check{t}_{\mathcal{L}\times\mathcal{R}}^n (a_1, a_2) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L}\times\mathcal{R}}^n ((b_1, b_2) + (a_1, a_2) - (b_1, b_2)) &= \check{i}_{\mathcal{L}\times\mathcal{R}}^n (b_1 + a_1 - b_1, b_2 + a_2 - b_2) \\ &\geq \min \{ \check{i}_{\mathcal{L}}^n (a_1), \check{i}_{\mathcal{R}}^n (a_2) \} \\ &\geq \check{i}_{\mathcal{L}\times\mathcal{R}}^n (a_1, a_2) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L}x\mathcal{R}}^n((b_1, b_2) + (a_1, a_2) - (b_1, b_2)) &= \check{f}_{\mathcal{L}x\mathcal{R}}^n(b_1 + a_1 - b_1, b_2 + a_2 - b_2) \\ &\geq \min\{\check{f}_{\mathcal{L}}^n(a_1), \check{f}_{\mathcal{R}}^n(a_2)\} \\ &\geq \check{f}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2) \end{aligned}$$

$$\begin{aligned} 3. \check{t}_{\mathcal{L}x\mathcal{R}}^n((\rho_1\tau((a_1, a_2) + \rho_2) - \rho_1\tau\rho_2)) & \\ &= \check{t}_{\mathcal{L}x\mathcal{R}}^n(\rho_1\tau(a_1 + \rho_2) - \rho_1\tau\rho_2, \rho_1\tau(a_2 + \rho_2) - \rho_1\tau\rho_2) \\ &= \max\{\check{t}_{\mathcal{L}}^n((\rho_1\tau(a_1 + \rho_2) - \rho_1\tau\rho_2)), \check{t}_{\mathcal{R}}^n((\rho_1\tau(a_2 + \rho_2) - \rho_1\tau\rho_2))\} \\ &\leq \max\{\check{t}_{\mathcal{L}}^n(a_1), \check{t}_{\mathcal{R}}^n(a_2)\} \\ &\leq \check{t}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L}x\mathcal{R}}^n((\rho_1\tau(a_1, a_2 + \rho_2) - \rho_1\tau\rho_2)) & \\ &= \check{i}_{\mathcal{L}x\mathcal{R}}^n(\rho_1\tau(a_1 + \rho_2) - \rho_1\tau\rho_2, \rho_1\tau(a_2 + \rho_2) - \rho_1\tau\rho_2) \\ &= \min\{\check{i}_{\mathcal{L}}^n((\rho_1\tau(a_1 + \rho_2) - \rho_1\tau\rho_2)), \check{i}_{\mathcal{R}}^n((\rho_1\tau(a_2 + \rho_2) - \rho_1\tau\rho_2))\} \\ &\geq \min\{\check{i}_{\mathcal{L}}^n(a_1), \check{i}_{\mathcal{R}}^n(a_2)\} \\ &\geq \check{i}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L}x\mathcal{R}}^n((\rho_1\tau(a_1, a_2 + \rho_2) - \rho_1\tau\rho_2)) & \\ &= \check{f}_{\mathcal{L}x\mathcal{R}}^n(\rho_1\tau(a_1 + \rho_2) - \rho_1\tau\rho_2, \rho_1\tau(a_2 + \rho_2) - \rho_1\tau\rho_2) \\ &= \min\{\check{f}_{\mathcal{L}}^n((\rho_1\tau(a_1 + \rho_2) - \rho_1\tau\rho_2)), \check{f}_{\mathcal{R}}^n((\rho_1\tau(a_2 + \rho_2) - \rho_1\tau\rho_2))\} \\ &\geq \min\{\check{f}_{\mathcal{L}}^n(a_1), \check{f}_{\mathcal{R}}^n(a_2)\} \\ &\geq \check{f}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2) \end{aligned}$$

∴ $\mathcal{L}x\mathcal{R}$ is also a anti neutrosophic multi fuzzy left ideal of M_1xM_2 .

4.3 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy right ideal of \mathcal{Y} near rings M_1 and M_2 then $\mathcal{L}\mathcal{X}\mathcal{R}$ is also a anti neutrosophic multi fuzzy right ideal of $M_1\mathcal{X}M_2$.

Proof:

Let \mathcal{L} and \mathcal{R} be anti neutrosophic fuzzy right ideals of $M_1\mathcal{X}M_2$ respectively

Let $(a_1, a_2), (b_1, b_2), (\rho_1, \rho_2) \in M_1\mathcal{X}M_2$

$$\begin{aligned} 1. \quad \check{t}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n((a_1, a_2) - (b_1, b_2)) &= \check{t}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(a_1 - b_1, a_2 - b_2) \\ &= \max(\check{t}_{\mathcal{L}}^n(a_1 - b_1), \check{t}_{\mathcal{R}}^n(a_2 - b_2)) \\ &\leq \max\{\max(\check{t}_{\mathcal{L}}^n(a_1), \check{t}_{\mathcal{L}}^n(b_1)), \max[\check{t}_{\mathcal{R}}^n(a_2), \check{t}_{\mathcal{R}}^n(b_2)]\} \\ &\leq \max\{\max[\check{t}_{\mathcal{L}}^n(a_1), \check{t}_{\mathcal{R}}^n(a_2)], \max[\check{t}_{\mathcal{L}}^n(b_1), \check{t}_{\mathcal{R}}^n(b_2)]\} \\ &\leq \max(\check{t}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(a_1, a_2), \check{t}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(b_1, b_2)) \end{aligned}$$

$$\begin{aligned} \check{i}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n((a_1, a_2) - (b_1, b_2)) &= \check{i}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(a_1 - b_1, a_2 - b_2) \\ &= \min(\check{i}_{\mathcal{L}}^n(a_1 - b_1), \check{i}_{\mathcal{R}}^n(a_2 - b_2)) \\ &\geq \min\{\min(\check{i}_{\mathcal{L}}^n(a_1), \check{i}_{\mathcal{L}}^n(b_1)), \min[\check{i}_{\mathcal{R}}^n(a_2), \check{i}_{\mathcal{R}}^n(b_2)]\} \\ &\geq \min\{\min[\check{i}_{\mathcal{L}}^n(a_1), \check{i}_{\mathcal{R}}^n(a_2)], \min[\check{i}_{\mathcal{L}}^n(b_1), \check{i}_{\mathcal{R}}^n(b_2)]\} \\ &\geq \min(\check{i}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(a_1, a_2), \check{i}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(b_1, b_2)) \end{aligned}$$

$$\begin{aligned} \check{f}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n((a_1, a_2) - (b_1, b_2)) &= \check{f}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(a_1 - b_1, a_2 - b_2) \\ &= \min(\check{f}_{\mathcal{L}}^n(a_1 - b_1), \check{f}_{\mathcal{R}}^n(a_2 - b_2)) \\ &\geq \min\{\min(\check{f}_{\mathcal{L}}^n(a_1), \check{f}_{\mathcal{L}}^n(b_1)), \min[\check{f}_{\mathcal{R}}^n(a_2), \check{f}_{\mathcal{R}}^n(b_2)]\} \\ &\geq \min\{\min[\check{f}_{\mathcal{L}}^n(a_1), \check{f}_{\mathcal{R}}^n(a_2)], \min[\check{f}_{\mathcal{L}}^n(b_1), \check{f}_{\mathcal{R}}^n(b_2)]\} \\ &\geq \min(\check{f}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(a_1, a_2), \check{f}_{\mathcal{L}\mathcal{X}\mathcal{R}}^n(b_1, b_2)) \end{aligned}$$

$$\begin{aligned}
 2. \quad \check{t}_{\mathcal{L}x\mathcal{R}}^n((b_1, b_2) + (a_1, a_2) - (b_1, b_2)) &= \check{t}_{\mathcal{L}x\mathcal{R}}^n(b_1 + a_1 - b_1, b_2 + a_2 - b_2) \\
 &= \max\{\check{t}_{\mathcal{L}}^n(b_1 + a_1 - b_1), \check{t}_{\mathcal{R}}^n(b_2 + a_2 - b_2)\} \\
 &\leq \max\{\check{t}_{\mathcal{L}}^n(a_1), \check{t}_{\mathcal{R}}^n(a_2)\} \\
 &\leq \check{t}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2)
 \end{aligned}$$

$$\begin{aligned}
 \check{i}_{\mathcal{L}x\mathcal{R}}^n((b_1, b_2) + (a_1, a_2) - (b_1, b_2)) &= \check{i}_{\mathcal{L}x\mathcal{R}}^n(b_1 + a_1 - b_1, b_2 + a_2 - b_2) \\
 &\geq \min\{\check{i}_{\mathcal{L}}^n(a_1), \check{i}_{\mathcal{R}}^n(a_2)\} \\
 &\geq \check{i}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2)
 \end{aligned}$$

$$\begin{aligned}
 \check{f}_{\mathcal{L}x\mathcal{R}}^n((b_1, b_2) + (a_1, a_2) - (b_1, b_2)) &= \check{f}_{\mathcal{L}x\mathcal{R}}^n(b_1 + a_1 - b_1, b_2 + a_2 - b_2) \\
 &\geq \min\{\check{f}_{\mathcal{L}}^n(a_1), \check{f}_{\mathcal{R}}^n(a_2)\} \\
 &\geq \check{f}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2)
 \end{aligned}$$

$$\begin{aligned}
 3. \quad \check{t}_{\mathcal{L}x\mathcal{R}}^n((a_1, a_2)\tau(\rho_1, \rho_2)) &= \check{t}_{\mathcal{L}x\mathcal{R}}^n\{(a_1 \tau\rho_1), (a_2 \tau\rho_2)\} \\
 &\leq \max\{\check{t}_{\mathcal{L}}^n(a_1), \check{t}_{\mathcal{R}}^n(a_2)\} \\
 &\leq \check{t}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2)
 \end{aligned}$$

$$\begin{aligned}
 \check{i}_{\mathcal{L}x\mathcal{R}}^n((a_1, a_2)\tau(\rho_1, \rho_2)) &= \check{i}_{\mathcal{L}x\mathcal{R}}^n\{(a_1 \tau\rho_1), (a_2 \tau\rho_2)\} \\
 &\geq \min\{\check{i}_{\mathcal{L}}^n(a_1), \check{i}_{\mathcal{R}}^n(a_2)\} \\
 &\geq \check{i}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2)
 \end{aligned}$$

$$\begin{aligned}
 \check{f}_{\mathcal{L}x\mathcal{R}}^n((a_1, a_2)\tau(\rho_1, \rho_2)) &= \check{f}_{\mathcal{L}x\mathcal{R}}^n\{(a_1 \tau\rho_1), (a_2 \tau\rho_2)\} \\
 &\geq \min\{\check{f}_{\mathcal{L}}^n(a_1), \check{f}_{\mathcal{R}}^n(a_2)\} \\
 &\geq \check{f}_{\mathcal{L}x\mathcal{R}}^n(a_1, a_2)
 \end{aligned}$$

$\therefore \mathcal{L}x\mathcal{R}$ is also a anti neutrosophic multi fuzzy right ideal of M_1xM_2 .

4.4 Theorem:

Let \mathcal{L} and \mathcal{R} anti neutrosophic multi fuzzy ideal of γ near rings M_1 and M_2 then $\mathcal{L}x\mathcal{R}$ is also a anti neutrosophic multi fuzzy ideal of M_1xM_2 .

Proof: It is clear

5. Conclusion

To conclude, the notion of neutrosophic multi fuzzy gamma near-ring, neutrosophic multi fuzzy ideals of gamma near-rings have been discussed. The proof for the theorem that states "Union and Intersection of two neutrosophic multi fuzzy ideals of gamma near-ring is also a Neutrosophic multi fuzzy ideal of gamma near-ring" has been provided.

References

1. Agboola A.A.A, Akwu A.D and Oyebo Y.T., Neutrosophic groups and subgroups, International J.Math.Combin.Vol.3(2012),1-9.
2. A. Solairaju and S. Thiruvén, Neutrosophic Fuzzy Ideals of Near-Rings, International Journal of Pure and Applied Mathematics, Volume 118 No. 6 2018, 527-539
3. Bh. Satyanarayana, A note on Γ -near rings Indian J. Math., 41:3(1999)427-433.
4. Bh. Satyanarayana, contribution to near ring theory, Doctoral Thesis, Nagarjuna University, India 1984.
5. Bh. Satyanarayana, Γ -near rings, proceedings of the national seminar on Algebra and its applications, july11-12, 2011, pp: 01-16.
6. Deli, I. Broumi, S. Smarandache, F. On neutrosophic multi sets and its application in medical diagnosis. J. New Theory 2015, 6, 88-98.
7. F. Smarandache, Neutrosophy, A new branch of Philosophy logic in multiple-valued logic, An international journal,8(3)(2002),297-384.
8. F. Smarandache, Neutrosophic set – a generalization of the Intuitionistic fuzzy sets, J. Pure Appl.Math.,24(2005),287-297
9. F. Smarandache, Neutrosophic Perspectives: Triplets, Duplets, Multi sets, Hybrid Operators, Modal Logic, Hedge Algebras And Applications, Pons Edition, Bruxelles,323p., 2017, 115-119.
10. G. L. Booth, A note on Γ -near rings, studia.sci.math.Hungar.23 (1988)471-475.

11. I. Arockiarani, J.MartinaJency , More on Fuzzy Neutrosophic sets and Fuzzy Neutrosophic Topological spaces , International journal of innovative research and studies ,May(2014),vol 3,Issue 5, 643-652.
12. J. Martina Jency, I. Arockiarani, Fuzzy Neutrosophic Sub groupoids, Asian Journal of Applied Sciences (ISSN:2321-0893), vol 04, Issue 01, February (2016)
13. K. Hemabala, Srinivasa kumar, Cartesian Product of multi L fuzzy ideals of Γ near ring, Advances in Mathematics: Scientific Journal 9(2020),no.7,5273-5281.
14. K. H. Kim and Y. B. Jun, On fuzzy R-subgroups of near-rings, J. Fuzzy Math. 8(3)(2008) 549-558.
15. R. Muthuraj and S. Balamurugan, multi fuzzy group, Gen.Math.Notes, vol17, No.1, July 2013, pp.74-81.
16. Rosenfield A, Fuzzy Groups, Journal of mathematical analysis and applications,35,512- 517(1971)
17. S. Abu zaid, On fuzzy sub near-rings and ideals, Fuzzy sets and systems 44(1991)139- 146.
18. S. D. Kim and H. S. Kim, On fuzzy ideals of near-rings, Bulletin Korean Mathematical society 33(1996) 593-601
19. S. Ragamayi, Y. Bhargavi, T. Eswarlal, L fuzzy ideals of a Γ near ring, International Journal of Pure and Applied Mathematics, Vol.118, No.2, 2018, 291- 307
20. S. Sabu and T.V. Ramakrishnan, Multi fuzzy sets, International mathematical Forum, 50(2010),2471-2476.
21. S. Sabu and T.V. Ramakrishnan, Multi fuzzy topology, Internatiional journal of Applied Mathematics.
22. T. K. Dutta and B. K. Biswas, Fuzzy ideal of a near-ring, Bull.Cal.Math.Soc.89 (1997), 447-456.
23. VakkasUlucay and MemetSahin, Neutrosophic Multi groups and Applications, new challenges in neutrosophic theory and applications , special issue, volume 7 , issue 1, 2019.
24. V. Chinnadurai and S.Kadalarasi, Direct product of fuzzy ideals of near-rings, Annals of Fuzzy Mathematics and Informatics,2016
25. Zadeh.L.A, Fuzzy sets, Information control, Vol.8,338-353(1965).

Received: Nov 10, 2021. Accepted: Feb 5, 2022