

4-12-2022

Tensor Product of Neutrosophic submodules of an R-module

Binu R.

Ursala Paul

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

R., Binu and Ursala Paul. "Tensor Product of Neutrosophic submodules of an R-module." *Neutrosophic Sets and Systems* 49, 1 (2022). https://digitalrepository.unm.edu/nss_journal/vol49/iss1/30

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



Tensor Product of Neutrosophic submodules of an R -module

Binu R.^{1,*} and Ursala Paul²

¹Department of Mathematics, Rajagiri School of Engineering and Technology (Autonomous), Ernakulam, India; 1984binur@gmail.com

²Department of Mathematics, St. Teresa's College (Autonomous), Ernakulam, India; ursalaj@gmail.com

*Correspondence: 1984binur@gmail.com; Tel.: (91-9946167116)

Abstract. In this paper, we develop a framework for tensor product with imprecise and indeterminate bounds as a neutrosophic submodule of an R -modules. The fundamental goal of this study is to extend the conventional tensor product in contemporary algebra to the most generalized domain of neutrosophic set algebraic structures. We discuss the construction of tensor product in neutrosophic submodules as a quotient space in this study and derives the universal uniqueness property of tensor product in neutrosophic domain .

Keywords: Neutrosophic set; Neutrosophic homomorphism; Direct product; Cartesian product of neutrosophic set; Neutrosophic R -bi additive ; Neutrosophic tensor product

1. Introduction

Algebra is a vital branch of Mathematics. The gist of Algebra lies in construction of fundamental mathematical structures and identification of relations between mathematical ideas. The group representation theory proposed by Frobenius [16] has a major role in advanced algebra and emphasises on the detailed study of symmetries in nature. The study of modules and group representation are inter-related and has several applications in different branches of physics and chemistry. Several researchers have looked into the algebraic structure underlying uncertainty in pure mathematics. When the novel set theories evolved as a result of the works of Zadeh [35], the concept of modules also underwent the subsequent transformation. The major inventions in this direction include the research contributions of Negoita and Ralescu [21] and Mashinchi and Zahedi [17, 18] towards fuzzy modules. In 1986, Atanassov [2] came up with the intuitionistic version of fuzzy sets. In 2011, P. Isaac, P.P. John [14] characterised an intuitionistic fuzzy submodule

The notion of neutrosophy originally appeared in philosophy [26], and then as a mathematical tool. In 1995, Smarandache [19, 23] invented the neutrosophic set with the main goal of

translating set theory into the the real world by bridging the gap with theoretical certainties and practical uncertainties. The intervention of neutrosophic sets to algebraic structures by Smarandache [34] paved the way to many innovative concepts in algebraic research field. Kandasamy and Smarandache [33] designed the fundamental algebraic neutrosophic systems. Smarandache and Ali [29] proposed the neutrosophic triplet group, a new algebraic structure that leads to evolutionary changes in the neutrosophic research domain. Gulistan and Naeem [11, 12] introduced the concept of neutrosophic triplet semi hypergroups and complex fuzzy hyper ideals in non-associative hyperrings. The works of Vidan Cetkin [5, 6] resulted in the creation of neutrosophic subgroups and modules. The idea of fuzzy G -modules and innovative group representations was proposed by Sherry [9]. The further study in this area was carried out by Ursala by infusing the concepts of rough and fuzzy sets with module theory [15, 24]. The notion of neutrosophic submodules of an R -module and the additional characteristics in methodology were introduced by Binu [3, 4]

In equivalences of module categories, homomorphism functions and tensor product play a major role, and it is the more flexible generalisation of free module. The study of bilinear operations and the extension of scalars is made possible by the tensor product of modules. The universal multiplication of modules is achieved by tensor algebra of modules. In a neutrosophic context, the principal application of the tensor product is the management of large amounts of ambiguous data and the reduction of object dimensions in mathematical modelling. The tensor product in neutrosophic modules can be used to set up computational uncertainty quantification in probabilistic and deterministic models, allowing for easier interpretation of latent information and extraction of more components of information. The concept of tensor product in neutrosophic submodules of an R -modules generalises the concept of tensor product in modules and gives the multilinear operation more strength. We design the tensor product in neutrosophic submodules and characterised its features in this paper. The objects or entities used in this study for the algebraic creation of tensor products are neutrosophic submodules, which allow us deal with ambiguous data in imprecise bounds.

The following is the organization of this work, with the first section serving as an introductory concept. The Section 2 of this research article deals with the pre-requisite definitions and results. The concept of tensor product between neutrosophic submodules and the characteristic properties are presented in Section 3. Section 4 provides an overview of the further research work in this particular area of neutrosophic set theory.

2. Preliminaries

In this section, we examine some of the preliminary definitions and outcomes that are necessary for a thorough understanding of the subsequent sections.

Definition 2.1. [1] Let R be a commutative ring with unity. A *module* M over R is an ‘Abelian’ group with a law of composition written ‘+’ and a scalar multiplication $R \times M \rightarrow M$, written $(r, x) \rightsquigarrow rx$, that satisfy these axioms

- (1) $1x = x$
- (2) $(rs)x = r(sx)$
- (3) $(r + s)x = rx + sx$
- (4) $r(x + y) = rx + ry \quad \forall r, s \in R \text{ and } x, y \in M.$

Definition 2.2. [8, 10] Let R be a ring and M be an R -module. Let N be a submodule of M . The (additive, abelian) quotient group M/N can be made into an R -module by defining an action of elements of R by

$$r(x + N) = (rx) + N, \forall r \in R, x + N \in M/N$$

Remark 2.1. $[m]$ represents the coset $m + N, \forall m \in M$.

Definition 2.3. [8] A homomorphism $\mathcal{Y} : M \rightarrow N$ of R -modules is a map compatible with the laws of composition

- (1) $\mathcal{Y}(x + y) = \mathcal{Y}(x) + \mathcal{Y}(y)$
- (2) $\mathcal{Y}(rx) = r\mathcal{Y}(x) \quad \forall x, y \in M, r \in R.$

Remark 2.2. $Hom_R(M, N)$ represent the set of all R -module homomorphisms of M into N .

Definition 2.4. [8] Let M_1, M_2, \dots, M_n be R -modules. Then the direct product is a collection of n -tuples, denoted and defined as $M_1 \times M_2 \times \dots \times M_n = (m_1, m_2, \dots, m_n)$, $m_i \in M_i, 1 \leq i \leq n$ with addition and action of R defined component wise is again an R -module.

Definition 2.5. [22] Let M be an R -module and $S \subseteq M$ the set of finite formal linear combinations $L(S)$ of elements of S is a submodule of M . A typical element of $L(S)$ is $r_1m_1 + r_2m_2 + \dots + r_nm_n$, $r_i \in R, m_i \in S, \forall i = 1, 2, \dots, n$.

Remark 2.3. If $S \subseteq M$ and $L(S)$ is the set of all finite linear combination of elements of S , then $L(S)$ is the smallest submodule that contains S .

Definition 2.6. [13] Let M, N and P be an R -modules. A map $\varphi : M \times N \rightarrow P$ is said to be R -bilinear if $\forall m_1, m_2 \in M, n_1, n_2 \in N$ and $r \in R$, the following conditions hold

- (1) $\varphi(m, n_1 + n_2) = \varphi(m, n_1) + \varphi(m, n_2) \quad \forall m \in M$
- (2) $\varphi(m_1 + m_2, n) = \varphi(m_1, n) + \varphi(m_2, n) \quad \forall n \in N$
- (3) $\varphi(rm, n) = \varphi(m, nr) = r\varphi(m, n) \quad \forall m \in M, n \in N, r \in R.$

Definition 2.7. [1, 13] The tensor product of R -modules M and N can be denoted and defined as $M \otimes N = (M \times N)/L(S)$ where S is the set of all formal sums of the following type

Binu R & Ursala Paul, Tensor Product of Neutrosophic submodules of an R -module




FIGURE 1. Tensor product

- (1) $(rm, n) - r(m, n)$
- (2) $(m, rn) - r(m, n)$
- (3) $(m_1 + m_2, n) - (m_1, n) - (m_2, n)$
- (4) $(m, n_1 + n_2) - (m, n_1) - (m, n_2), \forall m, m_1, m_2 \in M; n, n_1, n_2 \in N; r \in R.$

Remark 2.4. (1) Being the quotient of R -module by a submodule, the tensor product $M \otimes N$ is an another R -module.

- (2) \exists a map $\tau : M \times N \rightarrow M \otimes N$ such that $\tau(m, n) = (m, n) + L(S), \forall m \in M, n \in N$ and denote $\tau(m, n)$ by $m \otimes n$.

Definition 2.8. [1, 22] The tensor product $M \otimes N$ of R -modules M and N satisfies the following properties

- (1) $(rm) \otimes n = r(m \otimes n)$
- (2) $m \otimes (rn) = r(m \otimes n)$
- (3) $(m_1 + m_2) \otimes n = (m_1 \otimes n) + (m_2 \otimes n)$
- (4) $m \otimes (n_1 + n_2) = (m \otimes n_1) + (m \otimes n_2)$

where $\forall m, m_1, m_2 \in M, n, n_1, n_2 \in N, r \in R$

Definition 2.9. [8, 22] Let M and N be two R -modules. A tensor product of M and N over R is an R -module $M \otimes N$ which is equipped with an R - bilinear map

$$\tau : M \times N \rightarrow M \otimes N$$

such that for each R -module P and each R -bilinear map $\varphi : M \times N \rightarrow P$, there is a unique homomorphism $\phi : M \otimes N \rightarrow P$, that is $\varphi = \phi \circ \tau$ (Refer Fig. 1)

Proposition 2.1. [8] “Let M and N be two R -modules. Tensor products of M and N over R are unique upto isomorphism .

Definition 2.10. [27, 31] A neutrosophic set P of the universal set X is defined as $P = \{(\eta, t_P(\eta), i_P(\eta), f_P(\eta)) : \eta \in X\}$ where $t_P, i_P, f_P : X \rightarrow (-0, 1^+)$. The three components t_P, i_P and f_P represent membership value (Percentage of truth), indeterminacy (Percentage of indeterminacy) and non membership value (Percentage of falsity) respectively. These components are functions of non standard unit interval $(-0, 1^+)$ [25].

Remark 2.5. [3, 28, 32]

- (1) If $t_P, i_P, f_P : X \rightarrow [0, 1]$, then P is known as single valued neutrosophic set (SVNS).
- (2) The algebraic structure R -module with SVNS as the underlying set is discussed in this work. SVNS will be referred to as a neutrosophic set for the sake of convenience.
- (3) U^X denotes the set of all neutrosophic subset of X or neutrosophic power set of X .

Definition 2.11. [3, 20, 30] Let $P, Q \in U^X$. Then P is contained in Q , denoted as $P \subseteq Q$ if and only if $P(\eta) \leq Q(\eta) \forall \eta \in X$, this means that $t_P(\eta) \leq t_Q(\eta), i_P(\eta) \leq i_Q(\eta), f_P(\eta) \geq f_Q(\eta), \forall \eta \in X$.

Definition 2.12. [7] "Let M be an R module. Let $P \in U^M$ where U^M denotes the neutrosophic power set of R -module M . Then a neutrosophic subset $P = \{x, t_P(x), i_P(x), f_P(x) : x \in M\}$ in M is called neutrosophic submodule of M if it satisfies the following;

- (1) $t_P(0) = 1, i_P(0) = 1, f_P(0) = 0$
- (2) $t_P(x + y) \geq t_P(x) \wedge t_P(y)$
 $i_P(x + y) \geq i_P(x) \wedge i_P(y)$
 $f_P(x + y) \leq f_P(x) \vee f_P(y), \forall x, y \in M$
- (3) $t_P(rx) \geq t_P(x), i_P(rx) \geq i_P(x), f_P(rx) \leq f_P(x), \forall x \in M, \forall r \in R$

Remark 2.6. The set of all neutrosophic submodules of R -module M represented by $U(M)$.

Definition 2.13. [3] A homomorphism Υ of M into N is called a weak neutrosophic homomorphism of P onto Q if $\Upsilon(P) \subseteq Q$. If Υ is a **weak neutrosophic homomorphism** of P onto Q , then P is weakly homomorphic to Q and we write $P \sim Q$. A homomorphism Υ of M into N is called a **neutrosophic homomorphism** of P onto Q if $\Upsilon(P) = Q$ and we represent it as $P \approx Q$.

Definition 2.14. [3] If $P = \{m, t_P(m), i_P(m), f_P(m) : m \in M\} \in U(M)$ and N be a submodule of M , then define ω , a neutrosophic set in M/N as follows.

$$\omega = \{[m], t_\omega([m]), i_\omega([m]), f_\omega([m]) : m \in M\}$$

where

$$t_\omega([m]) = \vee \{t_P(u) : u \in [m]\}$$

$$i_\omega([m]) = \vee \{i_P(u) : u \in [m]\}$$

$$f_\omega([m]) = \wedge \{f_P(u) : u \in [m]\}$$

Then $\omega \in U(M/N)$.

3. Tensor Products in Neutrosophic submodules

The tensor product between R -modules M and N is a more general than the vector space tensor product. The construction of tensor products give a most characteristic strategy for joining two modules. This section describes the construction and properties of neutrosophic tensor products.

Definition 3.1. Let M and N be two R -modules. If $A \in U(M)$ and $B \in U(N)$, then the cartesian product $A \times B$ of A and B is a neutrosophic set of $M \times N$ drfined as $(A \times B)(x, y) = \{(x, y), t_{A \times B}(x, y), i_{A \times B}(x, y), f_{A \times B}(x, y) : (x, y) \in M \times N\}$ where

$$\begin{aligned} t_{A \times B}(x, y) &= t_A(x) \wedge t_B(y) \\ i_{A \times B}(x, y) &= i_A(x) \wedge i_B(y) \\ f_{A \times B}(x, y) &= f_A(x) \vee f_B(y). \end{aligned}$$

Proposition 3.1. If $A \in U(M)$ and $B \in U(N)$, then $A \times B \in U(M \times N)$.

Proof. We prove that $A \times B$ satisfies the following conditions :

- (1) $t_{A \times B}(0, 0) = 1$, $i_{A \times B}(0, 0) = 1$ & $f_{A \times B}(0, 0) = 0$
- (2) $t_{A \times B}((x_1, y_1) + (x_2, y_2)) \geq t_{A \times B}(x_1, y_1) \wedge t_{A \times B}(x_2, y_2)$
 $i_{A \times B}((x_1, y_1) + (x_2, y_2)) \geq i_{A \times B}(x_1, y_1) \wedge i_{A \times B}(x_2, y_2)$
 $f_{A \times B}((x_1, y_1) + (x_2, y_2)) \leq f_{A \times B}(x_1, y_1) \vee f_{A \times B}(x_2, y_2) \forall (x_1, y_1), (x_2, y_2) \in M \times N$
- (3) $t_{A \times B}(r(x, y)) \geq t_{A \times B}(x, y)$
 $i_{A \times B}(r(x, y)) \geq i_{A \times B}(x, y)$
 $f_{A \times B}(r(x, y)) \leq f_{A \times B}(x, y) \forall (x, y) \in M \times N, r \in R$

1. From the definition 3.1,

$$\begin{aligned} t_{A \times B}(0, 0) &= t_A(0) \wedge t_B(0) = 1 \\ i_{A \times B}(0, 0) &= i_A(0) \wedge i_B(0) = 1 \\ f_{A \times B}(0, 0) &= f_A(0) \vee f_B(0) = 0 \end{aligned}$$

2. Now $\forall (x_1, y_1), (x_2, y_2) \in M \times N$

$$\begin{aligned} t_{A \times B}((x_1, y_1) + (x_2, y_2)) &= t_{A \times B}(x_1 + x_2, y_1 + y_2) \\ &= t_A(x_1 + x_2) \wedge t_B(y_1 + y_2) \\ &\geq (t_A(x_1) \wedge t_A(x_2)) \wedge (t_B(y_1) \wedge t_B(y_2)) \\ &= (t_A(x_1) \wedge t_B(y_1)) \wedge (t_A(x_2) \wedge t_B(y_2)) \\ &= t_{A \times B}(x_1, y_1) \wedge t_{A \times B}(x_2, y_2) \end{aligned}$$

Similarly prove that $i_{A \times B}((x_1, y_1) + (x_2, y_2)) \geq i_{A \times B}(x_1, y_1) \wedge i_{A \times B}(x_2, y_2)$ and $f_{A \times B}((x_1, y_1) + (x_2, y_2)) \leq f_{A \times B}(x_1, y_1) \vee f_{A \times B}(x_2, y_2)$.

3. Consider $\forall (x, y) \in M \times N, r \in R$

$$\begin{aligned} t_{A \times B}(r(x, y)) &= t_{A \times B}(rx, ry) \\ &= t_A(rx) \wedge t_B(ry) \\ &\geq t_A(x) \wedge t_B(y) \\ &= t_{A \times B}(x, y) \end{aligned}$$

Similarly prove that $i_{A \times B}(r(x, y)) \geq i_{A \times B}(x, y)$ and $f_{A \times B}(r(x, y)) \leq f_{A \times B}(x, y)$. Hence $A \times B \in U(M \times N)$. \square

Definition 3.2. Let $A \in U(M), B \in U(N)$ and $C \in U(P)$ where M, N and P are R modules. A map $\varphi : M \times N \rightarrow P$ is called neutrosophic R bi additive if the following conditions are hold $\forall (m, n) \in M \times N, m \in M, n \in N$

- (1) The map $\varphi : M \times N \rightarrow P$ is R bi additive
- (2) $t_C(\varphi((m, n))) \geq t_{A \times B}((m, n))$
- (3) $i_C(\varphi((m, n))) \geq i_{A \times B}((m, n))$
- (4) $f_C(\varphi((m, n))) \leq f_{A \times B}((m, n))$

Definition 3.3. Let $A \in U(L(M \times N)) = U(Y)$ and $Y(S)$ be a submodule of $L(M \times N) = Y$. Then the neutrosophic tensor product of R -modules M and N , $(M \otimes N)$, is a neutrosophic set Q of $Y/Y(S)$ defined as follows

$$Q((m, n) + Y(S)) = \{(m, n) + Y(S), t_Q((m, n) + Y(S), i_Q((m, n) + Y(S), f_Q((m, n) + Y(S))\}$$

$\forall (m, n) \in M \times N, m \in M, n \in N$ where

$$\begin{aligned} t_Q((m, n) + Y(S)) &= \vee \{t_A((m, n) + y(S)) : y(S) \in Y(S)\} \\ i_Q((m, n) + Y(S)) &= \vee \{i_A((m, n) + y(S)) : y(S) \in Y(S)\} \\ f_Q((m, n) + Y(S)) &= \wedge \{f_A((m, n) + y(S)) : y(S) \in Y(S)\} \end{aligned}$$

Remark: The coset $(m, n) + Y(S)$ is represented by $[(m, n)]$

Theorem 3.1. Let Q be the neutrosophic tensor product of $M \otimes N$, then $Q \in U(Y/Y(S))$.

Proof. We have

$$Q([(m, n)]) = \{[(m, n)], t_Q([(m, n)]), i_Q([(m, n)]), f_Q([(m, n)]) : (m, n) \in M \otimes N, m \in M, n \in N\}$$

and $A \in U(L(M \times N)) = U(Y)$ and $Y(S)$ be a submodule of $L(M \times N) = Y$. Also

$$t_Q([(m, n)]) = \vee \{t_A(x, y) : (x, y) \in [m, n]\}$$

$$i_Q([(m, n)]) = \vee \{i_A(x, y) : (x, y) \in [m, n]\}$$

$$f_Q([(m, n)]) = \wedge \{f_A(x, y) : (x, y) \in [m, n]\}$$

We have $t_Q([0, 0]) = \vee \{t_A(x, y) : (x, y) \in [0, 0]\} = t_A(0) = 1$, similarly $i_Q([0]) = 1$ and

$$f_Q([0, 0]) = \wedge \{f_A(x, y) : (x, y) \in [0, 0]\} = f_A(0) = 0$$

Now for $(m_1, n_1), (m_2, n_2) \in M \otimes N$

$$\begin{aligned} t_Q([(m_1, n_1)] + [(m_2, n_2)]) &= \vee \{t_A(x, y) : (x, y) \in [(m_1, n_1)] + [(m_2, n_2)]\} \\ &= \vee \{t_A((x_1, y_1) + (x_2, y_2)) : (x_1, y_1) + (x_2, y_2) \in [(m_1, n_1)] + [(m_2, n_2)]\} \\ &\geq \vee \{t_A((x_1, y_1) + (x_2, y_2)) : (x_1, y_1) \in [(m_1, n_1)], (x_2, y_2) \in [(m_2, n_2)]\} \\ &\geq \vee \{t_A(x_1, y_1) \wedge t_A(x_2, y_2) : (x_1, y_1) \in [(m_1, n_1)], (x_2, y_2) \in [(m_2, n_2)]\} \\ &= (\vee \{t_A((x_1, y_1)) : (x_1, y_1) \in [m_1, n_1]\}) \wedge \\ &\quad (\vee \{t_A((x_2, y_2)) : (x_2, y_2) \in [m_2, n_2]\}) \\ &= t_Q([m_1, n_1]) + t_Q([m_2, n_2]) \end{aligned}$$

Similarly we can prove that

$$i_Q([m_1, n_1] + [m_2, n_2]) \geq i_Q([(m_1, n_1)]) \wedge i_Q([(m_2, n_2)])$$

and

$$f_Q([(m_1, n_1)] + [(m_2, n_2)]) \leq f_Q([(m_1, n_1)]) \vee f_Q([(m_2, n_2)])$$

Now for any $r \in R, (m, n) \in M \otimes N$,

$$\begin{aligned} t_Q(r[(m, n)]) &= t_Q([r(m, n)]) \\ &= \vee \{t_A(x, y) : (x, y) \in [r(m, n)]\} \\ &= \vee \{t_A(r(x, y) + y(s) : y(s) \in Y(S))\} \\ &\geq \vee \{t_A(r(x, y) + ry_1(S)) : y_1(S) \in Y(S)\} \\ &= \vee \{t_A(r((x, y) + y_1(S))) : y_1(S) \in Y(S)\} \\ &\geq \vee \{t_A((x, y) + y_1(S)) : y_1(S) \in Y(S)\} \\ &= \vee \{t_A(x_1, y_1) : (x_1, y_1) \in [m, n]\} \\ &= t_Q([m, n]) \end{aligned}$$

Similarly we can prove that

$$i_Q(r[(m, n)]) \geq i_Q([(m, n)]) \text{ and } f_Q(r[(m, n)]) \geq f_Q([(m, n)])$$

Thus $Q \in U(Y/Y(S))$. \square

Definition 3.4. A pair $(M \otimes N, \tau)$ or a map $\tau : M \times N \rightarrow M \otimes N$ is said to be the tensor product of A and B where $A \in U(M)$ and $B \in U(N)$ if for every neutrosophic R bi additive map $\varphi : M \times N \rightarrow P$ of $A \times B$ to C , there is unique neutrosophic homomorphism $\phi : M \otimes N \rightarrow P$ of $A \otimes B$ onto C such that $\phi \circ \tau = \varphi$ where

$$t_C(\phi(m \otimes n)) \geq t_{A \times B}(m, n)$$

$$i_C(\phi(m \otimes n)) \geq i_{A \times B}(m, n)$$

$$f_C(\phi(m \otimes n)) \leq f_{A \times B}(m, n)$$

Theorem 3.2. The tensor product of two neutrosophic R modules exists and it is unique up to isomorphism.

Proof. Let $A \in U(M)$, $B \in U(N)$ and $\tau : M \times N \rightarrow M \otimes N$ be the tensor product of R -modules M and N . Then by the definition of 3.4, for every neutrosophic R bi additive map $\varphi : M \times N \rightarrow P$ where P be an R module, there is unique neutrosophic homomorphism $\phi : M \otimes N \rightarrow P$ such that $\phi \circ \tau = \varphi$.

Now define a map $A \otimes B : M \otimes N \rightarrow [0, 1]$ by putting

$$A \otimes B(m \otimes n) = \{(m \otimes n), t_{A \otimes B}(m \otimes n), i_{A \otimes B}(m \otimes n), f_{A \otimes B}(m \otimes n)\}$$

where

$$t_{A \otimes B}(m \otimes n) = \bigvee \{t_{A \times B}(m', n') : (m' \otimes n') = (m \otimes n)\}$$

$$i_{A \otimes B}(m \otimes n) = \bigvee \{i_{A \times B}(m', n') : (m' \otimes n') = (m \otimes n)\}$$

$$f_{A \otimes B}(m \otimes n) = \bigwedge \{f_{A \times B}(m', n') : (m' \otimes n') = (m \otimes n)\}$$

Let $C \in U(P)$, then to prove that, $\forall m \otimes n \in M \otimes N$

$$t_C(\phi(m \otimes n)) \geq t_{A \otimes B}(m \otimes n)$$

$$i_C(\phi(m \otimes n)) \geq i_{A \otimes B}(m \otimes n)$$

$$f_C(\phi(m \otimes n)) \leq f_{A \otimes B}(m \otimes n)$$

Suppose $(m' \otimes n') = (m \otimes n) \in M \otimes N$

$$\begin{aligned} t_C(\phi(m' \otimes n')) &\geq \bigvee \{t_C(\phi \circ \tau)(m', n')\} \\ &= \bigvee \{t_{A \times B}(m, n')\} \\ &= t_{A \otimes B}(m \otimes n) \end{aligned}$$

□

Definition 3.5. Let A be left neutrosophic R -module of left R -module M and let B be right neutrosophic R -module of right R -module N . Let C be a neutrosophic abelian group and $\tilde{g} : A \times B \rightarrow C$ be neutrosophic biadditive. A pair (C, \tilde{g}) is called a tensor product of A and B if for every fuzzy biadditive $F : A \times B \rightarrow H$ where H is a neutrosophic abelian group, there is a unique neutrosophic map $\theta \in Hom(C, H)$ such that $\theta \circ \tilde{g} = F$.

Theorem 3.3. Let A be left neutrosophic R -module of left R -module M and let B be right neutrosophic R -module of right R -module N . The tensor product of the two neutrosophic R -modules A and B exist and it is unique up to isomorphism.

Proof. Let $\varphi : M \times N \rightarrow M \otimes N$ be the tensor products of R -modules A and B . Then we can define the maps $t, i, f : M \otimes N \rightarrow [0, 1]$ such that $\forall i$

$$\begin{aligned} t_{A \otimes B}(\sum (a_i \otimes b_i)) &= \bigvee \{t_{A \times B}(\sum (a_i', b_i')) : \sum (a_i' \otimes b_i') = \sum (a_i \otimes b_i)\} \\ i_{A \otimes B}(\sum (a_i \otimes b_i)) &= \bigvee \{i_{A \times B}(\sum (a_i', b_i')) : \sum (a_i' \otimes b_i') = \sum (a_i \otimes b_i)\} \\ f_{A \otimes B}(\sum (a_i \otimes b_i)) &= \bigwedge \{f_{A \times B}(\sum (a_i', b_i')) : \sum (a_i' \otimes b_i') = \sum (a_i \otimes b_i)\} \end{aligned}$$

Then $\varphi : A \times B \rightarrow A \otimes B$ is neutrosophic biadditive and \tilde{H} be a neutrosophic abelian group and $\psi : A \times B \rightarrow \tilde{H}$ be a neutrosophic biadditive. Thn by definition of tensor product, \exists a unique homomorphism $\theta : M \otimes N \rightarrow H$ such that $\theta \circ \psi = \varphi$.

Now we have to show that $\sum (a_i \otimes b_i) \in A \otimes B$ and

$$\begin{aligned} t_H(\theta(\sum (a_i \otimes b_i))) &\geq t_{A \otimes B}(\sum (a_i \otimes b_i)) \\ i_H(\theta(\sum (a_i \otimes b_i))) &\geq i_{A \otimes B}(\sum (a_i \otimes b_i)) \\ f_H(\theta(\sum (a_i \otimes b_i))) &\leq f_{A \otimes B}(\sum (a_i \otimes b_i)) \end{aligned}$$

Suppose $\sum (a_i' \otimes b_i') = \sum (a_i \otimes b_i) \in M \otimes N$

$$\begin{aligned} t_H(\theta \sum (a_i' \otimes b_i')) &= t_H(\sum (\theta(a_i' \otimes b_i'))) \\ &\geq \bigwedge \{t_H \theta(a_i' \otimes b_i')\} \\ &= \bigwedge \{t_H \theta \varphi(a_i', b_i')\} \\ &= \bigwedge \{t_H \psi(a_i', b_i')\} \\ &\geq \bigwedge \{t_{A \times B}(a_i', b_i')\} \\ &= t_{A \times B}(a_i', b_i') \\ &= t_{A \otimes B}(\sum (a_i' \otimes b_i')) \end{aligned}$$

Similarly ,

$$i_H(\theta(\sum (a_i \otimes b_i))) \geq i_{A \otimes B}(\sum (a_i \otimes b_i))$$

$$f_H(\theta(\sum(a_i \otimes b_i))) \leq f_{A \otimes B}(\sum(a_i \otimes b_i))$$

This concludes that $\sum(a_i \otimes b_i) \in A \otimes B$ and $A \otimes B$ is a tensor product of A and B . Also it is obvious that tensor product is unique up to isomorphism. \square

Remark 3.1 : Let A and B be two neutrosophic right and left R module, then $0_R \otimes A \cong A$ and $B \otimes 0_R \cong B$

4. Conclusion

The concept of tensor product is great significance in classical algebra, geometry and analysis. In the emerging algebraic research domain, the amalgamation of tensor product in a neutrosophic submodule context leads to the design of the most flexible version of algebraic product. In this research a neutrosophic quotient submodule of an R -module is constructed as a tensor product in neutrosophic submodules of an R -modules. It will definitely lead to the development of new theoretical and practical techniques for problem solving in the fields of classical and quantum mechanics, image processing and neural networks, artificial intelligence and machine learning. The concept of tensor product in neutrosophic submodules is an imminent tool for the vague multi dimensional real world big data processing and analysis. In our future research, we propose to extend the concept of exact sequences to use tensor factorization in the neutrosophic domain, as well as the associative property of the relationship between neutrosophic injective and projective modules. The above mentioned study leads to the homological properties of neutrosophic submodule tensor products and neutrosophic module category theory.

Conflicts of Interest: Declare conflicts of interest or state "The authors declare no conflict of interest."

References

1. ARTIN, M. *Algebra*. Pearson Prentice Hall, 2011.
2. ATANASSOV, K. Review and new results on intuitionistic fuzzy sets. *preprint Im-MFAIS-1-88, Sofia 5* (1988), 1.
3. BINU, R., AND ISAAC, P. Neutrosophic quotient submodules and homomorphisms. *Punjab University Journal of Mathematics*.
4. BINU, R., AND ISAAC, P. Some characterizations of neutrosophic submodules of an-module. *Applied Mathematics and Nonlinear Sciences 6*, 1 (2021), 359–372.
5. ÇETKIN, V., AND AYGÜN, H. An approach to neutrosophic subgroup and its fundamental properties. *Journal of Intelligent & Fuzzy Systems 29*, 5 (2015), 1941–1947.
6. ÇETKIN, V., AND AYGÜN, H. An approach to neutrosophic subrings. *Sakarya University Journal of Science 23*, 3 (2019), 472–477.
7. ÇETKIN, V., VAROL, B. P., AND AYGÜN, H. On neutrosophic submodules of a module. *Hacettepe Journal of Mathematics and Statistics 46*, 5 (2017), 791–799.

8. DUMMIT, D., AND FOOTE, R. *Abstract Algebra*. Wiley, 2003.
9. FERNADEZ, S. Fuzzy g -modules and fuzzy representations. *TAJOPAM 1* (2002), 107–114.
10. FUTA, Y., OKAZAKI, H., AND SHIDAMA, Y. Quotient module of z -module. *Formalized Mathematics 20*, 3 (2012), 205–214.
11. GULISTAN, M., NAWAZ, S., AND HASSAN, N. Neutrosophic triplet non-associative semihypergroups with application. *Symmetry 10*, 11 (2018), 613.
12. GULISTAN, M., YAQOUB, N., NAWAZ, S., AND AZHAR, M. A study of (α, β) -complex fuzzy hyperideals in non-associative hyperring. *Journal of Intelligent & Fuzzy Systems 36*, 6 (2019), 6025–6036.
13. HUANG, Y.-Z., AND LEPOWSKY, J. A theory of tensor products for module categories for a vertex operator algebra, ii. *Selecta Mathematica 1*, 4 (1995), 757.
14. ISAAC, P., AND JOHN, P. P. On intuitionistic fuzzy submodules of a module. *Int. J. of Mathematical Sciences and Applications 1*, 3 (2011), 1447–1454.
15. ISAAC, P., AND PAUL, U. Rough g -modules and their properties. *Advances in Fuzzy Mathematics 12*, 1 (2017), 93–100.
16. LAM, T. Y. Representations of finite groups: A hundred years, part i. *Notices of the AMS 45*, 3 (1998), 361–372.
17. LÓPEZ-PERMOUTH, S. R. Lifting morita equivalence to categories of fuzzy modules. *Information sciences 64*, 3 (1992), 191–201.
18. MASHINCHI, M., AND ZAHEDI, M. On l -fuzzy primary submodules. *Fuzzy sets and systems 49*, 2 (1992), 231–236.
19. MOI, S., BISWAS, S., AND PAL, S. Second-order neutrosophic boundary-value problem. *Complex & Intelligent Systems 7*, 2 (2021), 1079–1098.
20. MONDAL, K., AND PRAMANIK, S. Neutrosophic decision making model of school choice. *Neutrosophic Sets and Systems 7* (2015), 62–68.
21. NEGOIȚĂ, C. V., AND RALESCU, D. A. *Applications of fuzzy sets to systems analysis*. Springer, 1975.
22. OSTRIK, V. Module categories, weak hopf algebras and modular invariants. *Transformation Groups 8*, 2 (2003), 177–206.
23. PADILLA, R. Smarandache algebraic structures. *Bulletin of Pure and Applied Sciences, Delhi 17*, 1 (1998), 119–121.
24. PAUL, U., AND ISAAC, P. Fuzzy lattice ordered g -modules. *International Journal of Fuzzy System Applications 8*, 3 (2019), 94–107.
25. ROBINSON, A. *Non-standard analysis*. Princeton University Press, 2016.
26. SMARANDACHE, F. Neutrosophy: neutrosophic probability, set, and logic: analytic synthesis & synthetic analysis.
27. SMARANDACHE, F. Neutrosophic set—a generalization of the intuitionistic fuzzy set. *International journal of pure and applied mathematics 24*, 3 (2005), 287.
28. SMARANDACHE, F. Neutrosophic set—a generalization of the intuitionistic fuzzy set. *Journal of Defense Resources Management (JoDRM) 1*, 1 (2010), 107–116.
29. SMARANDACHE, F., AND ALI, M. Neutrosophic triplet group. *Neural Computing and Applications 29*, 7 (2018), 595–601.
30. SMARANDACHE, F., GÎFU, D., AND TEODORESCU, M. Neutrosophic elements in discourse. *Social Sciences and Education Research Review, 2* (1) (2015).
31. SMARANDACHE, F., AND PRAMANIK, S. *New trends in neutrosophic theory and applications*, vol. 1. Infinite Study, 2016.
32. SMARANDACHE, F., ZHANG, Y., AND SUNDERRAMAN, R. Single valued neutrosophic sets. *Neutrosophy: neutrosophic probability, set and logic 4* (2009), 126–129.

-
33. VASANTHA KANDASAMY, W. Smarandache neutrosophic algebraic structures. *arXiv Mathematics e-prints* (2006), math-0603708.
 34. WANG, H., SMARANDACHE, F., ZHANG, Y., AND SUNDERRAMAN, R. *Single valued neutrosophic sets*. Infinite study, 2010.
 35. ZADEH, L. A. Fuzzy sets. *Informtaion and control* 8, 3 (1965), 338–353.

Received: Nov. 7, 2021. Accepted: April 7, 2022.