

6-10-2022

On Characterizations of $(\$, \epsilon, \zeta)$ -Single Valued Neutrosophic Hyperrings and Hyperideals

Muhammad Shazib Hameed

Zaheer Ahmad

Said Broumi

Shahbaz Ali

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Hameed, Muhammad Shazib; Zaheer Ahmad; Said Broumi; and Shahbaz Ali. "On Characterizations of $(\$, \epsilon, \zeta)$ -Single Valued Neutrosophic Hyperrings and Hyperideals." *Neutrosophic Sets and Systems* 50, 1 (2022). https://digitalrepository.unm.edu/nss_journal/vol50/iss1/37

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



On Characterizations of $(\varpi, \varepsilon, \varsigma)$ -Single Valued Neutrosophic Hyperrings and Hyperideals

Muhammad Shazib Hameed¹, Zaheer Ahmad¹, Said Broumi² and Shahbaz Ali^{1,*}

¹Department of Mathematics, Khwaja Fareed University of Engineering and Information Technology, Rahim Yar Khan 64200, Pakistan; shazib.hameed@kfueit.edu.pk; zaheer@gmail.com; shahbaz.ali@kfueit.edu.pk

²Laboratory of Information Processing, Faculty of Science Ben MSik, University Hassan II, Casablanca, Morocco; broumisaid78@gmail.com

*Correspondence: shahbaz.ali@kfueit.edu.pk

Abstract. This study included concepts for $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic hyperring $((\varpi, \varepsilon, \varsigma)\text{-SVNHR})$ and $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic hyperideal $((\varpi, \varepsilon, \varsigma)\text{-SVNHI})$. $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic hyperrings $((\varpi, \varepsilon, \varsigma)\text{-SVNHRs})$ and $(\varpi, \varepsilon, \varsigma)$ -single valued hyperideals $((\varpi, \varepsilon, \varsigma)\text{-SVNHIs})$ are examined and validated in terms of algebraic features and of structural characteristics.

Keywords: Hyperring, Hyperideal, Single valued neutrosophic set, Single valued neutrosophic hyperring, Single valued neutrosophic hyperideal, $(\varpi, \varepsilon, \varsigma)$ -Single valued neutrosophic hyperring and $(\varpi, \varepsilon, \varsigma)$ -Single valued neutrosophic hyperideal.

1. Introduction

The theory of hyperstructure came into existence in 1934 when Marty [1] defined hypergroups as being generalized. To propose an overlying homomorphism, Corsini [2] developed the concept of hypering and general forms of hypering. The H_v -ring, the H_v -subring, and the H_v -ideals of the H_v -ring, all these are modifying the thoughts introduced by Corsini [2], have been invented by Vougiouklis [3, 4]. Generally, [5] offers a variety of rates in $[0, 1]$ stated by a single real number. In order to relieve ambiguities, a fuzzy set model was created by the Turksen [6], which was utilized to assess membership of the fuzzy set framework. An enhancement of fuzzy sets is intuitionistic sets, suggested in 1986 by Atanassov [7]. This approach was analogous to the interval-valued fuzzy sets described in [8]. Intuitionistic fuzzy sets can execute flawed data and not inexhaustible information, frequently in real-life [8]. Rosenfeld [9] launched the fuzzy algebra work, extending it to several fuzzy models such as intuitionistic fuzzy sets, fuzzy soft sets, and imprecise soft sets. Some artworks related to soft, fuzzy rings

and ideal vague soft groups, vague soft rings, and soft ideals are also found in [10–13]. In 1998, to attain these goals, Smarandache suggested the neutrosophic paradigm in [14]. In [15–20], numerous new neutrosophic theoretical fads were launched.

Wang et al. [8] pioneered the theory of a single-valued neutrosophic set (*SVNS*), whereas Smarandache plithogenic was presented into [21] as a refinement of neutrosophic structure. Hyperstructure theory is often used in numerous mathematical ideas. Algebraic hyperstructures have a wide range of applications, including fuzzy sets, design and data, artificial intelligence, lattices, automation, and combinatorics, and etc. As a result of fuzzy algebra research, fuzzy hyperalgebraic theory was produced. Liu [22] created the idea of fuzzy ideals of a ring. A lot of hyperstructure work has been done over the last two decades, such as fuzzy hyperalgebras [23], fuzzy hyperrings [24], fuzzy topological F-polygroups [25], Bipolar-valued fuzzy soft hyper BCK ideals [26], fuzzy hypergroup degree [27], fuzzy hypergraphs [28], hyper-spectral image analysis [29], fundamental relations on fuzzy hypermodules [30], and so on.

There are works available of hyperstructures related to hyperrings in these manuscripts: fuzzy hyperings [31], Γ -hyperrings [32], soft hyperrings [33], topological hyperrings [34], and topological structures of lower and upper rough subsets in a hyperring [35], etc. In [36], Davvaz initiated the generalization of fuzzy hyperideal. Bharathi and Vimala subsequently established the notions of fuzzy l -ideal in [37], and the fuzzy l -ideal was then expanded in [38]. In [39–41], Selvachandran et al. introduced the hypergroup and hyperring theory for imprecise soft sets, and some other important works on fuzzy sets are studied in [42–45].

In this paper, we focus on the theories of $(\varpi, \varepsilon, \varsigma)$ -*SVNHRs* and $(\varpi, \varepsilon, \varsigma)$ -*SVNHIs* in order to contribute to the advancement of the neutrosophic theory of hyperalgebraic.

2. Preliminaries

Let Ξ be a set of points where \hat{n} refers to a generic element of Ξ .

Definition 2.1. [8] A *SVNS* Υ neutrosophic set that is characterized by a truth membership function $\tau_{\Upsilon}(\hat{n})$, an indeterminacy-membership function $\iota_{\Upsilon}(\hat{n})$, and a falsity-membership function $F_{\Upsilon}(\hat{n})$, where $\tau_{\Upsilon}(\hat{n}), \iota_{\Upsilon}(\hat{n}), F_{\Upsilon}(\hat{n}) \in [0, 1]$. This set Υ can thus be written as:

$$\Upsilon = \{(\hat{n}, \tau_{\Upsilon}(\hat{n}), \iota_{\Upsilon}(\hat{n}), F_{\Upsilon}(\hat{n})) : \hat{n} \in \Xi\}.$$

The sum of $\tau_{\Upsilon}(\hat{n})$, $\iota_{\Upsilon}(\hat{n})$ and $F_{\Upsilon}(\hat{n})$ must fulfill the clause $0 \leq \tau_{\Upsilon}(\hat{n}) + \iota_{\Upsilon}(\hat{n}) + F_{\Upsilon}(\hat{n}) \leq 3$. For a *SVNS* Υ in Ξ , the triplet $(\tau_{\Upsilon}(\hat{n}), \iota_{\Upsilon}(\hat{n}), F_{\Upsilon}(\hat{n}))$ is referred to as a single valued neutrosophic number (SVNN). Let $\hat{n} = (\tau_{\hat{n}}, \iota_{\hat{n}}, F_{\hat{n}})$ stand for a SVNN.

Definition 2.2. [8] Assume Υ and Γ are two *SVNSs* in a universe Ξ .

- (1) Υ is contained in Γ , if $\tau_{\Upsilon}(\hat{n}) \leq \tau_{\Gamma}(\hat{n})$, $\iota_{\Upsilon}(\hat{n}) \leq \iota_{\Gamma}(\hat{n})$, and $F_{\Upsilon}(\hat{n}) \geq F_{\Gamma}(\hat{n})$, $\forall \hat{n} \in \Xi$.

This relationship is denoted as $\Upsilon \subseteq \Gamma$.

- (2) $\Upsilon = \Gamma$ if $\Upsilon \subseteq \Gamma$ and $\Gamma \subseteq \Upsilon$.
- (3) $\Upsilon^c = (\hat{n}, (F_{\Upsilon}(\hat{n}), 1 - \iota_{\Upsilon}(\hat{n}), \tau_{\Upsilon}(\hat{n})), \forall \hat{n} \in \Xi$.
- (4) $\Upsilon \cup \Gamma = (\hat{n}, (\bigvee(\tau_{\Upsilon}, \tau_{\Gamma}), \bigvee(\iota_{\Upsilon}, \iota_{\Gamma}), \bigwedge(F_{\Upsilon}, F_{\Gamma}))), \forall \hat{n} \in \Xi$.
- (5) $\Upsilon \cap \Gamma = (\hat{n}, (\bigwedge(\tau_{\Upsilon}, \tau_{\Gamma}), \bigwedge(\iota_{\Upsilon}, \iota_{\Gamma}), \bigvee(F_{\Upsilon}, F_{\Gamma}))), \forall \hat{n} \in \Xi$.

Definition 2.3. [1] A hypergroup $\langle H, \circ \rangle$ is a set H with an associative hyperoperation $(\circ) : H * H \rightarrow P(H)$ which satisfies $\hat{n} \circ H = H \circ \hat{n} = H, \forall \hat{n} \in H$ (reproduction axiom).

Definition 2.4. [36] If the following properties satisfy, a hyperstructure $\langle H, \circ \rangle$ is termed a H_v -group:

- (1) $\hat{n} \circ (\hat{o} \circ \hat{p}) \cap (\hat{n} \circ \hat{o}) \circ \hat{p} \neq \phi, \forall \hat{n}, \hat{o}, \hat{p} \in H, (H_v\text{-semigroup})$.
- (2) $\hat{n} \circ H = H \circ \hat{n} = H, \forall \hat{n} \in H$.

Definition 2.5. [1] A subset W of H is termed as subhypergroup if $\langle W, \circ \rangle$ is a hypergroup.

Definition 2.6. [2] A H_v -ring is a multi-valued system $(R, +, \circ)$ that satisfies the following axioms:

- (1) $(R, +)$ must a H_v -group,
- (2) (R, \circ) must a H_v -semigroup,
- (3) The hyperoperation “ \circ ” is weak distributive over the hyperoperation “ $+$ ”, that is for each $\hat{n}, \hat{o}, \hat{p} \in R$ the clauses $\hat{n} \circ (\hat{o} + \hat{p}) \cap ((\hat{n} \circ \hat{o}) + (\hat{n} \circ \hat{p})) \neq \phi$ and $(\hat{n} + \hat{o}) \circ \hat{p} \cap ((\hat{n} \circ \hat{p}) + (\hat{o} \circ \hat{p})) \neq \phi$ must satisfy.

Definition 2.7. [2] A nonempty subset R' of R is a subhyperring of $(R, +, \circ)$ if $(R', +)$ is a subhypergroup of $(R, +)$ and $\forall \hat{n}, \hat{o}, \hat{p} \in R', \hat{n} \circ \hat{o} \in P^*(R')$, where $P^*(R')$ denotes the set of all non-empty subsets of R' .

Definition 2.8. [2] Suppose H_v -ring be R . a nonempty subset I of R is called a left (resp. right) H_v -ideal if the following axioms hold:

- (1) $(I, +)$ be a H_v -subgroup of $(R, +)$,
- (2) $R \circ I \subseteq I$ (resp. $I \circ R \subseteq I$).

If I is both a left and right H_v -ideal of R , then I is called H_v -ideal of R .

3. $(\varpi, \varepsilon, \varsigma)$ -Single Valued Neutrosophic Hyperrings

We represent hyperring $(R, +, \circ)$ by R throughout this section.

Definition 3.1. If Υ be a single valued neutrosophic subset of Ξ then $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic Υ subset of Ξ is categorize as,

$$\Upsilon^{(\varpi, \varepsilon, \varsigma)} = \left\{ \langle \hat{n}, \tau_{\Upsilon}^{\varpi}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{n}) \mid \tau_{\Upsilon}^{\varpi}(\hat{n}) = \bigwedge \{ \tau_{\Upsilon}(\hat{n}), \varpi \}, \iota_{\Upsilon}^{\varepsilon}(\hat{n}) = \bigwedge \{ \iota_{\Upsilon}(\hat{n}), \varepsilon \}, F_{\Upsilon}^{\varsigma}(\hat{n}) = \bigvee \{ F_{\Upsilon}(\hat{n}), \varsigma \}, \hat{n} \in \Xi \right\},$$

and $0 \leq \tau_{\Upsilon}^{\varpi}(\hat{n}) + \iota_{\Upsilon}^{\varepsilon}(\hat{n}) + F_{\Upsilon}^{\varsigma}(\hat{n}) \leq 3$, where $\varpi, \varepsilon, \varsigma \in [0, 1]$ also $\tau, \iota, F : \Upsilon \rightarrow [0, 1]$, such that $\tau_{\Upsilon}^{\varpi}, \iota_{\Upsilon}^{\varepsilon}, F_{\Upsilon}^{\varsigma}$ represents the functions of truth, indeterminacy, and falsity-membership, respectively.

Definition 3.2. Let Υ be a $(\varpi, \varepsilon, \varsigma)$ -SVNS over R . Then Υ is called a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R , if,

- (1) $\forall \hat{k}, \hat{l} \in R,$
 $\bigwedge \{ \tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l}) \} \leq \inf \{ \tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l} \},$
 $\bigvee \{ \iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \} \geq \sup \{ \iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l} \},$ and
 $\bigvee \{ F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l}) \} \geq \sup \{ F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l} \}.$
- (2) $\forall \hat{n}, \hat{k} \in R, \exists \hat{l} \in R$ such that $\hat{k} \in \hat{n} + \hat{l}$ and
 $\bigwedge \{ \tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k}) \} \leq \tau_{\Upsilon}^{\varpi}(\hat{l}),$
 $\bigvee \{ \iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k}) \} \geq \iota_{\Upsilon}^{\varepsilon}(\hat{l}),$ and
 $\bigvee \{ F_{\Upsilon}^{\varsigma}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{k}) \} \geq F_{\Upsilon}^{\varsigma}(\hat{l})$
- (3) $\forall \hat{n}, \hat{k} \in R, \exists \hat{m} \in R$ such that $\hat{k} \in \hat{m} + \hat{n}$ and
 $\bigwedge \{ \tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k}) \} \leq \tau_{\Upsilon}^{\varpi}(\hat{m}),$
 $\bigvee \{ \iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k}) \} \geq \iota_{\Upsilon}^{\varepsilon}(\hat{m}),$ and
 $\bigvee \{ F_{\Upsilon}^{\varsigma}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{k}) \} \geq F_{\Upsilon}^{\varsigma}(\hat{m}).$
- (4) $\forall \hat{k}, \hat{l} \in R,$
 $\bigwedge \{ \tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l}) \} \leq \inf \{ \tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l} \},$
 $\bigvee \{ \iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \} \geq \sup \{ \iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l} \},$ and
 $\bigvee \{ F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l}) \} \geq \sup \{ F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l} \}.$

Example 3.3. The family of t -level sets of $(\varpi, \varepsilon, \varsigma)$ -SVNSs over R is a subhyperring of R is resulting below:

$$\Upsilon_t^{(\varpi, \varepsilon, \varsigma)} = \{ \hat{k} \in R : \tau_{\Upsilon}^{\varpi}(\hat{k}) \geq t, \iota_{\Upsilon}^{\varepsilon}(\hat{k}) \leq t, F_{\Upsilon}^{\varsigma}(\hat{k}) \leq t \}, \forall t \in [0, 1].$$

Then Υ over R is a $(\varpi, \varepsilon, \varsigma)$ -SVNHR.

Theorem 3.4. Υ is a $(\varpi, \varepsilon, \varsigma)$ -SVNS over R . Then Υ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R if and only if Υ is $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic semi hypergroup over (R, \circ) and also a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic hypergroup over $(R, +)$.

Proof. The definition 3.2 readily indicates this proof. \square

Proposition 3.5. If Υ and Γ be two $(\varpi, \varepsilon, \varsigma)$ -single-valued neutrosophic subset of ring R then $(\Upsilon \cap \Gamma)^{(\varpi, \varepsilon, \varsigma)} = \Upsilon^{(\varpi, \varepsilon, \varsigma)} \cap \Gamma^{(\varpi, \varepsilon, \varsigma)}$.

Proof. Assume that Υ and Γ are two $(\varpi, \varepsilon, \varsigma)$ -single-valued neutrosophic subset of ring R .

$$\begin{aligned} (\Upsilon \cap \Gamma)^{(\varpi, \varepsilon, \varsigma)}(\hat{n}) &= \left\{ \min\{\min\{\tau_{\Upsilon}(\hat{n}), \tau_{\Gamma}(\hat{n})\}, \varpi\}, \min\{\min\{\iota_{\Upsilon}(\hat{n}), \iota_{\Gamma}(\hat{n})\}, \varepsilon\}, \max\{\max\{F_{\Upsilon}(\hat{n}), F_{\Gamma}(\hat{n})\}, \varsigma\} \right\} \\ &= \left\{ \min\{\min\{\tau_{\Upsilon}(\hat{n}), \varpi\}, \min\{\tau_{\Gamma}(\hat{n}), \varpi\}\}, \min\{\min\{\iota_{\Upsilon}(\hat{n}), \varepsilon\}, \max\{\iota_{\Gamma}(\hat{n}), \varepsilon\}\}, \max\{\max\{F_{\Upsilon}(\hat{n}), \varsigma\}, \max\{F_{\Gamma}(\hat{n}), \varsigma\}\} \right\} \\ &= \left\{ \min(\{\tau_{\Upsilon}^{\varpi}(\hat{n})\}, \{\tau_{\Gamma}^{\varpi}(\hat{n})\}), \min(\{\iota_{\Upsilon}^{\varepsilon}(\hat{n})\}, \{\iota_{\Gamma}^{\varepsilon}(\hat{n})\}), \max(\{F_{\Upsilon}^{\varsigma}(\hat{n})\}, \{F_{\Gamma}^{\varsigma}(\hat{n})\}) \right\} = \Upsilon^{(\varpi, \varepsilon, \varsigma)}(\hat{n}) \cap \Gamma^{(\varpi, \varepsilon, \varsigma)}(\hat{n}), \forall \hat{n} \in R. \end{aligned}$$

□

Theorem 3.6. *Let Υ and Γ be $(\varpi, \varepsilon, \varsigma)$ -SVNHRs over R . Then $\Upsilon \cap \Gamma$ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R if it is non-null.*

Proof. Let Υ and Γ are $(\varpi, \varepsilon, \varsigma)$ -SVNHRs over R . By using Definition 3.2, and Proposition 3.5

$$(\Upsilon \cap \Gamma)^{(\varpi, \varepsilon, \varsigma)} = \Upsilon^{(\varpi, \varepsilon, \varsigma)} \cap \Gamma^{(\varpi, \varepsilon, \varsigma)} = \{ \langle \hat{k}, (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k}) \rangle : \hat{k} \in R \},$$

where

$$\begin{aligned} (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}) &= \wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k})), \\ (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}) &= \wedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k})), \\ (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k}) &= \vee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Gamma}^{\varsigma}(\hat{k})). \end{aligned}$$

Assuming $\forall \hat{k}, \hat{l} \in R$, we are only proven to include all four clauses for membership terms τ_{Υ}^{ϖ} , τ_{Γ}^{ϖ} and indeterminacy terms $\iota_{\Upsilon}^{\varepsilon}$, $\iota_{\Gamma}^{\varepsilon}$. Indications for falsity functions of F_{Υ}^{ς} , F_{Γ}^{ς} correspondingly derived.

$$\begin{aligned} (1) \quad \wedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l})\} &= \wedge\{\wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k})), \wedge(\tau_{\Upsilon}^{\varpi}(\hat{l}), \tau_{\Gamma}^{\varpi}(\hat{l}))\} \\ &\leq \wedge\{\wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})), \wedge(\tau_{\Gamma}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{l}))\} \\ &\leq \wedge\{\inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}, \inf\{\tau_{\Gamma}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}\} \\ &\leq \inf\{\wedge(\tau_{\Upsilon}^{\varpi}(\hat{m}), \tau_{\Gamma}^{\varpi}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\} \\ &= \inf\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \\ \Rightarrow \quad \wedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l})\} &\leq \inf\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \end{aligned}$$

Also

$$\begin{aligned} \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{l})\} &= \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k})), \bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{l}), \iota_{\Gamma}^{\varepsilon}(\hat{l}))\} \\ &\geq \bigwedge\{\bigvee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})), \bigvee(\iota_{\Gamma}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{l}))\} \\ &\geq \bigwedge\{\sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}, \sup\{\iota_{\Gamma}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}\} \\ &\geq \sup\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{m}), \iota_{\Gamma}^{\varepsilon}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\} \\ &= \sup\{(\iota_{\Upsilon}^{\varepsilon}(\hat{m}) \wedge \iota_{\Gamma}^{\varepsilon}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\}. \\ \Rightarrow \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{l})\} &\geq \sup\{(\iota_{\Upsilon}^{\varepsilon}(\hat{m}) \wedge \iota_{\Gamma}^{\varepsilon}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\}. \end{aligned}$$

Similarly,

$$\bigvee\{(F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k}), (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{l})\} \geq \sup\{(F_{\Upsilon}^{\varsigma}(\hat{m}) \vee F_{\Gamma}^{\varsigma}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\}.$$

(2) $\exists \forall \hat{n}, \hat{k} \in R$ such that $\hat{k} \in \hat{n} + \hat{l}$ then it argues that:

$$\begin{aligned} \bigwedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &= \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{n})), \bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})), \bigwedge(\tau_{\Gamma}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{l}), \tau_{\Gamma}^{\varpi}(\hat{l}))\} \\ &= (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l}). \\ \Rightarrow \bigwedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &\leq \{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l}) : \hat{k} \in \hat{n} + \hat{l}\}. \end{aligned}$$

Also, $\exists \forall \hat{n}, \hat{k} \in R$ such that $\hat{k} \in \hat{n} + \hat{l}$ then it argues that:

$$\begin{aligned} \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &= \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{n})), \bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})), \bigwedge(\iota_{\Gamma}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{l}), \iota_{\Gamma}^{\varepsilon}(\hat{l}))\} \\ &= \{(\iota_{\Upsilon}^{\varepsilon}(\hat{l}) \wedge \iota_{\Gamma}^{\varepsilon}(\hat{l}))\}. \\ \Rightarrow \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &\geq \{(\iota_{\Upsilon}^{\varepsilon}(\hat{l}) \wedge \iota_{\Gamma}^{\varepsilon}(\hat{l})) : \hat{k} \in \hat{n} + \hat{l}\} \end{aligned}$$

Similarly,

$$\bigvee\{(F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{n}), (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k})\} \geq \{(F_{\Upsilon}^{\varsigma}(\hat{l}) \vee F_{\Gamma}^{\varsigma}(\hat{l})) : \hat{k} \in \hat{n} + \hat{l}\}$$

(3) $\forall \hat{n}, \hat{k} \in R \exists \hat{m} \in R$ where $\hat{k} \in \hat{m} + \hat{n}$ can be readily proved that

$$\begin{aligned} \bigwedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &= \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{n})), \bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})), \bigwedge(\tau_{\Gamma}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{m}), \tau_{\Gamma}^{\varpi}(\hat{m})\} \\ &= (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}). \end{aligned}$$

$$\Rightarrow \bigwedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k})\} \leq \{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{k} \in \hat{m} + \hat{n}\}.$$

Also, $\forall \hat{n}, \hat{k} \in R \exists \hat{m} \in R$ where $\hat{k} \in \hat{m} + \hat{n}$ then it argues that:

$$\begin{aligned} \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &= \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{n})), \bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})), \bigwedge(\iota_{\Gamma}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \bigwedge\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}), \iota_{\Gamma}^{\varepsilon}(\hat{m})\} \\ &= \{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{m})\}. \end{aligned}$$

$$\Rightarrow \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} \geq \{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{m}) : \hat{k} \in \hat{m} + \hat{n}\}$$

Similarly,

$$\bigvee\{(F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{n}), (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k})\} \geq \{(F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{m}) : \hat{k} \in \hat{m} + \hat{n}\}$$

(4) $\forall \hat{k}, \hat{l} \in R$,

$$\begin{aligned} \bigwedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l})\} &\leq \inf\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}, \\ \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{l})\} &\geq \sup\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}, \\ \bigvee\{(F_{\Upsilon}^{\varsigma} \wedge F_{\Gamma}^{\varsigma})(\hat{k}), (F_{\Upsilon}^{\varsigma} \wedge F_{\Gamma}^{\varsigma})(\hat{l})\} &\geq \sup\{(F_{\Upsilon}^{\varsigma} \wedge F_{\Gamma}^{\varsigma})(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}. \end{aligned}$$

Hence, $\Upsilon \cap \Gamma$ is $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R . \square

Theorem 3.7. Let Υ be a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R . Then for every $t \in [0, 1]$, $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)} \neq \phi$ is a subhyperring over R .

Proof. Let Υ be a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R . $\forall t \in [0, 1]$, let $\hat{k}, \hat{l} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$.

Then $\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l}) \geq t$, $\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \leq t$ and $F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l}) \leq t$.

Since Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic subhypergroup of $(R, +)$, we have the following

$$\inf\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\} \geq \bigwedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l})\} \geq \bigwedge\{t, t\} = t,$$

$$\sup\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\} \leq \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{l})\} \leq \bigvee\{t, t\} = t,$$

and

$$\sup\{(F_{\Upsilon}^{\varsigma} \wedge F_{\Gamma}^{\varsigma})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\} \leq \bigvee\{(F_{\Upsilon}^{\varsigma} \wedge F_{\Gamma}^{\varsigma})(\hat{k}), (F_{\Upsilon}^{\varsigma} \wedge F_{\Gamma}^{\varsigma})(\hat{l})\} \leq \bigvee\{t, t\} = t.$$

This implies that $\hat{m} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ and then for every $\hat{m} \in \hat{k} + \hat{l}$, we obtain $\hat{k} + \hat{l} \subseteq \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$.

As such, for every $\hat{m} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$, we obtain $\hat{m} + \Upsilon_t^{(\varpi, \varepsilon, \varsigma)} \subseteq \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$.

Now let $\hat{k}, \hat{m} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$. Then $\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{m}) \geq t, \iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{m}) \leq t$ and $F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{m}) \leq t$. Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic subhypergroup of $(R, +)$, $\exists \hat{l} \in R$ such that $\hat{k} \in \hat{m} + \hat{l}$ and $\tau_{\Upsilon}^{\varpi}(\hat{l}) \geq \wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{m})) \geq t, \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \leq \vee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{m})) \leq t, F_{\Upsilon}^{\varsigma}(\hat{l}) \leq \vee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{m})) \leq t$, and this implies that $\hat{l} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$. Therefore, we obtain $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)} \subseteq \hat{m} + \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$.

As such, we obtain $\hat{m} + \Upsilon_t^{(\varpi, \varepsilon, \varsigma)} = \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$. As a result, $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ is a subhypergroup of $(R, +)$. Let $\hat{k}, \hat{l} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$, then $\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l}) \geq t, \iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \leq t$ and $F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l}) \leq t$. Since Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic sub-semihypergroup of (R, \circ) , then $\forall \hat{k}, \hat{l} \in R$, we have the following:

$$\inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\} \geq \wedge\{\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})\} = t,$$

$$\sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\} \leq \vee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})) = t,$$

and

$$\sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\} \leq \vee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l})) = t.$$

This implies that $\hat{m} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ and consequently $\hat{k} \circ \hat{l} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$.

Therefore, for every $\hat{k}, \hat{l} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ we obtain $\hat{k} \circ \hat{l} \in P^*(R)$. Hence $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ is a subhyperring over R . \square

Theorem 3.8. *Let Υ be a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic set over R . Then the following statements are equivalent:*

- (1) Υ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R .
- (2) $\forall t \in [0, 1]$, a non-empty $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ is a subhyperring over R .

Proof. (1) \Rightarrow (2) $\forall t \in [0, 1]$, by Theorem 3.7, $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ is subhyperring over R .

(2) \Rightarrow (1) Assume that $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ is a subhyperring over R . Let $\hat{k}, \hat{l} \in \Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ and therefore $\hat{k} + \hat{l} \subseteq \Upsilon_{t_0}^{(\varpi, \varepsilon, \varsigma)}$. Then for every $\hat{m} \in \hat{k} + \hat{l}$ we have $\tau_{\Upsilon}^{\varpi}(\hat{m}) \geq t_0, \iota_{\Upsilon}^{\varepsilon}(\hat{m}) \leq t_0$ and $F_{\Upsilon}^{\varsigma}(\hat{m}) \leq t_0$, which implies that:

$$\wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})) \leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\},$$

$$\vee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})) \geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\},$$

and

$$\vee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l})) \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}.$$

Thus, clause (1) of Definition 3.2 has been fulfilled.

Next, let $\hat{n}, \hat{k} \in \Upsilon_{t_1}^{(\varpi, \varepsilon, \varsigma)}$ for every $t_1 \in [0, 1]$ which means that $\exists \hat{l} \in \Upsilon_{t_1}^{(\varpi, \varepsilon, \varsigma)}$ such that $\hat{k} \in \hat{n} \circ \hat{l}$.

Since $\hat{l} \in \Upsilon_{t_1}^{(\varpi, \varepsilon, \varsigma)}$, we have $\tau_{\Upsilon}^{\varpi}(\hat{l}) \geq t_1$, $\iota_{\Upsilon}^{\varepsilon}(\hat{l}) \leq t_1$ and $F_{\Upsilon}^{\varsigma}(\hat{l}) \leq t_1$, and thus we have

$$\begin{aligned} \tau_{\Upsilon}^{\varpi}(\hat{l}) &\geq t_1 = \bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{m})), \\ \iota_{\Upsilon}^{\varepsilon}(\hat{l}) &\leq t_1 = \bigvee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{m})), \end{aligned}$$

and

$$F_{\Upsilon}^{\varsigma}(\hat{l}) \leq t_1 = \bigvee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{m})).$$

Thus, clause (2) of Definition 3.2 has been fulfilled.

Assurance of (3) of Definition 3.2 can be satisfied in a similar way.

As a result, Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic subhypergroup of $(R, +)$.

Now since $\Upsilon_t^{(\varpi, \varepsilon, \varsigma)}$ is a sub-semihypergroup of the semihypergroup (R, \circ) , we got the following.

Let $\hat{k}, \hat{l} \in \Upsilon_{t_2}^{(\varpi, \varepsilon, \varsigma)}$ and therefore we have $\hat{k} \circ \hat{l} \in \Upsilon_{t_2}^{(\varpi, \varepsilon, \varsigma)}$. Thus for every $\hat{m} \in \hat{k} \circ \hat{l}$, we obtain $\tau_{\Upsilon}^{\varpi}(\hat{m}) \geq t_2$, $\iota_{\Upsilon}^{\varepsilon}(\hat{m}) \leq t_2$ and $F_{\Upsilon}^{\varsigma}(\hat{m}) \leq t_2$, and therefore it follows that:

$$\begin{aligned} \bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})) &\leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}, \\ \bigvee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})) &\geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}, \end{aligned}$$

and

$$\bigvee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l})) \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}.$$

which reveals that clause (4) of Definition 3.2 is verified.

Hence Υ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHR over R . \square

4. $(\varpi, \varepsilon, \varsigma)$ -Single Valued Neutrosophic Hyperideals

Definition 4.1. Let Υ be a $(\varpi, \varepsilon, \varsigma)$ -SVNS over R . Then Υ is $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic left (resp. right) hyperideal over R , if,

- (1) $\forall \hat{k}, \hat{l} \in R$,
 - $\bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})\} \leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}$,
 - $\bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})\} \geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}$, and
 - $\bigvee\{F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l})\} \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}$.
- (2) $\forall \hat{n}, \hat{k} \in R, \exists \hat{l} \in R$ such that $\hat{k} \in \hat{n} + \hat{l}$, and
 - $\bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})\} \leq \tau_{\Upsilon}^{\varpi}(\hat{l})$,
 - $\bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})\} \geq \iota_{\Upsilon}^{\varepsilon}(\hat{l})$, and
 - $\bigvee\{F_{\Upsilon}^{\varsigma}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{k})\} \geq F_{\Upsilon}^{\varsigma}(\hat{l})$.
- (3) $\forall \hat{n}, \hat{k} \in R, \exists \hat{m} \in R$ such that $\hat{k} \in \hat{m} + \hat{n}$, and
 - $\bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})\} \leq \tau_{\Upsilon}^{\varpi}(\hat{m})$,
 - $\bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})\} \geq \iota_{\Upsilon}^{\varepsilon}(\hat{m})$, and
 - $\bigvee\{F_{\Upsilon}^{\varsigma}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{k})\} \geq F_{\Upsilon}^{\varsigma}(\hat{m})$.

$$(4) \forall \hat{k}, \hat{l} \in R, \\ \tau_{\Upsilon}^{\varpi}(\hat{l}) \leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\} \text{ (resp. } \tau_{\Upsilon}^{\varpi}(\hat{k}) \leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}), \\ \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\} \text{ (resp. } \iota_{\Upsilon}^{\varepsilon}(\hat{k}) \geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}), \text{ and} \\ F_{\Upsilon}^{\varsigma}(\hat{l}) \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\} \text{ (resp. } F_{\Upsilon}^{\varsigma}(\hat{k}) \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}).$$

If Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic left (resp. right) hyperideal of R then Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic subhypergroup of $(R, +)$ by clauses (1), (2) and (3).

Definition 4.2. Let Υ be a $(\varpi, \varepsilon, \varsigma)$ -SVNS over R . Then Υ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHI over R , if aforementioned clauses are met:

$$(1) \forall \hat{k}, \hat{l} \in R, \\ \bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})\} \leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}, \\ \bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})\} \geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}, \text{ and} \\ \bigvee\{F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l})\} \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \\ (2) \forall \hat{n}, \hat{k} \in R, \exists \hat{l} \in R \text{ such that } \hat{k} \in \hat{n} + \hat{l}, \text{ and} \\ \bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})\} \leq \tau_{\Upsilon}^{\varpi}(\hat{l}), \\ \bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})\} \geq \iota_{\Upsilon}^{\varepsilon}(\hat{l}), \text{ and} \\ \bigvee\{F_{\Upsilon}^{\varsigma}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{k})\} \geq F_{\Upsilon}^{\varsigma}(\hat{l}). \\ (3) \forall \hat{n}, \hat{k} \in R, \exists \hat{m} \in R \text{ such that } \hat{k} \in \hat{m} + \hat{n}, \text{ and} \\ \bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})\} \leq \tau_{\Upsilon}^{\varpi}(\hat{m}), \\ \bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})\} \geq \iota_{\Upsilon}^{\varepsilon}(\hat{m}), \text{ and} \\ \bigvee\{F_{\Upsilon}^{\varsigma}(\hat{n}), F_{\Upsilon}^{\varsigma}(\hat{k})\} \geq F_{\Upsilon}^{\varsigma}(\hat{m}). \\ (4) \forall \hat{k}, \hat{l} \in R, \\ \bigwedge\{\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})\} \leq \inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}, \\ \bigvee\{\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})\} \geq \sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}, \text{ and} \\ \bigvee\{F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l})\} \geq \sup\{F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l}\}.$$

Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic subhypergroup of $(R, +)$ by clauses (1), (2) and (3). Clause (4) indicate that Υ is both $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic left hyperideal and $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic right hyperideal.

$\Rightarrow \Upsilon$ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHI of R .

Theorem 4.3. Let Υ be a non-null $(\varpi, \varepsilon, \varsigma)$ -SVNS over R . Υ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHI over R if and only if Υ is a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic hypergroup over $(R, +)$ and also Υ is both a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic left hyperideal and a $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic right hyperideal of R .

Proof. With the help of Definitions 4.1 and 4.2, we get the required proof. \square

Theorem 4.4. *Let Υ and Γ be two $(\varpi, \varepsilon, \varsigma)$ -SVNHIs over R . Then $\Upsilon \cap \Gamma$ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHI over R if it is non-null.*

Proof. Let Υ and Γ are $(\varpi, \varepsilon, \varsigma)$ -SVNHIs over R . By using Definition 3.2, and Proposition 3.5

$$(\Upsilon \cap \Gamma)^{(\varpi, \varepsilon, \varsigma)} = \Upsilon^{(\varpi, \varepsilon, \varsigma)} \cap \Gamma^{(\varpi, \varepsilon, \varsigma)} = \{ \langle \hat{k}, (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k}) \rangle : \hat{k} \in R \},$$

where

$$\begin{aligned} (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}) &= \wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k})), \\ (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}) &= \wedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k})), \\ (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k}) &= \vee(F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Gamma}^{\varsigma}(\hat{k})). \end{aligned}$$

Assuming $\forall \hat{k}, \hat{l} \in R$, we are only proven to include all four clauses for membership terms τ_{Υ}^{ϖ} , τ_{Γ}^{ϖ} and indeterminacy terms $\iota_{\Upsilon}^{\varepsilon}$, $\iota_{\Gamma}^{\varepsilon}$.

$$\begin{aligned} (1) \quad \wedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l})\} &= \wedge\{\wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k})), \wedge(\tau_{\Upsilon}^{\varpi}(\hat{l}), \tau_{\Gamma}^{\varpi}(\hat{l}))\} \\ &\leq \wedge\{\wedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l})), \wedge(\tau_{\Gamma}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{l}))\} \\ &\leq \wedge\{\inf\{\tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}, \inf\{\tau_{\Gamma}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}\} \\ &\leq \inf\{\wedge(\tau_{\Upsilon}^{\varpi}(\hat{m}), \tau_{\Gamma}^{\varpi}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\} \\ &= \inf\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \\ \Rightarrow \wedge\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{k}), (\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{l})\} &\leq \inf\{(\tau_{\Upsilon}^{\varpi} \wedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \end{aligned}$$

Also

$$\begin{aligned} \vee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{l})\} &= \vee\{\wedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k})), \wedge(\iota_{\Upsilon}^{\varepsilon}(\hat{l}), \iota_{\Gamma}^{\varepsilon}(\hat{l}))\} \\ &\geq \wedge\{\vee(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l})), \vee(\iota_{\Gamma}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{l}))\} \\ &\geq \wedge\{\sup\{\iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}, \sup\{\iota_{\Gamma}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}\} \\ &\geq \sup\{\wedge(\iota_{\Upsilon}^{\varepsilon}(\hat{m}), \iota_{\Gamma}^{\varepsilon}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\} \\ &= \sup\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \\ \Rightarrow \vee\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{k}), (\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{l})\} &\geq \sup\{(\iota_{\Upsilon}^{\varepsilon} \wedge \iota_{\Gamma}^{\varepsilon})(\hat{m}) : \hat{m} \in \hat{k} + \hat{l}\}. \end{aligned}$$

Similarly,

$$\vee\{(F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{k}), (F_{\Upsilon}^{\varsigma} \vee F_{\Gamma}^{\varsigma})(\hat{l})\} \geq \sup\{(F_{\Upsilon}^{\varsigma}(\hat{m}) \vee F_{\Gamma}^{\varsigma}(\hat{m})) : \hat{m} \in \hat{k} + \hat{l}\}.$$

(2) $\exists \forall \hat{n}, \hat{k} \in R$ such that $\hat{k} \in \hat{n} + \hat{l}$ then it argues that:

$$\begin{aligned} \bigwedge\{(\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &= \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{n})), \bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})), \bigwedge(\tau_{\Gamma}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{l}), \tau_{\Gamma}^{\varpi}(\hat{l}))\} \\ &= (\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{l}). \\ \Rightarrow \bigwedge\{(\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &\leq \{(\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{l}) : \hat{k} \in \hat{n} + \hat{l}\}. \end{aligned}$$

Also, $\exists \forall \hat{n}, \hat{k} \in R$ such that $\hat{k} \in \hat{n} + \hat{l}$ then it argues that:

$$\begin{aligned} \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &= \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{n})), \bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})), \bigwedge(\iota_{\Gamma}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{l}), \iota_{\Gamma}^{\varepsilon}(\hat{l}))\} \\ &= \{(\iota_{\Upsilon}^{\varepsilon}(\hat{l}) \bigwedge \iota_{\Gamma}^{\varepsilon}(\hat{l}))\}. \\ \Rightarrow \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &\geq \{(\iota_{\Upsilon}^{\varepsilon}(\hat{l}) \bigwedge \iota_{\Gamma}^{\varepsilon}(\hat{l})) : \hat{k} \in \hat{n} + \hat{l}\} \end{aligned}$$

Similarly,

$$\bigvee\{(F_{\Upsilon}^{\varsigma} \bigvee F_{\Gamma}^{\varsigma})(\hat{n}), (F_{\Upsilon}^{\varsigma} \bigvee F_{\Gamma}^{\varsigma})(\hat{k})\} \geq \{(F_{\Upsilon}^{\varsigma}(\hat{l}) \bigvee F_{\Gamma}^{\varsigma}(\hat{l})) : \hat{k} \in \hat{n} + \hat{l}\}$$

(3) $\forall \hat{n}, \hat{k} \in R \exists \hat{m} \in R$ where $\hat{k} \in \hat{m} + \hat{n}$ can be readily proved that

$$\begin{aligned} \bigwedge\{(\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &= \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{n})), \bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \bigwedge\{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{n}), \tau_{\Upsilon}^{\varpi}(\hat{k})), \bigwedge(\tau_{\Gamma}^{\varpi}(\hat{n}), \tau_{\Gamma}^{\varpi}(\hat{k}))\} \\ &\leq \{\bigwedge(\tau_{\Upsilon}^{\varpi}(\hat{m}), \tau_{\Gamma}^{\varpi}(\hat{m}))\} \\ &= (\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{m}). \\ \Rightarrow \bigwedge\{(\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{n}), (\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{k})\} &\leq \{(\tau_{\Upsilon}^{\varpi} \bigwedge \tau_{\Gamma}^{\varpi})(\hat{m}) : \hat{k} \in \hat{m} + \hat{n}\}. \end{aligned}$$

Also, $\forall \hat{n}, \hat{k} \in R \exists \hat{m} \in R$ where $\hat{k} \in \hat{m} + \hat{n}$ then it argues that:

$$\begin{aligned} \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &= \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{n})), \bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \bigvee\{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{n}), \iota_{\Upsilon}^{\varepsilon}(\hat{k})), \bigwedge(\iota_{\Gamma}^{\varepsilon}(\hat{n}), \iota_{\Gamma}^{\varepsilon}(\hat{k}))\} \\ &\geq \{\bigwedge(\iota_{\Upsilon}^{\varepsilon}(\hat{m}), \iota_{\Gamma}^{\varepsilon}(\hat{m}))\} \\ &= \{(\iota_{\Upsilon}^{\varepsilon}(\hat{m}) \bigwedge \iota_{\Gamma}^{\varepsilon}(\hat{m}))\}. \\ \Rightarrow \bigvee\{(\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{n}), (\iota_{\Upsilon}^{\varepsilon} \bigwedge \iota_{\Gamma}^{\varepsilon})(\hat{k})\} &\geq \{(\iota_{\Upsilon}^{\varepsilon}(\hat{m}) \bigwedge \iota_{\Gamma}^{\varepsilon}(\hat{m})) : \hat{k} \in \hat{m} + \hat{n}\} \end{aligned}$$

Similarly,

$$\bigvee\{(F_{\Upsilon}^{\varsigma} \bigvee F_{\Gamma}^{\varsigma})(\hat{n}), (F_{\Upsilon}^{\varsigma} \bigvee F_{\Gamma}^{\varsigma})(\hat{k})\} \geq \{(F_{\Upsilon}^{\varsigma}(\hat{m}) \bigvee F_{\Gamma}^{\varsigma}(\hat{m})) : \hat{k} \in \hat{m} + \hat{n}\}$$

$$(4) \forall \hat{k}, \hat{l} \in R,$$

$$\begin{aligned} \bigvee \{ \tau_{\Upsilon}^{\varpi}(\hat{k}), \tau_{\Upsilon}^{\varpi}(\hat{l}) \} &\leq \inf \{ \tau_{\Upsilon}^{\varpi}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l} \}, \\ \bigvee \{ \iota_{\Upsilon}^{\varepsilon}(\hat{k}), \iota_{\Upsilon}^{\varepsilon}(\hat{l}) \} &\geq \sup \{ \iota_{\Upsilon}^{\varepsilon}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l} \}, \text{ and} \\ \bigvee \{ F_{\Upsilon}^{\varsigma}(\hat{k}), F_{\Upsilon}^{\varsigma}(\hat{l}) \} &\geq \sup \{ F_{\Upsilon}^{\varsigma}(\hat{m}) : \hat{m} \in \hat{k} \circ \hat{l} \}. \end{aligned}$$

Hence, it is verified that $\Upsilon \cap \Gamma$ is a $(\varpi, \varepsilon, \varsigma)$ -SVNHI over R . \square

5. Conclusions

This research has introduced the novel concepts of the $(\varpi, \varepsilon, \varsigma)$ -single valued neutrosophic theory of hyperrings and hyperideals through the introduction of a few hyperalgebraic structures and the analysis of some basic properties, outcomes, and structural characteristics of these concepts. We plan to meld more hyperalgebraic theory with real-world applications in the future for plithogenic sets for $(\varpi, \varepsilon, \varsigma)$ -single-valued neutrosophic sets and $(\varpi, \varepsilon, \varsigma)$ -interval-valued neutrosophic sets.

References

1. F. Marty. Sur unegeneralisation de la notion de groupe. Proceedings of the 8th Congress Mathematiciens Scandinaves, Stockholm, Sweden (1934) 45-49.
2. Corsini, P. (1993). Prolegomena of hypergroup theory. Aviani editore.
3. Vougiouklis, T. (1994). Hyperstructures and their representations. Hadronic Press.
4. Vougiouklis, T. (1995). A new class of hyperstructures. J. Combin. Inform. System Sci., 20, 229-235.
5. Zadeh, L. A. (1996). Fuzzy sets. Information and Control 8, 338-353.
6. Turksen, I. B. (1986). Interval valued fuzzy sets based on normal forms. Fuzzy sets and systems, 20(2), 191-210.
7. Atanassov, K. (1986). Intuitionistic fuzzy sets. fuzzy sets and systems 20 (1), 87-96.
8. Wang, H.; Smarandache, F.; Zhang, Y.; & Sunderraman, R. (2010). Single valued neutrosophic sets. Infinite study.
9. Rosenfeld, A. (1971). Fuzzy groups. Journal of mathematical analysis and applications, 35(3), 512-517.
10. Selvachandran, G.; & Salleh, A. R. (2016). Fuzzy soft ideals based on fuzzy soft spaces. Far East Journal of Mathematical Sciences, 99(3), 429.
11. Selvachandran, G.; & Salleh, A. R. (2016). Rings and ideals in a vague soft set setting. Far East Journal of Mathematical Sciences, 99(2), 279.
12. Selvachandran, G.; & Salleh, A. R. (2015). On normalistic vague soft groups and normalistic vague soft group homomorphism. Advances in Fuzzy Systems, 2015.
13. Khademan, S.; Zahedi, M. M.; Borzooei, R. A.; & Jun, Y. B. (2019). Neutrosophic hyper BCK-ideals. Infinite Study.
14. Smarandache, F. (1998). Neutrosophy: neutrosophic probability, set, and logic: analytic synthesis & synthetic analysis.
15. Abdel-Basset, M.; Mohamed, R.; Zaied, A. E. N. H.; & Smarandache, F. (2019). A hybrid plithogenic decision-making approach with quality function deployment for selecting supply chain sustainability metrics. Symmetry, 11(7), 903.

16. Abdel-Basset, M.; Saleh, M.; Gamal, A.; & Smarandache, F. (2019). An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number. *Applied Soft Computing*, 77, 438-452.
17. Abdel-Basset, M.; Chang, V.; Gamal, A.; & Smarandache, F. (2019). An integrated neutrosophic ANP and VIKOR method for achieving sustainable supplier selection: A case study in importing field. *Computers in Industry*, 106, 94-110.
18. Abdel-Basset, M.; Manogaran, G.; Gamal, A.; & Smarandache, F. (2019). A group decision making framework based on neutrosophic TOPSIS approach for smart medical device selection. *Journal of medical systems*, 43(2), 38.
19. Elavarasan, B.; Smarandache, F.; & Jun, Y. B. (2019). Neutrosophic ideals in semigroups. *Neutrosophic Sets and Systems*, 28(1), 21.
20. Smarandache, F.; & Neutrosophy, N. P. (1998). *Set, and Logic*, ProQuest Information & Learning. Ann Arbor, Michigan, USA, 105.
21. Smarandache, F. (2017). *Plithogeny, plithogenic set, logic, probability, and statistics*. Pons Publishing House, Brussels, Belgium, 141 p., arXiv.org (Cornell University).
22. Liu, W. J. (1982). Fuzzy invariant subgroups and fuzzy ideals. *Fuzzy sets and Systems*, 8(2), 133-139.
23. Ameri, R.; & Nozari, T. (2011). Fuzzy hyperalgebras. *Computers & Mathematics with Applications*, 61(2), 149-154.
24. Leoreanu-Fotea, V.; & Davvaz, B. (2009). Fuzzy hyperrings. *Fuzzy sets and systems*, 160(16), 2366-2378.
25. Davvaz, B.; & Abbasizadeh, N. (2017). Fuzzy topological F-polygroups. *Journal of Intelligent & Fuzzy Systems*, 33(6), 3433-3440.
26. Muhiuddin, G., Harizavi, H., & Jun, Y. B. (2020). Bipolar-valued fuzzy soft hyper BCK ideals in hyper BCK algebras. *Discrete Mathematics, Algorithms and Applications*, 12(02), 2050018.
27. Cristea, I. (2016). Fuzzy Subhypergroups Degree. *Journal of Multiple-Valued Logic & Soft Computing*, 27(1).
28. Akram, M.; & Luqman, A. (2020). *Fuzzy hypergraphs and related extensions*. Springer Singapore.
29. Van Pham, N.; Pham, L. T.; Pedrycz, W.; & Ngo, L. T. (2021). Feature-reduction fuzzy co-clustering approach for hyper-spectral image analysis. *Knowledge-Based Systems*, 216, 106549.
30. Davvaz, B.; & Firoozkouhi, N. (2019). Fundamental relation on fuzzy hypermodules. *Soft Computing*, 23(24), 13025-13033.
31. Leoreanu-Fotea, V.; & Davvaz, B. (2009). Fuzzy hyperrings. *Fuzzy sets and systems*, 160(16), 2366-2378.
32. Ersoy, B. A. (2020). Study of Γ -hyperrings by fuzzy hyperideals with respect to a t-norm. *Acta Universitatis Sapientiae, Mathematica*, 11(2), 306-331.
33. Ostadhadi-Dehkordi, S.; & Shum, K. P. (2019). Regular and strongly regular relations on soft hyperrings. *Soft Computing*, 23(10), 3253-3260.
34. Nodehi, M.; Norouzi, M.; & Dehghan, O. R. (2020). An introduction to topological hyperrings. *Caspian Journal of Mathematical Sciences (CJMS)*, 9(2), 210-223.
35. Abughazalah, N.; Yaqoob, N.; & Shahzadi, K. (2021). Topological Structures of Lower and Upper Rough Subsets in a Hyperring. *Journal of Mathematics*, 2021.
36. Davvaz, B. (1998). On H_v -rings and fuzzy H_v -ideal. *Journal of Fuzzy Mathematics*, 6(1), 33-42.
37. Bharathi, P.; & Vimala, J. (2016). The Role of Fuzzy l -ideals in a Commutative l -group. *Global Journal of Pure and Applied Mathematics*, 12(3), 2067-2074.
38. Vimala, J.; & ArockiaReeta, J. (2018). Ideal approach on lattice ordered fuzzy soft group and its application in selecting best mobile network coverage among travelling paths. *ARNP Journal of Engineering and Applied Sciences*, 13(7), 2414-2421.

39. Selvachandran, G.; & Salleh, A. R. (2014). Vague soft hypergroups and vague soft hypergroup homomorphism. *Advances in Fuzzy Systems*, 2014.
40. Selvachandran, G.; & Salleh, A. R. (2015). Algebraic hyperstructures of vague soft sets associated with hyperrings and hyperideals. *The Scientific World Journal*, 2015.
41. Selvachandran, G. (2015). Introduction to the theory of soft hyperrings and soft hyperring homomorphism. *JP Journal of Algebra, Number Theory and Applications*, 36(3), 279-294.
42. Hameed, M. S.; Ahmad, Z.; Mukhtar, S.; & Ullah, A. (2021). Some results on χ -single valued neutrosophic subgroups. *Indonesian Journal of Electrical Engineering and Computer Science*, 23(3), 1583-1589.
43. Ali, S.; Kousar, M.; Xin, Q.; Pamucar, D.; Hameed, M. S.; & Fayyaz, R. (2021). Belief and Possibility Belief Interval-Valued N-Soft Set and Their Applications in Multi-Attribute Decision-Making Problems. *Entropy*, 23(11), 1498.
44. Hameeda, M. S.; Alia, S.; Mukhtar, S.; Shoaib, M.; Ishaq, M. K.; & Mukhtiar, U. On characterization of χ -single valued neutrosophic sub-groups.
45. Hameed, M. S.; Mukhtar, S.; Khan, H. N.; Ali, S.; Mateen, M. H.; & Gulzar, M. (2021). Pythagorean fuzzy N-Soft groups. *Int. J. Electr. Comput. Eng*, 21, 1030-1038.

Received: Feb 8, 2022. Accepted: Jun 14, 2022