

10-5-2022

On Some New Concepts of Weakly Neutrosophic Crisp Separation Axioms

Qays Hatem Imran

Ali H. M. Al-Obaidi

Florentin Smarandache

Md. Hanif PAGE

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Imran, Qays Hatem; Ali H. M. Al-Obaidi; Florentin Smarandache; and Md. Hanif PAGE. "On Some New Concepts of Weakly Neutrosophic Crisp Separation Axioms." *Neutrosophic Sets and Systems* 51, 1 (2022). https://digitalrepository.unm.edu/nss_journal/vol51/iss1/21

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



On Some New Concepts of Weakly Neutrosophic Crisp Separation Axioms

Qays Hatem Imran^{1*}, Ali H. M. Al-Obaidi², Florentin Smarandache³ and Md. Hanif PAGE⁴

¹Department of Mathematics, College of Education for Pure Science, Al-Muthanna University, Samawah, Iraq.
E-mail: qays.imran@mu.edu.iq

²Department of Mathematics, College of Education for Pure Science, University of Babylon, Hillah, Iraq.
E-mail: aalobaidi@uobabylon.edu.iq

³Department of Mathematics, University of New Mexico 705 Gurley Ave. Gallup, NM 87301, USA.
E-mail: smarand@unm.edu

⁴Department of Mathematics, KLE Technological University, Hubballi-580031, Karnataka, India.
E-mail: mb_page@kletech.ac.in

*Correspondence: qays.imran@mu.edu.iq

Abstract: In this paper, we shall study some new concepts of weakly neutrosophic crisp separation axioms, which are called “neutrosophic crisp α -separation and neutrosophic crisp semi- α -separation axioms” such as neutrosophic crisp α - T_i and neutrosophic crisp semi- α - $T_i, \forall i = 0, 1, \dots, 4$. Moreover, we shall study the relationship between usual neutrosophic crisp separation axioms and these kinds of weakly neutrosophic crisp separation axioms.

Mathematics Subject Classification (2010): 03E72, 03F55, 54A40, 62C86.

Keywords: NC^α - T_i , $NC^{S\alpha}$ - $T_i, \forall i = 0, 1, \dots, 4$, NC^α -regular, $NC^{S\alpha}$ -regular, NC^α -normal and $NC^{S\alpha}$ -normal spaces.

1. Introduction

A. A. Salama et al. [1] give a concept of neutrosophic crisp topological space (briefly NCTS). A. A. Salama [2] provided some classes of neutrosophic crisp nearly open sets. A. H. M. Al-Obaidi et al. [3,4] give concepts of weakly neutrosophic crisp functions. Md. Hanif PAGE et al. [5] examined the view of neutrosophic generalized homeomorphism. Q. H. Imran et al. [6-8] established neutrosophic semi- α -open sets, new types of weakly neutrosophic crisp continuity and new concepts of neutrosophic crisp open sets. R. Dhavaseelan et al. [9] examined the view of neutrosophic α^m -continuity. R. K. Al-Hamido et al. [10] tendered the interpretation of neutrosophic crisp semi- α -closed sets. A. B. Al-Nafee et al. [11] demonstrated the principle of separation axioms in neutrosophic crisp topological spaces. R. K. Al-Hamido et al. [12] provided neutrosophic crisp semi separation axioms. The objective of this paper is to study some new concepts of weakly neutrosophic crisp separation axioms, which are called “neutrosophic crisp α -separation and neutrosophic crisp semi- α -separation axioms” such as neutrosophic crisp α - T_i and neutrosophic crisp semi- α - $T_i, \forall i = 0, 1, \dots, 4$. Moreover, we shall study the relationship between usual neutrosophic crisp separation axioms and these kinds of weakly neutrosophic crisp separation axioms.

2. Preliminaries

Throughout this paper, (\mathcal{S}, ζ) and (\mathcal{I}, η) (or simply \mathcal{S} and \mathcal{I}) always mean NCTSs. The complement of a neutrosophic crisp open set (briefly NC-OS) is called a neutrosophic crisp closed

set (briefly NC-CS) in (\mathcal{S}, ζ) . For a NCS \mathfrak{B} in a NCTS (\mathcal{S}, ζ) , $NCcl(\mathfrak{B})$, $NCint(\mathfrak{B})$ and \mathfrak{B}^c denote the NC-closure of \mathfrak{B} , the NC-interior of \mathfrak{B} and the NC-complement of \mathfrak{B} respectively.

Definition 2.1 [1]:

For any nonempty under-consideration set \mathcal{S} , a neutrosophic crisp set (in short NCS) \mathfrak{B} is an object holding the establish $\mathfrak{B} = \langle \mathfrak{B}_1, \mathfrak{B}_2, \mathfrak{B}_3 \rangle$ where $\mathfrak{B}_1, \mathfrak{B}_2$ and \mathfrak{B}_3 are mutually disjoint sets included in \mathcal{S} .

Definition 2.2:

A NC-subset \mathfrak{B} of a NCTS (\mathcal{S}, ζ) is said to be:

- (i) neutrosophic crisp α -open set (in short NC^α -OS) [2] if $\mathfrak{B} \subseteq NCint(NCcl(NCint(\mathfrak{B})))$. The family of all NC^α -OSs of \mathcal{S} is denoted by $NC^\alpha O(\mathcal{S})$. The complement of NC^α -OS is called a neutrosophic crisp α -closed set (in short NC^α -CS). The family of all NC^α -CSs of \mathcal{S} is denoted by $NC^\alpha C(\mathcal{S})$.
- (ii) neutrosophic crisp semi- α -open set (in short $NC^{S\alpha}$ -OS) [10] if there exists a NC^α -OS \mathfrak{D} in \mathcal{S} such that $\mathfrak{D} \subseteq \mathfrak{B} \subseteq NCcl(\mathfrak{D})$ or equivalently if $\mathfrak{B} \subseteq NCcl(NCint(NCcl(NCint(\mathfrak{B}))))$. The family of all $NC^{S\alpha}$ -OSs of \mathcal{S} is denoted by $NC^{S\alpha} O(\mathcal{S})$. The complement of $NC^{S\alpha}$ -OS is called a neutrosophic crisp semi- α -closed set (in short $NC^{S\alpha}$ -CS). The family of all $NC^{S\alpha}$ -CSs of \mathcal{S} is denoted by $NC^{S\alpha} C(\mathcal{S})$.

Example 2.3:

Let $\mathcal{S} = \{\mathfrak{k}_1, \mathfrak{k}_2, \mathfrak{k}_3, \mathfrak{k}_4\}$. Then $\zeta = \{\emptyset_N, \langle \{\mathfrak{k}_1\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_2\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_1, \mathfrak{k}_2\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_1, \mathfrak{k}_2, \mathfrak{k}_3\}, \emptyset, \emptyset \rangle, \mathcal{S}_N\}$ is a NCTS. The family of all NC^α -OSs of \mathcal{S} is : $NC^\alpha O(\mathcal{S}) = \zeta \sqcup \langle \{\mathfrak{k}_1, \mathfrak{k}_2, \mathfrak{k}_4\}, \emptyset, \emptyset \rangle$.

The family of all $NC^{S\alpha}$ -OSs of \mathcal{S} is : $NC^{S\alpha} O(\mathcal{S}) = NC^\alpha O(\mathcal{S}) \sqcup \langle \{\mathfrak{k}_1, \mathfrak{k}_3\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_1, \mathfrak{k}_4\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_2, \mathfrak{k}_3\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_2, \mathfrak{k}_4\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_1, \mathfrak{k}_3, \mathfrak{k}_4\}, \emptyset, \emptyset \rangle, \langle \{\mathfrak{k}_2, \mathfrak{k}_3, \mathfrak{k}_4\}, \emptyset, \emptyset \rangle$.

Remark 2.4 [10,14]:

In a NCTS (\mathcal{S}, ζ) , then the following statements hold, and the opposite of each statement is not true:

- (i) Every NC-OS (resp. NC-CS) is a NC^α -OS (resp. NC^α -CS) and $NC^{S\alpha}$ -OS (resp. $NC^{S\alpha}$ -CS).
- (ii) Every NC^α -OS (resp. NC^α -CS) is a $NC^{S\alpha}$ -OS (resp. $NC^{S\alpha}$ -CS).

Definition 2.5:

- (i) The NC^α -interior of a NCS \mathfrak{B} of a NCTS (\mathcal{S}, ζ) is the union of all NC^α -OSs contained in \mathfrak{B} and is denoted by $NC^\alpha int(\mathfrak{B})$ [3].
- (ii) The $NC^{S\alpha}$ -interior of a NCS \mathfrak{B} of a NCTS (\mathcal{S}, ζ) is the union of all $NC^{S\alpha}$ -OSs contained in \mathfrak{B} and is denoted by $NC^{S\alpha} int(\mathfrak{B})$ [10].

Definition 2.6:

- (i) The NC^α -closure of a NCS \mathfrak{B} of a NCTS (\mathcal{S}, ζ) is the intersection of all NC^α -CSs containing \mathfrak{B} and is denoted by $NC^\alpha cl(\mathfrak{B})$ [3].
- (ii) The $NC^{S\alpha}$ -closure of a NCS \mathfrak{B} of a NCTS (\mathcal{S}, ζ) is the intersection of all $NC^{S\alpha}$ -CSs containing \mathfrak{B} and is denoted by $NC^{S\alpha} cl(\mathfrak{B})$ [10].

Theorem 2.7:

Let (\mathcal{S}, ζ) and (\mathcal{I}, η) be two NCTSs. If $\mathfrak{B} \in NC^\alpha O(\mathcal{S})$ (resp. $\mathfrak{B} \in NC^{S\alpha} O(\mathcal{S})$), $\mathfrak{D} \in NC^\alpha O(\mathcal{I})$ (resp. $\mathfrak{D} \in NC^{S\alpha} O(\mathcal{I})$), then $\mathfrak{B} \times \mathfrak{D} \in NC^\alpha O(\mathcal{S} \times \mathcal{I})$ (resp. $\mathfrak{B} \times \mathfrak{D} \in NC^{S\alpha} O(\mathcal{S} \times \mathcal{I})$).

Proof:

Since $\mathfrak{B} \subseteq NCint(NCcl(NCint(\mathfrak{B})))$, $\mathfrak{D} \subseteq NCint(NCcl(NCint(\mathfrak{D})))$.

Hence $\mathfrak{B} \times \mathfrak{D} \subseteq NCint(NCcl(NCint(\mathfrak{B}))) \times NCint(NCcl(NCint(\mathfrak{D}))) = NCint(NCcl(NCint(\mathfrak{B} \times \mathfrak{D})))$.

Therefore $\mathfrak{B} \times \mathfrak{D} \subseteq NCint(NCcl(NCint(\mathfrak{B} \times \mathfrak{D}))) \Rightarrow \mathfrak{B} \times \mathfrak{D} \in NC^{\alpha}O(\mathcal{S} \times \mathcal{I})$. The second case is similar. ■

Corollary 2.8:

Let (\mathcal{S}, ζ) and (\mathcal{I}, η) be two NCTSs. If $\mathfrak{B} \in NC^{\alpha}C(\mathcal{S})$ (resp. $\mathfrak{B} \in NC^{S\alpha}C(\mathcal{S})$), $\mathfrak{D} \in NC^{\alpha}C(\mathcal{I})$ (resp. $\mathfrak{D} \in NC^{S\alpha}C(\mathcal{I})$), then $\mathfrak{B} \times \mathfrak{D} \in NC^{\alpha}C(\mathcal{S} \times \mathcal{I})$ (resp. $\mathfrak{B} \times \mathfrak{D} \in NC^{S\alpha}C(\mathcal{S} \times \mathcal{I})$).

Proof:

The proof of this is similar to that of theorem (2.6). ■

Proposition 2.9 [10]:

For any NC-subset \mathfrak{B} of a NCTS (\mathcal{S}, ζ) , then:

- (i) $NCint(\mathfrak{B}) \subseteq NC^{\alpha}int(\mathfrak{B}) \subseteq NC^{S\alpha}int(\mathfrak{B}) \subseteq NC^{S\alpha}cl(\mathfrak{B}) \subseteq NC^{\alpha}cl(\mathfrak{B}) \subseteq NCcl(\mathfrak{B})$.
- (ii) $NCint(NC^{S\alpha}int(\mathfrak{B})) = NC^{S\alpha}int(NCint(\mathfrak{B})) = NCint(\mathfrak{B})$.
- (iii) $NC^{\alpha}int(NC^{S\alpha}int(\mathfrak{B})) = NC^{S\alpha}int(NC^{\alpha}int(\mathfrak{B})) = NC^{\alpha}int(\mathfrak{B})$.
- (iv) $NCcl(NC^{S\alpha}cl(\mathfrak{B})) = NC^{S\alpha}cl(NCcl(\mathfrak{B})) = NCcl(\mathfrak{B})$.
- (v) $NC^{\alpha}cl(NC^{S\alpha}cl(\mathfrak{B})) = NC^{S\alpha}cl(NC^{\alpha}cl(\mathfrak{B})) = NC^{\alpha}cl(\mathfrak{B})$.
- (vi) $NC^{S\alpha}cl(\mathfrak{B}) = \mathfrak{B} \sqcup NCint(NCcl(NCint(NCcl(\mathfrak{B}))))$.
- (vii) $NC^{S\alpha}int(\mathfrak{B}) = \mathfrak{B} \cap NCcl(NCint(NCcl(NCint(\mathfrak{B}))))$.
- (viii) $NCint(NCcl(\mathfrak{B})) \subseteq NC^{S\alpha}int(NC^{S\alpha}cl(\mathfrak{B}))$

Definition 2.10 [1]:

Let $\rho: (\mathcal{S}, \zeta) \rightarrow (\mathcal{I}, \eta)$ be a function, then ρ is said to be NC-continuous (in short NC-CF) iff $\forall \mathfrak{B}$ NC-OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a NC-OS in \mathcal{S} .

Definition 2.11 [13]:

Let $\rho: (\mathcal{S}, \zeta) \rightarrow (\mathcal{I}, \eta)$ be a function, then ρ is said to be NC^{α} -continuous (in short NC^{α} -CF) iff $\forall \mathfrak{B}$ NC-OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a NC^{α} -OS in \mathcal{S} .

Definition 2.12 [10]:

Let $\rho: (\mathcal{S}, \zeta) \rightarrow (\mathcal{I}, \eta)$ be a function, then ρ is said to be:

- (i) $NC^{\alpha*}$ -continuous (in short $NC^{\alpha*}$ -CF) iff $\forall \mathfrak{B}$ NC^{α} -OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a NC^{α} -OS in \mathcal{S} .
- (ii) $NC^{\alpha**}$ -continuous (in short $NC^{\alpha**}$ -CF) iff $\forall \mathfrak{B}$ NC^{α} -OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a NC-OS in \mathcal{S} .

Definition 2.13 [10]:

Let $\rho: (\mathcal{S}, \zeta) \rightarrow (\mathcal{I}, \eta)$ be a function, then ρ is said to be:

- (i) $NC^{S\alpha}$ -continuous (in short $NC^{S\alpha}$ -CF) iff $\forall \mathfrak{B}$ NC-OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a $NC^{S\alpha}$ -OS in \mathcal{S} .
- (ii) $NC^{S\alpha*}$ -continuous (in short $NC^{S\alpha*}$ -CF) iff $\forall \mathfrak{B}$ $NC^{S\alpha}$ -OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a $NC^{S\alpha}$ -OS in \mathcal{S} .
- (iii) $NC^{S\alpha**}$ -continuous (in short $NC^{S\alpha**}$ -CF) iff $\forall \mathfrak{B}$ $NC^{S\alpha}$ -OS in \mathcal{I} , then $\rho^{-1}(\mathfrak{B})$ is a NC-OS in \mathcal{S} .

3. Some New Concepts of Weakly Neutrosophic Crisp Separation Axioms

Definition 3.1:

- (i) A NCTS (\mathcal{S}, ζ) is said to be a NC^{α} - T_0 -space if for each pair of distinct neutrosophic crisp points in (\mathcal{S}, ζ) there exists NC^{α} -OS of (\mathcal{S}, ζ) containing one neutrosophic crisp point but not the other.
- (ii) A NCTS (\mathcal{S}, ζ) is said to be a $NC^{S\alpha}$ - T_0 -space if for each pair of distinct neutrosophic crisp points in (\mathcal{S}, ζ) there exists $NC^{S\alpha}$ -OS of (\mathcal{S}, ζ) containing one neutrosophic crisp point but not the other.

Theorem 3.3:

A NCTS (\mathcal{S}, ζ) is $\text{NC}^\alpha\text{-}T_0$ -space ($\text{NC}^{\text{S}\alpha}\text{-}T_0$ -space respectively) iff $\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \neq \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)$ ($\text{NC}^{\text{S}\alpha}\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \neq \text{NC}^{\text{S}\alpha}\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)$ receptively) for each $u \neq v$ in \mathcal{S} .

Proof:

\Rightarrow Let $\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \neq \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)$, $\forall u \neq v \in \mathcal{S}$. Hence $\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \not\subseteq \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)$ or $\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle) \not\subseteq \text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle)$. Suppose that $\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \not\subseteq \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle) \Rightarrow u \notin \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle) \Rightarrow u \in (\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle))^c$ but $(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle))^c$ is a NC^α -OS and $v \notin (\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle))^c$. Therefore \mathcal{S} is a $\text{NC}^\alpha\text{-}T_0$ -space.

\Leftarrow Let \mathcal{S} be a $\text{NC}^\alpha\text{-}T_0$ -space, $\forall u \neq v \in \mathcal{S}$. Hence there exists a NC^α -OS \mathfrak{B} in \mathcal{S} such that $u \in \mathfrak{B}, v \notin \mathfrak{B}$ or $u \notin \mathfrak{B}, v \in \mathfrak{B}$. Then \mathfrak{B}^c is a NC^α -CS and $u \notin \mathfrak{B}^c, v \in \mathfrak{B}^c$. Therefore $u \notin \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)$ (since $u \notin \mathfrak{B}^c$). Hence $\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \not\subseteq \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)$. The second case is similar. ■

Theorem 3.4:

If (\mathcal{S}, ζ) is a $\text{NC}^\alpha\text{-}T_0$ -space ($\text{NC}^{\text{S}\alpha}\text{-}T_0$ -space respectively), then $\text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle)) \cap \text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)) = \emptyset_N$ ($\text{NC}^{\text{S}\alpha}\text{int}(\text{NC}^{\text{S}\alpha}\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle)) \cap \text{NC}^{\text{S}\alpha}\text{int}(\text{NC}^{\text{S}\alpha}\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)) = \emptyset_N$ receptively), $\forall u \neq v$ in \mathcal{S} .

Proof:

Let (\mathcal{S}, ζ) be a $\text{NC}^\alpha\text{-}T_0$ -space. Then there exists a NC^α -OS \mathfrak{B} such that $u \in \mathfrak{B}, v \notin \mathfrak{B}$ or $u \notin \mathfrak{B}, v \in \mathfrak{B}$. If $u \in \mathfrak{B}, v \notin \mathfrak{B} \Rightarrow u \notin \mathfrak{B}^c, v \in \mathfrak{B}^c$. Thus $\text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)) \subseteq \text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle) \subseteq \mathfrak{B}^c = \text{NC}^\alpha\text{cl}(\mathfrak{B}^c)$ (since \mathfrak{B}^c is a NC^α -CS). Hence $\text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)) \subseteq \mathfrak{B}^c \Rightarrow \text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)) \cap \mathfrak{B} = \emptyset_N$. Therefore, $u \in \mathfrak{B} \subseteq (\text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)))^c$. Hence $\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle) \subseteq (\text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)))^c \Rightarrow \text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle)) \subseteq (\text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)))^c \Rightarrow \text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{u\}, \emptyset, \emptyset \rangle)) \cap \text{NC}^\alpha\text{int}(\text{NC}^\alpha\text{cl}(\langle \{v\}, \emptyset, \emptyset \rangle)) = \emptyset_N$. The second case is similar. ■

Remark 3.5:

- (i) Every $\text{NC}\text{-}T_0$ -space is a $\text{NC}^\alpha\text{-}T_0$ -space and $\text{NC}^{\text{S}\alpha}\text{-}T_0$ -space.
- (ii) Every $\text{NC}^\alpha\text{-}T_0$ -space is a $\text{NC}^{\text{S}\alpha}\text{-}T_0$ -space.

Remark 3.6:

- (i) $\text{NC}^\alpha\text{-}T_0$ ($\text{NC}^{\text{S}\alpha}\text{-}T_0$ respectively) property is a $\text{NC}^{\alpha*}$ ($\text{NC}^{\text{S}\alpha*}$ respectively) topological property.
- (ii) $\text{NC}^\alpha\text{-}T_0$ ($\text{NC}^{\text{S}\alpha}\text{-}T_0$ respectively) property is a $\text{NC}^{\alpha**}$ ($\text{NC}^{\text{S}\alpha**}$ respectively) topological property.
- (iii) $\text{NC}^\alpha\text{-}T_0$ is a NC^α -hereditary property.

Proposition 3.7:

- (i) Let (\mathcal{S}, ζ) and (\mathcal{J}, η) be $\text{NC}^\alpha\text{-}T_0$ -spaces if and only if $\mathcal{S} \times \mathcal{J}$ is a $\text{NC}^\alpha\text{-}T_0$ -space.
- (ii) If (\mathcal{S}, ζ) and (\mathcal{J}, η) are $\text{NC}^{\text{S}\alpha}\text{-}T_0$ -spaces, then $\mathcal{S} \times \mathcal{J}$ is a $\text{NC}^{\text{S}\alpha}\text{-}T_0$ -space.

Proof:

(i) \Rightarrow Let \mathcal{S} and \mathcal{J} be $\text{NC}^\alpha\text{-}T_0$ -spaces. Let $(u_1, v_1) \neq (u_2, v_2)$ in $\mathcal{S} \times \mathcal{J}$. Then $u_1 \neq u_2$ in $\mathcal{S} \Rightarrow$ there exists $\mathfrak{B}_1 \in \text{NC}^\alpha\mathcal{O}(\mathcal{S})$ such that $u_1 \in \mathfrak{B}_1, u_2 \notin \mathfrak{B}_1$ or $u_1 \notin \mathfrak{B}_1, u_2 \in \mathfrak{B}_1$. Also $v_1 \neq v_2$ in $\mathcal{J} \Rightarrow$ there exists $\mathfrak{B}_2 \in \text{NC}^\alpha\mathcal{O}(\mathcal{J})$ such that $v_1 \in \mathfrak{B}_2, v_2 \notin \mathfrak{B}_2$ or $v_1 \notin \mathfrak{B}_2, v_2 \in \mathfrak{B}_2$. Then $(u_1, v_1) \in \mathfrak{B}_1 \times \mathfrak{B}_2, (u_2, v_2) \notin \mathfrak{B}_1 \times \mathfrak{B}_2$ or $(u_1, v_1) \notin \mathfrak{B}_1 \times \mathfrak{B}_2, (u_2, v_2) \in \mathfrak{B}_1 \times \mathfrak{B}_2$. But $\mathfrak{B}_1 \times \mathfrak{B}_2 \in \text{NC}^\alpha\mathcal{O}(\mathcal{S} \times \mathcal{J})$ (since by theorem (2.6)). Hence $\mathcal{S} \times \mathcal{J}$ is a $\text{NC}^\alpha\text{-}T_0$ -space.

\Leftarrow Let $\mathcal{S} \times \mathcal{I}$ be a $\text{NC}^\alpha\text{-}T_0$ -space, to prove that \mathcal{S} and \mathcal{I} are $\text{NC}^\alpha\text{-}T_0$ -spaces. Since $\mathcal{S} \times \mathcal{I}$ is a $\text{NC}^\alpha\text{-}T_0$ -space, then $\mathcal{S} \times \langle \{v_0\}, \emptyset, \emptyset \rangle$ and $\langle \{u_0\}, \emptyset, \emptyset \rangle \times \mathcal{I}$ are $\text{NC}^\alpha\text{-}T_0$ -spaces (since $\text{NC}^\alpha\text{-}T_0$ property is a NC^α -hereditary). Hence \mathcal{S} and \mathcal{I} are $\text{NC}^\alpha\text{-}T_0$ -spaces. The proof (ii) is evident for others. ■

Definition 3.8:

- (i) A NCTS (\mathcal{S}, ζ) is said to be a $\text{NC}^\alpha\text{-}T_1$ -space if for each pair of distinct NC- points u and v of \mathcal{S} , there exist two NC^α -OSs \mathfrak{B} and \mathfrak{D} containing u and v respectively, such that $u \in \mathfrak{B}$, $v \in \mathfrak{D}$.
- (ii) A NCTS (\mathcal{S}, ζ) is said to be a $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space if for each pair of distinct NC-points u and v of \mathcal{S} , there exist two $\text{NC}^{\text{S}\alpha}$ -OSs \mathfrak{B} and \mathfrak{D} containing u and v respectively, such that $u \in \mathfrak{B}$, $v \in \mathfrak{D}$.

Proposition 3.9:

A NCTS (\mathcal{S}, ζ) is $\text{NC}^\alpha\text{-}T_1$ -space ($\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space respectively) if and only if $\langle \{u\}, \emptyset, \emptyset \rangle$ is a $\text{NC}^\alpha\text{-CS}$ ($\text{NC}^{\text{S}\alpha}\text{-CS}$ respectively), $\forall u \in \mathcal{S}$.

Proof:

\Rightarrow Let \mathcal{S} be a $\text{NC}^\alpha\text{-}T_1$ -space. Let $w \in \mathcal{S}$, to prove that $\langle \{w\}, \emptyset, \emptyset \rangle$ is a $\text{NC}^\alpha\text{-CS}$. Let $u \in (\langle \{w\}, \emptyset, \emptyset \rangle)^c \Rightarrow u \neq w$ in \mathcal{S} . Hence there exists a NC^α -OS \mathfrak{B} such that $u \in \mathfrak{B}$, $w \notin \mathfrak{B}$ or $u \notin \mathfrak{B}$, $w \in \mathfrak{B}$. If $u \in \mathfrak{B}$, $w \notin \mathfrak{B} \Rightarrow u \in \mathfrak{B} \subseteq (\langle \{w\}, \emptyset, \emptyset \rangle)^c \Rightarrow (\langle \{w\}, \emptyset, \emptyset \rangle)^c$ is a $\text{NC}^\alpha\text{-OS} \Rightarrow \langle \{w\}, \emptyset, \emptyset \rangle$ is a $\text{NC}^\alpha\text{-CS}$.

\Leftarrow Let $\langle \{w\}, \emptyset, \emptyset \rangle$ be a $\text{NC}^\alpha\text{-CS}$, $\forall w \in \mathcal{S}$, to prove that \mathcal{S} is a $\text{NC}^\alpha\text{-}T_1$ -space. Let $u \neq v$ in \mathcal{S} . Hence $\langle \{u\}, \emptyset, \emptyset \rangle, \langle \{v\}, \emptyset, \emptyset \rangle$ are $\text{NC}^\alpha\text{-CSs} \Rightarrow (\langle \{u\}, \emptyset, \emptyset \rangle)^c, (\langle \{v\}, \emptyset, \emptyset \rangle)^c$ are $\text{NC}^\alpha\text{-OSs}$ and $v \in (\langle \{u\}, \emptyset, \emptyset \rangle)^c, u \notin (\langle \{u\}, \emptyset, \emptyset \rangle)^c, u \in (\langle \{v\}, \emptyset, \emptyset \rangle)^c, v \notin (\langle \{v\}, \emptyset, \emptyset \rangle)^c$. Therefore \mathcal{S} is a $\text{NC}^\alpha\text{-}T_1$ -space. The second case is similar. ■

Remark 3.10:

- (i) Every $\text{NC}\text{-}T_1$ -space is a $\text{NC}^\alpha\text{-}T_1$ -space and $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space.
- (ii) Every $\text{NC}^\alpha\text{-}T_1$ -space is a $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space.
- (iii) Every $\text{NC}^\alpha\text{-}T_1$ -space is a $\text{NC}^\alpha\text{-}T_0$ -space.
- (iv) Every $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space is a $\text{NC}^{\text{S}\alpha}\text{-}T_0$ -space.

Remark 3.11:

- (i) $\text{NC}^\alpha\text{-}T_1$ ($\text{NC}^{\text{S}\alpha}\text{-}T_1$ respectively) property is a $\text{NC}^{\alpha*}$ ($\text{NC}^{\text{S}\alpha*}$ respectively) topological property.
- (ii) $\text{NC}^\alpha\text{-}T_1$ ($\text{NC}^{\text{S}\alpha}\text{-}T_1$ respectively) property is a $\text{NC}^{\alpha**}$ ($\text{NC}^{\text{S}\alpha**}$ respectively) topological property.
- (iii) $\text{NC}^\alpha\text{-}T_1$ property is a NC^α -hereditary property.

Proposition 3.12:

- (i) Let \mathcal{S} and \mathcal{I} be $\text{NC}^\alpha\text{-}T_1$ -spaces if and only if $\mathcal{S} \times \mathcal{I}$ is a $\text{NC}^\alpha\text{-}T_1$ -space.
- (ii) If \mathcal{S} and \mathcal{I} are $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -spaces, then $\mathcal{S} \times \mathcal{I}$ is a $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space.

Proof:

The proof of this is similar to that of proposition (3.7). ■

Definition 3.13:

- (i) A NCTS (\mathcal{S}, ζ) is said to be a $\text{NC}^\alpha\text{-}T_2$ -space if for each pair of distinct NC-points u and v in \mathcal{S} , there exist two NC^α -OSs \mathfrak{D}_1 and \mathfrak{D}_2 such that $u \in \mathfrak{D}_1$, $v \in \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_{\mathcal{N}}$.
- (ii) A NCTS (\mathcal{S}, ζ) is said to be a $\text{NC}^{\text{S}\alpha}\text{-}T_2$ -space if for each pair of distinct NC-points u and v in \mathcal{S} , there exist two $\text{NC}^{\text{S}\alpha}$ -OSs \mathfrak{D}_1 and \mathfrak{D}_2 such that $u \in \mathfrak{D}_1$, $v \in \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_{\mathcal{N}}$.

Proposition 3.14:

If (\mathcal{S}, ζ) is a $\text{NC}^\alpha\text{-}T_2$ -space ($\text{NC}^{\text{S}\alpha}\text{-}T_2$ -space respectively), then $\mathfrak{B} = \{(u, v): u = v, u, v \in \mathcal{S}\}$ is a $\text{NC}^\alpha\text{-CS}$ ($\text{NC}^{\text{S}\alpha}\text{-CS}$ respectively).

Proof:

Let \mathcal{S} be a $\text{NC}^\alpha\text{-}T_2$ -space, to prove that \mathfrak{B} is a $\text{NC}^\alpha\text{-CS}$. Let $(u, v) \in \mathfrak{B}^c = \mathcal{S} \times \mathcal{S} - \mathfrak{B}$. Hence $u \neq v$ in $\mathcal{S} \Rightarrow$ there exist $\mathfrak{D}_1, \mathfrak{D}_2 \in \text{NC}^\alpha\mathcal{O}(\mathcal{S})$ such that $u \in \mathfrak{D}_1, v \in \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_{\mathcal{N}}$ (since \mathcal{S} is a $\text{NC}^\alpha\text{-}T_2$ -space). Hence $\mathfrak{D}_1 \times \mathfrak{D}_2 \in \text{NC}^\alpha\mathcal{O}(\mathcal{S} \times \mathcal{S})$ by theorem (2.7) $(u, v) \in \mathfrak{D}_1 \times \mathfrak{D}_2 \subseteq \mathfrak{B}^c$, hence \mathfrak{B}^c is a $\text{NC}^\alpha\text{-OS}$. Therefore \mathfrak{B} is a $\text{NC}^\alpha\text{-CS}$. The second case is similar. ■

Remark 3.15:

- (i) Every $\text{NC}\text{-}T_2$ -space is a $\text{NC}^\alpha\text{-}T_2$ -space and $\text{NC}^{\text{S}\alpha}\text{-}T_2$ -space.
- (ii) Every $\text{NC}^\alpha\text{-}T_2$ -space is a $\text{NC}^{\text{S}\alpha}\text{-}T_2$ -space.
- (iii) Every $\text{NC}^\alpha\text{-}T_2$ -space is a $\text{NC}^\alpha\text{-}T_1$ -space.
- (iv) Every $\text{NC}^{\text{S}\alpha}\text{-}T_2$ -space is a $\text{NC}^{\text{S}\alpha}\text{-}T_1$ -space.

Remark 3.16:

- (i) $\text{NC}^\alpha\text{-}T_2$ ($\text{NC}^{\text{S}\alpha}\text{-}T_2$ respectively) property is a NC^{α^*} ($\text{NC}^{\text{S}\alpha^*}$ respectively) topological property.
- (ii) $\text{NC}^\alpha\text{-}T_2$ ($\text{NC}^{\text{S}\alpha}\text{-}T_2$ respectively) property is a $\text{NC}^{\alpha^{**}}$ ($\text{NC}^{\text{S}\alpha^{**}}$ respectively) topological property.
- (iii) $\text{NC}^\alpha\text{-}T_2$ property is a NC^α -hereditary property.

Proposition 3.17:

- (i) Let \mathcal{S} and \mathcal{I} be $\text{NC}^\alpha\text{-}T_2$ -spaces if and only if $\mathcal{S} \times \mathcal{I}$ is a $\text{NC}^\alpha\text{-}T_2$ -space.
- (ii) If \mathcal{S} and \mathcal{I} are $\text{NC}^{\text{S}\alpha}\text{-}T_2$ -spaces, then $\mathcal{S} \times \mathcal{I}$ is a $\text{NC}^{\text{S}\alpha}\text{-}T_2$ -space.

Proof:

The proof of this is similar to that of proposition (3.12). ■

Proposition 3.18:

- (i) If $\rho, \mu: \mathcal{S} \rightarrow \mathcal{I}$ are $\text{NC}^{\alpha^*}\text{-CF}$ and \mathcal{I} is a $\text{NC}^\alpha\text{-}T_2$ -space, then the NC -set $\mathfrak{B} = \{u: u \in \mathcal{S}, \rho(u) = \mu(u)\}$ is a $\text{NC}^\alpha\text{-CS}$.
- (ii) If $\rho, \mu: \mathcal{S} \rightarrow \mathcal{I}$ are $\text{NC}^\alpha\text{-CF}$ and \mathcal{I} is a $\text{NC}\text{-}T_2$ -space, then the NC -set $\mathfrak{B} = \{u: u \in \mathcal{S}, \rho(u) = \mu(u)\}$ is a $\text{NC}^\alpha\text{-CS}$.

Proof:

(i) If $u \notin \mathfrak{B} \Rightarrow u \in \mathfrak{B}^c \Rightarrow \rho(u) \neq \mu(u)$ in \mathcal{I} . Hence there exist two $\text{NC}^\alpha\text{-OSs}$ \mathfrak{D}_1 and \mathfrak{D}_2 in \mathcal{I} such that $\rho(u) \in \mathfrak{D}_1, \mu(u) \in \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_{\mathcal{N}}$ (since \mathcal{I} is a $\text{NC}^\alpha\text{-}T_2$ -space). But $\rho^{-1}(\mathfrak{D}_1), \mu^{-1}(\mathfrak{D}_2) \in \text{NC}^\alpha\mathcal{O}(\mathcal{S})$ (since ρ, μ are $\text{NC}^{\alpha^*}\text{-CF}$). Therefore, $u \in \rho^{-1}(\mathfrak{D}_1)$ and $u \in \mu^{-1}(\mathfrak{D}_2)$. Hence $u \in \rho^{-1}(\mathfrak{D}_1) \cap \mu^{-1}(\mathfrak{D}_2)$. Let $\mathfrak{U} = \rho^{-1}(\mathfrak{D}_1) \cap \mu^{-1}(\mathfrak{D}_2) \in \text{NC}^\alpha\mathcal{O}(\mathcal{S})$. To prove $\mathfrak{U} \subseteq \mathfrak{B}^c$, i.e., $\mathfrak{U} \cap \mathfrak{B} = \emptyset_{\mathcal{N}}$. Suppose that $\mathfrak{U} \cap \mathfrak{B} \neq \emptyset_{\mathcal{N}} \Rightarrow \exists v \in \mathfrak{U} \cap \mathfrak{B} \Rightarrow v \in \mathfrak{U}$ and $v \in \mathfrak{B}$, i.e., $v \in \rho^{-1}(\mathfrak{D}_1)$ and $v \in \mu^{-1}(\mathfrak{D}_2)$ and $v \in \mathfrak{B}$. Hence $\rho(v) \in \mathfrak{D}_1, \mu(v) \in \mathfrak{D}_2$ and $v \in \mathfrak{B}$. Therefore $\rho(v) = \mu(v)$ (since $v \in \mathfrak{B}$). Hence $\mathfrak{D}_1 \cap \mathfrak{D}_2 \neq \emptyset_{\mathcal{N}}$ which is a contradiction. Therefore $\mathfrak{U} \subseteq \mathfrak{B}^c \Rightarrow \mathfrak{B}^c \in \text{NC}^\alpha\mathcal{O}(\mathcal{S}) \Rightarrow \mathfrak{B}$ is a $\text{NC}^\alpha\text{-CS}$. The proof (ii) is evident for others. ■

4. Some New Concepts of Weakly Neutrosophic Crisp Regularity

Definition 4.1:


Let (\mathcal{S}, ζ) be a NCTS, then \mathcal{S} is said to be:

- (i) NC^α -regular ($\text{NC}^{\text{S}\alpha}$ -regular respectively) if every $u \in \mathcal{S}$ and every \mathcal{Q} $\text{NC}\text{-CS}$ such that $u \notin \mathcal{Q}$,

- there exist two NC^α -OSs ($\text{NC}^{\text{S}\alpha}$ -OSs respectively) \mathfrak{B} and \mathfrak{D} such that $u \in \mathfrak{B}$, $Q \subseteq \mathfrak{D}$ and $\mathfrak{B} \cap \mathfrak{D} = \emptyset_{\mathcal{N}}$.
- (ii) NC^{α^*} -regular ($\text{NC}^{\text{S}\alpha^*}$ -regular respectively) if every $u \in \mathcal{S}$ and every Q NC^α -CS ($\text{NC}^{\text{S}\alpha}$ -CS respectively) such that $u \notin Q$, there exist two NC^α -OSs ($\text{NC}^{\text{S}\alpha}$ -OSs respectively) \mathfrak{B} and \mathfrak{D} such that $u \in \mathfrak{B}$, $Q \subseteq \mathfrak{D}$ and $\mathfrak{B} \cap \mathfrak{D} = \emptyset_{\mathcal{N}}$.
- (iii) $\text{NC}^{\alpha^{**}}$ -regular ($\text{NC}^{\text{S}\alpha^{**}}$ -regular respectively) if every $u \in \mathcal{S}$ and every Q NC^α -CS ($\text{NC}^{\text{S}\alpha}$ -CS respectively) such that $u \notin Q$, there exist two NC -OSs \mathfrak{B} and \mathfrak{D} such that $u \in \mathfrak{B}$, $Q \subseteq \mathfrak{D}$ and $\mathfrak{B} \cap \mathfrak{D} = \emptyset_{\mathcal{N}}$.

Remark 4.2:

The following diagram shows the relation between the different types of weakly NC-regular and weakly NC^α -regular ($\text{NC}^{\text{S}\alpha}$ -regular respectively) spaces:

**Fig. 4.1****Theorem 4.3:**

Let (\mathcal{S}, ζ) be a NCTS, then:

- (i) \mathcal{S} is a NC^α -regular if and only if for each \mathfrak{B} NC -OS containing u , there exists \mathfrak{D} NC^α -OS containing u such that $u \in \mathfrak{D} \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}) \subseteq \mathfrak{B}$.
- (ii) \mathcal{S} is a NC^{α^*} -regular if and only if for each \mathfrak{B} NC^α -OS contains u , there exists \mathfrak{D} NC^α -OS contains u such that $u \in \mathfrak{D} \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}) \subseteq \mathfrak{B}$.
- (iii) \mathcal{S} is a $\text{NC}^{\alpha^{**}}$ -regular if and only if for each \mathfrak{B} NC^α -OS contains u , there exists \mathfrak{D} NC -OS contains u such that $u \in \mathfrak{D} \subseteq \text{NCcl}(\mathfrak{D}) \subseteq \mathfrak{B}$.

Proof:

(i) \Rightarrow Let \mathcal{S} be a NC^α -regular space and let \mathfrak{B} be a NC -OS containing u . Hence \mathfrak{B}^c is a NC -CS and $u \notin \mathfrak{B}^c$. Then there exist $\mathfrak{D}_1, \mathfrak{D}_2$ NC^α -OSs in \mathcal{S} such that $u \in \mathfrak{D}_1$, $\mathfrak{B}^c \subseteq \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_{\mathcal{N}}$ (since \mathcal{S} is a NC^α -regular space). Hence $u \in \mathfrak{D}_1 \subseteq \mathfrak{D}_2^c \subseteq \mathfrak{B}$ (since $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_{\mathcal{N}} \Rightarrow \mathfrak{D}_1 \subseteq \mathfrak{D}_2^c$). Therefore $u \in \mathfrak{D}_1 \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_1) \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_2^c) \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{B})$. Therefore $u \in \mathfrak{D}_1 \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_1) \subseteq \mathfrak{D}_2^c \subseteq \mathfrak{B}$. This implies that $u \in \mathfrak{D}_1 \subseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_1) \subseteq \mathfrak{B}$, where \mathfrak{D}_1 is a NC^α -OS.

\Leftarrow Let \mathcal{Q} be a NC-CS such that $u \notin \mathcal{Q} \Rightarrow \mathcal{Q}^c$ is a NC-OS contains u . Hence there exists \mathcal{D} NC $^\alpha$ -OS contains u such that $u \in \mathcal{D} \subseteq NC^\alpha cl(\mathcal{D}) \subseteq \mathcal{Q}^c$. We get $\mathcal{Q} \subseteq (NC^\alpha cl(\mathcal{D}))^c$, so it is $(NC^\alpha cl(\mathcal{D}))^c$ is a NC $^\alpha$ -OS and contains \mathcal{Q} . Now, to prove $\mathcal{D} \cap (NC^\alpha cl(\mathcal{D}))^c = \emptyset_N$. Since $\mathcal{D} \subseteq NC^\alpha cl(\mathcal{D})$, but $NC^\alpha cl(\mathcal{D}) \cap (NC^\alpha cl(\mathcal{D}))^c = \emptyset_N \Rightarrow \mathcal{D} \cap (NC^\alpha cl(\mathcal{D}))^c = \emptyset_N$. Hence \mathcal{S} is a NC $^\alpha$ -regular space. The proofs (ii), (iii) are evident for others. ■

Theorem 4.4:

Let (\mathcal{S}, ζ) be a NCTS, then:

- (i) \mathcal{S} is a NC $^{S\alpha^*}$ -regular if and only if for each \mathcal{B} NC $^{S\alpha}$ -OS contains u , there exists \mathcal{D} NC $^{S\alpha}$ -OS contains u such that $u \in \mathcal{D} \subseteq NC^{S\alpha} cl(\mathcal{D}) \subseteq \mathcal{B}$.
- (ii) \mathcal{S} is a NC $^{S\alpha^{**}}$ -regular if and only if for each \mathcal{B} NC $^{S\alpha}$ -OS contains u , there exists \mathcal{D} NC-OS contains u such that $u \in \mathcal{D} \subseteq NC cl(\mathcal{D}) \subseteq \mathcal{B}$.

Proof:

The proof of this is similar to that of theorem (4.3). ■

Theorem 4.5:

Let (\mathcal{S}, ζ) be a NCTS, then:

- (i) \mathcal{S} is a NC $^\alpha$ -regular if and only if $u \notin \mathcal{Q}$ where \mathcal{Q} is a NC $^\alpha$ -CS, there exist two NC $^\alpha$ -OSs \mathcal{B} and \mathcal{D} such that $u \in \mathcal{B}, \mathcal{Q} \subseteq \mathcal{D}$ and $NC^\alpha cl(\mathcal{B}) \cap NC^\alpha cl(\mathcal{D}) = \emptyset_N$.
- (ii) \mathcal{S} is a NC $^{\alpha^{**}}$ -regular if and only if for each \mathcal{Q} NC $^\alpha$ -CS, such that $u \notin \mathcal{Q}$, there exist two NC-OSs \mathcal{B} and \mathcal{D} such that $u \in \mathcal{B}, \mathcal{Q} \subseteq \mathcal{D}$ and $NC cl(\mathcal{B}) \cap NC cl(\mathcal{D}) = \emptyset_N$.

Proof:

(i) Let \mathcal{S} be a NC $^\alpha$ -regular space and let \mathcal{Q} be a NC $^\alpha$ -CS, such that $u \notin \mathcal{Q}$. Then there exist two NC $^\alpha$ -OSs \mathcal{U} and \mathcal{V} such that $u \in \mathcal{U}, \mathcal{Q} \subseteq \mathcal{V}$ and $\mathcal{U} \cap \mathcal{V} = \emptyset_N$. Therefore \mathcal{U} is a NC $^\alpha$ -OS containing u in \mathcal{S} , where \mathcal{S} is a NC $^\alpha$ -regular space. Then there exists \mathcal{B} NC $^\alpha$ -OS containing u such that $u \in \mathcal{B} \subseteq NC^\alpha cl(\mathcal{B}) \subseteq \mathcal{U}$ (since by theorem (4.3) (ii)). Hence $NC^\alpha cl(\mathcal{B}) \subseteq \mathcal{U}$. Also, $\mathcal{Q} \subseteq \mathcal{V} \subseteq NC^\alpha cl(\mathcal{V})$, but $NC^\alpha cl(\mathcal{V}) \subseteq (NC^\alpha cl(\mathcal{U}))^c$ (since $\mathcal{U} \cap \mathcal{V} = \emptyset_N \Rightarrow \mathcal{V} \subseteq \mathcal{U}^c \Rightarrow NC^\alpha cl(\mathcal{V}) \subseteq NC^\alpha cl(\mathcal{U}^c)$). Hence $\mathcal{Q} \subseteq \mathcal{V} \subseteq NC^\alpha cl(\mathcal{V}) \subseteq NC^\alpha cl(\mathcal{U}^c) = \mathcal{U}^c$ (since \mathcal{U}^c is a NC $^\alpha$ -CS). Suppose that $\mathcal{V} = \mathcal{D}$, hence $\mathcal{Q} \subseteq \mathcal{D} \subseteq NC^\alpha cl(\mathcal{D}) \subseteq \mathcal{U}^c \Rightarrow NC^\alpha cl(\mathcal{D}) \subseteq \mathcal{U}^c$. Since $\mathcal{U} \cap \mathcal{U}^c = \emptyset_N$, hence $NC^\alpha cl(\mathcal{B}) \cap NC^\alpha cl(\mathcal{D}) = \emptyset_N$ (since $NC^\alpha cl(\mathcal{B}) \subseteq \mathcal{U}$ and $NC^\alpha cl(\mathcal{D}) \subseteq \mathcal{U}^c$). The other side is clear. The proof (ii) is evident for others. ■

Theorem 4.6:

Let (\mathcal{S}, ζ) be a NCTS, then:

- (i) \mathcal{S} is a NC $^{S\alpha^*}$ -regular if and only if $u \notin \mathcal{Q}$, where \mathcal{Q} is a NC $^{S\alpha}$ -CS, there exist two NC $^{S\alpha}$ -OSs \mathcal{B} and \mathcal{D} such that $u \in \mathcal{B}, \mathcal{Q} \subseteq \mathcal{D}$ and $NC^{S\alpha} cl(\mathcal{B}) \cap NC^{S\alpha} cl(\mathcal{D}) = \emptyset_N$.
- (ii) \mathcal{S} is a NC $^{S\alpha^{**}}$ -regular if and only if $u \notin \mathcal{Q}$, where \mathcal{Q} is a NC $^{S\alpha}$ -CS, there exist two NC-OSs \mathcal{B} and \mathcal{D} such that $u \in \mathcal{B}, \mathcal{Q} \subseteq \mathcal{D}$ and $NC cl(\mathcal{B}) \cap NC cl(\mathcal{D}) = \emptyset_N$.

Proof:

The proof of this is similar to that of theorem (4.5). ■

Remark 4.7:

- (i) NC $^\alpha$ -regular property is a NC $^{\alpha^{**}}$ -topological property.
- (ii) NC $^{\alpha^*}$ -regular property is a NC $^{\alpha^*}$ -topological property.
- (iii) NC $^{\alpha^{**}}$ -regular property is a NC $^{\alpha^{**}}$ -topological property.
- (iv) NC $^{S\alpha}$ -regular property is a NC $^{S\alpha^{**}}$ -topological property.
- (v) NC $^{S\alpha^*}$ -regular property is a NC $^{S\alpha^*}$ -topological property.

(vi) $NC^{S\alpha^{**}}$ -regular property is a $NC^{S\alpha^{**}}$ -topological property.

Proposition 4.8:

- (i) If $\mathcal{S} \times \mathcal{J}$ is a $NC^{\alpha^{**}}$ -regular, then both \mathcal{S} and \mathcal{J} are $NC^{\alpha^{**}}$ -regular spaces.
- (ii) If $\mathcal{S} \times \mathcal{J}$ is a $NC^{S\alpha^{**}}$ -regular, then both \mathcal{S} and \mathcal{J} are $NC^{S\alpha^{**}}$ -regular spaces.

Proof:

(i) Suppose that $\mathcal{S} \times \mathcal{J}$ is a $NC^{\alpha^{**}}$ -regular, to prove that \mathcal{S} and \mathcal{J} are $NC^{\alpha^{**}}$ -regular spaces. Let \mathcal{U} and \mathcal{V} be two NC^{α} -OSs in \mathcal{S} and \mathcal{J} containing u and v respectively. Hence $(u, v) \in \mathcal{U} \times \mathcal{V}$ where $\mathcal{U} \times \mathcal{V}$ is a NC^{α} -OS in $\mathcal{S} \times \mathcal{J}$ (by theorem (2.7)). Hence there exists NC -OS \mathcal{K} in $\mathcal{S} \times \mathcal{J}$ such that $(u, v) \in \mathcal{K} \subseteq NCcl(\mathcal{K}) \subseteq \mathcal{U} \times \mathcal{V}$ (since $\mathcal{S} \times \mathcal{J}$ is a $NC^{\alpha^{**}}$ -regular). Then there exist two NC -OSs \mathcal{B} and \mathcal{D} in \mathcal{S} and \mathcal{J} such that $(u, v) \in \mathcal{B} \times \mathcal{D} \subseteq NCcl(\mathcal{B} \times \mathcal{D}) = NCcl(\mathcal{B}) \times NCcl(\mathcal{D}) \subseteq \mathcal{U} \times \mathcal{V}$. Hence $u \in \mathcal{B} \subseteq NCcl(\mathcal{B}) \subseteq \mathcal{U} \Rightarrow \mathcal{S}$ is a $NC^{\alpha^{**}}$ -regular space. Also, $v \in \mathcal{D} \subseteq NCcl(\mathcal{D}) \subseteq \mathcal{V} \Rightarrow \mathcal{J}$ is a $NC^{\alpha^{**}}$ -regular space. The proof (ii) is evident for others. ■

Theorem 4.9:

If (\mathcal{S}, ζ) is a NC^{α^*} -regular ($NC^{\alpha^{**}}$ -regular respectively), then $\zeta = NC^{\alpha}O(\mathcal{S})$.

Proof:

It is clear that $\zeta \subseteq NC^{\alpha}O(\mathcal{S})$. Let \mathcal{B} be a NC^{α} -OS in \mathcal{S} containing u . Then there exists a NC^{α} -OS \mathcal{D} containing u such that $u \in \mathcal{D} \subseteq NC^{\alpha}cl(\mathcal{D}) \subseteq \mathcal{B}$ (\mathcal{S} is a NC^{α^*} -regular). Therefore $NC^{\alpha}int(\mathcal{D}) \subseteq NC^{\alpha}int(NC^{\alpha}cl(\mathcal{D})) \subseteq \mathcal{B}$. Thus $u \in \mathcal{D} \subseteq NCcl(NCint(\mathcal{D})) \subseteq \mathcal{B}$ (since by proposition (2.9)). Hence \mathcal{B} is a NC -OS $\Rightarrow NC^{\alpha}O(\mathcal{S}) \subseteq \zeta$. Therefore $\zeta = NC^{\alpha}O(\mathcal{S})$. ■

Proposition 4.10:

- (i) If $\rho: \mathcal{S} \rightarrow \mathcal{J}$ is a NC^{α} -CF and \mathcal{S} is a NC^{α^*} -regular, then ρ is a NC -CF.
- (ii) If $\rho: \mathcal{S} \rightarrow \mathcal{J}$ is a NC^{α} -CF and \mathcal{J} is a NC^{α^*} -regular, then ρ is a NC^{α^*} -CF.
- (iii) If $\rho: \mathcal{S} \rightarrow \mathcal{J}$ is a NC^{α^*} -CF and \mathcal{S} is a NC^{α^*} -regular, then ρ is a $NC^{\alpha^{**}}$ -CF.

Proof:

(i) Let $\rho: \mathcal{S} \rightarrow \mathcal{J}$ be a NC^{α} -CF, to prove that ρ is a NC -CF. Let \mathcal{B} be a NC -OS in \mathcal{J} , then $\rho^{-1}(\mathcal{B})$ is a NC^{α} -OS in \mathcal{S} (since ρ is a NC^{α} -CF). But \mathcal{S} is a NC^{α^*} -regular space (by hypothesis). Hence $\rho^{-1}(\mathcal{B})$ is a NC^{α} -OS in \mathcal{S} (since by theorem (4.9)). Therefore ρ is a NC -CF. The proofs (ii), (iii) are evident for others. ■

Definition 4.11:

Let (\mathcal{S}, ζ) be a NCTS, then \mathcal{S} is said to be:

- (i) $NC^{\alpha}-T_3$ -space if \mathcal{S} is a $NC^{\alpha}-T_1$ -space and NC^{α} -regular space.
- (ii) $NC^{\alpha^*}-T_3$ -space if \mathcal{S} is $NC^{\alpha}-T_1$ -space and NC^{α^*} -regular space.
- (iii) $NC^{\alpha^{**}}-T_3$ -space if \mathcal{S} is $NC^{\alpha}-T_1$ -space and $NC^{\alpha^{**}}$ -regular space.

Definition 4.12:

Let (\mathcal{S}, ζ) be a NCTS, then \mathcal{S} is said to be:

- (i) $NC^{S\alpha}-T_3$ -space if \mathcal{S} is a $NC^{S\alpha}-T_1$ -space and $NC^{S\alpha}$ -regular space.
- (ii) $NC^{S\alpha^*}-T_3$ -space if \mathcal{S} is $NC^{S\alpha}-T_1$ -space and $NC^{S\alpha^*}$ -regular space.
- (iii) $NC^{S\alpha^{**}}-T_3$ -space if \mathcal{S} is $NC^{S\alpha}-T_1$ -space and $NC^{S\alpha^{**}}$ -regular space.

Remark 4.13:

- (i) $NC^{\alpha^*}-T_3$ ($NC^{S\alpha^*}-T_3$ respectively) property is a NC^{α^*} ($NC^{S\alpha^*}$ respectively) topological property.
- (ii) $NC^{\alpha^{**}}-T_3$ ($NC^{S\alpha^{**}}-T_3$ respectively) property is a $NC^{\alpha^{**}}$ ($NC^{S\alpha^{**}}$ respectively) topological property.

Remark 4.14:

- (i) Every $\text{NC}-T_3$ -space is a $\text{NC}^\alpha-T_3$ -space and $\text{NC}^{S\alpha}-T_3$ -space.
- (ii) Every $\text{NC}^\alpha-T_3$ -space is a $\text{NC}^{S\alpha}-T_3$ -space.
- (iii) Every $\text{NC}^{\alpha^{**}}-T_3$ -space ($\text{NC}^{S\alpha^{**}}-T_3$ -space respectively) is a $\text{NC}^{\alpha^*}-T_3$ -space ($\text{NC}^{S\alpha^*}-T_3$ -space, respectively).
- (iv) Every $\text{NC}^{\alpha^*}-T_3$ -space ($\text{NC}^{S\alpha^*}-T_3$ -space respectively) is a $\text{NC}^\alpha-T_2$ -space ($\text{NC}^{S\alpha}-T_2$ -space, respectively).

Proposition 4.15:

$\mathcal{S} \times \mathcal{I}$ is a $\text{NC}^{\alpha^{**}}-T_3$ -space if and only if both \mathcal{S} and \mathcal{I} are $\text{NC}^{\alpha^{**}}-T_3$ -spaces.

Proof:

Follow directly from proposition (3.12) part (i) and proposition (4.8) part (i). ▀


5. Some New Concepts of Weakly Neutrosophic Crisp Normality**Definition 5.1:**

Let (\mathcal{S}, ζ) be a NCTS, then \mathcal{S} is said to be:

- (i) NC^α -normal ($\text{NC}^{S\alpha}$ -normal respectively) if for every two NC-CSs \mathcal{Q}_1 and \mathcal{Q}_2 such that $\mathcal{Q}_1 \cap \mathcal{Q}_2 = \emptyset_{\mathcal{N}}$ There exist two NC^α -OSs ($\text{NC}^{S\alpha}$ -OSs respectively) \mathfrak{B} and \mathfrak{D} such that $\mathcal{Q}_1 \subseteq \mathfrak{B}$ and $\mathcal{Q}_2 \subseteq \mathfrak{D}$ and $\mathfrak{B} \cap \mathfrak{D} = \emptyset_{\mathcal{N}}$.
- (ii) NC^{α^*} -normal ($\text{NC}^{S\alpha^*}$ -normal respectively) if for every two NC^α -CSs ($\text{NC}^{S\alpha}$ -CSs respectively) \mathcal{Q}_1 and \mathcal{Q}_2 such that $\mathcal{Q}_1 \cap \mathcal{Q}_2 = \emptyset_{\mathcal{N}}$ There exist two NC^α -OSs ($\text{NC}^{S\alpha}$ -OSs respectively) \mathfrak{B} and \mathfrak{D} such that $\mathcal{Q}_1 \subseteq \mathfrak{B}$ and $\mathcal{Q}_2 \subseteq \mathfrak{D}$ and $\mathfrak{B} \cap \mathfrak{D} = \emptyset_{\mathcal{N}}$.
- (iii) $\text{NC}^{\alpha^{**}}$ -normal ($\text{NC}^{S\alpha^{**}}$ -normal respectively) if for every two NC^α -CSs ($\text{NC}^{S\alpha}$ -CSs respectively) \mathcal{Q}_1 and \mathcal{Q}_2 such that $\mathcal{Q}_1 \cap \mathcal{Q}_2 = \emptyset_{\mathcal{N}}$, there exist two NC-OSs \mathfrak{B} and \mathfrak{D} such that $\mathcal{Q}_1 \subseteq \mathfrak{B}$ and $\mathcal{Q}_2 \subseteq \mathfrak{D}$ and $\mathfrak{B} \cap \mathfrak{D} = \emptyset_{\mathcal{N}}$.

Remark 5.2:

The following diagram shows the relation between the different types of weakly NC-normal and weakly NC^α -normal ($\text{NC}^{S\alpha}$ -normal respectively) spaces:

**Fig. 5.1****Theorem****5.3:**

Let (\mathcal{S}, ζ) be a NCTS, then:

- (i) \mathcal{S} is a NC^α -normal space if and only if for every NC-CS \mathcal{Q} and every NC-OS \mathfrak{B} containing \mathcal{Q} , there exists NC $^\alpha$ -OS say \mathfrak{D} , such that $\mathcal{Q} \sqsubseteq \mathfrak{D} \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}) \sqsubseteq \mathfrak{B}$.
- (ii) \mathcal{S} is a NC^{α^*} -normal space if and only if for every NC $^\alpha$ -CS \mathcal{Q} and every NC $^\alpha$ -OS \mathfrak{B} containing \mathcal{Q} , there exists NC $^\alpha$ -OS say \mathfrak{D} , such that $\mathcal{Q} \sqsubseteq \mathfrak{D} \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}) \sqsubseteq \mathfrak{B}$.
- (iii) \mathcal{S} is a $\text{NC}^{\alpha^{**}}$ -normal space if and only if for every NC $^\alpha$ -CS \mathcal{Q} and every NC $^\alpha$ -OS \mathfrak{B} containing \mathcal{Q} , there exists NC-OS say \mathfrak{D} , such that $\mathcal{Q} \sqsubseteq \mathfrak{D} \sqsubseteq \text{NCcl}(\mathfrak{D}) \sqsubseteq \mathfrak{B}$.

Proof:

(i) \Rightarrow Let \mathcal{S} be a NC^α -normal space. Let $\mathcal{Q} \sqsubseteq \mathfrak{B}$, where \mathcal{Q} is a NC-CS and \mathfrak{B} is a NC-OS $\Rightarrow \mathcal{Q} \cap \mathfrak{B}^c = \emptyset_N$, where \mathfrak{B}^c is a NC-CS. Hence there exist two NC-OSs $\mathfrak{D}_1, \mathfrak{D}_2$ such that $\mathcal{Q} \sqsubseteq \mathfrak{D}_1$ and $\mathfrak{B}^c \sqsubseteq \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_N$ (since \mathcal{S} is a NC^α -normal space). Therefore $\mathcal{Q} \sqsubseteq \mathfrak{D}_1 \sqsubseteq \mathfrak{D}_2^c \sqsubseteq \mathfrak{B} \Rightarrow \text{NC}^\alpha \text{cl}(\mathcal{Q}) \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_1) \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_2^c) = \mathfrak{D}_2^c \sqsubseteq \mathfrak{B}$. Hence $\mathcal{Q} \sqsubseteq \mathfrak{D}_1 \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}_1) \sqsubseteq \mathfrak{B}$, where \mathfrak{D}_1 is a NC $^\alpha$ -OS in \mathcal{S} .
 \Leftarrow To prove \mathcal{S} is a NC^α -normal space. Let \mathcal{Q}_1 and \mathcal{Q}_2 be NC-CSs in \mathcal{S} such that $\mathcal{Q}_1 \cap \mathcal{Q}_2 = \emptyset_N$. Hence $\mathcal{Q}_1 \sqsubseteq \mathcal{Q}_2^c$, where \mathcal{Q}_2^c is a NC-OS. Then there exists a NC $^\alpha$ -OS \mathfrak{D} such that $\mathcal{Q}_1 \sqsubseteq \mathfrak{D} \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D}) \sqsubseteq \mathcal{Q}_2^c$ (by hypothesis). Hence $\mathcal{Q}_1 \sqsubseteq \mathfrak{D}$, $\mathcal{Q}_2 \sqsubseteq (\text{NC}^\alpha \text{cl}(\mathfrak{D}))^c$. On the other hand $\text{NC}^\alpha \text{cl}(\mathfrak{D}) \cap (\text{NC}^\alpha \text{cl}(\mathfrak{D}))^c = \emptyset_N$. Hence $\mathfrak{D} \cap (\text{NC}^\alpha \text{cl}(\mathfrak{D}))^c = \emptyset_N$ (since $\mathfrak{D} \sqsubseteq \text{NC}^\alpha \text{cl}(\mathfrak{D})$). Therefore \mathcal{S} is a NC^α -normal space. The proofs (ii), (iii) are evident for others. ■

Theorem 5.4:

Let (\mathcal{S}, ζ) be a NCTS, then:

- (i) \mathcal{S} is a $\text{NC}^{\text{S}\alpha}$ -normal space if and only if for every NC-CS \mathcal{Q} and every NC-OS \mathfrak{B} containing \mathcal{Q} , there exists NC $^{\text{S}\alpha}$ -OS say \mathfrak{D} , such that $\mathcal{Q} \sqsubseteq \mathfrak{D} \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}) \sqsubseteq \mathfrak{B}$.
- (ii) \mathcal{S} is a $\text{NC}^{\text{S}\alpha^*}$ -normal space if and only if for every NC $^{\text{S}\alpha}$ -CS \mathcal{Q} and every NC $^{\text{S}\alpha}$ -OS \mathfrak{B} containing \mathcal{Q} , there exists NC $^\alpha$ -OS say \mathfrak{D} , such that $\mathcal{Q} \sqsubseteq \mathfrak{D} \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}) \sqsubseteq \mathfrak{B}$.
- (iii) \mathcal{S} is a $\text{NC}^{\text{S}\alpha^{**}}$ -normal space if and only if for every NC $^{\text{S}\alpha}$ -CS \mathcal{Q} and every NC $^{\text{S}\alpha}$ -OS \mathfrak{B} containing \mathcal{Q} , there exists NC-OS say \mathfrak{D} , such that $\mathcal{Q} \sqsubseteq \mathfrak{D} \sqsubseteq \text{NCcl}(\mathfrak{D}) \sqsubseteq \mathfrak{B}$.

Proof:

(i) \Rightarrow Let \mathcal{S} be a $\text{NC}^{\text{S}\alpha}$ -normal space. Let $\mathcal{Q} \sqsubseteq \mathfrak{B}$, where \mathcal{Q} is a NC-CS and \mathfrak{B} is a NC-OS $\Rightarrow \mathcal{Q} \cap \mathfrak{B}^c = \emptyset_N$, where \mathfrak{B}^c is a NC-CS. Hence there exist two NC $^{\text{S}\alpha}$ -OSs $\mathfrak{D}_1, \mathfrak{D}_2$ such that $\mathcal{Q} \sqsubseteq \mathfrak{D}_1$ and $\mathfrak{B}^c \sqsubseteq \mathfrak{D}_2$ and $\mathfrak{D}_1 \cap \mathfrak{D}_2 = \emptyset_N$ (since \mathcal{S} is a $\text{NC}^{\text{S}\alpha}$ -normal space). Therefore $\mathcal{Q} \sqsubseteq \mathfrak{D}_1 \sqsubseteq \mathfrak{D}_2^c \sqsubseteq \mathfrak{B} \Rightarrow \text{NC}^{\text{S}\alpha} \text{cl}(\mathcal{Q}) \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}_1) \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}_2^c) \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{B})$. Hence $\mathcal{Q} \sqsubseteq \mathfrak{D}_1 \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}_1) \sqsubseteq \mathfrak{B}$, where \mathfrak{D}_1 is a NC $^{\text{S}\alpha}$ -OS in \mathcal{S} .
 \Leftarrow To prove \mathcal{S} is a $\text{NC}^{\text{S}\alpha}$ -normal space. Let \mathcal{Q}_1 and \mathcal{Q}_2 be NC-CSs in \mathcal{S} , such that $\mathcal{Q}_1 \cap \mathcal{Q}_2 = \emptyset_N$. Hence $\mathcal{Q}_1 \sqsubseteq \mathcal{Q}_2^c$, where \mathcal{Q}_2^c is a NC-OS. Then there exists a NC $^{\text{S}\alpha}$ -OS \mathfrak{D} such that $\mathcal{Q}_1 \sqsubseteq \mathfrak{D} \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}) \sqsubseteq \mathcal{Q}_2^c$ (by hypothesis). Hence $\mathcal{Q}_1 \sqsubseteq \mathfrak{D}$, $\mathcal{Q}_2 \sqsubseteq (\text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}))^c$. On the other hand $\text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}) \cap (\text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}))^c = \emptyset_N$. Hence $\mathfrak{D} \cap (\text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D}))^c = \emptyset_N$ (since $\mathfrak{D} \sqsubseteq \text{NC}^{\text{S}\alpha} \text{cl}(\mathfrak{D})$). Therefore \mathcal{S} is a $\text{NC}^{\text{S}\alpha}$ -normal space. The proofs (ii), (iii) are evident for others. ■

Remark 5.5:

- (i) NC^α -normal property is a $\text{NC}^{\alpha^{**}}$ -topological property.
- (ii) NC^{α^*} -normal property is a NC^{α^*} -topological property.
- (iii) $\text{NC}^{\alpha^{**}}$ -normal property is a $\text{NC}^{\alpha^{**}}$ -topological property.
- (iv) $\text{NC}^{\text{S}\alpha}$ -normal property is a $\text{NC}^{\text{S}\alpha^*}$ -topological property.
- (v) $\text{NC}^{\text{S}\alpha^*}$ -normal property is a $\text{NC}^{\text{S}\alpha^*}$ -topological property.

(vi) $NC^{S\alpha^{**}}$ -normal property is a $NC^{S\alpha^{**}}$ -topological property.

Proposition 5.6:

- (i) If $\mathcal{S} \times \mathcal{J}$ is a $NC^{\alpha^{**}}$ -normal space, then both \mathcal{S} and \mathcal{J} are $NC^{\alpha^{**}}$ -normal spaces.
- (ii) If $\mathcal{S} \times \mathcal{J}$ is a $NC^{S\alpha^{**}}$ -normal space, then both \mathcal{S} and \mathcal{J} are $NC^{S\alpha^{**}}$ -normal spaces.

Proof:

(i) Suppose that $\mathcal{S} \times \mathcal{J}$ is a $NC^{\alpha^{**}}$ -normal space, to prove that \mathcal{S} and \mathcal{J} are $NC^{\alpha^{**}}$ -normal spaces.

Let \mathfrak{B}_1 and \mathfrak{B}_2 be two NC^{α} -OSs in \mathcal{S} and \mathcal{J} respectively, such that $Q_1 \sqsubseteq \mathfrak{B}_1$ and $Q_2 \sqsubseteq \mathfrak{B}_2$, where Q_1 and Q_2 are NC^{α} -CSs in \mathcal{S} and \mathcal{J} respectively. Hence $Q_1 \times Q_2 \sqsubseteq \mathfrak{B}_1 \times \mathfrak{B}_2$ where $Q_1 \times Q_2$ is a NC^{α} -CS and $\mathfrak{B}_1 \times \mathfrak{B}_2$ is a NC^{α} -OS in $\mathcal{S} \times \mathcal{J}$ (by theorem (2.7) and corollary (2.8)). But $\mathcal{S} \times \mathcal{J}$ is a $NC^{\alpha^{**}}$ -normal space. Then there exists a NC -OS say \mathfrak{D} in $\mathcal{S} \times \mathcal{J}$ such that $Q_1 \times Q_2 \sqsubseteq \mathfrak{D} \sqsubseteq NCcl(\mathfrak{D}) \sqsubseteq \mathfrak{B}_1 \times \mathfrak{B}_2$. Then there exist NC -OSs \mathcal{U}_1 and \mathcal{U}_2 in $\mathcal{S} \times \mathcal{J}$ such that $Q_1 \times Q_2 \sqsubseteq \mathcal{U}_1 \times \mathcal{U}_2 \sqsubseteq NCcl(\mathcal{U}_1 \times \mathcal{U}_2) = NCcl(\mathcal{U}_1) \times NCcl(\mathcal{U}_2) \sqsubseteq \mathfrak{B}_1 \times \mathfrak{B}_2$. Hence $Q_1 \sqsubseteq \mathcal{U}_1 \sqsubseteq NCcl(\mathcal{U}_1) \sqsubseteq \mathfrak{B}_1 \Rightarrow \mathcal{S}$ is a $NC^{\alpha^{**}}$ -normal space. Also, $Q_2 \sqsubseteq \mathcal{U}_2 \sqsubseteq NCcl(\mathcal{U}_2) \sqsubseteq \mathfrak{B}_2 \Rightarrow \mathcal{J}$ is a $NC^{\alpha^{**}}$ -normal space. The proof (ii) is evident for others. ■

Definition 5.7:

Let (\mathcal{S}, ζ) be a NCTS, then \mathcal{S} is said to be:

- (i) NC^{α} - T_4 -space if \mathcal{S} is a NC^{α} - T_1 -space and NC^{α} -normal space.
- (ii) NC^{α^*} - T_4 -space if \mathcal{S} is NC^{α} - T_1 -space and NC^{α^*} -normal space.
- (iii) $NC^{\alpha^{**}}$ - T_4 -space if \mathcal{S} is NC^{α} - T_1 -space and $NC^{\alpha^{**}}$ -normal space.

Definition 5.8:

Let (\mathcal{S}, ζ) be a NCTS, then \mathcal{S} is said to be:

- (i) $NC^{S\alpha}$ - T_4 -space if \mathcal{S} is a $NC^{S\alpha}$ - T_1 -space and $NC^{S\alpha}$ -normal space.
- (ii) $NC^{S\alpha^*}$ - T_4 -space if \mathcal{S} is $NC^{S\alpha}$ - T_1 -space and $NC^{S\alpha^*}$ -normal space.
- (iii) $NC^{S\alpha^{**}}$ - T_4 -space if \mathcal{S} is $NC^{S\alpha}$ - T_1 -space and $NC^{S\alpha^{**}}$ -normal space.

Remark 5.9:


- (i) NC^{α} - T_4 ($NC^{S\alpha}$ - T_4 respectively) property is a $NC^{\alpha^{**}}$ ($NC^{S\alpha^{**}}$ respectively) topological property.
- (ii) NC^{α^*} - T_4 ($NC^{S\alpha^*}$ - T_4 respectively) property is a NC^{α^*} ($NC^{S\alpha^*}$ respectively) topological property.
- (iii) $NC^{\alpha^{**}}$ - T_4 ($NC^{S\alpha^{**}}$ - T_4 respectively) property is a $NC^{\alpha^{**}}$ ($NC^{S\alpha^{**}}$ respectively) topological property.

Remark 5.10:

- (i) Every NC - T_4 -space is a NC^{α} - T_4 -space and $NC^{S\alpha}$ - T_4 -space.
- (ii) Every NC^{α} - T_4 -space is a $NC^{S\alpha}$ - T_4 -space.
- (iii) Every $NC^{\alpha^{**}}$ - T_4 -space is a NC^{α^*} - T_4 -space and $NC^{S\alpha}$ - T_4 -space.
- (iv) Every NC^{α^*} - T_4 -space ($NC^{S\alpha^*}$ - T_4 -space respectively) is a NC^{α^*} - T_3 -space ($NC^{S\alpha^*}$ - T_3 -space, respectively).
- (v) Every $NC^{\alpha^{**}}$ - T_4 -space ($NC^{S\alpha^{**}}$ - T_4 -space respectively) is a $NC^{\alpha^{**}}$ - T_3 -space ($NC^{S\alpha^{**}}$ - T_3 -space, respectively).

Remark 5.11:

The following diagram explains the relationships between usual NC -separation axioms, NC^{α} -separation axioms and $NC^{S\alpha}$ -separation axioms:



Also, we have the following diagram:




Fig. 5.2

6. Conclusions

We have provided some new concepts of weakly neutrosophic crisp separation axioms. Some characterizations have been provided to illustrate how far topological structures are conserved by the new neutrosophic crisp notion defined. Furthermore, some new concepts of weakly neutrosophic crisp regularity are also studied. The study demonstrated some new concepts of weakly neutrosophic crisp normality and proved some of their related attributes.

Funding: There is no external grant for this work.

Acknowledgments: The authors are appreciative to the Referees for their constructive comments.

Conflicts of Interest: There are no conflicts of interest declared by the authors.

References

- [1] A. A. Salama, F. Smarandache and V. Kroumov, Neutrosophic crisp sets and neutrosophic crisp topological spaces. *Neutrosophic Sets and Systems*, 2(2014), 25-30.
- [2] A. A. Salama, Basic structure of some classes of neutrosophic crisp nearly open sets & possible application to GIS topology. *Neutrosophic Sets and Systems*, 7(2015), 18-22.
- [3] A. H. M. Al-Obaidi and Q. H. Imran, On new types of weakly neutrosophic crisp open mappings. *Iraqi Journal of Science*, vol.62, No.8, (2021), 2660-2666.
- [4] A. H. M. Al-Obaidi, Q. H. Imran and M. M. Abdulkadhim, On new types of weakly neutrosophic crisp closed functions. *Communicated*.
- [5] Md. Hanif PAGE and Q. H. Imran, Neutrosophic generalized homeomorphism. *Neutrosophic Sets and Systems*, 35(2020), 340-346.
- [6] Q. H. Imran, F. Smarandache, R. K. Al-Hamido and R. Dhavaseelan, On neutrosophic semi- α -open sets. *Neutrosophic Sets and Systems*, 18(2017), 37-42.
- [7] Q. H. Imran, R. K. Al-Hamido and A. H. M. Al-Obaidi, On new types of weakly Neutrosophic crisp continuity. *Neutrosophic Sets and Systems*, 38(2020), 179-187.
- [8] Q. H. Imran, K. S. Tanak and A. H. M. Al-Obaidi, On new concepts of neutrosophic crisp open sets. *Journal of Interdisciplinary Mathematics*, 25(2)(2022), 563-572.
- [9] R. Dhavaseelan, R. Narmada Devi, S. Jafari and Q. H. Imran, Neutrosophic α^m -continuity. *Neutrosophic Sets and Systems*, 27(2019), 171-179.
- [10] R. K. Al-Hamido, Q. H. Imran, K. A. Alghurabi and T. Gharibah, On neutrosophic crisp semi- α -closed sets. *Neutrosophic Sets and Systems*, 21(2018), 28-35.
- [11] A. B. Al-Nafee, R. K. Al-Hamido and F. Smarandache, Separation axioms in neutrosophic crisp topological spaces. *Neutrosophic Sets and Systems*, 25(2019).
- [12] R. K. Al-Hamido, Luai Salha and Taleb Gharibah, Neutrosophic crisp semi separation axioms in neutrosophic crisp topological spaces. *International Journal of Neutrosophic Science*, 6(1)(2020), 32-40.
- [13] A. A. Salama, I. M. Hanafy, H. Elghawalby and M. S. Dabash, Some GIS Topological Concepts via Neutrosophic Crisp Set Theory. *New Trends in Neutrosophic Theory and Applications*, 2016.
- [14] W. Al-Omeri, Neutrosophic crisp sets via neutrosophic crisp topological spaces NCTS. *Neutrosophic Sets and Systems*, 13(2016), 96-104.

Received: Feb 5, 2022. Accepted: April 22, 2022.