

10-5-2022

## Dombi Aggregation Operators of Linguistic Neutrosophic Numbers for Multiple Attribute Group Decision-Making Problems in Landslide Treatment Schemes

Yingying Zhang

Rui Yong

Jun Ye

Zhen Zhong

Shuang Zheng

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Zhang, Yingying; Rui Yong; Jun Ye; Zhen Zhong; and Shuang Zheng. "Dombi Aggregation Operators of Linguistic Neutrosophic Numbers for Multiple Attribute Group Decision-Making Problems in Landslide Treatment Schemes." *Neutrosophic Sets and Systems* 51, 1 (2022). [https://digitalrepository.unm.edu/nss\\_journal/vol51/iss1/45](https://digitalrepository.unm.edu/nss_journal/vol51/iss1/45)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).



# Dombi Aggregation Operators of Linguistic Neutrosophic Numbers for Multiple Attribute Group Decision-Making Problems in Landslide Treatment Schemes

Yingying Zhang<sup>1</sup>, Rui Yong<sup>\*1</sup>, Jun Ye<sup>1</sup>, Zhen Zhong<sup>2</sup>, Shuang Zheng<sup>2</sup>

<sup>1</sup> School of Civil and Environmental Engineering, Ningbo University, Ningbo, Zhejiang, 315211, P.R. China;

<sup>2</sup> Key Laboratory of Rock Mechanics and Geohazards of Zhejiang Province, Shaoxing University, Shaoxing, Zhejiang, 32200, P.R. China

\* Correspondence: yongrui@nbu.edu.cn


**Abstract:** The landslide disaster caused huge losses to human lives and property, and the research on the selection of landslide treatment schemes has attracted global attention. Fuzzy multi-attribute decision-making methods are widely used for selecting the slope treatment schemes in engineering practice. But they do not take into account human linguistic arguments in the linguistic decision-making environment, which usually contains incomplete and uncertain information, and still lack a qualitative evaluation method. To deal with multiple attribute group decision-making (MAGDM) problems of landslide treatment schemes in the linguistic neutrosophic environment, a linguistic neutrosophic number Dombi weighted arithmetic averaging (LNNDWAA) operator and a linguistic neutrosophic number Dombi weighted geometric averaging (LNNDWGA) operator are developed to aggregate linguistic neutrosophic information. Then, a new MAGDM method using these aggregation operators is proposed in view of the Dombi operational flexibility. Finally, the proposed method is applied to select the optimal landslide treatment scheme under the linguistic neutrosophic environment. The results show that this method can effectively solve the decision-making problem of landslide treatment schemes and make the decision result more reasonable and flexible than other existing methods.

**Keywords:** landslide treatment scheme; multiple attribute decision; linguistic neutrosophic number; Dombi operation

## 1. Introduction

Landslides have caused immeasurable economic losses to human society. For example, China has an average of almost 30,000 landslides, rock falls, and debris flows every year, many of which have caused catastrophic disasters. On average, nearly 800 people are killed each year, and the direct economic loss exceeds 1 billion US dollars. In 2019, a total of 6,181 geological disasters occurred nationwide (Figure 1), where slope failure accounted for 68.27%, accounting for the vast majority [1]. Therefore, great attention has been paid to the prevention and treatment of landslides. At present, there are many prevention and control plans for slopes. To choose the most reasonable plan, some decision-making (DM) methods are needed. In recent years, research on DM methods of slope treatment schemes has received increasing attention. These DM methods include the empirical discriminant method [2], the analytic hierarchy process [3], the fuzzy multi-attribute DM method [4], the subjective and objective weighting method [5], and so on. However, most of these

methods are based on the analysis or subjective judgment of expert experience, which leads to unreasonable, uneconomical, and sometimes even huge waste in the final choice of the treatment schemes. Fuzzy multi-attribute DM methods have been widely used because they can deal with fuzzy evaluation and DM problems. But they does not take into account human linguistic arguments in the DM issue of the treatment schemes, which usually contains incomplete and uncertain information.



**Figure 1.** Types of geological disasters in 2019

Due to the indeterminacy and ambiguity of human cognition of objective things and the intricacy of multi-attribute group decision making (MAGDM) environment, linguistic variables can more effectively describe decision information than numerical values [6,7,8]. Therefore, to improve the DM effectiveness, many researchers performed extensive studies on DM challenges in linguistic environments. Zadeh [7] initially proposed the concept of linguistic variables (LVs), which can employ words or sentences to represent qualitative information. In order to solve DM problems with linguistic information, Herrera et al. [9, 10] created a technique for linguistic decision analysis. Later, Xu [11,12,13] developed linguistic aggregation operators and goal programming models to handle DM problems. To tackle incompleteness and ambiguity in DM situations more effectively, Merigó et al. [14,15,16] proposed some linguistic aggregation operators for the aggregation of LVs. Xu [17,18] proposed uncertain LVs given by interval values. Then, some scholars developed various aggregation operators of uncertain LVs for the MAGDM problems with uncertain linguistic information [19–23]. Chen et al. [24] proposed the concept of linguistic intuitionistic fuzzy numbers (LIFNs), which enables the direct description of real and false linguistic information using linguistic items. Liu et al. [25,26] put forward some LIFN aggregation operators for multi-attribute DM. However, LIFN cannot express uncertain and inconsistent linguistic information in DM problems. But the neutrosophic numbers (NNs) [27–30] and neutrosophic sets [28–30] proposed by Smarandache make up for the above shortcomings. Some scholars put forward new concepts focusing on the combination of neutrosophic set and linguistic set. Subsequently, Fang and Ye [8] introduced the linguistic neutrosophic number (LNN) which includes three independent LVs for describe true, false, and uncertain linguistic information. They also introduced the LNN weighted geometric and LNN weighted arithmetic averaging operators to handle MAGDM problems containing LNN information. However, this MAGDM method [8] can be better applicable to the expression and processing of inconsistent and uncertain linguistic information in DM problems. After that, some LNN aggregation operators were successively proposed, such as LNN normalized weighted geometric Bonferroni mean (LNNWGBM) and LNN normalized weighted Bonferroni

mean (LNNWBM) operators [31] and linguistic neutrosophic power weighted Heronian aggregation (LNPWHA) operators [32]. These aggregation operators can effectively deal with linguistic DM problems with inconsistent and uncertain linguistic information. Furthermore, Shi and Ye developed three correlation coefficients of LNNs [33] and two cosine similarity measures of LNNs [34] for MAGDM problems with LNN information. Cui and Ye [35] defined the concept of hesitant linguistic neutrosophic number (HLNN) sets and introduced the normalized generalized distance and similarity measures of HLNNs for DM problems with HLNN information.

In 1982, Dombi [36] developed Dombi t-conorm and Dombi t-norm operations, which contain the advantage of changeability by adjusting a parameter value. Hence, Liu et al. [37] introduced the Dombi operations of intuitionistic fuzzy sets (IFSs) and proposed some Dombi aggregation operators for the MAGDM problem with intuitionistic fuzzy information. Ye and Chen [38] introduced the Dombi operations of single-valued neutrosophic numbers (SVNNs), then presented the single-valued neutrosophic Dombi weighted geometric average (SVNDWGA) operator and the single-valued neutrosophic Dombi weighted arithmetic average (SVNDWAA) operator to handle DM problems with LNNs. Ye and Lu [39] extended the Dombi operations to the environment of linguistic cubic variables (LCVs) and developed the linguistic cubic variable Dombi weighted geometric average (LCVDWGA) operator and linguistic cubic variable Dombi weighted arithmetic average (LCVDWAA) operator for MAGDM problems. However, in the available research, the Dombi operations have not yet been extended to LNNs. Therefore, the main goals of this study are (1) to propose some Dombi operations of LNNs, (2) to propose the LNN Dombi weighted geometric averaging (LNNDWGA) and LNN Dombi weighted arithmetic averaging (LNNDWAA) operators, (3) to develop a DM method based on the LNNDWAA or LNNDWGA operator for performing MAGDM problems in the LNN information environment, and (4) to validate the viability of this method through a case study.

The following sections constitute the rest of this paper. Section 2 introduces some preliminaries. In Section 3, we define the Dombi operations of LNNs and propose the LNNDWAA and LNNDWGA operators and their properties. Section 4 introduces a new MAGDM method using the LNNDWAA or LNNDWGA operator. In Section 5, the application of the proposed method is demonstrated by an application example and then a comparative analysis is given to show its superiority over existing approaches. Section 6 gives the conclusions of this article.

## 2. Preliminaries

### 2.1 Several Concepts of LNNs

**Definition 1** [8]. Suppose that  $Fr^{Ro} = \{Fr_0^{Ro}, Fr_1^{Ro}, \dots, Fr_\Phi^{Ro}\}$  is a set of linguistic terms with an odd cardinality  $\Phi + 1$ . Then, LNN is defined as  $N = \langle Fr_x^{Ro}, Fr_y^{Ro}, Fr_z^{Ro} \rangle$  for  $Fr_x^{Ro}, Fr_y^{Ro}, Fr_z^{Ro} \in Fr^{Ro}$  and  $x, y, z \in [0, \Phi]$ , where  $Fr_x^{Ro}$ ,  $Fr_y^{Ro}$  and  $Fr_z^{Ro}$  independently represent truth, uncertainty, and falsity LVs, respectively.

**Definition 2** [8]. Set  $N = \langle Fr_x^{Ro}, Fr_y^{Ro}, Fr_z^{Ro} \rangle$  as LNN in  $Fr^{Ro}$ . The score and accuracy functions of  $N$  are determined by the following equations:

$$U(N) = (2\Phi + x - y - z) / (3\Phi) \text{ for } U(N) \in [0, 1], \quad (1)$$

$$V(N) = (x - z) / \Phi \text{ for } V(N) \in [-1, 1]. \quad (2)$$

**Definition 3** [8]. Let  $N_1 = \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle$  and  $N_2 = \langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \rangle$  be two LNNs in

$Fr^{Ro}$ , and they imply the following ranking relations:

- (1) When  $U(N_1) > U(N_2) \Rightarrow N_1 \succ N_2$ ;

- (2) When  $U(N_1) < U(N_2) \Rightarrow N_1 \prec N_2$ ;
- (3) When  $V(N_1) = V(N_2)$  and  $U(N_1) = U(N_2) \Rightarrow N_1 = N_2$ ;
- (4) When  $V(N_1) < V(N_2)$  and  $U(N_1) = U(N_2) \Rightarrow N_1 \prec N_2$ ;
- (5) When  $V(N_1) > V(N_2)$  and  $U(N_1) = U(N_2) \Rightarrow N_1 \succ N_2$ .

**Definition 4** [8]. Let  $N_1 = \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle$  and  $N_2 = \langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \rangle$  be two LNNs in  $Fr^{Ro}$ , and  $\lambda$  is a positive real number ( $\lambda > 0$ ). Their operational laws are introduced as follows:

- (1)  $N_1 \oplus N_2 = \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle \oplus \langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \rangle = \left\langle Fr_{x_1+x_2-\frac{x_1x_2}{\Phi}}^{Ro}, Fr_{\frac{y_1y_2}{\Phi}}^{Ro}, Fr_{\frac{z_1z_2}{\Phi}}^{Ro} \right\rangle$ ;
- (2)  $N_1 \otimes N_2 = \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle \otimes \langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \rangle = \left\langle Fr_{\frac{x_1x_2}{\Phi}}^{Ro}, Fr_{y_1+y_2-\frac{y_1y_2}{\Phi}}^{Ro}, Fr_{z_1+z_2-\frac{z_1z_2}{\Phi}}^{Ro} \right\rangle$ ;
- (3)  $\lambda N_1 = \lambda \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle = \left\langle Fr_{\Phi-\Phi\left(1-\frac{x_1}{\Phi}\right)^\lambda}^{Ro}, Fr_{\Phi\left(\frac{y_1}{\Phi}\right)^\lambda}^{Ro}, Fr_{\Phi\left(\frac{z_1}{\Phi}\right)^\lambda}^{Ro} \right\rangle$ ;
- (4)  $N_1^\lambda = \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle^\lambda = \left\langle Fr_{\Phi\left(\frac{x_1}{\Phi}\right)^\lambda}^{Ro}, Fr_{\Phi-\Phi\left(1-\frac{y_1}{\Phi}\right)^\lambda}^{Ro}, Fr_{\Phi-\Phi\left(1-\frac{z_1}{\Phi}\right)^\lambda}^{Ro} \right\rangle$ .

### 2.2 Weighted Aggregation Operators of LNNs

**Definition 5** [8]. Set  $N_g = \langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \rangle$  ( $g = 1, 2, \dots, h$ ) as an assemblage of LNNs in  $Fr^{Ro}$ . The LNNWAA operator is defined below:

$$LNNWAA(N_1, N_2, \dots, N_h) = \sum_{g=1}^h \gamma_g N_g, \tag{3}$$

where  $\gamma_g$  is the weight of  $N_g$  ( $g = 1, 2, \dots, h$ ) for  $0 \leq \gamma_g \leq 1$  and  $\sum_{g=1}^h \gamma_g = 1$ .

**Theorem 1** [8]. Let  $N_g = \langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \rangle$  ( $g = 1, 2, \dots, h$ ) as an assemblage of LNN in  $Fr^{Ro}$ , then the aggregation result is obtained based on the following aggregation equation:

$$LNNWAA(N_1, N_2, \dots, N_h) = \sum_{g=1}^h \gamma_g N_g = \left\langle Fr_{\Phi-\Phi\prod_{g=1}^h\left(1-\frac{x_g}{\Phi}\right)^{\gamma_g}}^{Ro}, Fr_{\Phi\prod_{g=1}^h\left(\frac{y_g}{\Phi}\right)^{\gamma_g}}^{Ro}, Fr_{\Phi\prod_{g=1}^h\left(\frac{z_g}{\Phi}\right)^{\gamma_g}}^{Ro} \right\rangle. \tag{4}$$

**Definition 6** [8]. Suppose that  $N_g = \langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \rangle$  ( $g = 1, 2, \dots, h$ ) is a group of LNNs in  $Fr^{Ro}$ , the LNNWGA operator is defined by

$$LNNWGA(N_1, N_2, \dots, N_h) = \prod_{g=1}^h N_g^{\gamma_g}, \tag{5}$$

where  $\gamma_g$  is the weight of  $N_g$  ( $g = 1, 2, \dots, h$ ) for  $0 \leq \gamma_g \leq 1$  and  $\sum_{g=1}^h \gamma_g = 1$ .

**Theorem 2** [8]. Let  $N_g = \langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \rangle$  ( $g = 1, 2, \dots, h$ ) as an assemblage of linguistic neutrosophic numbers in  $Fr^{Ro}$ , then the result of aggregation is obtained based on the following aggregation equation:

$$LNNWGA(N_1, N_2, \dots, N_h) = \prod_{g=1}^h N_g^{\gamma_g} = \left\langle Fr_{\Phi \prod_{g=1}^h \left(\frac{x_g}{\Phi}\right)^{\gamma_g}}^{Ro}, Fr_{\Phi - \Phi \prod_{g=1}^h \left(1 - \frac{y_g}{\Phi}\right)^{\gamma_g}}^{Ro}, Fr_{\Phi - \Phi \prod_{g=1}^h \left(1 - \frac{z_g}{\Phi}\right)^{\gamma_g}}^{Ro} \right\rangle. \quad (6)$$

### 3. Dombi Operations of LNNs

The Dombi operations contain the advantage of flexible aggregations by modifying the value of the parameter. In 1982, Dombi [36] proposed the Dombi T-norm and T-conorm operations for the first time. Although many researchers have introduced Dombi operations in various linguistic decision-making environments and decision-making methods [37–43], the Dombi operations have not yet expanded to LNNs. Therefore, this section proposes the Dombi T-norm and T-conorm operations of LNNs, then presents the LNNDWAA and LNNDWGA operators and their properties.

#### 3.1 Dombi Operational Laws of LNNs

**Definition 7** [36]. For any two real-values  $Th$  and  $Tj$ , the Dombi T-norm and T-conorm operations between  $Th$  and  $Tj$  are defined below:

$$O_D(Th, Tj) = \frac{1}{1 + \left\{ \left( \frac{1-Th}{Th} \right)^\rho + \left( \frac{1-Tj}{Tj} \right)^\rho \right\}^{1/\rho}}, \quad (7)$$

$$O_D^c(Th, Tj) = 1 - \frac{1}{1 + \left\{ \left( \frac{Th}{1-Th} \right)^\rho + \left( \frac{Tj}{1-Tj} \right)^\rho \right\}^{1/\rho}}, \quad (8)$$

where the parameter  $\rho \geq 1$  and  $(Th, Tj) \in [0, 1] \times [0, 1]$ .

**Definition 8.** Assume that  $N_1 = \langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \rangle$  and  $N_2 = \langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \rangle$  are two LNNs,  $\lambda > 0$ , and  $\rho > 0$ . Then, the Dombi T-norm and T-conorm operational laws of LNNs are expressed below:

$$\begin{aligned}
 N_1 \oplus N_2 &= \left\langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \right\rangle \oplus \left\langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \right\rangle \\
 &= \left( Fr_{\Phi \times \frac{1}{1 + \left\{ \left( \frac{x_1/\Phi}{1-x_1/\Phi} \right)^\rho + \left( \frac{x_2/\Phi}{1-x_2/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi \times \frac{1}{1 + \left\{ \left( \frac{1-y_1/\Phi}{y_1/\Phi} \right)^\rho + \left( \frac{1-y_2/\Phi}{y_2/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi \times \frac{1}{1 + \left\{ \left( \frac{1-z_1/\Phi}{z_1/\Phi} \right)^\rho + \left( \frac{1-z_2/\Phi}{z_2/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right), \quad (9) \\
 &= \left( Fr_{\frac{\Phi}{1 + \left\{ \left( \frac{x_1}{\Phi-x_1} \right)^\rho + \left( \frac{x_2}{\Phi-x_2} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \left( \frac{\Phi-y_1}{y_1} \right)^\rho + \left( \frac{\Phi-y_2}{y_2} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \left( \frac{\Phi-z_1}{z_1} \right)^\rho + \left( \frac{\Phi-z_2}{z_2} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right)
 \end{aligned}$$

$$\begin{aligned}
 N_1 \otimes N_2 &= \left\langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \right\rangle \otimes \left\langle Fr_{x_2}^{Ro}, Fr_{y_2}^{Ro}, Fr_{z_2}^{Ro} \right\rangle \\
 &= \left( Fr_{\Phi \times \frac{1}{1 + \left\{ \left( \frac{1-x_1/\Phi}{x_1/\Phi} \right)^\rho + \left( \frac{1-x_2/\Phi}{x_2/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi \times \frac{1}{1 + \left\{ \left( \frac{y_1/\Phi}{1-y_1/\Phi} \right)^\rho + \left( \frac{y_2/\Phi}{1-y_2/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi \times \frac{1}{1 + \left\{ \left( \frac{z_1/\Phi}{1-z_1/\Phi} \right)^\rho + \left( \frac{z_2/\Phi}{1-z_2/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right), \quad (10) \\
 &= \left( Fr_{\frac{\Phi}{1 + \left\{ \left( \frac{\Phi-x_1}{x_1} \right)^\rho + \left( \frac{\Phi-x_2}{x_2} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \left( \frac{y_1}{\Phi-y_1} \right)^\rho + \left( \frac{y_2}{\Phi-y_2} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \left( \frac{z_1}{\Phi-z_1} \right)^\rho + \left( \frac{z_2}{\Phi-z_2} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right)
 \end{aligned}$$

$$\begin{aligned}
 \lambda N_1 &= \lambda \left\langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \right\rangle \\
 &= \left( Fr_{\Phi \times \frac{1}{1 + \left\{ \lambda \left( \frac{x_1/\Phi}{1-x_1/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \lambda \left( \frac{1-y_1/\Phi}{y_1/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \lambda \left( \frac{1-z_1/\Phi}{z_1/\Phi} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right), \quad (11) \\
 &= \left( Fr_{\frac{\Phi}{1 + \left\{ \lambda \left( \frac{x_1}{\Phi-x_1} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \lambda \left( \frac{\Phi-y_1}{y_1} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\frac{\Phi}{1 + \left\{ \lambda \left( \frac{\Phi-z_1}{z_1} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right)
 \end{aligned}$$

$$\begin{aligned}
 N_1^\lambda &= \left\langle Fr_{x_1}^{Ro}, Fr_{y_1}^{Ro}, Fr_{z_1}^{Ro} \right\rangle^\lambda \\
 &= \left( \left( \frac{Fr^{Ro}}{\frac{\Phi}{1+\left\{\lambda\left(\frac{1-x_1/\Phi}{x_1/\Phi}\right)^\rho\right\}^{1/\rho}}}, Fr^{Ro}, Fr^{Ro} \right), \left( \frac{Fr^{Ro}}{\frac{\Phi}{1+\left\{\lambda\left(\frac{y_1/\Phi}{1-y_1/\Phi}\right)^\rho\right\}^{1/\rho}}}, Fr^{Ro}, Fr^{Ro} \right), \left( \frac{Fr^{Ro}}{\frac{\Phi}{1+\left\{\lambda\left(\frac{z_1/\Phi}{1-z_1/\Phi}\right)^\rho\right\}^{1/\rho}}}, Fr^{Ro}, Fr^{Ro} \right) \right) \\
 &= \left( \frac{Fr^{Ro}}{\frac{\Phi}{1+\left\{\lambda\left(\frac{\Phi-x_1}{x_1}\right)^\rho\right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\frac{\Phi}{1+\left\{\lambda\left(\frac{y_1}{\Phi-y_1}\right)^\rho\right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\frac{\Phi}{1+\left\{\lambda\left(\frac{z_1}{\Phi-z_1}\right)^\rho\right\}^{1/\rho}}} \right)
 \end{aligned} \tag{12}$$

However, the operational results of the equations (9)–(12) are also LNNs.

**Example 1.** Let  $N_1 = \langle Fr_2^{Ro}, Fr_1^{Ro}, Fr_3^{Ro} \rangle$  and  $N_2 = \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_4^{Ro} \rangle$  in  $Fr^{Ro} = \{Fr_0^{Ro}, Fr_1^{Ro}, \dots, Fr_6^{Ro}\}$  be two LNNs,  $\lambda = 0.5, \rho = 1$ . Based on the equations (9)–(12), we have the following operational results:

$$\begin{aligned}
 N_1 \oplus N_2 &= \left\langle Fr_2^{Ro}, Fr_1^{Ro}, Fr_3^{Ro} \right\rangle \oplus \left\langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_4^{Ro} \right\rangle \\
 &= \left( \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{\left(\frac{2}{6-2}\right)^1+\left(\frac{3}{6-3}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{\left(\frac{6-1}{1}\right)^1+\left(\frac{6-2}{2}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{\left(\frac{6-3}{3}\right)^1+\left(\frac{6-4}{4}\right)^1\right\}^{1/1}}}} \right) = \left\langle Fr_{3.6000}^{Ro}, Fr_{0.7500}^{Ro}, Fr_{2.4000}^{Ro} \right\rangle'
 \end{aligned}$$

$$\begin{aligned}
 N_1 \otimes N_2 &= \left\langle Fr_2^{Ro}, Fr_1^{Ro}, Fr_3^{Ro} \right\rangle \otimes \left\langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_4^{Ro} \right\rangle \\
 &= \left( \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{\left(\frac{6-2}{2}\right)^1+\left(\frac{6-3}{3}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{\left(\frac{1}{6-1}\right)^1+\left(\frac{2}{6-2}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{\left(\frac{3}{6-3}\right)^1+\left(\frac{4}{6-4}\right)^1\right\}^{1/1}}}} \right) = \left\langle Fr_{1.5000}^{Ro}, Fr_{2.4706}^{Ro}, Fr_{4.5000}^{Ro} \right\rangle'
 \end{aligned}$$

$$\begin{aligned}
 \lambda N_1 &= \lambda \left\langle Fr_2^{Ro}, Fr_1^{Ro}, Fr_3^{Ro} \right\rangle \\
 &= \left( \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{0.5 \times \left(\frac{6-2}{6-2}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{0.5 \times \left(\frac{6-1}{1}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{0.5 \times \left(\frac{6-3}{3}\right)^1\right\}^{1/1}}}} \right) = \left\langle Fr_{1.2000}^{Ro}, Fr_{1.7143}^{Ro}, Fr_{4.0000}^{Ro} \right\rangle'
 \end{aligned}$$

$$\begin{aligned}
 N_1^\lambda &= \left\langle Fr_2^{Ro}, Fr_1^{Ro}, Fr_3^{Ro} \right\rangle^\lambda \\
 &= \left( \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{0.5 \times \left(\frac{6-2}{2}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{0.5 \times \left(\frac{1}{6-1}\right)^1\right\}^{1/1}}}}, \frac{Fr^{Ro}}{6-\frac{6}{1+\left\{0.5 \times \left(\frac{3}{6-3}\right)^1\right\}^{1/1}}}} \right) = \left\langle Fr_{3.0000}^{Ro}, Fr_{0.5455}^{Ro}, Fr_{2.0000}^{Ro} \right\rangle'
 \end{aligned}$$



3.2 Dombi Weighted Aggregation Operators of LNNs

**Definition 9.** Set  $N_g = \left\langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \right\rangle$  ( $g = 1, 2, \dots, h$ ) as a group of LNNs. Let  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_h)$

be the weight vector of  $N_g$  such that  $\gamma_g \in [0, 1]$  and  $\sum_{g=1}^h \gamma_g = 1$ . The LNNDWAA and LNNDWGA operators are proposed below:

$$LNNDWAA(N_1, N_2, \dots, N_h) = \bigoplus_{g=1}^h \gamma_g N_g, \tag{13}$$

$$LNNDWGA(N_1, N_2, \dots, N_h) = \bigotimes_{g=1}^h N_g^{\gamma_g}. \tag{14}$$

**Theorem 3.** Let  $N_g = \left\langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \right\rangle$  ( $g = 1, 2, \dots, h$ ) be an assemblage of LNNs and  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_h)$  be the weight vector of  $N_g$  such that  $\gamma_g \in [0, 1]$  and  $\sum_{g=1}^h \gamma_g = 1$ . The aggregated result of the LNNDWAA operator is still an LNN, which can be expressed by

$$LNNDWAA(N_1, N_2, \dots, N_h) = \bigoplus_{g=1}^h \gamma_g N_g = \left\langle \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{s=1}^h \gamma_s \left( \frac{x_s}{\Phi - x_s} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{s=1}^h \gamma_s \left( \frac{\Phi - y_s}{y_s} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{s=1}^h \gamma_s \left( \frac{\Phi - z_s}{z_s} \right)^\rho \right\}^{1/\rho}}} \right\rangle. \tag{15}$$

Theorem 3 is proved through mathematical induction below.

**Proof:**

(a) Let  $h = 2$ . Based on Definition 8 we can obtain

$$\begin{aligned} LNNDWAA(N_1, N_2) &= N_1 \oplus N_2 \\ &= \left\langle \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \gamma_1 \left( \frac{x_1}{\Phi - x_1} \right)^\rho + \gamma_2 \left( \frac{x_2}{\Phi - x_2} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \gamma_1 \left( \frac{\Phi - y_1}{y_1} \right)^\rho + \gamma_2 \left( \frac{\Phi - y_2}{y_2} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \gamma_1 \left( \frac{\Phi - z_1}{z_1} \right)^\rho + \gamma_2 \left( \frac{\Phi - z_2}{z_2} \right)^\rho \right\}^{1/\rho}}} \right\rangle \\ &= \left\langle \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^2 \gamma_g \left( \frac{x_g}{\Phi - x_g} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^2 \gamma_g \left( \frac{\Phi - y_g}{y_g} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^2 \gamma_g \left( \frac{\Phi - z_g}{z_g} \right)^\rho \right\}^{1/\rho}}} \right\rangle \end{aligned}$$

(b) If  $h = k$ , we can keep the following result from the equation (15):

$$LNNDWAA(N_1, N_2, \dots, N_k) = \bigoplus_{g=1}^k \gamma_g N_g$$

$$= \left\langle Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{x_g}{\Phi-x_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{\Phi-y_g}{y_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{\Phi-z_g}{z_g}\right)^\rho\right\}^{1/\rho}}}^{Ro} \right\rangle.$$

(c) Set  $h = k + 1$ . Based on Definition 9 and the equation (15), there exists the following result:

$$LNNDWAA(N_1, N_2, \dots, N_k, N_{k+1}) = \bigoplus_{g=1}^{k+1} \gamma_g N_g$$

$$= \left\langle Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{x_g}{\Phi-x_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{\Phi-y_g}{y_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{\Phi-z_g}{z_g}\right)^\rho\right\}^{1/\rho}}}^{Ro} \right\rangle \oplus \gamma_{k+1} N_{k+1}.$$

$$= \left\langle Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^{k+1} \gamma_g \left(\frac{x_g}{\Phi-x_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^{k+1} \gamma_g \left(\frac{\Phi-y_g}{y_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^{k+1} \gamma_g \left(\frac{\Phi-z_g}{z_g}\right)^\rho\right\}^{1/\rho}}}^{Ro} \right\rangle.$$

In terms of the above results, the equation (15) can hold for all  $h$ .

Then, the LNNDWAA operator has some properties:

(1) Reducibility: When  $\gamma = (1/h, 1/h, \dots, 1/h)$ , there exists

$$LNNDWAA(N_1, N_2, \dots, N_h) = \bigoplus_{g=1}^h \gamma_g N_g$$

$$= \left\langle Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^h \frac{1}{h} \left(\frac{x_g}{\Phi-x_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^h \frac{1}{h} \left(\frac{\Phi-y_g}{y_g}\right)^\rho\right\}^{1/\rho}}}^{Ro}, Fr_{\Phi-\frac{\Phi}{1+\left\{\sum_{g=1}^h \frac{1}{h} \left(\frac{\Phi-z_g}{z_g}\right)^\rho\right\}^{1/\rho}}}^{Ro} \right\rangle.$$

(2) Idempotency: Let all LNNs be  $N_g = \left\langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \right\rangle = N$  ( $g = 1, 2, \dots, h$ ). Then,

$$LNNDWAA(N_1, N_2, \dots, N_h) = N.$$

(3) Commutativity: Let the LNN sequence  $(N_1', N_2', \dots, N_h')$  be an arbitrary arrangement of  $(N_1, N_2, \dots, N_h)$ . Then, there is  $LNNDWAA(N_1', N_2', \dots, N_h') = LNNDWAA(N_1, N_2, \dots, N_h)$ .

(4) Boundedness: If the maximum and minimum LNNs are  $N_{\max} = \left\langle Fr_{\max(x_g)}^{Ro}, Fr_{\min(y_g)}^{Ro}, Fr_{\min(z_g)}^{Ro} \right\rangle$  and  $N_{\min} = \left\langle Fr_{\min(x_g)}^{Ro}, Fr_{\max(y_g)}^{Ro}, Fr_{\max(z_g)}^{Ro} \right\rangle$ , then  $N_{\min} \leq LNNDWAA(N_1, N_2, \dots, N_h) \leq N_{\max}$ .

**Proof:**

(1) Based on the equation (15), we can see that the property (1) is valid.

(2) Since  $N_g = \langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \rangle = N$  ( $g = 1, 2, \dots, h$ ), by the equation (15) we can obtain the following result:

$$\begin{aligned} LNNDWAA(N_1, N_2, \dots, N_h) &= \bigoplus_{g=1}^h \gamma_g N_g \\ &= \left\langle Fr_{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{x_g}{\Phi - x_g} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - y_g}{y_g} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - z_g}{z_g} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right\rangle \\ &= \left\langle Fr_{\Phi - \frac{\Phi}{1 + \left\{ \left( \frac{x}{\Phi - x} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi - \frac{\Phi}{1 + \left\{ \left( \frac{\Phi - y}{y} \right)^\rho \right\}^{1/\rho}}}^{Ro}, Fr_{\Phi - \frac{\Phi}{1 + \left\{ \left( \frac{\Phi - z}{z} \right)^\rho \right\}^{1/\rho}}}^{Ro} \right\rangle \\ &= \left\langle Fr_{\Phi - \frac{\Phi}{1 + \left( \frac{x}{\Phi - x} \right)}}^{Ro}, Fr_{\Phi - \frac{\Phi}{1 + \left( \frac{\Phi - y}{y} \right)}}^{Ro}, Fr_{\Phi - \frac{\Phi}{1 + \left( \frac{\Phi - z}{z} \right)}}^{Ro} \right\rangle = \langle Fr_x^{Ro}, Fr_y^{Ro}, Fr_z^{Ro} \rangle = N. \end{aligned}$$

Hence,  $LNNDWAA(N_1, N_2, \dots, N_h) = N$  holds.

(3) The property (3) is obvious.

(4) Since  $\min(x_g) \leq x_g \leq \max(x_g), \max(y_g) \leq y_g \leq \min(y_g), \max(z_g) \leq z_g \leq \min(z_g)$ , there are the following inequalities:

$$\begin{aligned} \Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\min(x_g)}{\Phi - \min(x_g)} \right)^\rho \right\}^{1/\rho}} &= \min(x_g) \leq \Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{x_g}{\Phi - x_g} \right)^\rho \right\}^{1/\rho}} \leq \max(x_g) = \Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\max(x_g)}{\Phi - \max(x_g)} \right)^\rho \right\}^{1/\rho}}, \\ \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - \max(y_g)}{\max(y_g)} \right)^\rho \right\}^{1/\rho}} &= \max(y_g) \leq \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - y_g}{y_g} \right)^\rho \right\}^{1/\rho}} \leq \min(y_g) = \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - \min(y_g)}{\min(y_g)} \right)^\rho \right\}^{1/\rho}}, \\ \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - \max(z_g)}{\max(z_g)} \right)^\rho \right\}^{1/\rho}} &= \max(z_g) \leq \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - z_g}{z_g} \right)^\rho \right\}^{1/\rho}} \leq \min(z_g) = \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - \min(z_g)}{\min(z_g)} \right)^\rho \right\}^{1/\rho}}. \end{aligned}$$

Therefore,  $N_{\min} \leq LNNDWAA(N_1, N_2, \dots, N_h) \leq N_{\max}$  is true.

**Theorem 4.** Let  $N_g = \langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \rangle$  ( $g = 1, 2, \dots, h$ ) be a group of LNNs and  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_h)$  be the weight vector of  $N_g$  ( $g = 1, 2, \dots, h$ ) for  $\gamma_g \in [0, 1]$  and  $\sum_{g=1}^h \gamma_g = 1$ . The aggregated result of the LNNDWAA operator is still an LNN, which can be expressed by

$$LNNDWGA(N_1, N_2, \dots, N_h) = \bigotimes_{g=1}^h N_g^{\gamma_g} = \left\langle Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{\Phi - x_g}{x_g} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{y_g}{\Phi - y_g} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \gamma_g \left( \frac{z_g}{\Phi - z_g} \right)^\rho \right\}^{1/\rho}} \right\rangle. \tag{16}$$

Theorem 4 is also proved based on mathematical induction, which is given below.

**Proof:**

(a) Let  $h = 2$ . Based on Definition 8 we can obtain

$$LNNDWGA(N_1, N_2) = N_1 \otimes N_2 = \left\langle Fr^{Ro}, \frac{\Phi}{1 + \left\{ \gamma_1 \left( \frac{\Phi - x_1}{x_1} \right)^\rho + \gamma_2 \left( \frac{\Phi - x_2}{x_2} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \gamma_1 \left( \frac{y_1}{\Phi - y_1} \right)^\rho + \gamma_2 \left( \frac{y_2}{\Phi - y_2} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \gamma_1 \left( \frac{z_1}{\Phi - z_1} \right)^\rho + \gamma_2 \left( \frac{z_2}{\Phi - z_2} \right)^\rho \right\}^{1/\rho}} \right\rangle = \left\langle Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^2 \gamma_g \left( \frac{\Phi - x_g}{x_g} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^2 \gamma_g \left( \frac{y_g}{\Phi - y_g} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^2 \gamma_g \left( \frac{z_g}{\Phi - z_g} \right)^\rho \right\}^{1/\rho}} \right\rangle$$

(b) If  $h = k$ , we can get the following equation from the equation (16):

$$LNNDWGA(N_1, N_2, \dots, N_k) = \bigotimes_{g=1}^k N_g^{\gamma_g} = \left\langle Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^k \gamma_g \left( \frac{\Phi - x_g}{x_g} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^k \gamma_g \left( \frac{y_g}{\Phi - y_g} \right)^\rho \right\}^{1/\rho}}, Fr^{Ro}, \frac{\Phi}{1 + \left\{ \sum_{g=1}^k \gamma_g \left( \frac{z_g}{\Phi - z_g} \right)^\rho \right\}^{1/\rho}} \right\rangle$$

(c) If  $h = k + 1$ , there exists the following result:

$$\begin{aligned}
 LNNDWGA(N_1, N_2, \dots, N_k, N_{k+1}) &= \bigotimes_{g=1}^{k+1} N_g^{\gamma_g} \\
 &= \left\langle Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{\Phi-x_g}{x_g}\right)^\rho\right\}^{1/\rho}}}, Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{y_g}{\Phi-y_g}\right)^\rho\right\}^{1/\rho}}}, Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^k \gamma_g \left(\frac{z_g}{\Phi-z_g}\right)^\rho\right\}^{1/\rho}}} \right\rangle \otimes \gamma_{k+1} N_{k+1} . \\
 &= \left\langle Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^{k+1} \gamma_g \left(\frac{\Phi-x_g}{x_g}\right)^\rho\right\}^{1/\rho}}}, Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^{k+1} \gamma_g \left(\frac{y_g}{\Phi-y_g}\right)^\rho\right\}^{1/\rho}}}, Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^{k+1} \gamma_g \left(\frac{z_g}{\Phi-z_g}\right)^\rho\right\}^{1/\rho}}} \right\rangle
 \end{aligned}$$

In terms of the above results, the equation (16) is true for all  $h$ .

The LNNDWGA operator also contains some properties:

(1) Reducibility: When the weight vector is  $\gamma = (1/h, 1/h, \dots, 1/h)$ , the equation (16) yields the following result:

$$\begin{aligned}
 LNNDWGA(N_1, N_2, \dots, N_h) &= \bigotimes_{g=1}^h N_g^{\gamma_g} \\
 &= \left\langle Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^h \frac{1}{h} \left(\frac{\Phi-x_g}{x_g}\right)^\rho\right\}^{1/\rho}}}, Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^h \frac{1}{h} \left(\frac{y_g}{\Phi-y_g}\right)^\rho\right\}^{1/\rho}}}, Fr_{\frac{\Phi}{1+\left\{\sum_{g=1}^h \frac{1}{h} \left(\frac{z_g}{\Phi-z_g}\right)^\rho\right\}^{1/\rho}}} \right\rangle .
 \end{aligned}$$

(2) Idempotency: Let all LNNs be  $N_g = \left\langle Fr_{x_g}^{Ro}, Fr_{y_g}^{Ro}, Fr_{z_g}^{Ro} \right\rangle = N$  ( $g = 1, 2, \dots, h$ ). Then,  $LNNDWGA(N_1, N_2, \dots, N_h) = N$ .

(3) Commutativity: Let the LNN sequence  $(N_1', N_2', \dots, N_h')$  be any permutation of  $(N_1, N_2, \dots, N_h)$ . Then, there is  $LNNDWGA(N_1', N_2', \dots, N_h') = LNNDWGA(N_1, N_2, \dots, N_h)$ .

(4) Boundedness: If the maximum and minimum LNNs are  $N_{\max} = \left\langle Fr_{\max(x_g)}^{Ro}, Fr_{\min(y_g)}^{Ro}, Fr_{\min(z_g)}^{Ro} \right\rangle$  and  $N_{\min} = \left\langle Fr_{\min(x_g)}^{Ro}, Fr_{\max(y_g)}^{Ro}, Fr_{\max(z_g)}^{Ro} \right\rangle$ , then  $N_{\min} \leq LNNDWGA(N_1, N_2, \dots, N_h) \leq N_{\max}$ .

Since the characteristics of the LNNDWGA operator can be easily proved by the similar proof process of the characteristics of the LNNDWAA operator, it is omitted here.

#### 4. MAGDM Method based on the LNNDWAA and LNNDWGA Operators

This section proposed a new DM method based on the LNNDWAA and LNNDWGA operators to solve MAGDM problems in the LNN environment.

In a MAGDM problem, let  $P = \{P_1, P_2, \dots, P_u\}$  be a set of alternatives and  $\Psi = \{\Psi_1, \Psi_2, \dots, \Psi_h\}$  be a set of attributes. The weight vector of the attributes  $\Psi_g$  ( $g = 1, 2, \dots, h$ ) is  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_h)$ . Assume that there is a group of decision-makers  $\Omega = \{\Omega_1, \Omega_2, \dots, \Omega_r\}$  with their weight vector  $\eta = (\eta_1, \eta_2, \dots, \eta_r)$ .

Each decision maker evaluates the value of each attribute  $\Psi_g$  ( $g = 1, 2, \dots, h$ ) for each alternative  $P_i$  from the set of linguistic terms  $Fr^{Ro} = \{Fr_0^{Ro} = \text{very low}, Fr_1^{Ro} = \text{low}, Fr_2^{Ro} = \text{slightly low}, Fr_3^{Ro} = \text{medium}, Fr_4^{Ro} = \text{slightly high}, Fr_5^{Ro} = \text{high}, Fr_6^{Ro} = \text{very high}\}$ . According to the linguistic terms, each decision-maker can assign the three linguistic values of indeterminacy, falsity, and truth to each attribute  $\Psi_g$  for the alternative  $P_v$ . Thus, LNN is composed of the obtained linguistic values. Hence, the LNN assessment information of the attributes  $\Psi_g$  ( $g = 1, 2, \dots, h$ ) for the alternatives  $P_v$  ( $v = 1, 2, \dots, u$ ) provided by each decision-maker  $\Omega_s$  ( $s = 1, 2, \dots, r$ ) can establish the LNN decision matrix  $M_s = (N_{vg}^s)_{u \times h}$ , where  $N_{vg}^s = \langle Fr_{x_{vg}}^{Ro}, Fr_{y_{vg}}^{Ro}, Fr_{z_{vg}}^{Ro} \rangle$  ( $s = 1, 2, \dots, r; v = 1, 2, \dots, u; g = 1, 2, \dots, h$ ) are LNNs.

Then, we present a MAGDM method using the score function (accuracy function) and the LNNDWAA and LNNDWGA operators to perform the MAGDM problem with LNN information. Here, the MAGDM method is described by the specific decision-making steps below.

**Step 1:** Aggregate all  $M_s$  ( $s = 1, 2, \dots, r$ ) by using the following LNNDWAA or LNNDWGA operator:

$$\begin{aligned}
 N_{vg} &= LNNDWAA(N_{vg}^1, N_{vg}^2, \dots, N_{vg}^r) = \bigoplus_{s=1}^r \eta_s N_{vg}^s \\
 &= \left\langle \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{s=1}^r \gamma_s \left( \frac{x_{vg}^s}{\Phi - x_{vg}^s} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{s=1}^r \gamma_s \left( \frac{\Phi - y_{vg}^s}{y_{vg}^s} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{s=1}^r \gamma_s \left( \frac{\Phi - z_{vg}^s}{z_{vg}^s} \right)^\rho \right\}^{1/\rho}}} \right\rangle \tag{17}
 \end{aligned}$$

or

$$\begin{aligned}
 N_{vg} &= LNNDWGA(N_{vg}^1, N_{vg}^2, \dots, N_{vg}^r) = \bigotimes_{s=1}^r (N_{vg}^s)^{\eta_s} \\
 &= \left\langle \frac{Fr^{Ro}}{1 + \left\{ \sum_{s=1}^r \gamma_s \left( \frac{\Phi - x_{vg}^s}{x_{vg}^s} \right)^\rho \right\}^{1/\rho}}, \frac{Fr^{Ro}}{1 + \left\{ \sum_{s=1}^r \gamma_s \left( \frac{y_{vg}^s}{\Phi - y_{vg}^s} \right)^\rho \right\}^{1/\rho}}, \frac{Fr^{Ro}}{1 + \left\{ \sum_{s=1}^r \gamma_s \left( \frac{z_{vg}^s}{\Phi - z_{vg}^s} \right)^\rho \right\}^{1/\rho}} \right\rangle \tag{18}
 \end{aligned}$$

to obtain the integrated matrix  $R = (N_{vg})_{u \times h}$ , where  $N_{vg} = \langle Fr_{x_{vg}}^{Ro}, Fr_{y_{vg}}^{Ro}, Fr_{z_{vg}}^{Ro} \rangle$  ( $v = 1, 2, \dots, u; g = 1, 2, \dots, h$ ) are integrated LNNs.

**Step 2:** Use the following LNNDWAA or LNNDWGA operator to obtain the collective overall LNNs  $N_v$  for  $P_v$  ( $v = 1, 2, \dots, u$ ):

$$\begin{aligned}
 N_v &= LNNDWAA(N_{v1}, N_{v2}, \dots, N_{vh}) = \bigoplus_{g=1}^h \gamma_g N_{vg} \\
 &= \left\langle \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \eta_g \left( \frac{x_g}{\Phi - x_g} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \eta_g \left( \frac{\Phi - y_g}{y_g} \right)^\rho \right\}^{1/\rho}}}, \frac{Fr^{Ro}}{\Phi - \frac{\Phi}{1 + \left\{ \sum_{g=1}^h \eta_g \left( \frac{\Phi - z_g}{z_g} \right)^\rho \right\}^{1/\rho}}} \right\rangle \tag{19}
 \end{aligned}$$

$$N_v = LNNDWGA(N_{v1}, N_{v2}, \dots, N_{vh}) = \bigotimes_{g=1}^{v_g} N_{vg}^{\gamma_g}$$

or

$$= \left\langle Fr^{Ro}_{\frac{\Phi}{1 + \left\{ \sum_{g=1}^h \eta_g \left( \frac{\Phi - x_g}{x_g} \right)^\rho \right\}^{1/\rho}}}, Fr^{Ro}_{\frac{\Phi}{1 + \left\{ \sum_{g=1}^h \eta_g \left( \frac{y_g}{\Phi - y_g} \right)^\rho \right\}^{1/\rho}}}, Fr^{Ro}_{\frac{\Phi}{1 + \left\{ \sum_{g=1}^h \eta_g \left( \frac{z_g}{\Phi - z_g} \right)^\rho \right\}^{1/\rho}}} \right\rangle. \quad (20)$$

**Step 3:** Calculate the score values of  $U(N_v)$  (the accuracy values of  $V(N_v)$ ) ( $v = 1, 2, \dots, u$ ) through the equation (1) (the equation (2)).

**Step 4:** Rank all alternatives in decreasing order, then select the more reasonable one.

**Step 5:** End.

### 5. An Illustrative Example on Slope Treatment Scheme Selection

The application of the MAGDM method proposed in this paper is illustrated by the selection of slope treatment schemes. To avoid slope instability, a set of four slope treatment options  $P = \{P_1, P_2, P_3, P_4\}$  is proposed, where  $P_1$  is gravity retaining wall + lattice protection;  $P_2$  is anti-slide retaining wall + anti-slide pile;  $P_3$  is anchor retaining wall + lattice protection; and  $P_4$  is pile-plate retaining wall. The evaluation of the schemes should meet the following attribute requirements: (1)  $\Psi_1$  is the economic status; (2)  $\Psi_2$  is the security situation; (3)  $\Psi_3$  is the construction feasibility; and (4)  $\Psi_4$  is the environment situation. The weight vector of the four attributes is assigned as  $\gamma = (0.23, 0.28, 0.26, 0.23)$ . Assume that three experts are invited as a group of decision makers  $\Omega = \{\Omega_1, \Omega_2, \Omega_3\}$ , then the weight vector  $\eta = (0.29, 0.33, 0.38)$  is given to indicate the importance of the various decision makers.

Decision makers need to assess the four attributes on the four alternatives from the linguistic term set  $Fr^{Ro} = \{Fr_0^{Ro} = \text{very low}, Fr_1^{Ro} = \text{low}, Fr_2^{Ro} = \text{slightly low}, Fr_3^{Ro} = \text{medium}, Fr_4^{Ro} = \text{slightly high}, Fr_5^{Ro} = \text{high}, Fr_6^{Ro} = \text{very high}\}$  with  $\Phi = 6$ . Thus, the linguistic evaluation results of each decision-maker  $\Omega_s$  ( $s = 1, 2, 3$ ) can be established as the LNN decision matrices  $M_1, M_2$ , and  $M_3$ :

$$M_1 = \begin{bmatrix} \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_1^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_4^{Ro}, Fr_5^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_3^{Ro} \rangle \\ \langle Fr_5^{Ro}, Fr_2^{Ro}, Fr_1^{Ro} \rangle & \langle Fr_5^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_4^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_4^{Ro}, Fr_5^{Ro} \rangle \\ \langle Fr_4^{Ro}, Fr_4^{Ro}, Fr_5^{Ro} \rangle & \langle Fr_5^{Ro}, Fr_3^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_5^{Ro}, Fr_2^{Ro}, Fr_1^{Ro} \rangle & \langle Fr_5^{Ro}, Fr_4^{Ro}, Fr_3^{Ro} \rangle \\ \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_4^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle \end{bmatrix}$$

$$M_2 = \begin{bmatrix} \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_4^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle \\ \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_1^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_4^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_4^{Ro}, Fr_4^{Ro} \rangle \\ \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_4^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle \\ \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_1^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_4^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle \end{bmatrix}$$

$$M_3 = \begin{bmatrix} \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_1^{Ro} \rangle \\ \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_3^{Ro}, Fr_4^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_3^{Ro} \rangle \\ \langle Fr_3^{Ro}, Fr_4^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_4^{Ro}, Fr_3^{Ro}, Fr_2^{Ro} \rangle \\ \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_5^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_1^{Ro} \rangle & \langle Fr_3^{Ro}, Fr_2^{Ro}, Fr_3^{Ro} \rangle \end{bmatrix}$$

The decision procedures based on the LNNDWAA operator are indicated below.

**Step 1:** Aggregate the decision matrices  $M_1$ ,  $M_2$ , and  $M_3$  by the equation (17) for  $\rho = 1$  and obtain the integrated matrix  $R = (N_{vg})_{4 \times 4}$  :

$$R = \begin{bmatrix} \langle Fr_{3.4249}^{Ro}, Fr_{2.0000}^{Ro}, Fr_{2.5751}^{Ro} \rangle & \langle Fr_{4.0000}^{Ro}, Fr_{2.0000}^{Ro}, Fr_{2.0033}^{Ro} \rangle & \langle Fr_{3.4790}^{Ro}, Fr_{3.2345}^{Ro}, Fr_{2.8599}^{Ro} \rangle & \langle Fr_{3.4249}^{Ro}, Fr_{2.5751}^{Ro}, Fr_{1.7045}^{Ro} \rangle \\ \langle Fr_{4.4496}^{Ro}, Fr_{2.2472}^{Ro}, Fr_{1.2346}^{Ro} \rangle & \langle Fr_{4.2808}^{Ro}, Fr_{2.5751}^{Ro}, Fr_{2.2472}^{Ro} \rangle & \langle Fr_{3.0000}^{Ro}, Fr_{2.6201}^{Ro}, Fr_{4.0000}^{Ro} \rangle & \langle Fr_{4.0000}^{Ro}, Fr_{2.5751}^{Ro}, Fr_{3.7430}^{Ro} \rangle \\ \langle Fr_{3.3799}^{Ro}, Fr_{3.6036}^{Ro}, Fr_{2.7933}^{Ro} \rangle & \langle Fr_{4.3051}^{Ro}, Fr_{3.2698}^{Ro}, Fr_{2.2140}^{Ro} \rangle & \langle Fr_{4.4496}^{Ro}, Fr_{2.2472}^{Ro}, Fr_{1.7192}^{Ro} \rangle & \langle Fr_{4.4496}^{Ro}, Fr_{3.2345}^{Ro}, Fr_{2.2140}^{Ro} \rangle \\ \langle Fr_{3.3799}^{Ro}, Fr_{2.2140}^{Ro}, Fr_{3.0211}^{Ro} \rangle & \langle Fr_{3.4249}^{Ro}, Fr_{1.5038}^{Ro}, Fr_{2.2901}^{Ro} \rangle & \langle Fr_{3.0000}^{Ro}, Fr_{2.2472}^{Ro}, Fr_{1.8692}^{Ro} \rangle & \langle Fr_{3.3799}^{Ro}, Fr_{2.5210}^{Ro}, Fr_{2.2901}^{Ro} \rangle \end{bmatrix}$$

**Step 2:** Through the equation (19), obtain the collective overall LNNs of  $N_v$  for  $P_v$  ( $v = 1, 2, 3, 4$ ) below:

$$N_1 = \langle Fr_{3.6290}^{Ro}, Fr_{2.3546}^{Ro}, Fr_{2.1981}^{Ro} \rangle, N_2 = \langle Fr_{4.0502}^{Ro}, Fr_{2.6660}^{Ro}, Fr_{2.2865}^{Ro} \rangle, N_3 = \langle Fr_{4.2426}^{Ro}, Fr_{2.9738}^{Ro}, Fr_{2.1555}^{Ro} \rangle, \text{ and } N_4 = \langle Fr_{3.3043}^{Ro}, Fr_{2.0120}^{Ro}, Fr_{2.2835}^{Ro} \rangle.$$

**Step 3:** Calculate the score values of  $U(N_v)$  ( $v = 1, 2, 3, 4$ ) by the equation (1):

$$U(N_1) = 0.6154, U(N_2) = 0.6165, U(N_3) = 0.6174, \text{ and } U(N_4) = 0.6116.$$

**Step 4:** Rank the four alternatives:  $P_3 \succ P_2 \succ P_1 \succ P_4$ . It can be seen that  $P_3$  is the most reasonable option among the four ones.

Or the decision procedures based on the LNNDWGA operator are indicated below.

**Step 1:** Aggregate the decision matrices  $M_1$ ,  $M_2$ , and  $M_3$  by the equation (18) for  $\rho = 1$  and obtain the integrated matrix  $R = (N_{vg})_{4 \times 4}$  :

$$R = \begin{bmatrix} \langle Fr_{3.2698}^{Ro}, Fr_{2.0000}^{Ro}, Fr_{2.7302}^{Ro} \rangle & \langle Fr_{4.0000}^{Ro}, Fr_{2.0000}^{Ro}, Fr_{3.1401}^{Ro} \rangle & \langle Fr_{3.3149}^{Ro}, Fr_{3.3799}^{Ro}, Fr_{3.9967}^{Ro} \rangle & \langle Fr_{3.2698}^{Ro}, Fr_{2.7302}^{Ro}, Fr_{2.4623}^{Ro} \rangle \\ \langle Fr_{4.2463}^{Ro}, Fr_{2.3964}^{Ro}, Fr_{1.4338}^{Ro} \rangle & \langle Fr_{3.7430}^{Ro}, Fr_{2.7302}^{Ro}, Fr_{2.3964}^{Ro} \rangle & \langle Fr_{3.0000}^{Ro}, Fr_{2.7655}^{Ro}, Fr_{4.0000}^{Ro} \rangle & \langle Fr_{4.0000}^{Ro}, Fr_{3.7099}^{Ro}, Fr_{4.2808}^{Ro} \rangle \\ \langle Fr_{3.2345}^{Ro}, Fr_{3.7528}^{Ro}, Fr_{3.9798}^{Ro} \rangle & \langle Fr_{3.8023}^{Ro}, Fr_{3.4249}^{Ro}, Fr_{2.3526}^{Ro} \rangle & \langle Fr_{4.2463}^{Ro}, Fr_{2.3964}^{Ro}, Fr_{2.2570}^{Ro} \rangle & \langle Fr_{4.2463}^{Ro}, Fr_{3.3799}^{Ro}, Fr_{2.3526}^{Ro} \rangle \\ \langle Fr_{3.2345}^{Ro}, Fr_{2.3526}^{Ro}, Fr_{4.2222}^{Ro} \rangle & \langle Fr_{3.2698}^{Ro}, Fr_{1.7173}^{Ro}, Fr_{2.4497}^{Ro} \rangle & \langle Fr_{3.0000}^{Ro}, Fr_{2.3964}^{Ro}, Fr_{3.4093}^{Ro} \rangle & \langle Fr_{3.2345}^{Ro}, Fr_{2.6851}^{Ro}, Fr_{2.4497}^{Ro} \rangle \end{bmatrix}$$

**Step 2:** Through the equation (20), obtain the collective overall LNNs  $N_v$  for  $P_v$  ( $v = 1, 2, 3, 4$ ) below:

$$N_1 = \langle Fr_{3.4588}^{Ro}, Fr_{2.6338}^{Ro}, Fr_{3.2455}^{Ro} \rangle, N_2 = \langle Fr_{3.6611}^{Ro}, Fr_{2.9722}^{Ro}, Fr_{3.4480}^{Ro} \rangle, N_3 = \langle Fr_{3.8440}^{Ro}, Fr_{3.3047}^{Ro}, Fr_{2.9054}^{Ro} \rangle, \text{ and } N_4 = \langle Fr_{3.1795}^{Ro}, Fr_{2.2959}^{Ro}, Fr_{3.3218}^{Ro} \rangle.$$

**Step 3:** Obtain the score values of  $U(N_v)$  ( $v = 1, 2, 3, 4$ ) by the equation (1):

$$U(N_1) = 0.5322, U(N_2) = 0.5134, U(N_3) = 0.5352, \text{ and } U(N_4) = 0.5312.$$

**Step 4:** Rank the four alternatives:  $P_3 \succ P_1 \succ P_4 \succ P_2$ . It can be seen that  $P_3$  is the most reasonable choice among the four ones.



We can repeat the above decision process by changing the parameter  $\rho$  from 2 to 4. The sorting results obtained by using the LNNDWAA operator are shown in Figure 2. Then, the ranking orders based on the LNNDWGA operator are indicated in Figure 3.




Figure 2. Ranking orders of the four alternatives based on the LNNDWAA operator (I, II, III, IV are ranking numbers)

As shown in Figure 2, the sorting results obtained based on the LNNDWAA operator change with the change of the parameter values of  $\rho$ . With an increase of  $\rho$ , the score values of the four alternatives gradually increase. However, the ranking orders tend to robustness when  $\rho > 3$ .




Figure 3. Ranking orders of the four alternatives based on the LNNDWGA operator

For the evaluation results using the LNNDWGA operator in Figure 3, the ranking orders also change with the change of  $\rho$ . With an increase of  $\rho$ , the score values of the four alternatives gradually decrease. The ranking orders tend to robustness when the value of the parameter  $\rho$  exceeds 3.

Furthermore, a comparison is made between the new MAGDM method and the existing relative MAGDM methods by the operators of LNNWAA and LNNWGA proposed by Fan and Ye [24]. According to the calculational steps given by Fan and Ye [24], the alternatives are evaluated as follows.

**Step 1:** By the LNNWAA operator of the equation (4), we can obtain the integrated matrix

$$R = \left( N_{vg} \right)_{u \times h} :$$

$$R = \begin{bmatrix} \left\langle Fr_{3.3757}^{Ro}, Fr_{2.0000}^{Ro}, Fr_{2.6243}^{Ro} \right\rangle & \left\langle Fr_{4.0000}^{Ro}, Fr_{2.0000}^{Ro}, Fr_{2.3988}^{Ro} \right\rangle & \left\langle Fr_{3.4284}^{Ro}, Fr_{3.2610}^{Ro}, Fr_{3.0433}^{Ro} \right\rangle & \left\langle Fr_{3.3757}^{Ro}, Fr_{2.6243}^{Ro}, Fr_{1.9761}^{Ro} \right\rangle \\ \left\langle Fr_{4.3642}^{Ro}, Fr_{2.2863}^{Ro}, Fr_{1.3013}^{Ro} \right\rangle & \left\langle Fr_{4.0917}^{Ro}, Fr_{2.6243}^{Ro}, Fr_{2.2863}^{Ro} \right\rangle & \left\langle Fr_{3.0000}^{Ro}, Fr_{2.6672}^{Ro}, Fr_{4.0000}^{Ro} \right\rangle & \left\langle Fr_{4.0000}^{Ro}, Fr_{3.5858}^{Ro}, Fr_{3.8255}^{Ro} \right\rangle \\ \left\langle Fr_{3.3328}^{Ro}, Fr_{3.6377}^{Ro}, Fr_{2.9822}^{Ro} \right\rangle & \left\langle Fr_{4.1300}^{Ro}, Fr_{3.2988}^{Ro}, Fr_{2.2496}^{Ro} \right\rangle & \left\langle Fr_{4.3642}^{Ro}, Fr_{2.2863}^{Ro}, Fr_{1.9083}^{Ro} \right\rangle & \left\langle Fr_{4.3642}^{Ro}, Fr_{3.2610}^{Ro}, Fr_{2.2496}^{Ro} \right\rangle \\ \left\langle Fr_{3.3328}^{Ro}, Fr_{2.2496}^{Ro}, Fr_{3.3286}^{Ro} \right\rangle & \left\langle Fr_{3.3757}^{Ro}, Fr_{1.5911}^{Ro}, Fr_{2.3332}^{Ro} \right\rangle & \left\langle Fr_{3.0000}^{Ro}, Fr_{2.2863}^{Ro}, Fr_{2.3620}^{Ro} \right\rangle & \left\langle Fr_{3.3328}^{Ro}, Fr_{2.5716}^{Ro}, Fr_{2.3332}^{Ro} \right\rangle \end{bmatrix}.$$

**Step 2:** The collective overall linguistic neutrosophic numbers of  $N_v$  for  $P_v$  ( $v = 1, 2, 3, 4$ ) was determined below:

$$N_1 = \langle Fr_{3.5807}^{Ro}, Fr_{2.4175}^{Ro}, Fr_{2.4916}^{Ro} \rangle, N_2 = \langle Fr_{3.9258}^{Ro}, Fr_{2.7432}^{Ro}, Fr_{2.6147}^{Ro} \rangle, N_3 = \langle Fr_{4.0996}^{Ro}, Fr_{3.0590}^{Ro}, Fr_{2.2998}^{Ro} \rangle, \text{ and } N_4 = \langle Fr_{3.2625}^{Ro}, Fr_{2.1144}^{Ro}, Fr_{2.5240}^{Ro} \rangle.$$

**Step 3:** Calculate the score values of  $U(N_v)$  ( $v = 1, 2, 3, 4$ ) for the collective overall linguistic neutrosophic numbers of  $N_v$ :

$$U(N_1) = 0.5929, U(N_2) = 0.5860, U(N_3) = 0.5967, \text{ and } U(N_4) = 0.5902.$$

**Step 4:** We can get the ranking of the four alternatives:  $P_3 \succ P_1 \succ P_4 \succ P_2$ . It can be seen that  $P_3$  is the most reasonable choice among the four ones.

Or by the LNNWGA operator of the equation (5), the calculational steps are given below.

**Step 1:** This step is the same as Step 1 mentioned above.

**Step 2:** Through the equation (5), the collective overall LNNs of  $N_v$  for  $P_v$  ( $v = 1, 2, 3, 4$ ) below:

$$N_1 = \langle Fr_{3.5543}^{Ro}, Fr_{2.5138}^{Ro}, Fr_{2.5421}^{Ro} \rangle, N_2 = \langle Fr_{3.8110}^{Ro}, Fr_{2.8160}^{Ro}, Fr_{3.0491}^{Ro} \rangle, N_3 = \langle Fr_{4.0389}^{Ro}, Fr_{3.1457}^{Ro}, Fr_{2.3507}^{Ro} \rangle, \text{ and } N_4 = \langle Fr_{3.2545}^{Ro}, Fr_{2.1658}^{Ro}, Fr_{2.5717}^{Ro} \rangle.$$

**Step 3:** Calculate the score values of  $U(N_v)$  ( $v = 1, 2, 3, 4$ ):

$$U(N_1) = 0.5832, U(N_2) = 0.5525, U(N_3) = 0.5857, \text{ and } U(N_4) = 0.5843.$$

**Step 4:** We can get the ranking of the four alternatives:  $P_3 \succ P_4 \succ P_2 \succ P_1$ . It can be seen that  $P_3$  is the most reasonable option among the four ones.

Figure 4 shows the comparison of the decision results obtained using the LNNWGA and LNNWAA operators [24] and the proposed LNNDWAA and LNNDWGA operators in this study. The ranking orders in this MAGDM example are influenced by different aggregation operators and values of the parameter  $\rho$ . According to the results obtained using the LNNWGA and LNNWAA operators, the scheme  $P_3$  is the most reasonable option among the four alternatives. It is the same as the result based on the proposed LNNDWAA and LNNDWGA operators when  $\rho = 1$ . However, the

best alternative is  $P_2$  according to the proposed LNNDWAA operator when  $\rho = 2, 3, 4$ . According to the result of the proposed LNNDWGA operator, when  $\rho = 2, 3, 4$ , the best alternative is  $P_4$ .




Figure 4. Comparison of the decision results based on different aggregation operators and values of  $\rho$

### 6. Conclusion

In this study, the LNNDWAA and LNNDWGA operators and their properties were proposed in view of the Dombi operations in the LNN environment. A novel technique for MAGDM problems was proposed using the LNNDWAA or LNNDWGA operator. In the proposed MAGDM process, regarding the satisfactory assessment of alternatives over multiple attributes, we established a decision matrix based on the suitable evaluation results given by the decision makers. Then, we used the LNNDWAA/LNNDWGA operator to aggregate LNN information. Finally, the score values (accuracy values if necessary) was calculated and the ranking results of alternatives are given in a descending order to obtain the optimal choice. In the DM application, an illustrative example of the selection of landslide treatment schemes was presented to verify the feasibility of the proposed method. Compared with the related MAGDM methods in previous studies, this new method can influence the sorting order of alternatives by changing the parameter values of  $\rho$ . Thus, it can overcome the insufficiency of decision flexibility in the existing MAGDM method with LNNs. Therefore, we can more effectively deal with the DM problem of landslide treatment schemes by specifying various parameter values according to the preferences and demands of decision makers. It is obvious that this new method can better solve the DM problem of landslide treatment schemes and make the DM results more reasonable and flexible in the uncertainty and inconsistency of human linguistic judgments.

**Data Availability:** The data used to support the findings of the study are available in the article.

**Conflicts of Interest:** The authors declare no conflicts of interest.

**Acknowledgments:** The study was funded by the National Natural Science Foundation of China (Nos. 42177117, 41427802, and 41572299), Zhejiang Provincial Natural Science Foundation (No. LQ16D020001).

## References

1. National Bureau of Statistics of China. China statistical yearbook. *China Statistics Press*, Beijing, 2020. (in Chinese)
2. Wang, L. Study on treatment technique and scheme decision of slope disease of mountainous highway. Master thesis, Chongqing Jiaotong University, 2012. (in Chinese)
3. Park, S.; Choi, C.; Kim, B.; Kim, J. Landslide susceptibility mapping using frequency ratio, analytic hierarchy process, logistic regression, and artificial neural network methods at the Inje area, Korea. *Environmental earth sciences* 2013, 68, 1443–1464.
4. Wang, L.; Li, W.; Li, H. Decision-making for ecological landslide prevention in tropical rainforests. *Natural Hazards* 2020, 103, 985–1008.
5. Xu, S.; Zhang, M.; Ma, Y.; Liu, J.; Wang, Y.; Ma, X.; Chen, J. Multiclassification Method of Landslide Risk Assessment in Consideration of Disaster Levels: A Case Study of Xianyang City, Shaanxi Province. *ISPRS International Journal of Geo-Information* 2021, 10, 646.
6. Ma, Z.Z. Group decision-making method from a reliability perspective and its application. Ph.D. Thesis China: Nanjing University of Aeronautics and Astronautics, 2018.
7. Zadeh, L.A. The concept of a linguistic variable and its application to approximate reasoning-I. *Information sciences* 1975, 8, 199–249.
8. Fang, Z.; Ye, J. Multiple attribute group decision-making method based on linguistic neutrosophic numbers. *Symmetry* 2017, 9, 111.
9. Herrera, F.; Herrera-Viedma, E. A model of consensus in group decision making under linguistic assessments. *Fuzzy sets and Systems* 1996, 78, 73–87.
10. Herrera, F.; Herrera-Viedma, E. Linguistic decision analysis: steps for solving decision problems under linguistic information. *Fuzzy Sets and systems* 2000, 115, 67–82.
11. Xu, Z.S. A method based on linguistic aggregation operators for group decision making with linguistic preference relations. *Information sciences* 2004, 166, 19–30.
12. Xu, Z.S. A note on linguistic hybrid arithmetic averaging operator in multiple attribute group decision making with linguistic information. *Group Decision and Negotiation* 2006, 15, 593–604.
13. Xu, Z.S. Goal programming models for multiple attribute decision making under linguistic setting. *Journal of Management Sciences in China* 2006, 9, 9–17.
14. Merigó, J.M.; Casanovas, M.; Martínez, L. Linguistic aggregation operators for linguistic decision making based on the Dempster-Shafer theory of evidence. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems* 2010, 18, 287–304.
15. Merigó, J.M.; Gil-Lafuente, A.M. Induced 2-tuple linguistic generalized aggregation operators and their application in decision-making. *Information Sciences* 2013, 236, 1–16.
16. Merigó, J.M.; Palacios-Marques, D.; Zeng, S. Subjective and objective information in linguistic multi-criteria group decision making. *European Journal of Operational Research* 2016, 248, 522–531.
17. Xu, Z. Uncertain linguistic aggregation operators based approach to multiple attribute group decision making under uncertain linguistic environment. *Information sciences* 2004, 168, 171–184.
18. Xu, Z. Induced uncertain linguistic OWA operators applied to group decision making. *Information fusion* 2006, 7, 231–238.
19. Wei, G.W. Uncertain linguistic hybrid geometric mean operator and its application to group decision making under uncertain linguistic environment. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems* 2009, 17, 251–267.
20. Wei, G.; Zhao, X.; Lin, R. et al. Uncertain linguistic Bonferroni mean operators and their application to multiple attribute decision making. *Applied Mathematical Modelling* 2013, 37, 5277–5285.
21. Park, J.H.; Gwak, M.G.; Kwun, Y.C. Uncertain linguistic harmonic mean operators and their applications to multiple attribute group decision making. *Computing* 2011, 93, 47–64.
22. Peng, B.; Ye, C.; Zeng, S. Uncertain pure linguistic hybrid harmonic averaging operator and generalized interval aggregation operator based approach to group decision-making. *Knowledge-Based Systems* 2012, 36, 175–181.
23. Zhang, H. Uncertain linguistic power geometric operators and their use in multiattribute group decision making. *Mathematical Problems in Engineering* 2015, 2015, 1–10.

24. Chen, Z.; Liu, P.; Pei, Z. An approach to multiple attribute group decision making based on linguistic intuitionistic fuzzy numbers. *International Journal of Computational Intelligence Systems* **2015**, *8*, 747–760.
25. Liu, P.; Wang, P. Some improved linguistic intuitionistic fuzzy aggregation operators and their applications to multiple-attribute decision making. *International Journal of Information Technology & Decision Making* **2017**, *16*, 817–850.
26. Liu, P.; Liu, J.; Merigó, J.M. Partitioned Heronian means based on linguistic intuitionistic fuzzy numbers for dealing with multi-attribute group decision making. *Applied Soft Computing* **2018**, *62*, 395–422.
27. Smarandache, F. Introduction to Neutrosophic Measure, Neutrosophic Integral, and Neutrosophic Probability. Sitech & Education Publisher: Craiova, Romania, **2013**.
28. Smarandache F. Symbolic neutrosophic theory. *Infinite Study*, **2015**.
29. Smarandache, F. Introduction to Neutrosophic Statistics. Sitech & Education Publisher: Craiova, Romania, **2014**.
30. Smarandache, F. Neutrosophy: Neutrosophic Probability, Set, and Logic: Analytic Synthesis & Synthetic Analysis. American Research Press: Rehoboth, DE, USA, **1998**.
31. Fan, C.; Ye, J.; Hu, K.; Fan, E. Bonferroni mean operators of linguistic neutrosophic numbers and their multiple attribute group decision-making methods. *Information* **2017**, *8*, 107.
32. Liu, P.; Mahmood, T.; Khan, Q. Group decision making based on power Heronian aggregation operators under linguistic neutrosophic environment. *International Journal of Fuzzy Systems* **2018**, *20*, 970–985.
33. Shi, L.L.; Ye, J. Multiple attribute group decision-making method using correlation coefficients between linguistic neutrosophic numbers. *Journal of Intelligent & Fuzzy Systems* **2018**, *35*, 917–925.
34. Shi, L.; Ye, J. Cosine measures of linguistic neutrosophic numbers and their application in multiple attribute group decision-making. *Information* **2017**, *8*, 117.
35. Cui, W.H.; Ye, J. Multiple-attribute decision-making method using similarity measures of hesitant linguistic neutrosophic numbers regarding least common multiple cardinality. *Symmetry* **2018**, *10*, 330.
36. Dombi, J. A general class of fuzzy operators, the DeMorgan class of fuzzy operators and fuzziness measures induced by fuzzy operators. *Fuzzy sets and systems* **1982**, *8*, 149–163.
37. Liu, P.; Liu, J.; Chen, S.M. Some intuitionistic fuzzy Dombi Bonferroni mean operators and their application to multi-attribute group decision making. *Journal of the Operational Research Society* **2017**, 1–26.
38. Chen, J.; Ye, J. Some single-valued neutrosophic Dombi weighted aggregation operators for multiple attribute decision-making. *Symmetry* **2017**, *9*, 82.
39. Lu, X.; Ye, J. Dombi aggregation operators of linguistic cubic variables for multiple attribute decision making. *Information* **2018**, *9*, 188.
40. He, X. Typhoon disaster assessment based on Dombi hesitant fuzzy information aggregation operators. *Natural Hazards* **2018**, *90*, 1153–1175.
41. Jana, C.; Senapati, T.; Pal, M.; Yager, R.R. Picture fuzzy Dombi aggregation operators: Application to MADM process. *Applied Soft Computing* **2019**, *74*, 99–109.
42. Jana, C.; Pal, M. Multi-criteria decision making process based on some single-valued neutrosophic Dombi power aggregation operators. *Soft Computing* **2021**, *25*, 5055–5072.
43. Ayub, S.; Abdullah, S.; Ghani, F.; Qiyas, M.; Khan, M.Y. Cubic fuzzy Heronian mean Dombi aggregation operators and their application on multi-attribute decision-making problem. *Soft Computing* **2021**, *25*, 4175–4189.

Received: June 12, 2022. Accepted: September 21, 2022.