

4-11-2023

## On The 3-Refined Neutrosophic Analytical Structures and Number Theoretical Concepts

Hamiyet Merkepçi

Katy D. Ahmad

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Merkepçi, Hamiyet and Katy D. Ahmad. "On The 3-Refined Neutrosophic Analytical Structures and Number Theoretical Concepts." *Neutrosophic Sets and Systems* 54, 1 (). [https://digitalrepository.unm.edu/nss\\_journal/vol54/iss1/1](https://digitalrepository.unm.edu/nss_journal/vol54/iss1/1)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).



# On The 3-Refined Neutrosophic Analytical Structures and Number Theoretical Concepts

<sup>1</sup>Hamiyet Merkepçi,<sup>2</sup> Katy D. Ahmad

<sup>1</sup>-Gaziantep University, Department Of Mathematics, Gaziantep, Turkey; [hamiyet.merkeci@gmail.com](mailto:hamiyet.merkeci@gmail.com)

<sup>2</sup>-Islamic University Of Gaza, Department Of Mathematics, Gaza, Palestine; [Katyon765@gmail.com](mailto:Katyon765@gmail.com)

Co- Hamiyet Merkepçi: [hamiyet.merkeci@gmail.com](mailto:hamiyet.merkeci@gmail.com)

## Abstract:

n-refined neutrosophic structures are considered as generalizations of classical structures, and neutrosophic structures.

The main goal of this paper, is to study several structures generated by using 3-refined neutrosophic numbers, where we find the mathematical formulas of 3-refined neutrosophic real functions. Also, the inner products over 3-refined neutrosophic vector spaces and orthogonal properties. In addition, we present the foundations of 3-refined neutrosophic number theory, especially division, congruencies, and some related equations.

**Keywords:** 3-refined neutrosophic real function, 3-refined neutrosophic inner product, 3-refined neutrosophic vector space, 3-refined neutrosophic number theory

## Introduction and basic concepts

The concept of neutrosophic structures plays an important role in the theory of algebraic structures and analysis. Many concepts and structures were defined previously, such as neutrosophic vector spaces, neutrosophic matrices, and algebraic rings [1-3, 5-7, 14-16].

Laterally, refined neutrosophic structures were defined to generalize the neutrosophic structures, where refined neutrosophic rings, modules, and other structures were presented [4, 8-11, 24-28].

The concept of n-refined neutrosophic structure is considered as a generalization of refined structure [12, 29]. For each value of the integer n, we get a generalized structure.

This work will study some of 3-refined neutrosophic structures, where we present the formulas of 3-refined neutrosophic real functions, 3-refined neutrosophic inner products defined over 3-refined neutrosophic vector spaces, and 3-refined number theoretical concepts.

First, we recall some basic concepts.

**Definition:**

Let  $(R, +, \times)$  be a ring,  $R(I) = \{a + bI; a, b \in R\}$  is called the neutrosophic ring where I is a neutrosophic element with condition  $I^2 = I$ .

**Definition:**

Let  $(R, +, \times)$  be a ring,  $(R(I_1, I_2), +, \times)$  is called a refined neutrosophic ring generated by  $R, I_1, I_2$ .

**Definition:**

Let  $(R, +, \times)$  be a ring and  $I_k; 1 \leq k \leq n$  be n indeterminacies. We define  $R_n(I) = \{a_0 + a_1I + \dots + a_nI_n; a_i \in R\}$  to be n-refined neutrosophic ring.

Addition and multiplication on  $R_n(I)$  are defined as:

$$\sum_{i=0}^n x_i I_i + \sum_{i=0}^n y_i I_i = \sum_{i=0}^n (x_i + y_i) I_i, \sum_{i=0}^n x_i I_i \times \sum_{i=0}^n y_i I_i = \sum_{i,j=0}^n (x_i \times y_j) I_i I_j.$$

Where  $\times$  is the multiplication defined on the ring R.

For n=3, we get the 3-refined neutrosophic ring.

**Main Discussion**

**Definition.**

Let  $R_3(I)$  be the 3-refined neutrosophic ring of reals,  $f: R_3(I) \rightarrow R_3(I); f = f(X); X \in R_3(I)$ .  $f$  is called 3-refined neutrosophic real function with one variable.

**Theorem.**

$$R_3(I) \cong R^4.$$

**Proof.**

We define  $g: R_3(I) \rightarrow R^4; g(a + bI_1 + cI_2 + dI_3) = (a, a + b + c + d, a + c + d, a + d)$ .

It is clear that  $g$  is well defined function.

$\ker(g) = \{0\}$ , thus  $g$  is injective.

$\text{Im}(g) = R^4$ , thus  $g$  is surjective, so that  $g$  is one-to-one.

Now, let  $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3, B = b_0 + b_1I_1 + b_2I_2 + b_3I_3$ ,

$$A + B = (a_0 + b_0) + (a_1 + b_1)I_1 + (a_2 + b_2)I_2 + (a_3 + b_3)I_3$$

$$g(A + B) = g(A) + g(B).$$

$$\begin{aligned} A.B &= a_0b_0 + [(a_0 + a_1 + a_2 + a_3)(b_0 + b_1 + b_2 + b_3) - (a_0 + a_2 + a_3)(b_0 + b_2 + b_3)]I_1 \\ &\quad + [(a_0 + a_2 + a_3)(b_0 + b_2 + b_3) - (a_0 + a_3)(b_0 + b_3)]I_2 \\ &\quad + [(a_0 + a_3)(b_0 + b_3) - a_0b_0]I_3 \end{aligned}$$

$g(A.B) = g(A).g(B)$ , hence  $g$  is a ring isomorphism.

**Remark.**

Let  $f: R_3(I) \rightarrow R_3(I)$  be a 3-refined neutrosophic real function with one variable, then  $f$  can be represented by four classical real functions by taking the direct isomorphic image  $g(f(X))$ .

**Example.**

Take  $f(X) = (1 + I_1)X^2 + (2 - I_2 - I_3)X + 1 + 2I_1 + I_2 + I_3$ ,  $f$  can be represented as follows:

$$g(f(X)) = g(1 + I_1)(g(X))^2 + g(2 - I_2 - I_3)g(X) + g(1 + 2I_1 + I_2 + I_3)$$

$$g(f(X)) = (1,2,1,1)(x_0^2, (x_0 + x_1 + x_2 + x_3)^2, (x_0 + x_2 + x_3)^2, (x_0 + x_3)^2)$$

$$+ (2,0,0,1)(x_0, x_0 + x_1 + x_2 + x_3, x_0 + x_2 + x_3, x_0 + x_3) + (1,5,3,2)$$

$$\begin{aligned} g(f(X)) &= (x_0^2 + 2x_0 + 1, 2(x_0 + x_1 + x_2 + x_3)^2 + 5, (x_0 + x_2 + x_3)^2 + 3, (x_0 + x_3)^2 \\ &\quad + (x_0 + x_3) + 2) \end{aligned}$$

The four classical functions that represent  $(f)$  are:

$$f_1: R \rightarrow R; f_1(x_0) = x_0^2 + 2x_0 + 1$$

$$f_2: R \rightarrow R; f_2(x_0 + x_1 + x_2 + x_3) = 2(x_0 + x_1 + x_2 + x_3)^2 + 5$$

$$f_3: R \rightarrow R; f_3(x_0 + x_2 + x_3) = (x_0 + x_2 + x_3)^2 + 3$$

$$f_4: R \rightarrow R; f_4(x_0 + x_3) = (x_0 + x_3)^2 + (x_0 + x_3) + 2$$

**Theorem.**

Let  $g: R_3(I) \rightarrow R^4$  be the isomorphism defined above, then:

$$g^{-1}: R^4 \rightarrow R_3(I); g^{-1}(a, b, c, d) = a + (b - c)I_1 + (c - d)I_2 + (d - a)I_3.$$

The proof is easy.

**Remark.**

To find the formula of a 3-refined neutrosophic real function  $f: R_3(I) \rightarrow R_3(I)$ , we went compute:

$$g^{-1}(g(f(X))).$$

**Example.**

For the function  $f(X) = (1 + I_1)X^2 + (2 - I_2 - I_3)X + 1 + 2I_1 + I_2 + I_3$ , we compute:

$$\begin{aligned} g^{-1}(g(f(X))) &= (x_0^2 + 2x_0 + 1) + [2(x_0 + x_1 + x_2 + x_3)^2 + 5 - (x_0 + x_2 + x_3)^2 - 3]I_1 \\ &\quad + [(x_0 + x_2 + x_3)^2 + 3 - (x_0 + x_3)^2 - (x_0 + x_3) - 2]I_2 \\ &\quad + [(x_0 + x_3)^2 + (x_0 + x_3) + 2 - x_0^2 - 2x_0 - 1]I_3 \\ &= x_0^2 + 2x_0 + 1 + [2(x_0 + x_1 + x_2 + x_3)^2 - (x_0 + x_2 + x_3)^2 + 2]I_1 \\ &\quad + [(x_0 + x_2 + x_3)^2 - (x_0 + x_3)^2 - (x_0 + x_3) + 1]I_2 \\ &\quad + [(x_0 + x_3)^2 + (x_0 + x_3) - x_0^2 - 2x_0 + 1]I_3 \end{aligned}$$

**Definition.**

Let  $f: R_3(I) \rightarrow R_3(I)$  be a 3-refined neutrosophic real function, and  $g(f(X)) = (f_1, f_2, f_3, f_4)$ , with  $f_i: R \rightarrow R; 1 \leq i \leq 4$ , we say:

- a).  $f$  is differentiable if and only if  $f_i$  are differentiable.
- b).  $f$  is integrable if and only if  $f_i$  are integrable.

We mean by differentiable/integrable on all  $R$  not only for sub-domains  $]a, b[ \subseteq R$ .

**Example on famous functions.**

1.  $f: R_3(I) \rightarrow R_3(I), f(X) = \sin(X)$ .

It's formula is  $f(X) = g^{-1}(g(f(X))) = \sin(x_0) + [\sin(x_0 + x_1 + x_2 + x_3) - \sin(x_0 + x_2 + x_3)]I_1 + [\sin(x_0 + x_2 + x_3) - \sin(x_0 + x_3)]I_2 + [\sin(x_0 + x_3) - \sin(x_0)]I_3$ .

2.  $f(X) = \cos(X) = g^{-1}(g(f(X))) = \cos(x_0) + [\cos(x_0 + x_1 + x_2 + x_3) - \cos(x_0 + x_2 + x_3)]I_1 + [\cos(x_0 + x_2 + x_3) - \cos(x_0 + x_3)]I_2 + [\cos(x_0 + x_3) - \cos(x_0)]I_3$ .

3.  $f(X) = \tan(X) = \tan(x_0) + [\tan(x_0 + x_1 + x_2 + x_3) - \tan(x_0 + x_2 + x_3)]I_1 + [\tan(x_0 + x_2 + x_3) - \tan(x_0 + x_3)]I_2 + [\tan(x_0 + x_3) - \tan(x_0)]I_3$

4.  $f(X) = \cot(X) = \cot(x_0) + [\cot(x_0 + x_1 + x_2 + x_3) - \cot(x_0 + x_2 + x_3)]I_1 + [\cot(x_0 + x_2 + x_3) - \cot(x_0 + x_3)]I_2 + [\cot(x_0 + x_3) - \cot(x_0)]I_3$

5.  $f(X) = e^X = e^{x_0} + [e^{x_0+x_1+x_2+x_3} - e^{x_0+x_2+x_3}]I_1 + [e^{x_0+x_2+x_3} - e^{x_0+x_3}]I_2 + [e^{x_0+x_3} - e^{x_0}]I_3$

6.  $f(X) = \ln(X) = \ln(x_0) + [\ln(x_0 + x_1 + x_2 + x_3) - \ln(x_0 + x_2 + x_3)]I_1 + [\ln(x_0 + x_2 + x_3) - \ln(x_0 + x_3)]I_2 + [\ln(x_0 + x_3) - \ln(x_0)]I_3$ , with  $X > 0$ .
7.  $f(X) = X^n = x_0^n + [(x_0 + x_1 + x_2 + x_3)^n - (x_0 + x_2 + x_3)^n]I_1 + [(x_0 + x_2 + x_3)^n - (x_0 + x_3)^n]I_2 + [(x_0 + x_3)^n - x_0^n]I_3; n \in \mathbb{N}$ .

**Definition.**

Let  $V$  be vector space over  $R$ , the 3-refined neutrosophic vector space is defined as follows:

$$V_3(I) = V + VI_1 + VI_2 + VI_3 = \{x + yI_1 + zI_2 + tI_3; x, y, z, t \in R\}$$

**Remark.**

Addition on  $V_3(I)$  is defined:

$$\begin{aligned} (x_0 + y_0I_1 + z_0I_2 + t_0I_3) + (x_1 + y_1I_1 + z_1I_2 + t_1I_3) \\ = (x_0 + x_1) + (y_0 + y_1)I_1 + (z_0 + z_1)I_2 + (t_0 + t_1)I_3 \end{aligned}$$

Where  $x_i, y_i, z_i, t_i \in V; 0 \leq i \leq 1$ .

Multiplication on  $V_3(I)$  is defined:

$$\begin{aligned} (a + bI_1 + cI_2 + dI_3) \cdot (x + yI_1 + zI_2 + tI_3) \\ = (a \cdot x) + (ay + bx + by + bz + bt + cy + dy)I_1 + (az + cx + cz + ct + dz)I_2 \\ + (at + dx + dt)I_3 \end{aligned}$$

Where  $a, b, c, d \in R, x, y, z, t \in V$

**Remark.**

$(V_3(I), +, \cdot)$  Is a module over  $R_3(I)$ .

**Definition.**

Let  $f: V_3(I) \times V_3(I) \rightarrow R_3(I)$  be a well defined mapping, we call  $f$  a 3-refined neutrosophic real inner product if and only if the following conditions hold:

- 1).  $f(X, X) \geq 0; \forall X \in V_3(I)$ .
- 2).  $f(X, X) = 0 \Leftrightarrow X = 0$ .
- 3).  $f(X, Y) = f(Y, X); \forall X, Y \in V_3(I)$ .
- 4).  $f(\alpha X + \beta Y, Z) = \alpha f(X, Z) + \beta f(Y, Z); \forall X, Y, Z \in V_3(I), \alpha, \beta \in R_3(I)$ .

**Theorem.**

Let  $g: V \times V \rightarrow R$  be an inner production on  $V$ , then for  $X = x_0 + x_1I_1 + x_2I_2 + x_3I_3, Y = y_0 + y_1I_1 + y_2I_2 + y_3I_3 \in V_3(I)$ , the mapping  $f: V_3(I) \times V_3(I) \rightarrow R_3(I)$  such that:

$$\begin{aligned}
f(X, Y) &= g(x_0, y_0) \\
&\quad + (g(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3) - g(x_0 + x_2 + x_3, y_0 + y_2 + y_3))I_1 \\
&\quad + (g(x_0 + x_2 + x_3, y_0 + y_2 + y_3) - g(x_0 + x_3, y_0 + y_3))I_2 \\
&\quad + (g(x_0 + x_3, y_0 + y_3) - g(x_0, y_0))I_3
\end{aligned}$$

Is a 3-refined neutrosophic inner product.

**Proof.**

$$\begin{aligned}
f(X, X) &= g(x_0, x_0) \\
&\quad + (g(x_0 + x_1 + x_2 + x_3, x_0 + x_1 + x_2 + x_3) - g(x_0 + x_2 + x_3, x_0 + x_2 + x_3))I_1 \\
&\quad + (g(x_0 + x_2 + x_3, x_0 + x_2 + x_3) - g(x_0 + x_3, x_0 + x_3))I_2 \\
&\quad + (g(x_0 + x_3, x_0 + x_3) - g(x_0, x_0))I_3 \\
&= \|x_0\|^2 + (\|x_0 + x_1 + x_2 + x_3\|^2 - \|x_0 + x_2 + x_3\|^2)I_1 + (\|x_0 + x_2 + x_3\|^2 - \|x_0 + x_3\|^2)I_2 \\
&\quad + (\|x_0 + x_3\|^2 - \|x_0\|^2)I_3 \geq 0
\end{aligned}$$

According to the concept of partial ordering on  $R_3(I)$ .

$$f(X, X) = 0 \Leftrightarrow \|x_0\|^2 = \|x_0 + x_1 + x_2 + x_3\|^2 = \|x_0 + x_2 + x_3\|^2 = \|x_0 + x_3\|^2 = 0$$

Thus  $x_0 = x_1 = x_2 = x_3 = 0$  and  $X = 0$ .

It is clear that  $f(X, Y) = f(Y, X)$ .

Now, let  $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3, B = b_0 + b_1I_1 + b_2I_2 + b_3I_3 \in R_3(I)$  and  $Z = z_0 + z_1I_1 + z_2I_2 + z_3I_3 \in V_3(I)$ , we have:

$$\begin{aligned}
AX + BY &= (a_0x_0 + b_0y_0) + ((a_0 + a_1 + a_2 + a_3)(x_0 + x_1 + x_2 + x_3) - (a_0 + a_2 + a_3)(x_0 + \\
&\quad x_2 + x_3) + (b_0 + b_1 + b_2 + b_3)(y_0 + y_1 + y_2 + y_3) - (b_0 + b_2 + b_3)(y_0 + y_2 + y_3))I_1 + \\
&\quad ((a_0 + a_2 + a_3)(x_0 + x_2 + x_3) - (a_0 + a_3)(x_0 + x_3) + (b_0 + b_2 + b_3)(y_0 + y_2 + y_3) - \\
&\quad (b_0 + b_3)(y_0 + y_3))I_2 + ((a_0 + a_3)(x_0 + x_3) - a_0x_0 + (b_0 + b_3)(y_0 + y_3) - b_0y_0)I_3
\end{aligned}$$

$$\begin{aligned}
f(AX + BY, Z) &= g(a_0x_0 + b_0y_0, z_0) \\
&+ \left( g((a_0 + a_1 + a_2 + a_3)(x_0 + x_1 + x_2 + x_3) \right. \\
&+ (b_0 + b_1 + b_2 + b_3)(y_0 + y_1 + y_2 + y_3), z_0 + z_1 + z_2 + z_3) \\
&- g((a_0 + a_2 + a_3)(x_0 + x_2 + x_3) + (b_0 + b_2 + b_3)(y_0 + y_2 + y_3), z_0 + z_2 \\
&+ z_3) \Big) I_1 \\
&+ \left( g((a_0 + a_2 + a_3)(x_0 + x_2 + x_3) + (b_0 + b_2 + b_3)(y_0 + y_2 + y_3), z_0 + z_2 \right. \\
&+ z_3) - g((a_0 + a_3)(x_0 + x_3) + (b_0 + b_3)(y_0 + y_3), z_0 + z_3) \Big) I_2 \\
&+ \left( g((a_0 + a_3)(x_0 + x_3) + (b_0 + b_3)(y_0 + y_3), z_0 + z_3) \right. \\
&- g(a_0x_0 + b_0y_0, z_0) \Big) I_3 \\
&= (a_0 + a_1I_1 + a_2I_2 + a_3I_3)f(X, Z) + (b_0 + b_1I_1 + b_2I_2 + b_3I_3)f(X, Z)
\end{aligned}$$

**Theorem.**

Let  $f: V_3(I) \times V_3(I) \rightarrow R_3(I)$  be a 3-refined neutrosophic real inner product, then  $g: V \times V \rightarrow R$  such that:

$$g(x, y) = f(x + 0I_1 + 0I_2 + 0I_3, y + 0I_1 + 0I_2 + 0I_3) \text{ is a classical inner product on } V.$$

The proof is clear.

**Definition.**

Let  $X = x_0 + x_1I_1 + x_2I_2 + x_3I_3 \in V_3(I)$  and  $f: V_3(I) \times V_3(I) \rightarrow R_3(I)$  be a 3-refined neutrosophic real inner product, then:

1. If  $Y = y_0 + y_1I_1 + y_2I_2 + y_3I_3 \in V_3(I)$ , then  $X \perp Y$  if and only if  $f(X, Y) = 0$ .
2.  $\|X\|^2 = f(X, X)$ .

**Theorem.**

Let  $f$  be a 3-refined neutrosophic real inner product on  $V_3(I)$  and  $X = x_0 + x_1I_1 + x_2I_2 + x_3I_3, Y = y_0 + y_1I_1 + y_2I_2 + y_3I_3 \in V_3(I)$ , then:

- 1).  $X \perp Y$  if and only if  $\begin{cases} x_0 \perp y_0, x_0 + x_3 \perp y_0 + y_3 \\ x_0 + x_2 + x_3 \perp y_0 + y_2 + y_3 \\ x_0 + x_1 + x_2 + x_3 \perp y_0 + y_1 + y_2 + y_3 \end{cases}$
- 2).  $\|X\| = \|x_0\| + (\|x_0 + x_1 + x_2 + x_3\| - \|x_0 + x_2 + x_3\|)I_1 + (\|x_0 + x_2 + x_3\| - \|x_0 + x_3\|)I_2 + (\|x_0 + x_3\| - \|x_0\|)I_3$

**Proof.**

- 1).  $X \perp Y \Leftrightarrow f(X, Y) = 0 \Leftrightarrow g(x_0, y_0) = g(x_0 + x_3, y_0 + y_3) = g(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3) = g(x_0 + x_2 + x_3, y_0 + y_2 + y_3) = 0$ , hence the proof holds.



2). We put  $T = \|x_0\| + (\|x_0 + x_1 + x_2 + x_3\| - \|x_0 + x_2 + x_3\|)I_1 + (\|x_0 + x_2 + x_3\| - \|x_0 + x_3\|)I_2 + (\|x_0 + x_3\| - \|x_0\|)I_3$

We compute  $T^2 = \|x_0\|^2 + (\|x_0 + x_1 + x_2 + x_3\|^2 - \|x_0 + x_2 + x_3\|^2)I_1 + (\|x_0 + x_2 + x_3\|^2 - \|x_0 + x_3\|^2)I_2 + (\|x_0 + x_3\|^2 - \|x_0\|^2)I_3 = f(X, X) = \|X\|^2$ , thus  $T = \|X\|$ .

**Example.**

Let  $X = 3 + 2I_1 - I_2 - I_3$ ,  $x_0 = 3, x_1 = 2, x_2 = -1, x_3 = -1$ , then:

$$\|X\| = |3| + (|3| - |1|)I_1 + (|1| - |2|)I_2 + (|2| - |3|)I_3 = 3 + 2I_1 - I_2 - I_3$$

**Example.**

Let  $V = R^2$ ,  $V_3(I) = R_3^2(I)$ , take  $X = (1,1) + (2,1)I_1 + (3,-1)I_2 + (-1,4)I_3$ .

$x_0 = (1,1), \|x_0\| = \sqrt{2}$ ,  $x_0 + x_3 = (0,5), \|x_0 + x_3\| = 5$ ,  $x_0 + x_2 + x_3 = (3,4), \|x_0 + x_2 + x_3\| = 5$ ,  $x_0 + x_1 + x_2 + x_3 = (5,5), \|x_0 + x_1 + x_2 + x_3\| = 5\sqrt{2}$

$$\|X\| = \sqrt{2} + (5\sqrt{2} - 5)I_1 + (5 - 5)I_2 + (5 - \sqrt{2})I_3 = \sqrt{2} + (5\sqrt{2} - 5)I_1 + (5 - \sqrt{2})I_3$$

**Remark.**

$\forall X, Y \in V_3(I)$ , then:  $\|X\| \geq 0$ ,  $\|X + Y\| \leq \|X\| + \|Y\|$ .

**Theorem.**

Let  $X, Y \in V_3(I)$ , then  $|f(X, Y)| \leq \|X\| \cdot \|Y\|$ .

**Proof.**

$$\begin{aligned} f(X, Y) &= g(x_0, y_0) \\ &\quad + (g(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3) - g(x_0 + x_2 + x_3, y_0 + y_2 + y_3))I_1 \\ &\quad + (g(x_0 + x_2 + x_3, y_0 + y_2 + y_3) - g(x_0 + x_3, y_0 + y_3))I_2 \\ &\quad + (g(x_0 + x_3, y_0 + y_3) - g(x_0, y_0))I_3 \end{aligned}$$

According to Cauchy-Shwartz inequality on the space  $V$ , we have:

$$|g(x_0, y_0)| \leq \|x_0\| \cdot \|y_0\|, |g(x_0 + x_2 + x_3, y_0 + y_2 + y_3)| \leq \|x_0 + x_2 + x_3\| \cdot \|y_0 + y_2 + y_3\|$$

$$|g(x_0 + x_3, y_0 + y_3)| \leq \|x_0 + x_3\| \cdot \|y_0 + y_3\|$$

$$|g(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3)| \leq \|x_0 + x_1 + x_2 + x_3\| \cdot \|y_0 + y_1 + y_2 + y_3\|$$

Thus,  $|f(X, Y)| \leq \|X\| \cdot \|Y\|$ , according to the definition of partial order relation.

**Example.**

Take  $V_3(I) = R_3^2(I)$ ,  $X = (1,1) + (1,0)I_1 + (0,1)I_2, Y = (2,0) + (0,3)I_1 + (1,0)I_2 + (0,1)I_3$

$x_0 = (1,1), y_0 = (2,0), g(x_0, y_0) = 2, \|x_0\| = \sqrt{2}, \|y_0\| = 2$ ,  $x_0 + x_3 = (1,1), \|x_0 + x_3\| = \sqrt{2}, y_0 + y_3 = (2,1), \|y_0 + y_3\| = \sqrt{5}, g(x_0 + x_3, y_0 + y_3) = 3$ ,  $x_0 + x_2 + x_3 = (1,2), \|x_0 + x_2 + x_3\| = \sqrt{3}, y_0 + y_2 + y_3 = (3,1), \|y_0 + y_2 + y_3\| = \sqrt{10}$ ,  $x_0 + x_1 + x_2 + x_3 =$

$$(2,2), \|x_0 + x_1 + x_2 + x_3\| = 2\sqrt{2}, y_0 + y_1 + y_2 + y_3 = (3,4), \|y_0 + y_1 + y_2 + y_3\| = 5,$$

$$g(x_0 + x_2 + x_3, y_0 + y_2 + y_3) = 5, g(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3) = 14.$$

$$f(X, Y) = 2 + (14 - 5)I_1 + (5 - 3)I_2 + (3 - 2)I_3 = 2 + 9I_1 + 2I_2 + I_3$$

$$|f(X, Y)| = 2 + 9I_1 + 2I_2 + I_3$$

$$\begin{aligned} \|X\| &= \sqrt{2} + (2\sqrt{2} - \sqrt{5})I_1 + (\sqrt{5} - \sqrt{2})I_2 + (\sqrt{2} - \sqrt{2})I_3 \\ &= \sqrt{2} + (2\sqrt{2} - \sqrt{5})I_1 + (\sqrt{5} - \sqrt{2})I_2 \end{aligned}$$

$$\|X\| = 2 + (5 - \sqrt{10})I_1 + (\sqrt{10} - \sqrt{5})I_2 + (\sqrt{5} - \sqrt{2})I_3$$

$$\begin{aligned} \|X\| \cdot \|Y\| &= 2\sqrt{2} + (10\sqrt{2} - 5\sqrt{2})I_1 + (5\sqrt{2} - \sqrt{10})I_2 + (\sqrt{10} - 2\sqrt{2})I_3 \\ &= 2\sqrt{2} + 5\sqrt{2}I_1 + (5\sqrt{2} - \sqrt{10})I_2 + (\sqrt{10} - 2\sqrt{2})I_3 \end{aligned}$$

On the other hand, we have:

$$2 \leq 2\sqrt{2}, 2 + 1 = 3 \leq \sqrt{10}, 5 \leq 5\sqrt{2}, 14 \leq 10\sqrt{2}, \text{ hence } |f(X, Y)| \leq \|X\| \cdot \|Y\|$$

### The Foundations 3-Refined Number Theory

#### Definition.

Let  $Z_3(I) = \{a + bI_1 + cI_2 + dI_3; a, b, c, d \in \mathbb{Z}\}$  be a set. It is called the ring of 3-refined neutrosophic integers if  $I_i \cdot I_j = I_{\min(i,j)}$ ,  $I_i^2 = I_i$ ;  $1 \leq i \leq 3$ .

It is a special case of the n-refined neutrosophic ring with  $n = 3$ .

#### Definition.

Let  $X = x_0 + x_1I_1 + x_2I_2 + x_3I_3, Y = y_0 + y_1I_1 + y_2I_2 + y_3I_3, Z = z_0 + z_1I_1 + z_2I_2 + z_3I_3 \in Z_3(I)$ , we define:

1).  $X \setminus Y$  if there exists  $T = t_0 + t_1I_1 + t_2I_2 + t_3I_3 \in Z_3(I)$  such that  $T \cdot X = Y$ .

2).  $X \equiv Y \pmod{Z}$  if and only if  $Z \setminus X - Y$ .

3).  $X \geq Y$  if and only if 
$$\begin{cases} x_0 \geq y_0 \\ x_0 + x_1 + x_2 + x_3 \geq y_0 + y_1 + y_2 + y_3 \\ x_0 + x_2 + x_3 \geq y_0 + y_2 + y_3 \\ x_0 + x_3 \geq y_0 + y_3 \end{cases}$$

#### Theorem.

Let  $X, Y, Z$  be the previous 3-refined neutrosophic integers, then:

1.  $X \setminus Y$  if and only if  $x_0 \setminus y_0, x_0 + x_1 + x_2 + x_3 \setminus y_0 + y_1 + y_2 + y_3, x_0 + x_2 + x_3 \setminus y_0 + y_2 + y_3, x_0 + x_3 \setminus y_0 + y_3$ .
2. If  $X \setminus Y$ , then  $X \leq Y$ .

3.  $X \equiv Y(\text{mod } Z)$  if and only if

$$\begin{cases} x_0 \equiv y_0(\text{mod } z_0) \\ x_0 + x_1 + x_2 + x_3 \equiv y_0 + y_1 + y_2 + y_3(\text{mod } z_0 + z_1 + z_2 + z_3) \\ x_0 + x_2 + x_3 \equiv y_0 + y_2 + y_3(\text{mod } z_0 + z_2 + z_3) \\ x_0 + x_3 \equiv y_0 + y_3(\text{mod } z_0 + z_3) \end{cases}$$

**Proof.**

1. Assume that  $X \setminus Y$ , this is true if and only if there exists  $T = t_0 + t_1I_1 + t_2I_2 + t_3I_3 \in Z_3(I)$  such that  $Y = X.T$ .

We have:

$$\begin{aligned} X.T &= (x_0 + x_1I_1 + x_2I_2 + x_3I_3)(t_0 + t_1I_1 + t_2I_2 + t_3I_3) \\ &= x_0t_0 + (x_0t_1 + x_1t_0 + x_1t_1 + x_1t_2 + x_1t_3 + x_2t_1 + x_3t_1)I_1 \\ &\quad + (x_0t_2 + x_2t_0 + x_2t_2 + x_3t_2)I_2 + (x_0t_3 + x_3t_0 + x_3t_3)I_3 \\ &= y_0 + y_1I_1 + y_2I_2 + y_3I_3 \end{aligned}$$

Thus:

$$\begin{cases} y_0 = x_0t_0 \dots (1) \\ y_1 = x_0t_1 + x_1t_0 + x_1t_1 + x_1t_2 + x_1t_3 + x_2t_1 + x_3t_1 \dots (2) \\ y_2 = x_0t_2 + x_2t_0 + x_2t_2 + x_3t_2 \dots (3) \\ y_3 = x_0t_3 + x_3t_0 + x_3t_3 \dots (4) \end{cases}$$

We add (1) to (4), (1) to (2) to (4), (1) to (2) to (3) to (4).

$$\begin{cases} y_0 = x_0t_0 \\ y_0 + y_3 = (x_0 + x_3)(t_0 + t_3) \\ y_0 + y_2 + y_3 = (x_0 + x_2 + x_3)(t_0 + t_2 + t_3) \\ y_0 + y_1 + y_2 + y_3 = (x_0 + x_1 + x_2 + x_3)(t_0 + t_1 + t_2 + t_3) \end{cases}$$

Thus, the proof of (1) is complete.

2. If  $X \setminus Y$ , then  $x_0 \setminus y_0$ , so  $x_0 \leq y_0$ .

Also:

$$\begin{cases} x_0 + x_3 \setminus y_0 + y_3, \text{ so } x_0 + x_3 \leq y_0 + y_3 \\ x_0 + x_2 + x_3 \setminus y_0 + y_2 + y_3, \text{ so } x_0 + x_2 + x_3 \leq y_0 + y_2 + y_3 \\ x_0 + x_1 + x_2 + x_3 \setminus y_0 + y_1 + y_2 + y_3, \text{ so } x_0 + x_1 + x_2 + x_3 \leq y_0 + y_1 + y_2 + y_3 \end{cases}$$

Thus  $X \leq Y$ .

3.  $X \equiv Y(\text{mod } Z)$  if and only if  $Z \setminus X - Y$ , thus:

$$\begin{cases} z_0 \setminus x_0 - y_0 \\ z_0 + z_3 \setminus (x_0 + x_3) - (y_0 + y_3) \\ z_0 + z_2 + z_3 \setminus (x_0 + x_2 + x_3) - (y_0 + y_2 + y_3) \\ z_0 + z_1 + z_2 + z_3 \setminus (x_0 + x_1 + x_2 + x_3) - (y_0 + y_1 + y_2 + y_3) \end{cases}$$

Thus  $x_0 \equiv y_0(\text{mod } z_0)$ ,  $x_0 + x_3 \equiv y_0 + y_3(\text{mod } z_0 + z_3)$ ,  $x_0 + x_2 + x_3 \equiv y_0 + y_2 + y_3(\text{mod } z_0 + z_2 + z_3)$ ,  $x_0 + x_1 + x_2 + x_3 \equiv y_0 + y_1 + y_2 + y_3(\text{mod } z_0 + z_1 + z_2 + z_3)$ .

**Example.**

Take  $X = 3 + 2I_1 + I_2 - I_3, Y = 3 + 4I_1 + 2I_2 + I_3$ , we have  $X \setminus Y$  that is because:

$$3 \setminus 3, 3 - 1 = 2 \setminus 3 + 1 = 4, 3 + 1 - 1 = 3 \setminus 3 + 2 + 1 = 6, 3 + 2 + 1 - 1 \setminus 3 + 4 + 2 + 1 = 10.$$

**Example.**

Take  $X = 7 + 3I_1 + I_2 + 5I_3, Y = 4 + I_1 + I_2 + I_3, Z = 3 + 2I_1 + I_3$ , we have  $7 \equiv 4(\text{mod } 3)$ ,  $7 + 5 = 12 \equiv 4 + 1(\text{mod } 3 + 4)$ ,  $7 + 1 + 5 = 13 \equiv 4 + 1 + 1(\text{mod } 3 + 0 + 4)$ ,  $7 + 3 + 1 + 5 = 16 \equiv 4 + 1 + 1 + 1(\text{mod } 9)$  thus,  $X \equiv Y(\text{mod } Z)$ .

**Theorem.**

The relation  $(\leq)$  is a partial order relation.

**Proof.**

$X \leq Y$  clearly.

If  $X \leq Y$  and  $Y \leq Z$ , then  $x_0 \leq y_0 \leq z_0, x_0 + x_3 \leq y_0 + y_3 \leq z_0 + z_3, x_0 + x_2 + x_3 \leq y_0 + y_2 + y_3 \leq z_0 + z_2 + z_3, x_0 + x_1 + x_2 + x_3 \leq y_0 + y_1 + y_2 + y_3 \leq z_0 + z_1 + z_2 + z_3$ .

Thus  $X \leq Z$ .

If  $X \leq Y$  and  $Y \leq X$ , then  $x_0 = y_0, x_0 + x_3 = y_0 + y_3, x_0 + x_2 + x_3 = y_0 + y_2 + y_3, x_0 + x_1 + x_2 + x_3 = y_0 + y_1 + y_2 + y_3$ , thus  $X = Y$ .

**Theorem.**

Let  $X = x_0 + x_1I_1 + x_2I_2 + x_3I_3, Y = y_0 + y_1I_1 + y_2I_2 + y_3I_3, Z = z_0 + z_1I_1 + z_2I_2 + z_3I_3, T = t_0 + t_1I_1 + t_2I_2 + t_3I_3, S = s_0 + s_1I_1 + s_2I_2 + s_3I_3 \in Z_3(I)$ , then:

1). If  $X \equiv Y(\text{mod } Z)$ ,  $T \equiv S(\text{mod } Z)$ , then  $X + T \equiv Y + S(\text{mod } Z)$  and  $X - T \equiv Y - S(\text{mod } Z), X.T \equiv Y.S(\text{mod } Z)$ .

2).  $X^n = x_0^n + [(x_0 + x_1 + x_2 + x_3)^n - (x_0 + x_2 + x_3)^n]I_1 + [(x_0 + x_2 + x_3)^n - (x_0 + x_3)^n]I_2 + [(x_0 + x_3)^n - x_0^n]I_3; n \in N$ .

3).  $X^n \equiv Y^n(\text{mod } Z^n); n \in N$ .

**Proof.**

1).  $X + T = (x_0 + t_0) + (x_1 + t_1)I_1 + (x_2 + t_2)I_2 + (x_3 + t_3)I_3$ .

$Y + S = (y_0 + s_0) + (y_1 + s_1)I_1 + (y_2 + s_2)I_2 + (y_3 + s_3)I_3$ .

Since  $z_0 \setminus x_0 - y_0, z_0 \setminus t_0 - s_0$ , then  $z_0 \setminus (x_0 + t_0) - (y_0 + s_0)$  and  $x_0 + t_0 \equiv y_0 + s_0(\text{mod } z_0)$ .

$z_0 + z_3 \setminus (x_0 + x_3) - (y_0 + y_3), z_0 \setminus (t_0 + t_3) - (s_0 + s_3)$ , then:

$z_0 + z_3 \setminus (x_0 + x_3 + t_0 + t_3) - (y_0 + y_3 + s_0 + s_3)$ , thus:

$$(x_0 + x_3) + (t_0 + t_3) \equiv (y_0 + y_3) + (s_0 + s_3) \pmod{z_0 + z_3}$$

By a similar discussion, we get:

$$z_0 + z_2 + z_3 \setminus (x_0 + x_2 + x_3 + t_0 + t_2 + t_3) - (y_0 + y_2 + y_3 + s_0 + s_2 + s_3)$$

$$z_0 + z_1 + z_2 + z_3 \setminus (x_0 + x_1 + x_2 + x_3 + t_0 + t_1 + t_2 + t_3) \\ - (y_0 + y_1 + y_2 + y_3 + s_0 + s_1 + s_2 + s_3)$$

So that  $X + T \equiv Y + S \pmod{Z}$ .

It is easy check that  $X - T \equiv Y - S \pmod{Z}, X.T \equiv Y.S \pmod{Z}$ .

2). For  $n = 1$  it is true clearly.

Assume that it is true for  $n = k$ , we must prove it for  $n = k + 1$ .

$$\begin{aligned} X^{k+1} &= X.X^k = [x_0 + x_1I_1 + x_2I_2 + x_3I_3][x_0^n + [(x_0 + x_1 + x_2 + x_3)^n - (x_0 + x_2 + x_3)^n]I_1 \\ &\quad + [(x_0 + x_2 + x_3)^n - (x_0 + x_3)^n]I_2 + [(x_0 + x_3)^n - x_0^n]I_3] \\ &= x_0^{n+1} + [x_0(x_0 + x_1 + x_2 + x_3)^n - x_0(x_0 + x_2 + x_3)^n + x_1(x_0 + x_1 + x_2 + x_3)^n \\ &\quad - x_1(x_0 + x_2 + x_3)^n + x_1x_0^n + x_1(x_0 + x_2 + x_3)^n - x_1(x_0 + x_3)^n \\ &\quad + x_1(x_0 + x_3)^n - x_1x_0^n + x_2(x_0 + x_1 + x_2 + x_3)^n - x_2(x_0 + x_2 + x_3)^n \\ &\quad + x_3(x_0 + x_1 + x_2 + x_3)^n - x_3(x_0 + x_2 + x_3)^n]I_1 \\ &\quad + [x_0(x_0 + x_2 + x_3)^n - x_0(x_0 + x_3)^n + x_2x_0^n + x_2(x_0 + x_2 + x_3)^n \\ &\quad - x_2(x_0 + x_3)^n + x_2(x_0 + x_3)^n - x_2x_0^n + x_3(x_0 + x_2 + x_3)^n - x_3(x_0 + x_3)^n]I_2 \\ &\quad + [x_0(x_0 + x_3)^n - x_0^{n+1} + x_3x_0^n + x_3(x_0 + x_3)^n - x_3x_0^n]I_3 \\ &= x_0^{n+1} + [(x_0 + x_1 + x_2 + x_3)^{n+1} - (x_0 + x_2 + x_3)^{n+1}]I_1 \\ &\quad + [(x_0 + x_2 + x_3)^{n+1} - (x_0 + x_3)^{n+1}]I_2 + [(x_0 + x_3)^{n+1} - x_0^{n+1}]I_3 \end{aligned}$$

This implies that is true by induction.

3). It holds directly from (1) and (2).

**Example.**

Take  $X = 1 + 2I_1 - I_2 + I_3, n = 2$ , then:

$$X^2 = 1 + [(3)^2 - 1]I_1 + [1 - (2)^2]I_2 + [(2)^2 - 1]I_3 = 1 + 8I_1 - 3I_2 + 3I_3$$

**Theorem.**

Let  $X, Y \in Z_3(I)$ , then  $gcd(X, Y) = gcd(x_0, y_0) + [gcd(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3) - gcd(x_0 + x_2 + x_3, y_0 + y_2 + y_3)]I_1 + [gcd(x_0 + x_2 + x_3, y_0 + y_2 + y_3) - gcd(x_0 + x_3, y_0 + y_3)]I_2 + [gcd(x_0 + x_3, y_0 + y_3) - gcd(x_0, y_0)]I_3$

**Example.**

Take  $X = 4 + 3I_1 + 5I_2 - I_3, Y = 7 + I_1 + I_2 + 3I_3$ .

$gcd(4,7) = 1, gcd(11,13) = 1, gcd(8,11) = 1, gcd(3,10) = 1$ , thus  $gcd(X,Y) = 1$ .

**Remark.**

$X, Y$  are called coprime (relatively prime) if and only if  $gcd(X, Y) = 1$ , which is equivalent to:

$$gcd(x_0, y_0) = gcd(x_0 + x_3, y_0 + y_3) = gcd(x_0 + x_2 + x_3, y_0 + y_2 + y_3) = gcd(x_0 + x_1 + x_2 + x_3, y_0 + y_1 + y_2 + y_3) = 1.$$

**Definition.**

Let  $X = x_0 + x_1I_1 + x_2I_2 + x_3I_3 \in Z_3(I)$ , with  $X > 0$ , we define:

$$\begin{aligned} \varphi_s(X) = & \varphi(x_0) + [\varphi(x_0 + x_1 + x_2 + x_3) - \varphi(x_0 + x_2 + x_3)]I_1 \\ & + [\varphi(x_0 + x_2 + x_3) - \varphi(x_0 + x_3)]I_2 + [\varphi(x_0 + x_3) - \varphi(x_0)]I_3 \end{aligned}$$

where  $\varphi$  is the ordinary Euler's function,  $\varphi_s$  is called the special 3-refined neutrosophic Euler's function.

**Example.**

Take  $X = 3 + I_1 + I_2 + I_3 > 0$ ;  $x_0 = 3, x_1 = 1, x_2 = x_3 = 1$ .

$$\varphi(x_0) = 2, \varphi(x_0 + x_1 + x_2 + x_3) = 2, \varphi(x_0 + x_2 + x_3) = 4, \varphi(x_0 + x_3) = 2.$$

Thus

$$\varphi_s(X) = 2 + [2 - 4]I_1 + [4 - 2]I_2 + [2 - 2]I_3 = 2 - 2I_1 + 2I_2 + 0I_3.$$

It is clear that  $\varphi_s(X) > 0$ ;  $\forall X > 0$ .

**Theorem.**

Let  $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3, M = m_0 + m_1I_1 + m_2I_2 + m_3I_3 \in Z_3(I)$ , such that:

$A > 0, M > 0$  and  $gcd(A, M) = 1$ , then:

- 1).  $A^{\varphi_s(M)} \equiv 1 \pmod{M}$ .
- 2).  $A^{-1} \pmod{M} \equiv a_0^{-1} \pmod{m_0} + [(a_0 + a_1 + a_2 + a_3)^{-1} \pmod{m_0 + m_1 + m_2 + m_3} - (a_0 + a_2 + a_3)^{-1} \pmod{m_0 + m_2 + m_3}]I_1 + [(a_0 + a_2 + a_3)^{-1} \pmod{m_0 + m_2 + m_3} - (a_0 + a_3)^{-1} \pmod{m_0 + m_3}]I_2 + [(a_0 + a_3)^{-1} \pmod{m_0 + m_3} - a_0^{-1} \pmod{m_0}]I_3$ .

**Proof.**

- 1). 
$$A^{\varphi_s(M)} = a_0^{\varphi(m_0)} + [(a_0 + a_1 + a_2 + a_3)^{\varphi(m_0+m_1+m_2+m_3)} - (a_0 + a_2 + a_3)^{\varphi(m_0+m_2+m_3)}]I_1 + [(a_0 + a_2 + a_3)^{\varphi(m_0+m_2+m_3)} - (a_0 + a_3)^{\varphi(m_0+m_3)}]I_2 + [(a_0 + a_3)^{\varphi(m_0+m_3)} - a_0^{\varphi(m_0)}]I_3 \equiv 1 \pmod{M}.$$

- 2). It holds directly by computing the product  $AA^{-1}$ .

**3-refined Diophantine equations:**

**Definition.**

Let  $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3, B = b_0 + b_1I_1 + b_2I_2 + b_3I_3, C = c_0 + c_1I_1 + c_2I_2 + c_3I_3, X = x_0 + x_1I_1 + x_2I_2 + x_3I_3, Y = y_0 + y_1I_1 + y_2I_2 + y_3I_3$ , where  $a_i, b_i, c_i, x_i, y_i \in Z_3(I)$ .

We define the 3-refined neutrosophic linear Diophantine equation with two variables as follows:

$$AX + BY = C.$$

**Example.**

Consider the following 3-refined neutrosophic linear Diophantine equation:

$$(3 + 2I_1 + I_2 + I_3)X + (2 + 4I_2)Y = 3 + 9I_1 - 7I_3$$

**Theorem.**

Let  $AX + BY = C$  be a 3-refined neutrosophic linear Diophantine equation, then it is equivalent to:

$$\begin{cases} a_0x_0 + b_0y_0 = c_0 \\ (a_0 + a_3)(x_0 + x_3) + (b_0 + b_3)(y_0 + y_3) = c_0 + c_3 \\ (a_0 + a_2 + a_3)(x_0 + x_2 + x_3) + (b_0 + b_2 + b_3)(y_0 + y_2 + y_3) = c_0 + c_2 + c_3 \\ (a_0 + a_1 + a_2 + a_3)(x_0 + x_1 + x_2 + x_3) + (b_0 + b_1 + b_2 + b_3)(y_0 + y_1 + y_2 + y_3) = c_0 + c_1 + c_2 + c_3 \end{cases}$$

**Proof.**

We compute  $AX = a_0x_0 + [(a_0 + a_1 + a_2 + a_3)(x_0 + x_1 + x_2 + x_3) - (a_0 + a_2 + a_3)(x_0 + x_2 + x_3)]I_1 + [(a_0 + a_2 + a_3)(x_0 + x_2 + x_3) - (a_0 + a_3)(x_0 + x_3)]I_2 + [(a_0 + a_3)(x_0 + x_3) - a_0x_0]I_3$

On the other hand, we have:

$$\begin{aligned} BY &= b_0y_0 + [(b_0 + b_1 + b_2 + b_3)(y_0 + y_1 + y_2 + y_3) - (b_0 + b_2 + b_3)(y_0 + y_2 + y_3)]I_1 \\ &\quad + [(b_0 + b_2 + b_3)(y_0 + y_2 + y_3) - (b_0 + b_3)(y_0 + y_3)]I_2 \\ &\quad + [(b_0 + b_3)(y_0 + y_3) - b_0y_0]I_3 \end{aligned}$$

The equation  $AX + BY = C$  equivalents:

$$\begin{cases} a_0x_0 + b_0y_0 = c_0 \\ (a_0 + a_3)(x_0 + x_3) + (b_0 + b_3)(y_0 + y_3) = c_0 + c_3 \\ (a_0 + a_2 + a_3)(x_0 + x_2 + x_3) + (b_0 + b_2 + b_3)(y_0 + y_2 + y_3) = c_0 + c_2 + c_3 \\ (a_0 + a_1 + a_2 + a_3)(x_0 + x_1 + x_2 + x_3) + (b_0 + b_1 + b_2 + b_3)(y_0 + y_1 + y_2 + y_3) = c_0 + c_1 + c_2 + c_3 \end{cases}$$

**Example.**

Find a solution of the equation:

$$(3 + 2I_1 + I_2 + I_3)X + (2 + 4I_2)Y = 3 + 9I_1 - 7I_3$$

We have  $a_0 = 3, a_1 = 2, a_2 = 1, a_3 = 1, b_0 = 2, b_1 = 0, b_2 = 4, b_3 = 0, c_0 = 3, c_1 = 9, c_2 = 0, c_3 = -7$

The equivalent system is:

$$\begin{cases} 3x_0 + 2y_0 = 3 \dots (1) \\ 4(x_0 + x_3) + 2(y_0 + y_3) = -4 \dots (2) \\ 5(x_0 + x_2 + x_3) + 7(y_0 + y_2 + y_3) = -4 \dots (3) \\ 7(x_0 + x_1 + x_2 + x_3) + 6(y_0 + y_1 + y_2 + y_3) = 5 \dots (4) \end{cases}$$

The equation (1) has a solution  $x_0 = 1, y_0 = 0$ .

The equation (2) has a solution  $x_0 + x_3 = -1, y_0 + y_3 = 0$ , thus  $x_3 = -2, y_3 = 0$ .

The equation (3) has a solution  $x_0 + x_2 + x_3 = 9, y_0 + y_2 + y_3 = -7$ , thus  $x_2 = 10, y_2 = -7$ .

The equation (4) has a solution  $x_0 + x_1 + x_2 + x_3 = 5, y_0 + y_1 + y_2 + y_3 = 5$ , thus  $x_1 = -4, y_1 = 12$ .

This means that  $X = 1 - 4I_1 + 10I_2 - 2I_3, Y = 12I_1 - 7I_2$ .

### Future research directions and suggestions

3-refined neutrosophic number as generalizations of classical real numbers and integers, may have a great impact on many areas of scientific knowledge.

In the following, we suggest many possible applications of 3-refined neutrosophic real numbers.

- 1-) How can we build a crypto-system from 3-refined neutrosophic integers which generalize RSA algorithm. [19]
- 2-) How can we build a crypto-system from 3-refined neutrosophic integers which generalize El-Gamal algorithm. [20-21]
- 3-) How can we solve 3-refined neutrosophic differential equations, and integral equations.
- 4-) How can we define Hillbert and Banach 3-refined neutrosophic spaces, and do classical functional inequalities still true in this case.

### References

- [1] Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona 2006.
- [2] Celik, M., and Olgun, N., " An Introduction To Neutrosophic Real Banach And Hillbert Spaces", Galoitica Journal Of Mathematical Structures And Applications, 2022.
- [3] Celik, M., and Olgun, N., " On The Classification Of Neutrosophic Complex Inner Product Spaces", Galoitica Journal Of Mathematical Structures And Applications, 2022.



- 
- [4] Celik, M., and Hatip, A., " On The Refined AH-Isometry And Its Applications In Refined Neutrosophic Surfaces", *Galoitica Journal Of Mathematical Structures And Applications*, 2022
- [5] Merkepci, H., and Ahmad, K., " On The Conditions Of Imperfect Neutrosophic Duplets and Imperfect Neutrosophic Triplets", *Galoitica Journal Of Mathematical Structures And Applications*, Vol.2, 2022.
- [6] Agboola, A.A.A., Akwu, A.D., and Oyebo, Y.T., " Neutrosophic Groups and Subgroups", *International J .Math. Combin*, Vol. 3, pp. 1-9. 2012.
- [7] Smarandache, F., "  $n$ -Valued Refined Neutrosophic Logic and Its Applications in Physics", *Progress in Physics*, 143-146, Vol. 4, 2013.
- [8] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, *Journal Of Mathematics*, Hindawi, 2021
- [9] Olgun, N., Hatip, A., Bal, M., and Abobala, M., " A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", *International Journal of neutrosophic Science*, Vol. 16, pp. 72-79, 2021.
- [10] Abobala, M., Hatip, A., and Bal, M., " A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol.17, 2021.
- [11] Abobala, M., Ziena, B,M., Doewes, R., and Hussein, Z., "The Representation Of Refined Neutrosophic Matrices By Refined Neutrosophic Linear Functions", *International Journal Of Neutrosophic Science*, 2022.
- [12] Ahmad, K., Thjeel, N., AbdulZahra, M., and Jaleel, R., " On The Classification of  $n$ -Refined Neutrosophic Rings And Its Applications In Matrix Computing Algorithms And Linear Systems", *International Journal Of Neutrosophic Science*, 2022.
- [13] Hatip, A., "An Introduction To Weak Fuzzy Complex Numbers ", *Galoitica Journal Of Mathematical Structures and Applications*, Vol.3, 2023.

- 
- [14] Merkepci, H., and Abobala, M., " The Application of AH-isometry In The Study Of Neutrosophic Conic Sections", *Galoitica Journal Of Mathematical Structures And Applications*, Vol.2, 2022.
- [15] Abobala, M., and Zeina, M.B., " A Study Of Neutrosophic Real Analysis By Using One Dimensional Geometric AH-Isometry", *Galoitica Journal Of Mathematical Structures And Applications*, Vol.3, 2023.
- [16] Khaldi, A., " A Study On Split-Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [17] Ahmad, K., " On Some Split-Complex Diophantine Equations", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [18] Ali, R., " On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [19] Merkepci, M., Abobala, M., and Allouf, A., " The Applications of Fusion Neutrosophic Number Theory in Public Key Cryptography and the Improvement of RSA Algorithm ", *Fusion: Practice and Applications*, 2023.
- [20] Merkepci, M.; Sarkis, M. An Application of Pythagorean Circles in Cryptography and Some Ideas for Future Non Classical Systems. *Galoitica Journal of Mathematical Structures and Applications* **2022**.
- [21] Merkepci, M., and Abobala, M., " Security Model for Encrypting Uncertain Rational Data Units Based on Refined Neutrosophic Integers Fusion and El Gamal Algorithm ", *Fusion: Practice and Applications*, 2023.
- [22] Hatip, A., " On Intuitionistic Fuzzy Subgroups of (M-N) Type and Their Algebraic Properties", *Galoitica Journal Of Mathematical Structures and Applications*, Vol.4, 2023.
- [23] Abobala, M., "Neutrosophic Real Inner Product Spaces", *Neutrosophic Sets and Systems*, Vol. 43, 2021
- [24] Agboola, A.A.A., "On Refined Neutrosophic Algebraic Structures", *Neutrosophic Sets and Systems*, Vol.10, pp. 99-101, 2015.

- 
- [25] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., " Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81, 2020.
- [26] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., " Refined Neutrosophic Rings II", International Journal of Neutrosophic Science, Vol. 2(2), pp. 89-94, 2020.
- [27] Abobala, M., and Hatip, A., " An Algebraic Approach To Neutrosophic Euclidean Geometry", Neutrosophic Sets and Systems, 2021.
- [28] Hatip, A., and Olgun, N., " On Refined Neutrosophic R-Module", International Journal of Neutrosophic Science, Vol. 7, pp.87-96, 2020.
- [29] Smarandache, F., and Abobala, M., "n-Refined Neutrosophic Rings", International Journal Of Neutrosophic Science, 2020.
- [30] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [31] Sarkis, M., " On The Solutions Of Fermat's Diophantine Equation In 3-refined Neutrosophic Ring of Integers", Neoma Journal of Mathematics and Computer Science, 2023.

**Received:** December 16, 2022. **Accepted:** April 03, 2023