

4-11-2023

An Introduction to Symbolic 2-Plithogenic Vector Spaces Generated from The Fusion of Symbolic Plithogenic Sets and Vector Spaces

Nader Mahmoud Taffach

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Taffach, Nader Mahmoud. "An Introduction to Symbolic 2-Plithogenic Vector Spaces Generated from The Fusion of Symbolic Plithogenic Sets and Vector Spaces." *Neutrosophic Sets and Systems* 54, 1 (). https://digitalrepository.unm.edu/nss_journal/vol54/iss1/4

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



An Introduction to Symbolic 2-Plithogenic Vector Spaces Generated from The Fusion of Symbolic Plithogenic Sets and Vector Spaces

Nader Mahmoud Taffach

1. Faculty Of Science, Department Of Mathematics, Idleb University, Syria

ntaffash77@windowslive.com

Abstract:

The fusion of symbolic plithogenic sets with algebraic structures generates novel algebraic neutrosophic structures that generalize the classical known structures. The objective of this paper is to define the concept of symbolic 2-plithogenic vector space over a symbolic 2-plithogenic field.

Concepts such as AH-subspace and AH-linear transformation will be presented and discussed in terms of theorems.

Keywords: 2-plithogenic symbolic set, 2-plithogenic vector space, 2-plithogenic dimension

Introduction

The concept of symbolic plithogenic sets was defined by Smarandache in [13-17,30], and he suggested an algebraic approach of these sets. Laterally, the concept of symbolic 2-plithogenic rings [31], where the concepts such as symbolic AH-ideals, and AH-homomorphisms were presented and discussed.

In general, we can say that symbolic plithogenic structures are very close to neutrosophic algebraic structures with many differences in the definition of multiplication operation [1-10].

Let R be a ring, the symbolic 2-plithogenic ring is defined as follows:

$$2 - SP_R = \{a_0 + a_1P_1 + a_2P_2; a_i \in R, P_j^2 = P_j, P_1 \times P_2 = P_{\max(1,2)} = P_2\}.$$

Smarandache has defined algebraic operations on $2 - SP_R$ as follows:

Addition:

$$[a_0 + a_1P_1 + a_2P_2] + [b_0 + b_1P_1 + b_2P_2] = (a_0 + b_0) + (a_1 + b_1)P_1 + (a_2 + b_2)P_2.$$

Multiplication:

$$[a_0 + a_1P_1 + a_2P_2] \cdot [b_0 + b_1P_1 + b_2P_2] = a_0b_0 + a_0b_1P_1 + a_0b_2P_2 + a_1b_0P_1^2 + a_1b_2P_1P_2 + a_2b_0P_2 + a_2b_1P_1P_2 + a_2b_2P_2^2 + a_1b_1P_1P_1 = a_0b_0 + (a_0b_1 + a_1b_0 + a_1b_1)P_1 + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)P_2.$$

It is clear that $(2 - SP_R)$ is a ring.

Also, if R is commutative, then $2 - SP_R$ is commutative, and if R has a unity (1), then $2 - SP_R$ has the same unity (1).

If R is a field, then $2 - SP_R$ is called a symbolic 2-plithogenic field.

In this paper, we study the symbolic 2-plithogenic vector spaces according to many points of view, where substructures such as AH-subspaces, and AH-linear transformations will be presented in terms of theorems. In addition, many examples will be illustrated to explain the novelty of these ideas.

Main Discussion

Definition.

Let V be a vector space over the field F , let $2 - SP_F$ be the corresponding symbolic 2-plithogenic field.

$$2 - SP_F = \{x + yP_1 + zP_2; x, y, z \in F, P_i^2 = P_i, P_1P_2 = P_2P_1 = P_2\}.$$

We define the symbolic 2-plithogenic vector space as follows:

$$2 - SP_V = V + VP_1 + VP_2 = \{a + bP_1 + cP_2; a, b, c \in V\}.$$

Operations on $2 - SP_V$ can be defined as follows:

Addition: (+): $2 - SP_V \rightarrow 2 - SP_V$, such that:

$$[x_0 + x_1P_1 + x_2P_2] + [y_0 + y_1P_1 + y_2P_2] = (x_0 + y_0) + (x_1 + y_1)P_1 + (x_2 + y_2)P_2$$

Multiplication: (.): $2 - SP_F \times 2 - SP_V \rightarrow 2 - SP_V$, such that:

$$\begin{aligned} [a + bP_1 + cP_2] \cdot [x_0 + x_1P_1 + x_2P_2] \\ = ax_0 + (ax_1 + bx_0 + bx_1)P_1 + (ax_2 + bx_2 + cx_0 + cx_1 + cx_2)P_2 \end{aligned}$$

where $x_i, y_i \in V, a, b, c \in F$

Theorem.

Let $(2 - SP_V, +, \cdot)$ Is a module over the ring $2 - SP_F$.

Proof.

Let $X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2 \in 2 - SP_V$, $A = a_0 + a_1P_1 + a_2P_2, B = b_0 + b_1P_1 + b_2P_2 \in 2 - SP_F$ we have:

$$1. X = X, (X + Y) + Z = X + (Y + Z), X + (-X) = -X + X = 0, X + 0 = 0 + X = X$$

Also

$$\begin{aligned} A(X + Y) &= (a_0 + a_1P_1 + a_2P_2)[(x_0 + y_0) + (x_1 + y_1)P_1 + (x_2 + y_2)P_2] \\ &= a_0(x_0 + y_0) + (a_0(x_1 + y_1) + a_1(x_0 + y_0) + a_1(x_1 + y_1))P_1 \\ &\quad + (a_0(x_2 + y_2) + a_1(x_2 + y_2) + a_2(x_0 + y_0) + a_2(x_1 + y_1) + a_2(x_2 + y_2))P_2 \\ &= A.X + A.Y \end{aligned}$$

$$\begin{aligned} (A + B)X &= [(a_0 + b_0) + (a_1 + b_1)P_1 + (a_2 + b_2)P_2](x_0 + x_1P_1 + x_2P_2) \\ &= (a_0 + b_0)x_0 + ((a_0 + b_0)x_1 + (a_1 + b_1)x_0 + (a_1 + b_1)x_1)P_1 \\ &\quad + ((a_0 + b_0)x_2 + (a_1 + b_1)x_2 + (a_2 + b_2)x_0 + (a_2 + b_2)x_1 + (a_2 + b_2)x_2)P_2 \\ &= A.X + B.X \end{aligned}$$

$$\begin{aligned} (A.B).X &= [a_0b_0 + (a_0b_1 + a_1b_0 + a_1b_1)P_1 + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)P_2](x_0 \\ &\quad + x_1P_1 + x_2P_2) \\ &= a_0b_0x_0 + [a_0b_0x_1 + (a_0b_1 + a_1b_0 + a_1b_1)x_0 + (a_0b_1 + a_1b_0 + a_1b_1)x_1]P_1 \\ &\quad + [a_0b_0x_2 + (a_0b_2 + a_2b_0 + a_1b_1)x_2 + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)x_0 \\ &\quad + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)x_1 \\ &\quad + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)x_2]P_2 = A(B.X) \end{aligned}$$

Example.

Let $V = R^3$ be the Euclidean space over the field $F = R$.

The corresponding symbolic 2-plithogenic vector space over $2 - SP_F$ is:

$$2 - SP_{R^3} = \{(x_0, y_0, z_0) + (x_1, y_1, z_1)P_1 + (x_2, y_2, z_2)P_2; x_i, y_i, z_i \in R\}$$

Consider $X = (1,1,0) + (2, -1,1)P_1 + (0,1, -1)P_2 \in 2 - SP_{R^3}, A = 2 + P_1 + P_2 \in 2 - SP_R$. We have:

$$\begin{aligned} A.X &= (2,2,0) + [(4, -2,2) + (1,1,0) + (2, -1,1)]P_1 \\ &\quad + [(0,2,2) + (0,1,1) + (1,1,0) + (2, -1,1) + (0,1,1)]P_2 \\ &= (2,2,0) + (7, -2,3)P_1 + (3,4,5)P_2 \end{aligned}$$

Definition.

Let $2 - SP_V$ be a symbolic 2-plithogenic vector space over $2 - SP_F$, let V_0, V_1, V_2 be the three subspaces of V , we define the AH-subspace as follows:

$$W = V_0 + V_1P_1 + V_2P_2 = \{x + yP_1 + zP_2; x \in V_0, y \in V_1, z \in V_2\}$$

If $V_0 = V_1 = V_2$, then W is called an AHS-subspace.

Example.

Consider $2 - SP_{R^3}$, we have $V_0 = \{(a, 0, 0); a \in R\}, V_1 = \{(0, b, 0); b \in R\}, V_2 = \{(0, 0, c); c \in R\}$ are three subspaces of $V = R^3$.

$W = V_0 + V_1P_1 + V_2P_2 = \{(a, 0, 0) + (0, b, 0)P_1 + (0, 0, c)P_2; a, b, c \in R\}$ is an AH-subspace of $2 - SP_{R^3}$.

$T = V_1 + V_1P_1 + V_1P_2 = \{(0, a, 0) + (0, b, 0)P_1 + (0, c, 0)P_2; a, b, c \in R\}$ is an AHS-subspace.

Theorem.

Let $2 - SP_V$ be a symbolic 2-plithogenic vector space over $2 - SP_F$, let W be an AHS-subspace of $2 - SP_V$, then W is a submodule of $2 - SP_V$.

Proof.

Suppose that W is an AHS-subspace, then there exists a subspace $V_0 \leq V$, such that

$$W = V_0 + V_0P_1 + V_0P_2 = \{x + yP_1 + zP_2; x, y, z \in V_0\}.$$

Let $X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2 \in W$, then:

$$X - Y = (x_0 - y_0) + (x_1 - y_1)P_1 + (x_2 - y_2)P_2 \in W$$

$\forall A = a_0 + a_1P_1 + a_2P_2 \in 2 - SP_F$, then:

$A.X = a_0x_0 + (a_0x_1 + a_1x_0 + a_1x_1)P_1 + (a_0x_2 + a_1x_2 + a_2x_0 + a_2x_1 + a_2x_2)P_2 \in W$, that is because $a_0x_0 \in V_0, a_0x_1 + a_1x_0 + a_1x_1 \in V_0, a_0x_2 + a_1x_2 + a_2x_0 + a_2x_1 + a_2x_2 \in V_0$, this implies the proof.

Definition.

Let V, W be two vector spaces over the field F . Let $2 - SP_V, 2 - SP_W$ be the corresponding symbolic 2-plithogenic vector spaces over $2 - SP_F$.

Let $L_0, L_1, L_2: V \rightarrow W$ be three linear transformations, we define the AH-linear transformation as follows:

$$L: 2 - SP_V \rightarrow 2 - SP_W, L = L_0 + L_1P_1 + L_2P_2; L(x + yP_1 + zP_2) = L_0(x) + L_1(y)P_1 + L_2(z)P_2.$$

If $L_0 = L_1 = L_2$, then L is called AHS-linear transformation.

Definition.

Let $L = L_0 + L_1P_1 + L_2P_2: 2 - SP_V \rightarrow 2 - SP_W$ be an AH-linear transformation, we define:

1. $AH - \ker(L) = \ker(L_0) + \ker(L_1)P_1 + \ker(L_2)P_2 = \{x + yP_1 + zP_2; x \in \ker(L_0), y \in \ker(L_1), z \in \ker(L_2)\}.$

2. $AH - Im(L) = Im(L_0) + Im(L_1)P_1 + Im(L_2)P_2 = \{a + bP_1 + cP_2\}; a \in Im(L_0), b \in Im(L_1), c \in Im(L_2)$

If L is AHS-linear transformation, then we get $AHS - kernel$, $AHS - Image$.

Theorem.

Let $L = L_0 + L_1P_1 + L_2P_2: 2 - SP_V \rightarrow 2 - SP_W$ be an AH-linear transformation, then:

1. $AH - ker(L)$ is AH-subspace of $2 - SP_V$.
2. $AH - Im(L)$ is AH-subspace of $2 - SP_W$.

Proof.

1. Since $ker(L_0), ker(L_1), ker(L_2)$ are subspaces of V , then $AH - ker(L)$ is an AH-subspace of $2 - SP_V$.
2. It is holds by the same.

Remark.

If L_0, L_1, L_2 are isomorphism, then $ker(L_0) = ker(L_1) = ker(L_2) = \{0\}, Im(L_0) = Im(L_1) = Im(L_2) = W$, thus $AH - ker(L) = \{0\}, AH - Im(L) = 2 - SP_W$.

Example.

Take $V = R^3, W = R^3, L_0, L_1, L_2: V \rightarrow W$ such that:

$$L_0(x, y, z) = (x, y), L_1(x, y, z) = (2x, z), L_2(x, y, z) = (x - y, y - z)$$

The corresponding AH-linear transformation is:

$$L = L_0 + L_1P_1 + L_2P_2: 2 - SP_{R^3} \rightarrow 2 - SP_{R^2}$$

$$\begin{aligned} L[(x_0, y_0, z_0) + (x_1, y_1, z_1)P_1 + (x_2, y_2, z_2)P_2] \\ = L_0(x_0, y_0, z_0) + L_1(x_1, y_1, z_1)P_1 + L_2(x_2, y_2, z_2)P_2 \\ = (x_0, y_0) + (2x_1, z_1)P_1 + (x_2 - y_2, y_2 - z_2)P_2 \end{aligned}$$

For example, take $X = (1, 2, 1) + (4, 3, -5)P_1 + (1, 1, 1)P_2$, then:

$$L(X) = (1, 2) + (8, -5)P_1 + (0, 0)P_2 = (1, 2) + (8, -5)P_1.$$

$$\left\{ \begin{array}{l} ker(L_0) = \{(0, 0, z_0); z_0 \in R\} \\ ker(L_1) = \{(0, y_1, 0); y_1 \in R\} \\ ker(L_2) = \{(x_2, x_2, x_2); x_2 \in R\} \\ AH - ker(L) = \{(0, 0, z_0) + (0, y_1, 0)P_1 + (x_2, x_2, x_2)P_2; z_0, y_1, x_2 \in R\} \end{array} \right.$$

Also,

$$\left\{ \begin{array}{l} Im(L_0) = R^2 \\ Im(L_1) = R^2 \\ Im(L_2) = R^2 \\ AH - Im(L) = R^2 + R^2P_1 + R^2P_2 = 2 - SP_W \end{array} \right.$$

Example.

Take $W = V = R^2$, $L_0, L_1, L_2: V \rightarrow W$ such that:

$$L_0(x, y) = (3x, -2x), L_1(x, y) = (x - y, 2x), L_2(x, y, z) = (x + 2y, y)$$

The corresponding AH-linear transformation is $L = L_0 + L_1P_1 + L_2P_2: 2 - SP_V \rightarrow 2 - SP_W$;

$$\begin{aligned} L[(x_0, y_0) + (x_1, y_1)P_1 + (x_2, y_2)P_2] &= L_0(x_0, y_0) + L_1(x_1, y_1)P_1 + L_2(x_2, y_2)P_2 \\ &= (3x_0, -2x_0) + (x_1 - y_1, 2x_1)P_1 + (x_2 + 2y_2, y_2)P_2 \end{aligned}$$

For example $X = (1, 4) + (2, 8)P_1 + (3, -1)P_2$

$$L(X) = (1, 4) + (2, 8)P_1 + (3, -1)P_2.$$

$$\left\{ \begin{array}{l} \ker(L_0) = \{(0, y_0); y_0 \in R\} \\ \ker(L_1) = \{0\} \\ \ker(L_2) = \{0\} \\ AH - \ker(L) = \{(0, y_0) + 0P_1 + 0P_2; y_0 \in R\} \end{array} \right.$$

Also,

$$\left\{ \begin{array}{l} \text{Im}(L_0) = \{(a, 0); a \in R\} \\ \text{Im}(L_1) = R^2 \\ \text{Im}(L_2) = R^2 \\ AH - \text{Im}(L) = \{(a, 0) + (a_1, b_1)P_1 + (a_2, b_2)P_2; a, a_1, a_2, b_1, b_2 \in R\} \end{array} \right.$$

Theorem.

Let $L = f + fP_1 + fP_2: 2 - SP_V \rightarrow 2 - SP_W$ be an AHS-linear transformation, then L is a module homomorphism.

Proof.

Let $X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2 \in 2 - SP_V$, then:

$$\begin{aligned} L(X + Y) &= f(x_0 + y_0) + f(x_1 + y_1)P_1 + f(x_2 + y_2)P_2 \\ &= [f(x_0) + f(x_1)P_1 + f(x_2)P_2] + [f(y_0) + f(y_1)P_1 + f(y_2)P_2] = L(X) + L(Y) \end{aligned}$$

Let $A = a_0 + a_1P_1 + a_2P_2 \in 2 - SP_F$, then:

$$\begin{aligned} L(A.X) &= f(a_0x_0) + f(a_0x_1 + a_1x_0 + a_1x_1)P_1 + f(a_0x_2 + a_2x_0 + a_2x_2 + a_1x_2 + a_2x_1)P_2 \\ &= a_0f(x_0) + (a_0f(x_1) + a_1f(x_0) + a_1f(x_1))P_1 \\ &\quad + (a_0f(x_2) + a_2f(x_0) + a_2f(x_2) + a_1f(x_2) + a_2f(x_1))P_2 \\ &= [a_0 + a_1P_1 + a_2P_2]. [f(x_0) + f(x_1)P_1 + f(x_2)P_2] = A.L(X) \end{aligned}$$

Thus, L is a module homomorphism.

The algebraic relations between symbolic 2-plithogenic vector spaces and neutrosophic vector spaces .

Theorem.

Let V be a vector space over the field F , consider $V(I) = V + VI = \{x + yI; x, y \in V\}$ is the corresponding neutrosophic vector space over the neutrosophic field $F(I) = \{a + bI; a, b \in F\}$.

$V(I_1, I_2) = V + VI_1 + VI_2 = \{x + yI_1 + zI_2; x, y, z \in V\}$ is the corresponding refined neutrosophic vector space over the refined neutrosophic field $F(I_1, I_2) = \{a + bI_1 + cI_2; a, b, c \in F\}$.

$2 - SP_V = V + VP_1 + VP_2 = \{x + yP_1 + zP_2; x, y, z \in V\}$ is the corresponding symbolic 2-plithogenic vector space over $2 - SP_F$, then:

1. $2 - SP_V$ is semi homomorphic to $V(I)$.
2. $2 - SP_V$ is semi isomorphic to $V(I_1, I_2)$.

Proof.

1. We define $f: 2 - SP_V \rightarrow V(I), g: 2 - SP_F \rightarrow F(I)$ such that:

$$f(x + yP_1 + zP_2) = x + yI; x, y, z \in V$$

$$g(a + bP_1 + cP_2) = a + bI; a, b, c \in F$$

We have the following:

g is a ring homomorphism, that is because:

$$A = a_0 + a_1P_1 + a_2P_2, B = b_0 + b_1P_1 + b_2P_2; a_i, b_i \in F, \text{ then:}$$

$$\text{If } A = B, \text{ then } a_i = b_i \text{ for all } i, \text{ thus } a_0 + a_1I = b_0 + b_1I, \text{ i.e. } g(A) = g(B).$$

$$g(A + B) = g[(a_0 + b_0) + (a_1 + b_1)P_1 + (a_2 + b_2)P_2] = a_0 + b_0 + (a_1 + b_1)I = g(A) + g(B).$$

$$\begin{aligned} g(A \cdot B) &= g[a_0b_0 + (a_0b_1 + a_1b_0 + a_1b_1)P_1 + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)P_2] \\ &= a_0b_0 + (a_0b_1 + a_1b_0 + a_1b_1)I = (a_0 + a_1I)(b_0 + b_1I) = g(A) \cdot g(B) \end{aligned}$$

On the other hand, f is well defined, that is because:

$$\text{If } X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2, \text{ then } x_i = y_i \text{ for all } i, \text{ hence } a_0 + a_1I = b_0 + b_1I, \text{ thus } f(X) = f(Y).$$

f preserves addition, that is because:

$$\text{For } X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2, \text{ we have:}$$

$$f(X + Y) = f[(x_0 + y_0) + (x_1 + y_1)P_1 + (x_2 + y_2)P_2] = x_0 + y_0 + (x_1 + y_1)I = f(X) + f(Y).$$

f preserves multiplication, that is because:

$$\text{For } A = a_0 + a_1P_1 + a_2P_2 \in 2 - SP_V, \text{ we have:}$$

$$f(A \cdot X) = a_0x_0 + (a_0x_1 + a_1x_0 + a_1x_1)I = (a_0 + a_1I)(x_0 + x_1I) = g(A) \cdot f(X)$$

Thus f is a semi module homomorphism.

We define $f: 2 - SP_V \rightarrow V(I_1, I_2)$, $g: 2 - SP_F \rightarrow F(I_1, I_2)$, where $f(x + yP_1 + zP_2) = x + zI_1 + yI_2$, and $g(a + bP_1 + cP_2) = a + cI_1 + bI_2; x, y, z \in V, a, b, c \in F$.

(g) is well defined, that is because:

If $A = a_0 + a_1P_1 + a_2P_2, B = b_0 + b_1P_1 + b_2P_2$, then:

$a_0 = a_1, b_0 = b_1, c_0 = c_1$, hence: $a_0 + c_0I_1 + b_0I_2 = a_1 + c_1I_1 + b_1I_2$, so that $g(A) = g(B)$.

(f) is well defined by a similar discussion.

(g) is one-to-one mapping, that is because:

$$\ker(g) = \{a + bP_1 + cP_2; g(a + bP_1 + cP_2) = 0\} = 0$$

$$\begin{aligned} \text{Im}(g) &= \{a + cI_1 + bI_2; g(a + bP_1 + cP_2) \in F(I_1, I_2); \exists A \in 2 - SP_F, g(A) = a + cI_1 + bI_2\} \\ &= F(I_1, I_2) \end{aligned}$$

(f) is one-to-one mapping, it can be proved by the same.

(g) and (f) preserve addition, that is because:

Consider $A = a_0 + a_1P_1 + a_2P_2, B = b_0 + b_1P_1 + b_2P_2 \in 2 - SP_F$, $X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2 \in 2 - SP_V$, then:

$$\begin{aligned} g(A + B) &= g[(a_0 + b_0) + (a_1 + b_1)P_1 + (a_2 + b_2)P_2] = a_0 + b_0 + (a_1 + b_1)I_1 + (a_2 + b_2)I_2 \\ &= g(A) + g(B) \end{aligned}$$

$f(X + Y) = f(X) + f(Y)$ by a similar discussion.

(g) preserves multiplication, that is because:

$$\begin{aligned} g(A.B) &= a_0b_0 + (a_0b_2 + a_2b_0 + a_2b_2 + a_1b_2 + a_2b_1)I_1 + (a_0b_1 + a_1b_0 + a_1b_1)I_2 = \\ &= g(A).g(B). \end{aligned}$$

(f) is semi module homomorphism, that is because:

$$\begin{aligned} f(A.X) &= a_0x_0 + (a_0x_2 + a_2x_0 + a_2x_2 + a_1x_2 + a_2x_1)I_1 + (a_0x_1 + a_1x_0 + a_1x_1)I_2 \\ &= (a_0 + a_1I_1 + a_2I_2)(x_0 + x_2I_1 + x_1I_2) = g(A).f(X) \end{aligned}$$

The basis of a symbolic 2-plithogenic vector spaces:

Theorem.

Let $T = \{t_1, \dots, t_n\}$ be a basis of the vector space V over the field F , then the set:

$T_P = \{t_i + (t_j - t_i)P_1 + (t_k - t_j)P_2; 1 \leq i, j, k \leq n\}$ is a basis of $2 - SP_V$.

Proof.

Let $X = x_0 + x_1P_1 + x_2P_2 \in 2 - SP_V, x_0, x_1, x_2 \in V$.

$$x_0 = \sum_{i=1}^n \alpha_i t_i, x_0 + x_1 = \sum_{j=1}^n \beta_j t_j, x_0 + x_1 + x_2 = \sum_{k=1}^n \gamma_k t_k; \alpha_i, \beta_j, \gamma_k \in F.$$

We put $A_{i,j,k} = \alpha_i + (\beta_j - \alpha_i)P_1 + (\gamma_k - \beta_j)P_2; 1 \leq i, j, k \leq n$

$$T_{i,j,k} = t_i + (t_j - t_i)P_1 + (t_k - t_j)P_2; 1 \leq i, j, k \leq n$$

$$\begin{aligned}
& \sum_{i,j,k=1}^n A_{i,j,k} T_{i,j,k} \\
&= \sum_{i=1}^n [\alpha_i t_i + [\beta_j t_j - \beta_j t_i - \alpha_i t_j + \alpha_i t_i + \beta_j t_i - \alpha_i t_i + \alpha_i t_j - \alpha_i t_i] P_1 \\
&\quad + [\alpha_i t_k - \alpha_i t_j + \gamma_k t_i - \beta_j t_i - \gamma_k t_j + \gamma_k t_i - \beta_j t_j + \beta_j t_i + \gamma_k t_k - \gamma_k t_j - \beta_j t_k \\
&\quad + -\beta_j t_j + \beta_j t_k - \beta_j t_j - \alpha_i t_k + \alpha_i t_j] P_2] \\
& \sum_{i=1}^n \alpha_i t_i + P_1 \left[\sum_{j=1}^n \beta_j t_j - \sum_{i=1}^n \alpha_i t_i \right] + P_2 \left[\sum_{k=1}^n \gamma_k t_k - \sum_{j=1}^n \beta_j t_j \right] \\
&= x_0 + P_1 [x_0 + x_1 - x_0] + P_2 [x_0 + x_1 + x_2 - (x_0 + x_1)] = x_0 + x_1 P_1 + x_2 P_2 \\
&= X
\end{aligned}$$

Thus T generates $2 - SP_V$.

On the other hand, T is linearly independent, that is because:

If $\sum_{i,j,k=1}^n A_{i,j,k} \cdot X = 0$, then:

$\sum_{i=1}^n \alpha_i t_i = 0, \sum_{j=1}^n \beta_j t_j = 0, \sum_{k=1}^n \gamma_k t_k = 0$, hence $\alpha_i = \beta_j = \gamma_k = 0$ for all i, j, k , thus $A_{i,j,k} = 0$.

This implies that T is a basis of $2 - SP_V$.

Example.

Find a basis of $2 - SP_{R^2}$.

Solution.

First of all, we have $\{u_1 = (1,0), u_2 = (0,1)\}$ is a basis of R^2 .

The corresponding basis of $2 - SP_{R^2}$ is:

$T = \{T_1, T_2, T_3, T_4, T_5, T_6, T_7, T_8\}$ such that:

$$T_1 = (1,0), T_2 = (0,1), T_3 = u_1 + (u_2 - u_1)P_1 + (u_2 - u_2)P_2 = (1,0) + (-1,1)P_1$$

$$T_4 = u_1 + (u_2 - u_1)P_1 + (u_1 - u_2)P_2 = (1,0) + (-1,1)P_1 + (1,-1)P_2$$

$$T_5 = u_2 + (u_2 - u_1)P_1 + (u_1 - u_1)P_2 = (0,1) + (1,-1)P_1$$

$$T_6 = u_2 + (u_2 - u_1)P_1 + (u_2 - u_1)P_2 = (0,1) + (1,-1)P_1 + (-1,1)P_2$$

$$T_7 = u_1 + (u_1 - u_1)P_1 + (u_2 - u_1)P_2 = (1,0) + (-1,1)P_2$$

$$T_8 = u_2 + (u_2 - u_2)P_1 + (u_1 - u_2)P_2 = (0,1) + (1,-1)P_2$$

Remark.

$$\dim(2 - SP_V) = (\dim V)^3.$$

Conclusion

In this paper we have defined the concept of symbolic 2-plithogenic vector spaces over a symbolic 2-plithogenic field, where we have presented some of their elementary properties such as basis, linear transformations, and AH-subspaces. On the other hand, we have suggested many examples to clarify the validity of our work.

References

- [1] Abobala, M., "AH-Subspaces in Neutrosophic Vector Spaces", International Journal of Neutrosophic Science, Vol. 6 , pp. 80-86. 2020.
- [2] Abobala, M.,. "A Study of AH-Substructures in n -Refined Neutrosophic Vector Spaces", International Journal of Neutrosophic Science", Vol. 9, pp.74-85. 2020.
- [3] Abobala, M., Hatip, A., Bal,M.," A Study Of Some Neutrosophic Clean Rings", International journal of neutrosophic science, 2022.
- [4] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.
- [5] Celik, M., and Hatip, A., " On The Refined AH-Isometry And Its Applications In Refined Neutrosophic Surfaces", Galoitica Journal Of Mathematical Structures And Applications, 2022.
- [6] Adeleke, E.O., Agboola, A.A.A.,and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81. 2020.
- [7] M. Ibrahim. A. Agboola, B.Badmus and S. Akinleye. On refined Neutrosophic Vector Spaces . International Journal of Neutrosophic Science, Vol. 7, 2020, pp. 97-109.
- [8] Von Shtawzen, O., " On A Novel Group Derived From A Generalization Of Integer Exponents and Open Problems", Galoitica journal Of Mathematical Structures and Applications, Vol 1, 2022.
- [9] Hatip, A., "An Introduction To Weak Fuzzy Complex Numbers ", Galoitica Journal Of Mathematical Structures and Applications, Vol.3, 2023.
- [10] Merkepci, H., and Ahmad, K., " On The Conditions Of Imperfect Neutrosophic Duplets and Imperfect Neutrosophic Triplets", Galoitica Journal Of Mathematical Structures And Applications, Vol.2, 2022.
- [11] Von Shtawzen, O., " Conjectures For Invertible Diophantine Equations Of 3-Cyclic and 4-Cyclic Refined Integers", Journal Of Neutrosophic And Fuzzy Systems, Vol.3, 2022.

-
- [12] Abobala, M., and Zeina, M.B., " A Study Of Neutrosophic Real Analysis By Using One Dimensional Geometric AH-Isometry", *Galoitica Journal Of Mathematical Structures And Applications*, Vol.3, 2023.
- [13] F. Smarandache, Neutrosophic Quadruple Numbers, Refined Neutrosophic Quadruple Numbers, Absorbance Law, and the Multiplication of Neutrosophic Quadruple Numbers. In *Symbolic Neutrosophic Theory*, Chapter 7, pages 186-193, Europa Nova, Brussels, Belgium, 2015.
- [14] F. Smarandache, Plithogeny, Plithogenic Set, Logic, Probability, and Statistics, 141 pages, Pons Editions, Brussels, Belgium, 2017. arXiv.org (Cornell University), Computer Science - Artificial Intelligence, 03Bxx:
- [15] Florentin Smarandache, Physical Plithogenic Set, 71st Annual Gaseous Electronics Conference, Session LW1, Oregon Convention Center Room, Portland, Oregon, USA, November 5–9, 2018.
- [16] Florentin Smarandache: Plithogenic Set, an Extension of Crisp, Fuzzy, Intuitionistic Fuzzy, and Neutrosophic Sets – Revisited, *Neutrosophic Sets and Systems*, vol. 21, 2018, pp. 153-166.
- [17] Florentin Smarandache, Plithogenic Algebraic Structures. Chapter in "Nidus idearum Scilogs, V: joining the dots" (third version), Pons Publishing Brussels, pp. 123-125, 2019.
- [18] P. K. Singh, Data with Turiyam Set for fourth dimension Quantum Information Processing. *Journal of Neutrosophic and Fuzzy Systems*, Vol. 1, Issue 1, pp. 9-23, 2021.
- [19] Khaldi, A., " A Study On Split-Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [20] Ahmad, K., " On Some Split-Complex Diophantine Equations", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [21]. Ali, R., " On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.

-
- [22] A. Alrida Basher, Katy D. Ahmad, Rosina Ali, An Introduction to the Symbolic Turiyam Groups and AH-Substructures,,*Journal of Neutrosophic and Fuzzy Systems*, Vol. 03, No. 02, pp. 43-52, 2022.
- [23] Agboola, A.A.A., "On Refined Neutrosophic Algebraic Structures," *Neutrosophic Sets and Systems*, Vol.10, pp. 99-101. 2015.
- [24] T.Chalapathi and L. Madhavi. "Neutrosophic Boolean Rings", *Neutrosophic Sets and Systems*, Vol. 33, pp. 57-66, 2020.
- [25] G. Shahzadi, M. Akram and A. B. Saeid, "An Application of Single-Valued Neutrosophic Sets in Medical Diagnosis," *Neutrosophic Sets and Systems*, vol. 18, pp. 80-88, 2017.
- [26] Sarkis, M., " On The Solutions Of Fermat's Diophantine Equation In 3-refined Neutrosophic Ring of Integers", *Neoma Journal of Mathematics and Computer Science*, 2023.
- [27] J. Anuradha and V. S, "Neutrosophic Fuzzy Hierarchical Clustering for Dengue Analysis in Sri Lanka," *Neutrosophic Sets and Systems*, vol. 31, pp. 179-199, 2020.
- [28] Celik, M., and Olgun, N., " An Introduction To Neutrosophic Real Banach And Hillbert Spaces", *Galoitica Journal Of Mathematical Structures And Applications*, 2022.
- [29] Celik, M., and Olgun, N., " On The Classification Of Neutrosophic Complex Inner Product Spaces", *Galoitica Journal Of Mathematical Structures And Applications*, 2022.
- [30] Smarandache, F., " Introduction to the Symbolic Plithogenic Algebraic Structures (revisited)", *Neutrosophic Sets and Systems*, vol. 53, 2023.
- [31] Merkepçi, H., and Abobala, M., " On The Symbolic 2-Plithogenic Rings", *International Journal of Neutrosophic Science*, 2023.
- [32] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", *Neutrosophic Sets and Systems*, Vol. 43, 2021.
- [33] Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", *Mathematical Problems in Engineering*, Hindawi, 2021
- [34] Agboola, A.A.A., Akinola, A.D., and Oyebola, O.Y., " Neutrosophic Rings I" , *International J.Mathcombin*, Vol 4, pp 1-14. 2011

-
- [35] Smarandache, F., and Ali, M., "Neutrosophic Triplet Group", Neural. Compute. Appl. 2019.
- [36] Chellamani, P., and Ajay, D., "Pythagorean neutrosophic Fuzzy Graphs", International Journal of Neutrosophic Science, Vol. 11, 2021.
- [37] Giorgio, N, Mehmood, A., and Broumi, S.," Single Valued neutrosophic Filter", International Journal of Neutrosophic Science, Vol. 6, 2020.
- [38] Hatip, A., " On Intuitionistic Fuzzy Subgroups of (M-N) Type and Their Algebraic Properties", Galoitica Journal Of Mathematical Structures and Applications, Vol.4, 2023.

Received: December 25, 2022. **Accepted:** March 28, 2023