

4-11-2023

## On The Classification of The Group of Units of Rational and Real 2-Cyclic Refined Neutrosophic Rings

Hasan Sankari

Mohammad Abobala

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Sankari, Hasan and Mohammad Abobala. "On The Classification of The Group of Units of Rational and Real 2-Cyclic Refined Neutrosophic Rings." *Neutrosophic Sets and Systems* 54, 1 ().  
[https://digitalrepository.unm.edu/nss\\_journal/vol54/iss1/8](https://digitalrepository.unm.edu/nss_journal/vol54/iss1/8)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).



# On The Classification of The Group of Units of Rational and Real 2-Cyclic Refined Neutrosophic Rings

<sup>1</sup>Hasan Sankari , <sup>2</sup>Mohammad Abobala

<sup>1,2</sup>Tishreen University, Faculty of Science, Department Of Mathematics, Latakia, Syria

Co-[Mohammadabobala777@gmail.com](mailto:Mohammadabobala777@gmail.com)

**Abstract:** The objective of this paper is to solve two open problems about the group of units of some 2-cyclic refined neutrosophic rings asked by Sadiq. Where it provides a classification theorem for these rings, and uses this classification property to give a full answer of these open questions.

Also, this work presents a novel algorithm to find all imperfect neutrosophic duplets and triplets in many numerical 2-cyclic refined neutrosophic rings by using the classification isomorphisms.

## 1. Introduction

Neutrosophic logic as a new generalization of fuzzy logic concerns with indeterminacy in science and real life problems [1]. Neutrosophy was proposed by Smarandache [6] for these logical purposes.

Laterally, neutrosophy was applied to algebra and algebraic structures, where we find many algebraic structures defined by using an indeterminacy element (I) such as neutrosophic rings, neutrosophic spaces, neutrosophic modules, and matrices [2-5].

The concept of n-cyclic refined neutrosophic ring was presented firstly in [7], and studied widely in [8-9].

In [10], Sadiq has studied the group of units problem for 2-CRNR rings, where he proved that it is isomorphic to 3 times direct product of  $Z_2$ . Also, he presented the following open research problems: [10]:

**Open problem 1:** If the ring  $R$  with no zero divisors, then is the group of units of  $R_2(I)$  is isomorphic to  $U(R) \times U(R) \times U(R)$ .

**Open problem 2:** Find a homomorphism between  $R_2(I)$  and the direct product  $\times R \times R$ .

**Open problem 3:** Is the group of units of the 2-cyclic refined ring of real numbers isomorphic to  $R^* \times R^* \times R^*$ .

This motivates us to continue these efforts to classify the group of units of 2-cyclic refined rings, and to prove the validity of Sadiq's open problems.

On the other hand, we classify all imperfect duplets and triplets in the ring of 2-cyclic refined neutrosophic integers by solving many related Diophantine equations.

We denote the 2-cyclic refined ring by 2-CRNR.

## 2. Preliminaries

### Definition 1.2:

Let  $(R, +, \times)$  be a ring and  $I_k; 1 \leq k \leq n$  be  $n$  sub-indeterminacies. We define  $R_n(I) = \{a_0 + a_1I + \dots + a_nI_n; a_i \in R\}$  to be  $n$ -cyclic refined neutrosophic ring.

Operations on  $R_n(I)$  are defined as:

$$\begin{aligned} \sum_{i=0}^n p_i I_i + \sum_{i=0}^n q_i I_i &= \sum_{i=0}^n (p_i + q_i) I_i, \sum_{i=0}^n p_i I_i \\ \times \sum_{i=0}^n q_i I_i &= \sum_{i,j=0}^n (p_i \times q_j) I_i I_j = \sum_{i,j=0}^n (p_i \times q_j) I_{(i+j \bmod n)} \end{aligned}$$

### Example 2.2:

(a) The 2-CRNR of integers is defined as follows:

$$Z_2(I) = \{t_0 + t_1 I_1 + t_2 I_2; t_i \in Z\}.$$

(b) Addition on  $Z_2(I)$  :

$$(a + bI_1 + cI_2) + (m + nI_1 + tI_2) = (a + m) + I_1(b + n) + I_2(c + t).$$

(c) Multiplication on  $Z_2(I)$  :

$$(a + bI_1 + cI_2)(m + nI_1 + tI_2) = am + anI_1 + atI_2 + bml_1 + bnI_2 + btI_1 + cml_2 + cnI_1 + ctI_2$$

$$= am + I_1(an + bm + bt + cn) + I_2(at + bn + cm + ct) .$$

Where  $I_1I_1 = I_{(1+1 \bmod 2)} = I_2, I_2I_2 = I_{(2+2 \bmod 2)} = I_2, I_1I_2 = I_{(1+2 \bmod 2)} = I_1$ .

### Definition 3.2:

Let  $R$  be a ring, a duplet  $(x, y)$  is called an imperfect duplet with  $x$  acts as an identity if and only if  $xy = yx = y$ .

A triple  $(x, y, z)$  is called an imperfect triplet with  $x$  acts as an identity if and only if  $xy = yx = y, xz = zx = z, zy = yz = x$ .

### 3. Main discussion

**Theorem 1.3 :** Let  $Z$  be the ring of integers, and  $S = \{(b_0, b_1, b_2); b_i \in Z \text{ and } b_1 - b_2 \in 2Z\}$ , then  $(S, +, \cdot)$  Is a subring of  $Z \times Z \times Z$ .

Proof: It is clear that  $S \neq \emptyset$

$\forall x, y \in S, x = (a_0, a_1, a_2), y = (b_0, b_1, b_2)$ , where  $b_1 - b_2, a_1 - a_2 \in 2Z$

$$x + y = (a_0 + b_0, a_1 + b_1, a_2 + b_2), xy = (a_0b_0, a_1b_1, a_2b_2)$$

We have:  $(a_1 + b_1) - (a_2 + b_2) = (a_1 - a_2) + (b_1 - b_2) \in 2Z$ , thus  $x + y \in S$

Also,  $a_1b_1 - a_2b_2 = a_1b_1 + a_1b_2 - a_1b_2 - a_2b_2 = a_1(b_1 + b_2) - b_2(a_1 + a_2)$  . By the assumption, we have  $b_1 - b_2, a_1 - a_2 \in 2Z$ , hence

$b_1 + b_2, a_1 + a_2 \in 2Z$ , this implies  $a_1(b_1 + b_2) - b_2(a_1 + a_2) \in 2Z$  and  $x \cdot y \in S$ .

**Theorem 2.3:** Let  $Z_2(I)$  be the 2-CRNR of integers, then  $Z_2(I) \cong S$ .

Proof:

Define  $f: Z_2(I) \rightarrow S; f(a_0 + a_1I_1 + a_2I_2) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2)$ .

It's clear that  $f$  is well defined. On the other hand we have:

(a).  $f$  is injective,  $\ker f = \{a_0 + a_1I_1 + a_2I_2 \in Z_2(I); f(a_0 + a_1I_1 + a_2I_2) = (0,0,0)\}$ , hence.

$a_0 = 0, a_0 + a_1 + a_2 = 0, a_0 - a_1 + a_2 = 0$ , thus  $a_0 = a_1 = a_2$ , this means that  $\ker f = \{0_S\}$ .

(b).  $f$  is surjective,  $\forall y = (a_0, a_1, a_2) \in S$ , we have:  $a_1 - a_2 \in 2Z$ , hence  $x = a_0 +$

$$I_1\left(\frac{a_1 - a_2}{2}\right) + I_2\left(\frac{a_1 + a_2 - 2a_0}{2}\right) \in Z_2(I).$$

This is because  $a_1 - a_2, a_1 + a_2 - 2a_0 \in 2Z$ .

Now, we compute  $f(x) = \left( a_0, a_0 + \frac{a_1 - a_2}{2} + \frac{a_1 + a_2 - 2a_0}{2}, a_0 - \frac{a_1 - a_2}{2} + \frac{a_1 + a_2 - 2a_0}{2} \right) = (a_0, a_1, a_2) = y$

(c).  $f$  is a homomorphism because clearly  $f$  preserves addition and multiplication, thus  $S \cong Z_2(I)$ .

**Theorem 3.3:** Let  $R$  be a of real numbers,  $R_2(I)$  be the corresponding 22-CRNR of real numbers, then  $R_2(I) \cong R^3$ .

Proof. Define  $f: R_2(I) \rightarrow R^3; f(a_0 + a_1I_1 + a_2I_2) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2)$ .

$f$  is well defined and bijective. (the proof is exactly similar to the previous theorem).

$f$  is a homomorphism.  $\forall x, y \in R_2(I), x = a_0 + a_1I_1 + a_2I_2, y = b_0 + b_1I_1 + b_2I_2$ .

$$x + y = a_0 + b_0 + (a_1 + b_1)I_1 + (a_2 + b_2)I_2.$$

$$f(x + y) = (a_0 + b_0, a_0 + b_0 + a_1 + b_1 + a_2 + b_2, a_0 + b_0 - (a_1 + b_1) + a_2 + b_2)$$

$$f(x + y) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2) + (b_0, b_0 + b_1 + b_2, b_0 - b_1 + b_2) = f(x) + f(y).$$

$$xy = a_0b_0 + (a_1b_0 + a_0b_1 + a_2b_1 + a_1b_2)I_1 + (a_1b_0 + a_0b_2 + a_1b_1 + a_2b_2)I_2$$

$$f(xy) = (a_0b_0, a_0b_0 + a_1b_0 + a_0b_1 + a_1b_1 + a_2b_0 + a_0b_2 + a_1b_1 + a_2b_2, a_0b_0$$

$$- (a_1b_0 + a_0b_1 + a_2b_1 + a_1b_2) + a_2b_0 + a_0b_2 + a_1b_1 + a_2b_2)$$

$$f(xy) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2) \cdot (b_0, b_0 + b_1 + b_2, b_0 - b_1 + b_2) = f(x) \cdot f(y) ,$$

hence  $R_2(I) \cong R^3$ .

### Answers to the open questions

The following theorem answers the open question 3.

**Theorem 4.3:** Let  $\cup (R_2(I))$  be the group of units of the 2-CRNR  $R_2(I)$ , then  $\cup (R_2(I)) \cong R^{*3}$ .

Proof.

According to the previous theorem,  $R_2(I) \cong R \times R \times R$ , hence.  $\cup (R_2(I)) \cong \cup (R) \times \cup (R) \times \cup (R) = R^{*3}$ .

The following remark answers the open question 2.

**Remark 5.3:** If  $R$  is a ring, and  $R_2(I)$  is the corresponding 2-CRNR, hence the map

$$f: R_2(I) \rightarrow R \times R \times R; f(a_0 + a_1I_1 + a_2I_2) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2),$$

Is a ring homomorphism (the proof is similar to Theorem 3.2). Thus the answer to the open question 2 is yes. Remark that  $f$  is not supposed to be an isomorphism, check Theorem 1.3 for example.

The first question is still open, but we can solve the problem in a special case for the ring of integers modulo  $n$ , with odd  $n$ .

**Theorem 6.3:** Let  $R$  be the ring of integers modulo  $n$ , with an odd integer  $n$ , then  $R_2(I) \cong Z_n \times Z_n \times Z_n$

Proof. . Define  $f: R_2(I) \rightarrow Z_n \times Z_n \times Z_n; f(a_0 + a_1I_1 + a_2I_2) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2)$ .

It's clear that  $f$  is a well defined homomorphism, by a similar argument of the previous theorem, we should prove that  $f$  is a bijective map.

$\ker f = \{a_0 + a_1I_1 + a_2I_2 \in R_2(I); f(a_0 + a_1I_1 + a_2I_2) = (0,0,0)\}$ , hence.

$$a_0 = 0 \dots (1)$$

$$a_0 + a_1 + a_2 = 0 \dots (2)$$

$$a_0 - a_1 + a_2 = 0 \dots (3)$$

From equation (2) and (3), we get  $2a_2 = 0$ , By the proposition of the theorem,  $n$  is odd, this means that  $\gcd(2,n)=1$  and 2 cannot be a zero divisor, thus  $2a_2 = 0 \Rightarrow a_2 = 0$ .

This implies that  $a_1 = 0$ , and  $\ker f = \{(0,0,0)\}$ .

$f$  is surjective:

$\forall y = (a_0, a_1, a_2) \in Z_n \times Z_n \times Z_n$  , we have  $x = a_0 + I_1((a_1 + a_2)2^{-1}) + I_2((a_1 + a_2 - 2a_0)2^{-1}) \in R_2(I)$ .

That is because 2 is a unit in  $Z_n$  and  $2^{-1}$  is existed.

Now, we compute

$$\begin{aligned} f(x) &= (a_0, (a_1 - a_2)2^{-1} + (a_1 + a_2 + 2a_0)2^{-1} + a_0, a_0 + (a_1 - a_2)2^{-1} \\ &\quad + (a_1 + a_2 + 2a_0)2^{-1}) \\ &= (a_0, a_12^{-1} - a_22^{-1} + a_12^{-1} + a_22^{-1} - 2a_02^{-1} + a_0, a_0 - a_12^{-1} + a_22^{-1} + a_12^{-1} + a_22^{-1} \\ &\quad - 2a_02^{-1}) \\ &= (a_0, 2a_12^{-1}, 2a_22^{-1}) = (a_0, a_1, a_2). \end{aligned}$$

So that,  $R_2(I) \cong Z_n \times Z_n \times Z_n$ .

**Theorem 7.3:** If  $R = Z_n$  the ring of integers modulo  $n$  with an odd integer  $n$ , we have:

$$\cup (R_2(I)) \cong \cup (Z_n) \times \cup (Z_n) \times \cup (Z_n).$$

The proof holds directly by the previous result.

**Theorem 8.3:** If  $R = Z$  the ring of integers,  $Z_2(I)$  be the corresponding 2-CRNR, then  $Z_2(I)$  has exactly 8 forms of imperfect duplets.

Proof. We have  $Z_2(I) \cong S$ ;  $S = \{(a_0, a_1, a_2); a_i \in Z \text{ and } a_1 - a_2 \in 2Z\}$ .

To find imperfect duplets in  $Z_2(I)$ , it is sufficient to compute duplets in  $S$ :

Let  $x = (a_0, a_1, a_2), y = (b_0, b_1, b_2)$  be an imperfect duplet in  $S$ , with  $y$  acts as an identity, we have.

$$x \cdot y = x \Rightarrow \begin{cases} a_0 b_0 = a_0 \\ a_1 b_1 = a_1 \\ a_2 b_2 = a_2 \end{cases} \Rightarrow \begin{cases} a_0 = 0 \text{ or } b_0 = 0 \\ a_1 = 0 \text{ or } b_1 = 0 \\ a_2 = 0 \text{ or } b_2 = 0 \end{cases}$$

The possible imperfect duplets are:

(1).  $x = (0, 0, 0), y = (b_0, b_1, b_2)$

(With  $b_1 - b_2 \in 2Z$ )

(2).  $x = (0, a_1, a_2), y = (b_0, 1, 1)$

(With  $a_1 - a_2 \in 2Z$ )

(3).  $x = (0, 0, a_2), y = (b_0, b_1, 1)$

(With  $a_2$  is even and  $b_1$  is odd)

(4).  $x = (a_0, 0, a_2), y = (1, b_1, 1)$

(With  $a_2$  is even and  $b_1$  is odd)

(5).  $x = (a_0, a_1, 0), y = (1, 1, b_2)$

(With  $a_1$  is even and  $b_2$  is odd)

(6).  $x = (a_0, a_1, a_2), y = (1, 1, 1)$

(With  $a_1 - a_2 \in 2Z$ )

(7).  $x = (a_0, 0, 0), y = (1, b_1, b_2)$

(With  $b_1 - b_2 \in 2Z$ )

(8).  $x = (0, a_1, 0), y = (b_0, 1, b_2)$

(With  $a_1$  is even and  $b_2$  is odd)

Thus, the imperfect duplets in  $Z_2(I)$  are the converse image of the duplets in  $S$ , according to the isomorphism

$$f(a_0 + a_1I_1 + a_2I_2) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2).$$

$f^{-1}(a_0 + a_1I_1 + a_2I_2) = \left(a_0 + \frac{a_1 - a_2}{2}I_1 + \frac{a_1 + a_2 - 2a_0}{2}I_2\right)$ , so that the duplets of  $Z_2(I)$  are:

$$(1). x = 0, y = b_0 + b_1I_1 + b_2I_2$$

(With  $b_1 - b_2 \in 2Z$ )

$$(2). x = \frac{a_1 - a_2}{2}I_1 + \frac{a_1 + a_2}{2}I_2, y = b_0 + \frac{2 - 2b_0}{2}I_2$$

(With  $a_1 - a_2 \in 2Z$ )

$$(3). x = \frac{-a_2}{2}I_1 + \frac{a_2}{2}I_2, y = b_0 + \frac{b_1 - 1}{2}I_1 + \frac{b_1 + 1 - 2b_0}{2}I_2$$

(With  $a_2$  is even and  $b_1$  is odd)

$$(4). x = a_0 + \frac{-a_2}{2}I_1 + \frac{a_2 - 2a_0}{2}I_2, y = 1 + \frac{b_1 - 1}{2}I_1 + \frac{b_1 + 1 - 2(1)}{2}I_2$$

(With  $a_2$  is even and  $b_1$  is odd)

$$(5). x = a_0 + \frac{a_1}{2}I_1 + \frac{a_1 - 2a_0}{2}I_2, y = 1 + \frac{1 - b_2}{2}I_1 + \frac{1 + b_2 - 2(1)}{2}I_2$$

(With  $a_1$  is even and  $b_2$  is odd)

$$(6). x = a_0 + a_1I_1 + a_2I_2, y = 1$$

(With  $a_1 - a_2 \in 2Z$ )

$$(7). x = a_0 - a_0I_2, y = 1 + \frac{b_1 - b_2}{2}I_1 + \frac{b_1 + b_2 - 2}{2}I_2$$

(With  $b_1 - b_2 \in 2Z$ )

$$(8). x = \frac{a_1}{2}I_1 + \frac{a_1}{2}I_2, y = b_0 + \frac{1 - b_2}{2}I_1 + \frac{1 + b_2 - 2b_0}{2}I_2$$

(With  $a_1$  is even and  $b_2$  is odd)

**Theorem 9.3:** Let  $R$  be the ring of real numbers,  $R_2(I)$  be its 2-CRNR, then  $R_2(I)$  has exactly 8 forms of imperfect duplets.

Proof. We have  $R_2(I) \cong R \times R \times R$  with the isomorphism:

$$f: R_2(I) \rightarrow R \times R \times R; f(a_0 + a_1I_1 + a_2I_2) = (a_0, a_0 + a_1 + a_2, a_0 - a_1 + a_2).$$



For determining the imperfect duplets in  $R_2(I)$ , it is sufficient to find duplets in  $R \times R \times R$  and go back to  $R_2(I)$  by the inverse isomorphism.

The imperfect duplets in  $R \times R \times R$  are:

$$(1). x = (0, 0, 0), y = (b_0, b_1, b_2)$$

$$(2). x = (0, a_1, a_2), y = (b_0, 1, 1)$$

$$(3). x = (0, 0, a_2), y = (b_0, b_1, 1)$$

$$(4). x = (a_0, 0, a_2), y = (1, b_1, 1)$$

$$(5). x = (a_0, a_1, 0), y = (1, 1, b_2)$$

$$(6). x = (a_0, a_1, a_2), y = (1, 1, 1)$$

$$(7). x = (a_0, 0, 0), y = (1, b_1, b_2)$$

$$(8). x = (0, a_1, 0), y = (b_0, 1, b_2)$$

Thus  $R_2(I)$  has 8 forms of imperfect duplets.

**Remark 10.3:** To find any imperfect duplets in  $R_2(I)$ , we should compute the inverse image of the corresponding duplet in  $R \times R \times R$  as follows:

$$f^{-1}(a_0, a_1, a_2) = a_0 + \frac{a_1 - a_2}{2}I_1 + \frac{a_1 + a_2 - 2a_0}{2}I_2$$

**Example 11.3:** Let's, take a duplet with form:  $x = (2, 0, 3), y = (1, 5, 1)$ , it is clear  $x \cdot y = x$ .

The corresponding duplet in  $R_2(I)$  is:

$$x_1 = f^{-1}(x) = 2 + \frac{-3}{2}I_1 + \frac{-1}{2}I_2, y_1 = f^{-1}(y) = 1 + 2I_1 + 2I_2.$$

**Remark 12.3:** that  $x_1 \cdot y_1 = 2 + 4I_1 + 4I_2 = \frac{-3}{2}I_1 - 3I_2 - 3I_1 - \frac{1}{2}I_2 - I_1 - I_2 = 2 - \frac{3}{2}I_1 -$

$$\frac{1}{2}I_2 = x_1.$$

**Theorem 13.3:** Let  $Z_2(I)$  be the 2-CRNR of integers, then it has exactly 14 forms of imperfect triplets.

Proof.

Let  $x, y, z$  be a triplet in  $S$ , then we have:

$xy = yx = x$ ,  $yz = zy = z$ ,  $xz = zx = y$ , so that,  $(x, y), (y, z)$  are imperfect duplets in  $S$ .

We discuss the 8 forms of imperfect duplets to find the desired imperfect duplets:

Form 1:  $x = (0, 0, 0), y = (b_0, b_1, b_2), z = (0, 0, 0)$  it is a triplet if and only if  $xy = z$ , thus  $y = (0, 0, 0)$ .

(the first triplet is  $x = y = z = (0, 0, 0)$ ).

Form 2:  $x = (0, a_1, a_2), y = (b_0, 1, 1), z = (0, c_1, c_2)$  it is a triplet if and only if  $xz = y$ , thus  $b_0 = 0, a_1c_1 = 1, a_2c_2 = 1$ .

the possible triplets are:

$$x = (0, 1, 1), y = (0, 1, 1), z = (0, 1, 1).$$

$$x = (0, 1, -1), y = (0, 1, 1), z = (0, 1, -1)$$

$$x = (0, 1, -1), y = (0, 1, 1), z = (0, 1, -1)$$

$$x = (0, -1, 1), y = (0, 1, 1), z = (0, -1, -1)$$

Form 3:  $x = (0, 0, a_2), y = (b_0, b_1, 1), z = (0, 0, c_2)$  it is a triplet if and only if  $xz = y$ , thus  $b_0 = b_1 = 0, a_2 = c_2 = 1$ .

the possible triplets are:

$$x = (0, 0, 1), y = (0, 0, 1), z = (0, 1, 1).$$

$$x = (0, 0, -1), y = (0, 0, 1), z = (0, 1, -1)$$

Form 4:  $x = (a_0, 0, a_1), y = (1, b_1, 1), z = (c_0, 0, c_1)$  it is a triplet if and only if  $xz = y$ , thus  $a_0c_0 = a_1c_1 = 1, b_1 = 0$ .

the possible triplets are:

$$x = (1, 0, 1), y = (1, 0, 1), z = (1, 0, 1)$$

$$x = (-1, 0, 1), y = (1, 0, 1), z = (-1, 0, 1).$$

$$x = (1, 0, -1), y = (1, 0, 1), z = (1, 0, -1)$$

$$x = (-1, 0, -1), y = (1, 0, 1), z = (-1, 0, -1)$$

Form 5:  $x = (a_0, a_1, 0), y = (1, 1, b_2), z = (c_0, c_1, 0)$  it is a triplet if and only if  $xz = y$ , thus  $a_0c_0 = 1, a_1c_1 = 1, b_2 = 0$ .

the possible triplets are:

$$x = (1, 1, 0), y = (1, 1, 0), z = (1, 1, 0)$$

$$x = (-1, -1, 0), y = (1, 1, 0), z = (-1, -1, 0).$$

$$x = (-1, 1, 0), y = (1, 1, 0), z = (-1, 1, 0)$$

$$x = (1, 1, 0), y = (1, 1, -1), z = (1, -1, 0)$$

Form 6:  $x = (a_0, a_1, a_2), y = (1, 1, 1), z = (c_0, c_1, c_2)$  it is a triplet if and only if  $xz = y$ ,

$$\text{thus, } a_0c_0 = a_1c_1 = a_2c_2 = 1.$$

the possible triplets are:

$$x = (1, 1, 1), y = (1, 1, 1), z = (1, 1, 1)$$

$$x = (1, 1, -1), y = (1, 1, 1), z = (1, 1, -1)$$

$$x = (1, -1, 1), y = (1, 1, 1), z = (1, -1, 1)$$

$$x = (-1, 1, 1), y = (1, 1, 1), z = (-1, 1, 1)$$

$$x = (-1, -1, 1), y = (1, 1, 1), z = (-1, -1, 1)$$

$$x = (1, -1, -1), y = (1, 1, 1), z = (1, -1, -1)$$

$$x = (-1, 1, -1), y = (1, 1, 1), z = (-1, 1, -1)$$

$$x = (-1, -1, -1), y = (1, 1, 1), z = (-1, -1, -1)$$

Form 7:  $x = (a_0, 0, 0), y = (1, b_1, b_2), z = (c_0, 0, 0)$  it is a triplet if and only if  $xz = y$ ,

$$\text{thus, } a_0c_0 = 1, b_1 = b_2 = 0.$$

the possible triplets are:

$$x = (1, 0, 0), y = (1, 0, 0), z = (1, 0, 0)$$

$$x = (-1, 0, 0), y = (1, 0, 0), z = (-1, 0, 0)$$

Form 8:  $x = (0, a_1, 0), y = (b_0, 1, b_2), z = (0, c_1, 0)$  it is a triplet if and only if  $xz = y$ ,

$$\text{thus, } a_1c_1 = 1, b_0 = b_2 = 0.$$

the possible triplets are:

$$x = (0, 1, 0), y = (0, 1, 0), z = (0, 1, 0)$$

$$x = (0, -1, 0), y = (0, -1, 0), z = (0, 1, 0).$$

## References

1. Smarandache, F., "  $n$ -Valued Refined Neutrosophic Logic and Its Applications in Physics", Progress in Physics, 143-146, Vol. 4, 2013.
2. M. Ibrahim. A. Agboola, B.Badmus and S. Akinleye. On refined Neutrosophic Vector Spaces . International Journal of Neutrosophic Science, Vol. 7, 2020, pp. 97-109.

3. Abobala, M., "A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", International Journal of Mathematics and Mathematical Sciences, hindawi, 2021.
4. Celik, M., and Olgun, N., " On The Classification Of Neutrosophic Complex Inner Product Spaces", Galoitica Journal Of Mathematical Structures And Applications, 2022.
5. Celik, M., and Olgun, N., " An Introduction To Neutrosophic Real Banach And Hillbert Spaces", Galoitica Journal Of Mathematical Structures And Applications, 2022.
6. Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona, 2006.
7. Abobala, M, " $n$ -Cyclic Refined Neutrosophic Algebraic Systems Of Sub-Indeterminacies, An Application To Rings and Modules", International Journal of Neutrosophic Science, Vol. 12, pp. 81-95 . 2020.
8. Von Shtawzen, O., " Conjectures For Invertible Diophantine Equations Of 3-Cyclic and 4-Cyclic Refined Integers", Journal Of Neutrosophic And Fuzzy Systems, Vol.3, 2022.
9. Basheer, A., Ahmad, K., and Ali, R., " Examples On Some Novel Diophantine Equations Derived From The Group Of Units Problem In  $n$ -Cyclic Refined Neutrosophic Rings Of Integers", Galoitica Journal Of Mathematical Structures and Applications, 2022.
10. Sadiq, B., " A Contribution To The Group Of Units' Problem In Some 2-Cyclic Refined Neutrosophic Rings", International Journal Of Neutrosophic Science, 2022.
11. Khaldi, A., " A Study On Split-Complex Vector Spaces", Neoma Journal Of Mathematics and Computer Science, 2023.
12. Ahmad, K., " On Some Split-Complex Diophantine Equations", Neoma Journal Of Mathematics and Computer Science, 2023.
13. Ali, R., " On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces", Neoma Journal Of Mathematics and Computer Science, 2023.
14. Von Shtawzen, O., " On A Novel Group Derived From A Generalization Of Integer Exponents and Open Problems", Galoitica journal Of Mathematical Structures and Applications, Vol 1, 2022.

15. Hatip, A., "An Introduction To Weak Fuzzy Complex Numbers ", Galoitica Journal Of Mathematical Structures and Applications, Vol.3, 2023.
16. Merkepci, H., and Ahmad, K., " On The Conditions Of Imperfect Neutrosophic Duplets and Imperfect Neutrosophic Triplets", Galoitica Journal Of Mathematical Structures And Applications, Vol.2, 2022.

**Received:** December 24, 2022. **Accepted:** April 03, 2023