

4-11-2023

On Some Algorithms for Solving Different Types of Symbolic 2-Plithogenic Algebraic Equations

Ahmad Khaldi

Khadija Ben Othman

Oliver Von Shtawzen

Rozina Ali

Sarah Jalal Mosa

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Khaldi, Ahmad; Khadija Ben Othman; Oliver Von Shtawzen; Rozina Ali; and Sarah Jalal Mosa. "On Some Algorithms for Solving Different Types of Symbolic 2-Plithogenic Algebraic Equations." *Neutrosophic Sets and Systems* 54, 1 (). https://digitalrepository.unm.edu/nss_journal/vol54/iss1/9

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



On Some Algorithms for Solving Different Types of Symbolic 2-Plithogenic Algebraic Equations

¹Ahmad Khaldi,² Khadija Ben Othman,³ Oliver Von Shtawzen,⁴ Rozina Ali,⁵ Sarah Jalal Mosa

¹ Mutah University, Faculty of Science, Jordan, khaldiahmad1221@gmail.com

² Umm Al-Qura University, Mekka, Saudi Arabia, khagijabenothonman33@gmail.com

³ University Of Nizwa, Sultanate Of Oman, Vonshtawzen1970abc@gmail.com

⁴ Cairo University, Cairo, Egypt, rozyyy123n@gmail.com

⁵ College Of Engineering Techniques, Al-Farahidi University, Baghdad, Iraq
s.mosa@uoalfarahidi.edu.iq

Abstract:

The main goal of this paper is to study three different types of algebraic symbolic 2-plithogenic equations. The symbolic 2-plithogenic linear Diophantine equations, symbolic 2-plithogenic quadratic equations, and linear system of symbolic 2-plithogenic equations will be discussed and handled, where algorithms to solve the previous types will be presented and proved by transforming them to classical algebraic systems of equations.

Keywords: symbolic 2-plithogenic Diophantine equation, symbolic 2-plithogenic quadratic equation, linear system, symbolic 2-plithogenic field

Introduction and preliminaries

The process of extending classical algebraic structures by using logical symbols and elements can be considered as a novel approach to generalize algebraic structures, where many algebraic structures were generalized by using neutrosophic elements, fuzzy elements, and refined neutrosophic elements [1-15].

Smarandache has defined the concept of Symbolic 2-plithogenic sets and structures [16-20] as new generalizations of classical structures. Also, he has presented many open research problems [20].

In [21], Smarandache ideas was discussed in a special case of $n=2$, where the symbolic 2-plithogenic rings were defined and studied with many elementary interesting substructures and properties.

Let R be a ring, the symbolic 2-plithogenic ring is defined as follows:

$$2 - SP_R = \{a_0 + a_1P_1 + a_2P_2; a_i \in R, P_j^2 = P_j, P_1 \times P_2 = P_{\max(1,2)} = P_2\}.$$

Smarandache has defined algebraic operations on $2 - SP_R$ as follows:

Addition:

$$[a_0 + a_1P_1 + a_2P_2] + [b_0 + b_1P_1 + b_2P_2] = (a_0 + b_0) + (a_1 + b_1)P_1 + (a_2 + b_2)P_2.$$

Multiplication:

$$[a_0 + a_1P_1 + a_2P_2] \cdot [b_0 + b_1P_1 + b_2P_2] = a_0b_0 + a_0b_1P_1 + a_0b_2P_2 + a_1b_0P_1^2 + a_1b_2P_1P_2 + a_2b_0P_2 + a_2b_1P_1P_2 + a_2b_2P_2^2 + a_1b_1P_1P_1 = a_0b_0 + (a_0b_1 + a_1b_0 + a_1b_1)P_1 + (a_0b_2 + a_1b_2 + a_2b_0 + a_2b_1 + a_2b_2)P_2.$$

It is clear that $(2 - SP_R)$ is a ring.

If R is a field, then $2 - SP_R$ is called a symbolic 2-plithogenic field [22].

Also, the following open problems were asked in [22]:

Problem (3):

If F is a field then $2 - SP_F$ is called a 2-plithogenic symbolic field. Now, can we find a strong algorithm that explains how can we solve the previous equations by turning it into the classical equations.

Problem (4):

If Z is the ring of integers ring then $2 - SP_Z$ is called a 2-plithogenic symbolic ring of integers. Can we find a strong algorithm that explains how can we solve the previous equations by turning it into the classical Diophantine equations.

In this paper, we solve the previous two open problems by suggesting effective algorithms that help us to transform symbolic 2-plithogenic equations to classical algebraic equations.

Main Results

Definition.

Let $2 - SP_Z = \{a + bP_1 + cP_2; a, b, c \in Z\}$ be the symbolic 2-plithogenic ring of integers, the Diophantine equation with two variables is defined as follows:

$$AX + BY = C; A = a_0 + a_1P_1 + a_2P_2, B = b_0 + b_1P_1 + b_2P_2, C = c_0 + c_1P_1 + c_2P_2,$$

$$X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2, a_i, b_i, c_i, x_i, y_i \in 2 - SP_Z.$$

The following theorem describes an algorithm to solve the symbolic 2-plithogenic linear Diophantine equation with two variables.

Theorem.

Let $AX + BY = C$ be the symbolic 2-plithogenic linear Diophantine equation with two variables, it is solvable if and only if the following linear Diophantine equations are solvable.

$$\begin{cases} a_0x_0 + b_0y_0 = c_0 \\ (a_0 + a_1)(x_0 + x_1) + (b_0 + b_1)(y_0 + y_1) = c_0 + c_1 \\ (a_0 + a_1 + a_2)(x_0 + x_1 + x_2) + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2) = c_0 + c_1 + c_2 \end{cases}$$

Proof.

The equation $AX + BY = C$ equivalents:

$$\begin{aligned} & a_0x_0 + b_0y_0 + (a_0x_1 + a_1x_0 + a_1x_1 + b_0y_1 + b_1y_0 + b_1y_1)P_1 \\ & \quad + (a_0x_2 + a_2x_0 + a_2x_2 + a_1x_2 + a_2x_1 + b_0y_2 + b_2y_0 + b_2y_2 + b_1y_2 + b_2y_1)P_2 \\ & = c_0 + c_1P_1 + c_2P_2 \end{aligned}$$

$$\begin{cases} a_0x_0 + b_0y_0 = c_0 \dots (1) \\ a_0x_1 + a_1x_0 + a_1x_1 + b_0y_1 + b_1y_0 + b_1y_1 = c_1 \dots (2) \\ a_0x_2 + a_2x_0 + a_2x_2 + a_1x_2 + a_2x_1 + b_0y_2 + b_2y_0 + b_2y_2 + b_1y_2 + b_2y_1 = c_2 \dots (3) \end{cases}$$

We add (1) to (2), and (1) to (2) to (3), we get:

$$\begin{cases} a_0x_0 + b_0y_0 = c_0 \\ (a_0 + a_1)(x_0 + x_1) + (b_0 + b_1)(y_0 + y_1) = c_0 + c_1 \\ (a_0 + a_1 + a_2)(x_0 + x_1 + x_2) + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2) = c_0 + c_1 + c_2 \end{cases}$$

And the proof is complete.

The description of the algorithm.

To solve $AX + BY = C$ in $2 - SP_Z$, we must follow these steps.

Step1.

We compute $gcd(a_0, b_0), gcd(a_0 + a_1, b_0 + b_1), gcd(a_0 + a_1 + a_2, b_0 + b_1 + b_2)$.

If $gcd(a_0, b_0)/c_0, gcd(a_0 + a_1, b_0 + b_1)/c_0 + c_1, gcd(a_0 + a_1 + a_2, b_0 + b_1 + b_2)/c_0 + c_1 + c_2$, then it is solvable.

Step2.

We solve the equivalent system and get the values of $x_i, y_i; 0 \leq i \leq 2$.

Example.

Consider the following symbolic 2-plithogenic linear Diophantine equation:

$$(2 + P_1 + P_2)X + (3 + 2P_1 - P_2)Y = 8 + 5P_1 + 7P_2.$$

$$\gcd(a_0, b_0) = \gcd(2, 3) = 1/8.$$

$$\gcd(a_0 + a_1, b_0 + b_1) = \gcd(3, 5) = 1/c_0 + c_1 = 13.$$

$$\gcd(a_0 + a_1 + a_2, b_0 + b_1 + b_2) = \gcd(4, 4) = 4/c_0 + c_1 + c_2 = 20.$$

So that, the equation is solvable.

The equivalent system of linear Diophantine equations is:

$$\begin{cases} 2x_0 + 3y_0 = 8 \dots (1) \\ 3(x_0 + x_1) + 5(y_0 + y_1) = 13 \dots (2) \\ 4(x_0 + x_1 + x_2) + 4(y_0 + y_1 + y_2) = 20 \dots (3) \end{cases}$$

The equation (1) has a solution $(x_0 = 1, y_0 = 2)$.

The equation (2) has a solution $(x_0 + x_1 = 1, y_0 + y_1 = 2)$, there for $(x_1 = 0, y_1 = 0)$.

The equation (3) has a solution $(x_0 + x_1 + x_2 = 2, y_0 + y_1 + y_2 = 3)$, there for $(x_2 = 1, y_2 = 1)$.

This implies a solution $X = 1 + P_2, Y = 2 + P_2$.

Example.

Consider the following:

$$(3 + P_1 + 5P_2)X + (6 - 2P_1 + 10P_2)Y = 5 + P_1 + P_2.$$

$$\gcd(a_0, b_0) = \gcd(3, 6) = 3 \nmid 5, \text{ there for it is not solvable.}$$

2-symbolic plithogenic Quadratic equation.

Let $2 - SP_F$ be a symbolic 2-plithogenic field, the formula

$$AX^2 + BY^2 + C = 0; A = a_0 + a_1P_1 + a_2P_2, B = b_0 + b_1P_1 + b_2P_2,$$

$$C = c_0 + c_1P_1 + c_2P_2, X = x_0 + x_1P_1 + x_2P_2, Y = y_0 + y_1P_1 + y_2P_2, a_i, b_i, c_i, x_i, y_i \in 2 - SP_F.$$

Is called the symbolic 2-plithogenic quadratic equation.

Theorem.

Let $AX^2 + BY^2 + C = 0$ be a symbolic 2-plithogenic quadratic equation over $2 - SP_F$, then it is solvable if and only if the following system is solvable:

$$\begin{cases} a_0x_0^2 + b_0y_0^2 + c_0 = 0 \dots (1) \\ (a_0 + a_1)(x_0 + x_1)^2 + (b_0 + b_1)(y_0 + y_1)^2 + (c_0 + c_1) = 0 \dots (2) \\ (a_0 + a_1 + a_2)(x_0 + x_1 + x_2)^2 + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2)^2 + (c_0 + c_1 + c_2) = 0 \dots (3) \end{cases}$$

Proof.

We have $X^2 = x_0^2 + P_1[(x_0 + x_1)^2 - x_0^2] + P_2[(x_0 + x_1 + x_2)^2 - (x_0 + x_1)^2]$, see [].

So that:

$$\begin{aligned} AX^2 &= a_0x_0^2 + P_1[a_0(x_0 + x_1)^2 - a_0x_0^2 + a_1(x_0 + x_1)^2 - a_1x_0^2 + a_1x_0^2] \\ &\quad + P_2[a_0(x_0 + x_1 + x_2)^2 - a_0(x_0 + x_1)^2 + a_1(x_0 + x_1 + x_2)^2 - a_1(x_0 + x_1)^2 \\ &\quad + a_2(x_0 + x_1 + x_2)^2 - a_2(x_0 + x_1)^2 + a_2x_0^2 + a_2(x_0 + x_1)^2 - a_2x_0^2] \end{aligned}$$

$$\begin{aligned} AX^2 &= a_0x_0^2 + P_1[(a_0 + a_1)(x_0 + x_1)^2 - a_0x_0^2] \\ &\quad + P_2[(a_0 + a_1 + a_2)(x_0 + x_1 + x_2)^2 - (a_0 + a_1)(x_0 + x_1)^2] \end{aligned}$$

There for, the equation $AX^2 + BY^2 + C = 0$ is equivalent to:

$$\begin{cases} a_0x_0^2 + b_0y_0^2 + c_0 = 0 \dots (1) \\ (a_0 + a_1)(x_0 + x_1)^2 + (b_0 + b_1)(y_0 + y_1)^2 + (c_0 + c_1) = 0 \dots (2) \\ (a_0 + a_1 + a_2)(x_0 + x_1 + x_2)^2 + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2)^2 + (c_0 + c_1 + c_2) = 0 \dots (3) \end{cases}$$

The description of algorithm.

To solve $AX^2 + BY^2 + C = 0$ in $2 - SP_F$, follow these steps:

Step1.

Solve the equivalent classical system of quadratic equations. If (1),(2), and (3) are solvable in the field F , then the symbolic 2-plithogenic quadratic equation is solvable.

Step2.

Discuss all possible cases of x_0, x_1, x_2 .

Remark.

If $AX^2 + BY^2 + C = 0$ is solvable in $2 - SP_F$, then it has at most 8 solutions.

Example.

Consider the following:

$$(1 + P_1 + P_2)X^2 + (3 - P_1)X - 4 - 12P_2 = 0$$

We have:

$$\begin{cases} a_0 = 1, a_1 = 1, a_2 = 1 \\ b_0 = 3, b_1 = -1, b_2 = 0 \\ c_0 = -4, c_1 = 0, c_2 = -12 \end{cases}$$

The equivalent system is:

$$\begin{cases} x_0^2 + 3x_0 - 4 = 0 \dots (1) \\ 2(x_0 + x_1)^2 + 2(x_0 + x_1) - 4 = 0 \dots (2) \\ 3(x_0 + x_1 + x_2)^2 + 2(x_0 + x_1 + x_2) - 16 = 0 \dots (3) \end{cases}$$

The solutions of (1): $x_0 = 1, x_0 = -4$.

The solutions of (2): $x_0 + x_1 = 1, x_0 + x_1 = -2$.

The solutions of (3): $x_0 + x_1 + x_2 = 2, x_0 + x_1 + x_2 = -\frac{8}{3}$.

Case1.

If $x_0 = 1, x_0 + x_1 = 1, x_0 + x_1 + x_2 = 2$, then $x_1 = 0, x_2 = 1$, and $X = 1 + P_2$.

Case2.

If $x_0 = 1, x_0 + x_1 = 1, x_0 + x_1 + x_2 = -\frac{8}{3}$, then $x_1 = 0, x_2 = -\frac{11}{3}$, and $X = 1 - \frac{11}{3}P_2$.

Case3.

If $x_0 = 1, x_0 + x_1 = -2, x_0 + x_1 + x_2 = 2$, then $x_1 = -3, x_2 = 4$, and $X = 1 - 3P_1 + 4P_2$.

Case4.

If $x_0 = 1, x_0 + x_1 = -2, x_0 + x_1 + x_2 = -\frac{8}{3}$, then $x_1 = -3, x_2 = -\frac{2}{3}$, and $X = 1 - 3P_1 - \frac{2}{3}P_2$.

Case5.

If $x_0 = -4, x_0 + x_1 = 1, x_0 + x_1 + x_2 = 2$, then $x_1 = 5, x_2 = 1$, and $X = -4 + 5P_1 + P_2$.

Case6.

If $x_0 = -4, x_0 + x_1 = 1, x_0 + x_1 + x_2 = -\frac{8}{3}$, then $x_1 = 5, x_2 = -\frac{11}{3}$, and $X = -4 + 5P_1 - \frac{11}{3}P_2$.

Case7.

If $x_0 = -4, x_0 + x_1 = -2, x_0 + x_1 + x_2 = 2$, then $x_1 = 2, x_2 = 4$, and $X = -4 + 2P_1 + 4P_2$.

Case8.

If $x_0 = -4, x_0 + x_1 = -2, x_0 + x_1 + x_2 = -\frac{8}{3}$, then $x_1 = 2, x_2 = -\frac{2}{3}$, and $X = -4 + 2P_1 - \frac{2}{3}P_2$.

So that, the solutions of the original symbolic 2-plithogenic quadratic equation are:

$$X \in \left\{ -4 + 2P_1 - \frac{2}{3}P_2, -4 + 2P_1 + 4P_2, -4 + 5P_1 - \frac{11}{3}P_2, -4 + 5P_1 + P_2, 1 - 3P_1 - \frac{2}{3}P_2, 1 - 3P_1 + 4P_2, 1 - \frac{11}{3}P_2, 1 + P_2 \right\}$$

Example.

Consider the following:

$$(2 + 3P_1 - P_2)X^2 + (4 + P_1 + P_2)X - 6 - 4P_1 = 0$$

We have:

$$\begin{cases} a_0 = 2, a_1 = 3, a_2 = -1 \\ b_0 = 4, b_1 = 1, b_2 = 1 \\ c_0 = -6, c_1 = -4, c_2 = 0 \end{cases}$$

The equivalent system is:

$$\begin{cases} 2x_0^2 + 4x_0 - 6 = 0 \dots (1) \\ 5(x_0 + x_1)^2 + 5(x_0 + x_1) - 10 = 0 \dots (2) \\ 4(x_0 + x_1 + x_2)^2 + 6(x_0 + x_1 + x_2) - 10 = 0 \dots (3) \end{cases}$$

The solutions of (1): $x_0 = 1, x_0 = -3$.

The solutions of (2): $x_0 + x_1 = 1, x_0 + x_1 = -2$.

The solutions of (3): $x_0 + x_1 + x_2 = 1, x_0 + x_1 + x_2 = -\frac{5}{2}$.

Case1.

If $x_0 = 1, x_0 + x_1 = 1, x_0 + x_1 + x_2 = 1$, then $x_1 = x_2 = 0$, and $X = 1$.

Case2.

If $x_0 = 1, x_0 + x_1 = 1, x_0 + x_1 + x_2 = -\frac{5}{2}$, then $x_1 = 0, x_2 = -\frac{7}{2}$, and $X = 1 - \frac{5}{2}P_2$.

Case3.

If $x_0 = 1, x_0 + x_1 = -2, x_0 + x_1 + x_2 = 1$, then $x_1 = -3, x_2 = 3$, and $X = 1 - 3P_1 + 3P_2$.

Case4.

If $x_0 = 1, x_0 + x_1 = -2, x_0 + x_1 + x_2 = -\frac{5}{2}$, then $x_1 = -3, x_2 = -\frac{1}{2}$, and $X = 1 - 3P_1 - \frac{1}{2}P_2$.

Case5.

If $x_0 = -3, x_0 + x_1 = 1, x_0 + x_1 + x_2 = 1$, then $x_1 = 4, x_2 = 0$, and $X = -3 + 4P_1$.

Case6.

If $x_0 = -3, x_0 + x_1 = 1, x_0 + x_1 + x_2 = -\frac{5}{2}$, then $x_1 = 4, x_2 = -\frac{7}{2}$, and $X = -3 + 4P_1 - \frac{7}{2}P_2$.

Case7.

If $x_0 = -3, x_0 + x_1 = -2, x_0 + x_1 + x_2 = 1$, then $x_1 = 1, x_2 = 3$, and $X = -3 + P_1 + 3P_2$.

Case8.

If $x_0 = -3, x_0 + x_1 = -2, x_0 + x_1 + x_2 = -\frac{5}{2}$, then $x_1 = 1, x_2 = -\frac{1}{2}$, and $X = -3 + P_1 - \frac{1}{2}P_2$.

So that, the solutions of the original symbolic 2-plithogenic quadratic equation are:

$$X \in \left\{ 1, 1 - \frac{5}{2}P_2, 1 - 3P_1 + 3P_2, 1 - 3P_1 - \frac{1}{2}P_2, -3 + 4P_1, -3 + 4P_1 - \frac{7}{2}P_2, -3 + P_1 + 3P_2, -3 + P_1 - \frac{1}{2}P_2 \right\}$$

2-plithogenic Linear equations.

We begin the simplest case, a symbolic 2-plithogenic linear equation with one variable

$$A.X = B.$$

This equation is solvable uniquely if and only if A is invertible and $X = A^{-1}B$.

According to [31], $A^{-1} = a_0^{-1} + P_1[(a_0 + a_1)^{-1} - a_0^{-1}] + P_2[(a_0 + a_1 + a_2)^{-1} - (a_0 + a_1)^{-1}]$.

Example.

Consider the equation $(2 + P_1 + P_2)X = 3 - P_1$ over $2 - SP_R$.

$a_0 = 2, a_0^{-1} = \frac{1}{2}, a_0 + a_1 = 3, (a_0 + a_1)^{-1} = \frac{1}{3}, a_0 + a_1 + a_2 = 4, (a_0 + a_1 + a_2)^{-1} = \frac{1}{4}$, thus:

$A^{-1} = \frac{1}{2} - \frac{1}{6}P_1 - \frac{1}{12}P_2$, there for:

$$X = \left(\frac{1}{2} - \frac{1}{6}P_1 - \frac{1}{12}P_2\right)(3 - P_1) = \frac{3}{2} - \frac{1}{2}P_1 - \frac{1}{2}P_1 + \frac{1}{6}P_1 - \frac{1}{4}P_2 + \frac{1}{12}P_2 = \frac{3}{2} - \frac{5}{6}P_1 - \frac{1}{6}P_2$$

The general case is about a linear system of n symbolic 2-plithogenic equations $A_i \cdot X_i = B_i; 1 \leq i \leq n$.

To solve a system like that, we must transform it to an equivalent classical system. We present the following algorithm.

To solve the symbolic 2-plithogenic linear system:

$$\begin{cases} A_{11} \cdot X_1 + A_{12} \cdot X_2 + \dots + A_{1n} \cdot X_n = B_{1n} \\ A_{21} \cdot X_1 + A_{22} \cdot X_2 + \dots + A_{2n} \cdot X_n = B_{2n} \\ \vdots \\ A_{n1} \cdot X_1 + A_{n2} \cdot X_2 + \dots + A_{nn} \cdot X_n = B_{nn} \end{cases}$$

Where: $A_{ij} = a_{ij}^{(0)} + a_{ij}^{(1)}P_1 + a_{ij}^{(2)}P_2, X_i = X_i^{(0)} + X_i^{(1)}P_1 + X_i^{(2)}P_2, B_{ij} = b_{ij}^{(0)} + b_{ij}^{(1)}P_1 + b_{ij}^{(2)}P_2 \in 2 - SP_F$.

Follow these steps:

Step1.

Find the classical equivalent system as follows:

$$\begin{cases} \sum_{i,j=1}^n a_{ij}^{(0)} X_i^{(0)} = \sum_{i,j=1}^n b_{ij}^{(0)} \\ \sum_{i,j=1}^n (a_{ij}^{(0)} + a_{ij}^{(1)}) (X_i^{(0)} + X_i^{(1)}) = \sum_{i,j=1}^n (b_{ij}^{(0)} + b_{ij}^{(1)}) \\ \sum_{i,j=1}^n (a_{ij}^{(0)} + a_{ij}^{(1)} + a_{ij}^{(2)}) (X_i^{(0)} + X_i^{(1)} + X_i^{(2)}) = \sum_{i,j=1}^n (b_{ij}^{(0)} + b_{ij}^{(1)} + b_{ij}^{(2)}) \end{cases}$$

step2.

Solve each system and remark that:

The first system gives the values of $X_i^{(0)}; 1 \leq i \leq n$.

The second one gives the values of $X_i^{(0)} + X_i^{(1)}; 1 \leq i \leq n$.

The third one gives values of $X_i^{(0)} + X_i^{(1)} + X_i^{(2)}; 1 \leq i \leq n$.

Step3.

If each system is solvable, then the original 2-plithogenic system is solvable, and if the number of solutions of every classical system is k , then the number of solutions for the 2-plithogenic system is k^3 .

Example.

Consider the following symbolic 2-plithogenic system of three linear equations with three variables:

$$\begin{cases} (1 + P_2)X_1 + (3 - P_1)X_2 + (1 + P_1 - P_2)X_3 = 5 \\ P_2X_1 + P_1X_2 + (P_1 - P_2)X_3 = 2P_1 + 2P_2 \\ (1 + P_1 - P_2)X_1 + (4 + 3P_1 - P_2)X_2 + (5 + 2P_2)X_3 = 11 + 4P_2 \end{cases}$$

the equivalent classical systems are:

$$\begin{cases} X_1^{(0)} + 3X_2^{(0)} + X_3^{(0)} = 5 \\ 0X_1^{(0)} + 0X_2^{(0)} + 0X_3^{(0)} = 0 \quad \dots \text{system(1)} \\ 2X_1^{(0)} + 4X_2^{(0)} + 5X_3^{(0)} = 11 \end{cases}$$

$$\begin{cases} (X_1^{(0)} + X_1^{(1)}) + 2(X_2^{(0)} + X_2^{(1)}) + 2(X_3^{(0)} + X_3^{(1)}) = 5 \\ 0(X_1^{(0)} + X_1^{(1)}) + (X_2^{(0)} + X_2^{(1)}) + (X_3^{(0)} + X_3^{(1)}) = 2 \quad \dots \text{system(2)} \\ 3(X_1^{(0)} + X_1^{(1)}) + 7(X_2^{(0)} + X_2^{(1)}) + 5(X_3^{(0)} + X_3^{(1)}) = 15 \end{cases}$$

$$\begin{cases} 2(X_1^{(0)} + X_1^{(1)} + X_1^{(2)}) + 2(X_2^{(0)} + X_2^{(1)} + X_2^{(2)}) + (X_3^{(0)} + X_3^{(1)} + X_3^{(2)}) = 5 \\ (X_1^{(0)} + X_1^{(1)} + X_1^{(2)}) + (X_2^{(0)} + X_2^{(1)} + X_2^{(2)}) + 2(X_3^{(0)} + X_3^{(1)} + X_3^{(2)}) = 4 \quad \dots \text{system(3)} \\ 2(X_1^{(0)} + X_1^{(1)} + X_1^{(2)}) + 6(X_2^{(0)} + X_2^{(1)} + X_2^{(2)}) + 7(X_3^{(0)} + X_3^{(1)} + X_3^{(2)}) = 15 \end{cases}$$

The system(1) has infinite solutions, thus the 2-plithogenic system has infinite solutions.

We will find some solutions to clarify the algorithm.

For example system(1) has a solution $X_1^{(0)} = X_2^{(0)} = X_3^{(0)} = 1$.

The system (2) has a solution $X_1^{(0)} + X_1^{(1)} = X_2^{(0)} + X_2^{(1)} = X_3^{(0)} + X_3^{(1)} = 1$, thus $X_1^{(1)} = X_2^{(1)} = X_3^{(1)} = 0$.

The system (3) has a solution $X_1^{(0)} + X_1^{(1)} + X_1^{(2)} = X_2^{(0)} + X_2^{(1)} + X_2^{(2)} = X_3^{(0)} + X_3^{(1)} + X_3^{(2)} = 1$, there for $X_1^{(2)} = X_2^{(2)} = X_3^{(2)} = 0$, and $X_1 = X_1^{(0)} + X_1^{(1)}P_1 + X_1^{(2)}P_2 = 1$, $X_2 = 1, X_3 = 1$ is a solution for the 2-plithogenic system.

Also, the system (1) has a solution $X_1^{(0)} = \frac{13}{2}, X_2^{(0)} = -\frac{1}{2}, X_3^{(0)} = 0$.

The system (2) has a solution $X_1^{(0)} + X_1^{(1)} = X_2^{(0)} + X_2^{(1)} = X_3^{(0)} + X_3^{(1)} = 1$.

The system (3) has a solution $X_1^{(0)} + X_1^{(1)} + X_1^{(2)} = X_2^{(0)} + X_2^{(1)} + X_2^{(2)} = X_3^{(0)} + X_3^{(1)} + X_3^{(2)} = 1$.

There for $X_1^{(1)} = 1 - \frac{13}{2} = -\frac{11}{2}, X_2^{(1)} = 1 + \frac{1}{2} = \frac{3}{2}, X_3^{(1)} = 1, X_1^{(2)} = X_2^{(2)} = X_3^{(2)} = 0$.

This implies that:

$X_1 = \frac{13}{2} - \frac{11}{2}P_1, X_2 = -\frac{1}{2} + \frac{3}{2}P_1, X_3 = P_1$ is a solution of the 2-plithogenic system.

Conclusion

In this paper, we have presented novel algorithms to solve many different types of 2-plithogenic algebraic equations (quadratic, linear, and linear Diophantine equations) by transforming them to classical systems of algebraic equations. Also, many examples were illustrated to explain the validity of our work.

References

- [1] Abobala, M., Hatip, A., Bal, M., "A Study Of Some Neutrosophic Clean Rings", International journal of neutrosophic science, 2022.
- [2] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.
- [3] Celik, M., and Hatip, A., "On The Refined AH-Isometry And Its Applications In Refined Neutrosophic Surfaces", Galoitica Journal Of Mathematical Structures And Applications, 2022.
- [4] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81. 2020.
- [5] Ahmad, K., Thjeel, N., AbdulZahra, M., and Jaleel, R., "On The Classification of n-Refined Neutrosophic Rings And Its Applications In Matrix Computing Algorithms And Linear Systems", International Journal Of Neutrosophic Science, 2022.
- [6] Hatip, A., "An Introduction To Weak Fuzzy Complex Numbers ", Galoitica Journal Of Mathematical Structures and Applications, Vol.3, 2023.
- [7] Merkepçi, H., and Ahmad, K., "On The Conditions Of Imperfect Neutrosophic Duplets and Imperfect Neutrosophic Triplets", Galoitica Journal Of Mathematical Structures And Applications, Vol.2, 2022.
- [8] Von Shtawzen, O., "Conjectures For Invertible Diophantine Equations Of 3-Cyclic and 4-Cyclic Refined Integers", Journal Of Neutrosophic And Fuzzy Systems, Vol.3, 2022.

-
- [9] Abobala, M., and Zeina, M.B., " A Study Of Neutrosophic Real Analysis By Using One Dimensional Geometric AH-Isometry", *Galoitica Journal Of Mathematical Structures And Applications*, Vol.3, 2023.
- [10] Khaldi, A., " A Study On Split-Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [11] Ahmad, K., " On Some Split-Complex Diophantine Equations", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [12]. Ali, R., " On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [13] Sarkis, M., " On The Solutions Of Fermat's Diophantine Equation In 3-refined Neutrosophic Ring of Integers", *Neoma Journal of Mathematics and Computer Science*, 2023.
- [14] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, *Journal Of Mathematics*, Hindawi, 2021
- [15] Abobala, M., Ziena, B,M., Doewes, R., and Hussein, Z., "The Representation Of Refined Neutrosophic Matrices By Refined Neutrosophic Linear Functions", *International Journal Of Neutrosophic Science*, 2022.
- [16] F. Smarandache, *Plithogeny, Plithogenic Set, Logic, Probability, and Statistics*, 141 pages, Pons Editions, Brussels, Belgium, 2017. arXiv.org (Cornell University), Computer Science - Artificial Intelligence, 03Bxx:
- [17] Florentin Smarandache, *Physical Plithogenic Set*, 71st Annual Gaseous Electronics Conference, Session LW1, Oregon Convention Center Room, Portland, Oregon, USA, November 5–9, 2018.
- [18] Florentin Smarandache: *Plithogenic Set, an Extension of Crisp, Fuzzy, Intuitionistic Fuzzy, and Neutrosophic Sets – Revisited*, *Neutrosophic Sets and Systems*, vol. 21, 2018, pp. 153-166.
- [19] Florentin Smarandache, *Plithogenic Algebraic Structures*. Chapter in "Nidus idearum Scilogs, V: joining the dots" (third version), Pons Publishing Brussels, pp. 123-125, 2019.

-
- [20] Smarandache, F., " Introduction to the Symbolic Plithogenic Algebraic Structures (revisited)", *Neutrosophic Sets and Systems*, vol. 53, 2023.
- [21] Merkepci, H., and Abobala, M., " On The Symbolic 2-Plithogenic Rings", *International Journal of Neutrosophic Science*, 2023.
- [22] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", *Neutrosophic Sets and Systems*, Vol. 43, 2021.
- [23] Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", *Mathematical Problems in Engineering*, Hindawi, 2021
- [24] Agboola, A.A.A., Akinola, A.D., and Oyebola, O.Y., " Neutrosophic Rings I" , *International J.Mathcombin*, Vol 4,pp 1-14. 2011
- [25] Smarandache, F., and Ali, M., "Neutrosophic Triplet Group", *Neural. Compute. Appl.* 2019.
- [26] Chellamani, P., and Ajay, D., "Pythagorean neutrosophic Fuzzy Graphs", *International Journal of Neutrosophic Science*, Vol. 11, 2021.
- [27] Giorgio, N, Mehmood, A., and Broumi, S.," Single Valued neutrosophic Filter", *International Journal of Neutrosophic Science*, Vol. 6, 2020.

Received: December 31, 2022. **Accepted:** March 30, 2023