

5-1-2023

Neutrosophic n-Valued Refined Sets and Topologies

Murad Ara

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Ara, Murad. "Neutrosophic n-Valued Refined Sets and Topologies." *Neutrosophic Sets and Systems* 55, 1 (2023). https://digitalrepository.unm.edu/nss_journal/vol55/iss1/19

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



Neutrosophic n -Valued Refined Sets and Topologies

Murad Arar *

Mathematics Department, Prince Sattam Bin Abdulaziz University-Al Kharj, Saudi Arabia;

m.arar@psau.edu.sa , muradshhada@gmail.com

*Correspondence: m.arar@psau.edu.sa

Abstract. In n -Valued refined logic truth value T can be split into many types of truths: T_1, T_2, \dots, T_p and I into many types of indeterminacies: I_1, I_2, \dots, I_r and F into many types of falsities: F_1, F_2, \dots, F_s , where p, r and s are integers greater than 1, and $p + r + s = n$. Importance of n -valued refined logic and sets appeared in different applications specially in medical diagnosis. In this paper we post a condition on neutrosophic n -valued refined sets to make them functional to be applied in different mathematical branches. We define and study n -valued refined topological spaces. We defined neutrosophic n -valued refined α -open, β -open, pre-open and semi-open sets and studied their properties. We constructed different counter examples to clarify the relations between these different types of neutrosophic n -valued refined generalized open sets.

Keywords: n -valued refined topology; refined logic; refined sets; n -valued refined α -open; semi-open sets; n -valued refined generalized open sets.)

1. INTRODUCTION

Neutrosophic sets are, first, introduced in 2005 by [26,27] as a generalization of intuitionistic fuzzy sets [13], where any element $x \in X$ we have three degrees; the degree of membership(T), indeterminacy(I), and non-membership(F). Neutrosophic vague sets are introduced in 2015 by [30]. Neutrosophic vague topological spaces introduced in [21] we are many different notations are introduced and studied such as neutrosophic vague continuity and compactness.

Neutrosophic topologies are defined and studied by Smarandache [27], Lupianez [19,20] and Salama [?]. Open and closed neutrosophic sets, interior, exterior, closure and boundary of neutrosophic sets can be found in [29].

Neutrosophic sets applied to generalize many notations about soft topology and applications [18], [23], [16], generalized open and closed sets [31] , fixed point theorems [18] , graph theory

[17]and rough topology and applications [22]. Neutrosophy has many applications especially in decision making, for more details about new trends of neutrosophic applications one can consult [1]- [7].

Generalized topology and continuity introduced in 2002 in [?] which is a generalization of topological spaces and has different properties than general topology, see for example [8], [11] and [12]. Neutrosophic generalized sets and topologies are introduced and studies by Murad M. Arar in 2020 see [9] and [10]. In n -valued refined logic truth value T can be split into many types of truths: T_1, T_2, \dots, T_p and I into many types of indeterminacies: I_1, I_2, \dots, I_r and F into many types of falsities: F_1, F_2, \dots, F_s , where p, r and s are integers greater than 1, and $p + r + s = n$ see [28]. Importance of n -valued refined logic and sets appeared in different applications specially in medial diagnosis see [25] and [14], where a strong assumption is assumed to make them functional; that is $p = r = s$.

Definition 1.1. [26]: We say that the set A is *neutrosophic* on X if

$$A = \{ \langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle; x \in X \}; \mu, \sigma, \nu : X \rightarrow]-0, 1^+[\text{ and } -0 \leq \mu(x) + \sigma(x) + \nu(x) \leq 3^+.$$

The class of all neutrosophic sets on the universe X will be denoted by $\mathcal{N}(X)$. The basic neutrosophic operations (inclusion, union, and intersection) were first introduced by [24].

Definition 1.2 (*Neutrosophic sets operations*). Let $A, A_\alpha, B \in \mathcal{N}(X)$ such that $\alpha \in \Delta$. Then we define the neutrosophic:

- (1) (*Inclusion*): $A \sqsubseteq B$ If $\mu_A(x) \leq \mu_B(x)$, $\sigma_A(x) \geq \sigma_B(x)$ and $\nu_A(x) \geq \nu_B(x)$.
- (2) (*Equality*): $A = B \Leftrightarrow A \sqsubseteq B$ and $B \sqsubseteq A$.
- (3) (*Intersection*) $\bigcap_{\alpha \in \Delta} A_\alpha(x) = \{ \langle x, \bigwedge_{\alpha \in \Delta} \mu_{A_\alpha}(x), \bigvee_{\alpha \in \Delta} \sigma_{A_\alpha}(x), \bigvee_{\alpha \in \Delta} \nu_{A_\alpha}(x) \rangle; x \in X \}$.
- (4) (*Union*) $\bigcup_{\alpha \in \Delta} A_\alpha(x) = \{ \langle x, \bigvee_{\alpha \in \Delta} \mu_{A_\alpha}(x), \bigwedge_{\alpha \in \Delta} \sigma_{A_\alpha}(x), \bigwedge_{\alpha \in \Delta} \nu_{A_\alpha}(x) \rangle; x \in X \}$.
- (5) (*Complement*) $A^c = \{ \langle x, \nu_A(x), 1 - \sigma_A(x), \mu_A(x) \rangle; x \in X \}$
- (6) (*Universal set*) $1_X = \{ \langle x, 1, 0, 0 \rangle; x \in X \}$; called the *neutrosophic universal set*.
- (7) (*Empty set*) $0_X = \{ \langle x, 0, 1, 1 \rangle; x \in X \}$; called the *neutrosophic empty set*.

Proposition 1.3. [24] For $A, A_\alpha \in \mathcal{N}(X)$ for every $\alpha \in \Delta$ we have:

- (1) $A \cap (\bigcup_{\alpha \in \Delta} A_\alpha) = \bigcup_{\alpha \in \Delta} (A \cap A_\alpha)$.
- (2) $A \cup (\bigcap_{\alpha \in \Delta} A_\alpha) = \bigcap_{\alpha \in \Delta} (A \cup A_\alpha)$.

Definition 1.4. [24] [*Neutrosophic Topology*] $\tau \subseteq \mathcal{N}(X)$ is called a *neutrosophic topology* for X if

- (1) $0_X, 1_X \in \tau$.
- (2) If $A_\alpha \in \tau$ for every $\alpha \in \Delta$, then $\bigcup_{\alpha \in \Delta} A_\alpha \in \tau$,
- (3) For every $A, B \in \tau$, we have $A \cap B \in \tau$.

The ordered pair (X, τ) will be said a *neutrosophic space* over X . The elements of τ will be called *neutrosophic open sets*. For any $A \in \mathcal{N}(X)$, If $A^c \in \tau$, then we say A is *neutrosophic closed*.

2. NEUTROSOPHIC n -VALUED REFINED SETS AND TOPOLOGY

In neutrosophic n -valued refined logic (see [28]) the membership degree refined (split) into r values $\mu_1, \mu_2, \dots, \mu_r$, the indeterminacy refined into s values $\sigma_1, \sigma_2, \dots, \sigma_s$ and the nonmembership refined into t values $\nu_1, \nu_2, \dots, \nu_t$ such that $n = r + s + t$ and

$$-0 \leq \sum_{i=1}^r \mu_i + \sum_{i=1}^s \sigma_i + \sum_{i=1}^t \nu_i \leq n^+$$

Some authors assumes that $r = s = t$ see for example [14]. Actually, there is no guarantee that the membership, indeterminacy and nonmembership degrees refined or split into the same number of values, and we will not get a functional system of Neutrosophic n -valued refined sets if no more restrictions are assumed on r, s and t . This occurs when we define the basic set operations on the neutrosophic n -valued refined sets, especially when we try to define the neutrosophic n -valued refined complement of a given neutrosophic n -valued refined set; where r plays the role of t and vice versa. We will be back to this discussion after stating some definitions and theorems.

Definition 2.1. [26]: A is called a *neutrosophic n -valued refined set* on a universe X if $A = \{ \langle x, \mu_A^1(x), \mu_A^2(x), \dots, \mu_A^r(x); \sigma_A^1(x), \sigma_A^2(x), \dots, \sigma_A^s(x); \nu_A^1(x), \nu_A^2(x), \dots, \nu_A^t(x) \rangle; x \in X \}; \mu_A^i, \sigma_A^j, \nu_A^k : X \rightarrow]-0, 1^+[$ for every $i = 1, \dots, r, j = 1, \dots, s, k = 1, \dots, t$ such that $r + s + t = n$ and

$$-0 \leq \sum_{i=1}^r \mu_A^i(x) + \sum_{j=1}^s \sigma_A^j + \sum_{k=1}^t \nu_A^k \leq n^+.$$

The class of all neutrosophic n -valued refined sets on the universe X will be denoted by $\mathcal{R}_n(X)$.

The following is the definition of the basic operations (inclusion, union, intersection and complement) on neutrosophic n -valued refined sets.

Definition 2.2. [*Neutrosophic n -valued refined sets operations*] Let $A, A_\alpha, B \in \mathcal{R}_n(X)$ such that $\alpha \in \Delta$. Then we define the neutrosophic n -valued refined:

- (1) (*Inclusion*): $A \sqsubseteq_R B$ If $\mu_A^i(x) \leq \mu_B^i(x), \sigma_A^j(x) \geq \sigma_B^j(x)$ and $\nu_A^k(x) \geq \nu_B^k(x)$ for every $i = 1, \dots, r, j = 1, \dots, s, k = 1, \dots, t$.
- (2) (*Equality*): $A = B \Leftrightarrow A \sqsubseteq_R B$ and $B \sqsubseteq_R A$.
- (3) (*Intersection*) $\bigcap_{\alpha \in \Delta_R} A_\alpha(x) = \{ \langle x, \bigwedge_{\alpha \in \Delta} \mu_{A_\alpha}^1(x), \dots, \bigwedge_{\alpha \in \Delta} \mu_{A_\alpha}^r(x); \bigvee_{\alpha \in \Delta} \sigma_A^1(x), \dots, \bigvee_{\alpha \in \Delta} \sigma_A^s(x); \bigvee_{\alpha \in \Delta} \nu_A^1(x), \dots, \bigvee_{\alpha \in \Delta} \nu_A^t(x) \rangle; x \in X \}$.

- (4) (Union) $\sqcup_{\alpha \in \Delta_R} A_\alpha(x) = \{\langle x, \bigvee_{\alpha \in \Delta} \mu_{A_\alpha}^1(x), \dots, \bigvee_{\alpha \in \Delta} \mu_{A_\alpha}^r(x); \bigwedge_{\alpha \in \Delta} \sigma_A^1(x), \dots, \bigwedge_{\alpha \in \Delta} \sigma_A^s(x); \bigwedge_{\alpha \in \Delta} \nu_A^1(x), \dots, \bigwedge_{\alpha \in \Delta} \nu_A^t(x) \rangle; x \in X\}$.
- (5) (Complement) $A^c = \{\langle x, \nu_A^1(x), \dots, \nu_A^t(x); 1 - \sigma_A^1(x), \dots, 1 - \sigma_A^s(x); \mu_A^1(x), \dots, \mu_A^r(x) \rangle; x \in X\}$
- (6) (Universal set) $1_X = \{\langle x, 1, \dots, 1; 0, \dots, 0; 0, \dots, 0 \rangle; x \in X\}$; called the *neutrosophic n-valued refined universal set*.
- (7) (Empty set) $0_X = \{\langle x, 0, \dots, 0; 1, \dots, 1; 1, \dots, 1 \rangle; x \in X\}$; called the *neutrosophic n-valued refined empty set*.

Theorem 2.3. Let $A_\alpha, A, B \in \mathcal{R}_n(X)$ such that $\alpha \in \Delta$. Then we have

- (1) If $A \sqsubseteq_R B \sqsubseteq_R C$, then $A \sqsubseteq_R C$.
 - (2) If $A \sqsubseteq_R B$, then $B^c \sqsubseteq_R A^c$.
 - (3) $(\sqcup_{\alpha \in \Delta_R} A_\alpha) \sqcap_R A = \sqcup_{\alpha \in \Delta_R} (A_\alpha \sqcap_R A)$
 - (4) $(\sqcap_{\alpha \in \Delta_R} A_\alpha) \sqcup_R A = \sqcap_{\alpha \in \Delta_R} (A_\alpha \sqcup_R A)$
- [Demorgan's Laws]
- (5) $(A \sqcup_R B)^c = A^c \sqcap_R B^c$
 - (6) $(A \sqcap_R B)^c = A^c \sqcup_R B^c$

Proof. (1) and (2) are Straight forward! (3) and (4) can be proved using the following two propositions:

$$\begin{aligned}
 & - (\bigvee_{\alpha \in \Delta} a_\alpha) \wedge b = \bigvee_{\alpha \in \Delta} (a_\alpha \wedge b) \\
 & - (\bigwedge_{\alpha \in \Delta} a_\alpha) \vee b = \bigwedge_{\alpha \in \Delta} (a_\alpha \vee b)
 \end{aligned}$$

Now, we prove (3) and (4) can be proved by duality:

$$\begin{aligned}
 (A \sqcup_R B)^c &= (\{\langle x, \mu_A^1(x) \vee \mu_B^1(x), \dots, \mu_A^r(x) \vee \mu_B^r(x); \sigma_A^1(x) \wedge \sigma_B^1(x), \dots, \sigma_A^s(x) \wedge \sigma_B^s(x); \nu_A^1(x) \wedge \nu_B^1(x), \dots, \nu_A^t(x) \wedge \nu_B^t(x) \rangle; x \in X\})^c \\
 &= \{\langle x, \nu_A^1(x) \wedge \nu_B^1(x), \dots, \nu_A^t(x) \wedge \nu_B^t(x); 1 - (\sigma_A^1(x) \wedge \sigma_B^1(x)), \dots, 1 - (\sigma_A^s(x) \wedge \sigma_B^s(x)); \mu_A^1(x) \vee \mu_B^1(x), \dots, \mu_A^r(x) \vee \mu_B^r(x) \rangle; x \in X\} \\
 &= \{\langle x, \nu_A^1(x) \wedge \nu_B^1(x), \dots, \nu_A^t(x) \wedge \nu_B^t(x); (1 - \sigma_A^1(x)) \vee (1 - \sigma_B^1(x)), \dots, (1 - \sigma_A^s(x)) \vee (1 - \sigma_B^s(x)); \mu_A^1(x) \vee \mu_B^1(x), \dots, \mu_A^r(x) \vee \mu_B^r(x) \rangle; x \in X\} \\
 &= \{\langle x, \nu_A^1(x), \dots, \nu_A^t(x); 1 - \sigma_A^1(x), \dots, 1 - \sigma_A^s(x); \mu_A^1(x), \dots, \mu_A^r(x) \rangle; x \in X\} \sqcap_R \\
 &\{\langle x, \nu_B^1(x), \dots, \nu_B^t(x); 1 - \sigma_B^1(x), \dots, 1 - \sigma_B^s(x); \mu_B^1(x), \dots, \mu_B^r(x) \rangle; x \in X\} = A^c \sqcap_R B^c \quad \square
 \end{aligned}$$

So, as the above theorem shows, the system defined in Definition 2.2 is rich to a certain extent, but it still needs to be stronger to deal with some situations: for example $A \sqcap_R A^c$ is not well-defined if $r \neq t$. The concept *True* (membership) and *False* (nonmembership) are related, it is reasonable to discuss them in any world simultaneously, so we can assume $r = t$, and this is what F. Smarandache did in [28] when he discussed the relative (absolute)

truth and falsity simultaneously. The condition $r = s = t$ mentioned in [14] is very strong and will not add any value to us, actually it implies that n is divisible by 3, since $n = r + s + t$, so it does not include some worlds, for example a world of seven and five-valued logic which discussed in [28]. On the other hand if we, only, assume $r = t$, then n can be any value since we have not assumed any condition on s and worlds of any n -valued logic will be included.

Definition 2.4. : Let A be a *neutrosophic n -valued refined set* on a universe X . If $r = s$, then we call A a *homogeneous neutrosophic n -valued refined set*. n will be called the dimension of A , and r, s will be called the sub-dimensions of A . The class of all *homogeneous neutrosophic n -valued refined sets* on the universe X with sub-dimensions r, s will be denoted by $\mathcal{R}_{(n,r,s)}(X)$.

The following is obvious:

Proposition 2.5. Let $A, B \in \mathcal{R}_{(n,r,s)}(X)$. Then

- (1) $A \sqcap_R B \in \mathcal{R}_{(n,r,s)}(X)$.
- (2) $A \sqcup_R B \in \mathcal{R}_{(n,r,s)}(X)$.
- (3) $A^c \in \mathcal{R}_{(n,r,s)}(X)$.

Example 2.6. Let $X = \{a, b\}$, and let $A, B \in \mathcal{R}_{(5,2,1)}(X)$ such that $A = \{\langle a, 0.2, 0.1; 0.7; 0.1, 0.4 \rangle, \langle b, 0.5, 0.3; 0.2; 0.9, 0.5 \rangle\}$ and $B = \{\langle a, 0.4, 0.01; 0.3; 0.4, 0.3 \rangle, \langle b, 0.4, 0.2; 0.1; 0.7, 0.7 \rangle\}$. Then we have:
 $A \sqcap_R B = \{\langle a, 0.2, 0.01; 0.7; 0.4, 0.4 \rangle, \langle b, 0.4, 0.2; 0.2; 0.9, 0.7 \rangle\} \in \mathcal{R}_{(5,2,1)}$
 $A \sqcup_R B = \{\langle a, 0.4, 0.1; 0.3; 0.1, 0.3 \rangle, \langle b, 0.5, 0.3; 0.1; 0.7, 0.5 \rangle\} \in \mathcal{R}_{(5,2,1)}$
 $A^c = \{\langle a, 0.1, 0.4; 0.3; 0.2, 0.1 \rangle, \langle b, 0.9, 0.5; 0.8; 0.5, 0.3 \rangle\} \in \mathcal{R}_{(5,2,1)}$

Definition 2.7 (*Neutrosophic n -valued Refined Topology*). $\tau \subseteq \mathcal{R}_{(n,r,s)}(X)$ is called a *neutrosophic n -valued refined topology* on X if

- (1) $0_X, 1_X \in \tau$.
- (2) For every $A, B \in \tau$, we have $A \sqcap_R B \in \tau$.
- (3) If $A_\alpha \in \tau$ for every $\alpha \in \Delta$, then $\sqcup_R A_\alpha \in \tau$,

Elements of τ are called *neutrosophic n -valued refined open sets*. $A \in \mathcal{R}_{(n,r,s)}(X)$ is said *neutrosophic n -valued refined closed set* if $A^c \in \tau$.

The class of all neutrosophic n -valued refined topologies on X with sub-dimensions r, s will be denoted by $TOP_{(n,r,s)}(X)$.

Definition 2.8. Let $\tau \subseteq \mathcal{R}_{(n,r,s)}(X)$ be a neutrosophic n -valued refined topology on X and let $A \in \mathcal{R}_{(n,r,s)}(X)$. Then:

- (1) The neutrosophic n -valued refined interior of A is defined to be $Int_R(A) = \sqcup_R \{O \in \tau; O \sqsubseteq_R A\}$.

(2) The neutrosophic n -valued refined closure of A is defined to be

$$Cl_R(A) = \sqcap_R \{C \in \mathcal{R}_{(n,r,s)}(X); C^c \in \tau \text{ and } A \sqsubseteq_R C\}$$

Example 2.9. Let $X = \{a, b\}$, and let $\tau = \{0_X, 1_X, A, B, C, D\} \subset \mathcal{R}_{(5,2,1)}(X)$ where

$$A = \{\langle a, 0.2, 0.1; 0.7; 0.1, 0.4 \rangle, \langle b, 0.5, 0.3; 0.2; 0.9, 0.5 \rangle\},$$

$$B = \{\langle a, 0.4, 0.01; 0.3; 0.4, 0.3 \rangle, \langle b, 0.4, 0.2; 0.1; 0.7, 0.7 \rangle\},$$

$$C = \{\langle a, 0.2, 0.01; 0.7; 0.4, 0.4 \rangle, \langle b, 0.4, 0.2; 0.2; 0.9, 0.7 \rangle\}$$

$$D = \{\langle a, 0.4, 0.1; 0.3; 0.1, 0.3 \rangle, \langle b, 0.5, 0.3; 0.1; 0.7, 0.5 \rangle\}$$

Then τ is a Neutrosophic 5-valued refined topology on X . All closed set are: $0_X, 1_X, A^c, B^c, C^c, D^c$ where

$$A^c = \{\langle a, 0.1, 0.4; 0.3; 0.2, 0.1 \rangle, \langle b, 0.9, 0.5; 0.8; 0.5, 0.3 \rangle\}$$

$$B^c = \{\langle a, 0.4, 0.3; 0.7; 0.4, 0.01 \rangle, \langle b, 0.7, 0.7; 0.9; 0.4, 0.2 \rangle\},$$

$$C^c = \{\langle a, 0.4, 0.4; 0.3; 0.2, 0.01 \rangle, \langle b, 0.9, 0.7; 0.8; 0.4, 0.2 \rangle\}$$

$$D^c = \{\langle a, 0.1, 0.3; 0.7; 0.4, 0.1 \rangle, \langle b, 0.7, 0.5; 0.9; 0.5, 0.3 \rangle\}$$

Let $K = \{\langle a, 0.43, 0.09; 0.2; 0.1, 0.2 \rangle, \langle b, 0.5, 0.25; 0.1; 0.5, 0.6 \rangle\}$. Then the open sets in τ contained in K are only $0_X, B, C$, so that $Int_R(K) = 0_X \sqcup_R B \sqcup_R C = B$. Now; we consider the set $K^c = \{\langle a, 0.1, 0.2; 0.8; 0.43, 0.09 \rangle, \langle b, 0.5, 0.6; 0.9; 0.5, 0.25 \rangle\}$ and compute $Cl_R(K^c)$; the only closed sets containing K^c are $1_X, B^c$ and C^c , so that $Cl_R(K^c) = 1_X \sqcap_R B^c \sqcap_R C^c = B^c$. Which means $Cl_R(K^c) = B^c$ and so $(Cl_R(K^c))^c = B = Int_R(K)$; that is $Int_R(K) = (Cl_R(K^c))^c$ and this leads us to the following theorem:

Theorem 2.10. Let (X, τ) be an n -valued refined topological space with sub-dimensions r, s and let $A \in \mathcal{R}_{(n,r,s)}(X)$. Then we have:

- (1) $Int_R(A) = (Cl_R(A^c))^c$
- (2) $Cl_R(K) = (Int_R(K^c))^c$

Proof. Since \vee and \wedge has duality, we will, only, proof part (1).

Let $A = \{\langle x, \mu_A^1(x), \dots, \mu_A^r(x); \sigma_A^1(x), \dots, \sigma_A^s(x); \nu_A^1(x), \dots, \nu_A^r(x) \rangle; x \in X\}$. Then

$$A^c = \{\langle x, \nu_A^1(x), \dots, \nu_A^r(x); 1 - \sigma_A^1(x), \dots, 1 - \sigma_A^s(x); \mu_A^1(x), \dots, \mu_A^r(x) \rangle; x \in X\}, \text{ so}$$

$Cl_R(A^c) = \sqcap_R \{C \in \mathcal{R}_{(n,r,s)}(X); C^c \in \tau \text{ and } A^c \sqsubseteq_R C\}$. We apply Demorgan's Laws in Theorem 2.3 to get: $(Cl_R(A^c))^c = \sqcup_R \{C^c \in \mathcal{R}_{(n,r,s)}(X); C^c \in \tau \text{ and } C^c \sqsubseteq_R A\} = \sqcup_R \{O \in \mathcal{R}_{(n,r,s)}(X); O \in \tau \text{ and } O \sqsubseteq_R A\} = Int_R(A)$.

□

Theorem 2.11. Let (X, τ) be an n -valued refined topological space with sub-dimensions r, s and let $A, B \in \mathcal{R}_{(n,r,s)}(X)$. Then we have:

- (1) $Int_R(A) \sqsubseteq_R A$.
- (2) If A is a neutrosophic n -valued refined open set, then $Int_R(A) = A$.

- (3) $Int_R(Int_R(A)) = Int_R(A)$.
- (4) If $A \sqsubseteq_R B$, then $Int_R(A) \sqsubseteq_R Int_R(B)$.
- (5) $Int_R(A \sqcap_R B) = Int_R(A) \sqcap_R Int_R(B)$
- (6) $Int_R(A \sqcup_R B) \supseteq_R Int_R(A) \sqcup_R Int_R(B)$
- (7) $Int_R(\bigsqcup_{\alpha \in \Delta} A_\alpha) \supseteq_R \bigsqcup_{\alpha \in \Delta} Int_R(A_\alpha)$
- (8) $A \sqsubseteq_R Cl_R(A)$.
- (9) If A is a neutrosophic n -valued refined closed set, then $Cl_R(A) = A$.
- (10) $Cl_R(Cl_R(A)) = Cl_R(A)$.
- (11) If $A \sqsubseteq_R B$, then $Int_R(A) \sqsubseteq_R Int_R(B)$.
- (12) $Cl_R(A \sqcup_R B) = Cl_R(A) \sqcup_R Cl_R(B)$
- (13) $Cl_R(A \sqcap_R B) \sqsubseteq_R Cl_R(A) \sqcap_R Cl_R(B)$
- (14) $Cl_R(\bigsqcup_{\alpha \in \Delta} A_\alpha) \supseteq_R \bigsqcup_{\alpha \in \Delta} Cl_R(A_\alpha)$

Proof. (1) Let $O \in \tau$ such that $O \sqsubseteq_R A$. Then for every $x \in X$ we have $\mu_O^i(x) \leq \mu_A^i(x)$ for every $i = 1, \dots, r$, $\sigma_O^i(x) \geq \sigma_A^i(x)$ for every $i = 1, \dots, s$ and $\nu_O^i(x) \geq \nu_A^i(x)$ for every $i = 1, \dots, r$, which implies that $\bigvee_{O \in \tau, O \sqsubseteq_R A} \mu_O^i(x) \leq \mu_A^i(x)$ for every $i = 1, \dots, r$, $\bigwedge_{O \in \tau, O \sqsubseteq_R A} \sigma_O^i(x) \geq \sigma_A^i(x)$ for every $i = 1, \dots, s$ and $\bigwedge_{O \in \tau, O \sqsubseteq_R A} \nu_O^i(x) \geq \nu_A^i(x)$ for every $i = 1, \dots, r$; that is $Int_R(A) \sqsubseteq A$.

- (2) Since A is open, then, from the definition of $Int_R(A)$, we have $A \sqsubseteq_R Int_R(A)$, and from part (1) we have the converse, and we done.
- (3) Since $Int_R(A)$ is a neutrosophic n -valued refined open set, we have (from part (2)) $Int_R(Int_R(A)) = Int_R(A)$.
- (4) Let O be a neutrosophic n -valued refined open set such that $O \sqsubseteq_R A$. Then since $A \sqsubseteq_R B$, we have $O \sqsubseteq_R B$, that is $Int_R(A) \sqsubseteq_R Int_R(B)$
- (5) From part (4) we have $Int_R(A \sqcap_R B) \sqsubseteq_R Int_R(A) \sqcap_R Int_R(B)$. On the other hand, $Int_R(A) \sqcap_R Int_R(B)$ is a neutrosophic n -valued refined open set contained in A and B , so that $Int_R(A) \sqcap_R Int_R(B) \sqsubseteq_R Int_R(A \sqcap_R B)$, and we done.
- (6) Since $Int_R(A) \sqsubseteq_R A$ and $Int_R(B) \sqsubseteq_R B$, we have $Int_R(A) \sqcup_R Int_R(B)$ is a neutrosophic n -valued refined open set contained in $A \sqcup_R B$, which implies that $Int_R(A) \sqcup_R Int_R(B) \sqsubseteq_R Int_R(A \sqcup_R B)$.
- (7) Since $A_\alpha \sqsubseteq_R \bigsqcup_{\alpha \in \Delta} A_\alpha$ for every $\alpha \in \Delta$, $Int_R(A_\alpha) \sqsubseteq_R Int_R(\bigsqcup_{\alpha \in \Delta} A_\alpha)$ for every $\alpha \in \Delta$, that is $\bigsqcup_{\alpha \in \Delta} Int_R(A_\alpha) \sqsubseteq_R Int_R(\bigsqcup_{\alpha \in \Delta} A_\alpha)$.

The remaining 5 parts can be proved by duality. \square

Equality in parts (7) and (13) of Theorem 2.11 does not hold.

Example 2.12. Consider the neutrosophic 5-valued refined topological space (X, τ) defined in Example 2.9 and let $K = \{\langle a, 0, 1; 0; 1, 1 \rangle, \langle b, 1, 1; 0; 0, 1 \rangle\}$, and $L = \{\langle a, 1, 0; 1; 0, 0 \rangle, \langle b, 0, 0; 1; 1, 0 \rangle\}$. Then $K \sqcup_R L = \{\langle a, 1, 1; 0; 0, 0 \rangle, \langle b, 1, 1; 0; 0, 0 \rangle\} = 1_X$. So we have $Int_R(K \sqcup_R L) = 1_X$, and since K and L contains no neutrosophic n -valued refined open set except 0_X we have $Int_R(K) = Int_R(L) = 0_X$, which means $Int_R(K) \sqcup_R Int_R(L) = 0_X$, hence equality in parts (7) and (8) of Theorem 2.11 does not hold. For part (13) let $K = \{\langle a, 0.1, 0.4; 0.6; 0.5, 0.1 \rangle, \langle b, 0.7, 0.5; 0.9; 0.5, 0.3 \rangle\}$, $L = \{\langle a, 0.1, 0.3; 0.7; 0.3, 0.1 \rangle, \langle b, 0.7, 0.5; 0.9; 0.5, 0.3 \rangle\}$. Then $K \sqcap_R L = \{\langle a, 0.1, 0.3; 0.7; 0.5, 0.1 \rangle, \langle b, 0.7, 0.5; 0.9; 0.5, 0.3 \rangle\}$. The only neutrosophic 5-valued Refined closed sets containing K are: $1_X, A^c$ and C^c , so that we have $Cl_R(K) = 1_X \sqcap_R A^c \sqcap_R C^c = A^c$. Again the only neutrosophic 5-valued Refined closed sets containing L are: $1_X, A^c$ and C^c , so that we have $Cl_R(L) = 1_X \sqcap_R A^c \sqcap_R C^c = A^c$, and $Cl_R(K) \sqcap_R Cl_R(L) = A^c \sqcap_R A^c = A^c$, on the other hand the only neutrosophic 5-valued Refined closed sets containing $K \sqcap_R L$ are: $1_X, A^c, B^c$ and D^c , so that we have $Cl_R(K \sqcap_R L) = 1_X \sqcap_R A^c \sqcap_R B^c \sqcap_R D^c = D^c$. Note that D^c is a proper subset of A^c , so equality in Theorem 2.11 part (13) does not hold.

Question 2.13. *Is there a neutrosophic n -valued refined topological space (X, τ) shows that equality in part (14) of Theorem 2.11 does not hold.*

Definition 2.14 (*Neutrosophic n -valued refined pre-open and pre-closed sets*). Let $\tau \in TOP_{(n,r,s)}(X)$ and $A \in \mathcal{R}_{(n,r,s)}(X)$. Then A is said to be:

- (1) A *neutrosophic n -valued refined semi-open set*, if $A \sqsubseteq_R Cl_R(Int_R(A))$. The complement of a neutrosophic n -valued refined semi-open set is called a *neutrosophic n -valued refined semi-closed set*.
- (2) A *neutrosophic n -valued refined pre-open set*, if $A \sqsubseteq_R Int_R(Cl_R(A))$. The complement of a neutrosophic n -valued refined pre-open set is called a *neutrosophic n -valued refined pre-closed set*.
- (3) A *neutrosophic n -valued refined α -open set*, if $A \sqsubseteq_R Int_R(Cl_R(Int_R(A)))$. The complement of a neutrosophic n -valued refined α -open set is called a *neutrosophic n -valued refined α -closed set*.
- (4) A *neutrosophic n -valued refined β -open set*, if $A \sqsubseteq_R Cl_R(Int_R(Cl_R(A)))$. The complement of a neutrosophic n -valued refined β -open set is called a *neutrosophic n -valued refined β -closed set*.

Theorem 2.15. *Let $\tau \in TOP_{(n,r,s)}(X)$ and $A \in \mathcal{R}_{(n,r,s)}(X)$. Then:*

- (1) *Every Neutrosophic n -valued refined open (closed) set, is neutrosophic n -valued refined α -open (closed) set.*

- (2) Every Neutrosophic n -valued refined α -open (α -closed) set, is neutrosophic n -valued refined pre-open (pre-closed) set and neutrosophic n -valued refined semi-open (semi-closed) set.
- (3) Every Neutrosophic n -valued refined pre-open (pre-closed) or semi-open (semi-closed) set, is a neutrosophic n -valued refined β -open (β -closed) set.

Proof. (1) Let A be a Neutrosophic n -valued refined open set. Then, from Theorem 2.11 part (2) and (8), we have $Int_R(A) = A$ and $A \sqsubseteq_R Cl_R(A)$. So $Int_R(Cl_R(int_R(A))) \supseteq_R Int_R(Cl_R(A)) \supseteq_R Int_R(A) = A$. That is A is a neutrosophic n -valued refined α -open set. Now, suppose that A is a Neutrosophic n -valued refined closed set. Then A^c is a Neutrosophic n -valued refined open set, which implies A^c is a neutrosophic n -valued refined α -open set, and so A is a neutrosophic n -valued refined α -closed set.

(2) Obvious! we only use Theorem 2.11 part (1).

(3) Obvious! we only use Theorem 2.11 part (8) .

□

None of the above implications reverse. The following is an example of a neutrosophic 5-valued refined α -open set which is not open, and another example of a neutrosophic 5-valued refined pre-open (so it is β -open) set which is neither semi-open nor α -open.

Example 2.16. Consider $\tau = \{0_X, 1_X, A, B, C, D\}$ in Example 2.9 and let

$$H = \{\langle a, 0.5, 0.1; 0.3; 0.1, 0.3 \rangle, \langle b, 0.5, 0.3; 0.1; 0.7, 0.5 \rangle\}.$$

Then the neutrosophic 5-valued refined open sets contained in H are $0_X, A, B, C, D$; so we have $Int_R(H) = 0_X \sqcup_R A \sqcup_R B \sqcup_R C \sqcup_R D = D$, and since the only neutrosophic 5-valued refined close set containing D is 1_X , we have $Cl_R(Int_R(H)) = 1_X$, which implies $Int_R(Cl_R(int_R(H))) = 1_X$, hence $A \sqsubseteq_R Int_R(Cl_R(int_R(A)))$ and H is a neutrosophic 5-valued refined α -open set but not a neutrosophic 5-valued refined open set.

Consider, again, the set $K = \{\langle a, 0.1, 0.4; 0.6; 0.1, 0.3 \rangle, \langle b, 0.9, 0.2; 0.4; 0.1, 0.5 \rangle\}$. Since $\mu_K^1(a) < \mu_O^1(a)$ for every $O \in \tau - \{0_X\}$, we have the only Neutrosophic 5-valued refined open set contained in K is 0_X and $Int_R(K) = 0_X$, which implies $Cl_R(Int_R(K)) = 0_X$ and $Int_R(Cl_R(Int_R(K))) = 0_X$, so K is not a neutrosophic 5-valued refined semi-open nor α -open set; on the other hand, $\mu_K^1(b) > \mu_D^1(b)$ for every neutrosophic 5-valued refined closed set D in τ except for 1_X , that means $Cl_R(K) = 1_X$ and $int_R(Cl_R(A)) = 1_X$, hence $K \sqsubseteq_R Int_R(Cl_R(A))$ and K is a neutrosophic 5-valued refined pre-open set but not α -open. Since every neutrosophic 5-valued refined pre-open set is a neutrosophic 5-valued refined β -open set, K is, also, and example of a neutrosophic 5-valued refined β -open set which is not neutrosophic 5-valued refined semi-open.

Here we give an example of a a neutrosophic 5-valued refined *semi-open* (so it is β -open) set which is neither *pre-open* nor α -open.

Example 2.17. Let $X = \{a\}$, and let $\tau = \{0_X, 1_X, A, B\} \subset \mathcal{R}_{(5,2,1)}(X)$ where $A = \{\langle a, 0.2, 0.1; 0.7; 0.3, 0.4 \rangle\}$, $B = \{\langle a, 0.3, 0.2; 0.5; 0.2, 0.3 \rangle\}$. Since $A \sqcap_R B = A$ and $A \sqcup_R B = B$, τ is a neutrosophic 5-valued refined topology on X . The 5-valued refined closed sets in (X, τ) are: $0_X, 1_X, A^c, B^c$ where $A^c = \{\langle a, 0.3, 0.4; 0.3; 0.2, 0.1 \rangle\}$ and $B^c = \{\langle a, 0.2, 0.3; 0.5; 0.3, 0.2 \rangle\}$. Consider the neutrosophic 5-valued refined set $L = \{\langle a, 0.2, 0.2; 0.5; 0.3, 0.3 \rangle\}$. Then the only neutrosophic 5-valued refined open sets contained in K are $0_X, A$, so that $Int_R(L) = 0_X \sqcup_R A = A$. To find $Cl_R(Int_R(L))$ we note that the neutrosophic 5-valued refined closed sets containing $Int_R(L)$ are $1_X, A^c, B^c$, so $Cl_R(Int_R(L)) = 1_X \sqcap_R A^c \sqcap_R B^c = B^c$, and since $L \sqsubseteq_R B^c$, L is a neutrosophic 5-valued refined semi-open sets. Now, we will show that L is not α -open. First note that the neutrosophic 5-valued refined open sets contained in $Cl_R(Int_R(K)) = B^c$ are 0_X and A , so we have $Int_R(Cl_R(Int_R(L))) = A$, and since L is not contained in A , L is not a neutrosophic α -open set.

We will show L is not a neutrosophic 5-valued refined *pre-open* set. The only neutrosophic 5-valued refined closed sets containing L are $1_X, A^c$ and B^c , so $Cl_R(L) = 1_X \sqcap_R A^c \sqcap_R B^c = B^c$, and since the neutrosophic 5-valued refined open sets contained in B^c are 0_X and A , we have $Int_R(Cl_R(L)) = A$ which not containing L , that is L is not a neutrosophic 5-valued refined pre-open set. So L is, also, an example of a neutrosophic 5-valued refined *semi-open* set which is not pre-open. And since every neutrosophic 5-valued refined *semi-open* set is β -open set, K is an example of a neutrosophic 5-valued refined β -open set which is not *pre-open*.

Finally we will give an example of a a neutrosophic 5-valued refined β -open set which is neither *pre-open* nor *semi-open*.

Example 2.18. Let (X, τ) as in Example 2.17 and consider the neutrosophic 5-valued refined set $M = \{\langle a, 0.2, 0.1; 0.9; 0.3, 0.5 \rangle\}$. Then the only neutrosophic 5-valued refined open sets in τ contained in K is 0_X , so $Int_R(M) = 0_X$, which implies $Cl_R(Int_R(M)) = 0_X$, and since M is not contained in 0_X , we have M is not neutrosophic 5-valued refined semi-open set; on the other hand the neutrosophic 5-valued refined closed sets containing M are $1_X, A^c$ and B^c , so that $Cl_R(M) = B^c$, and since the only neutrosophic 5-valued refined open sets contained in B^c are 0_X and A we have $Int_R(Cl_R(M)) = A$. Since $Int_R(Cl_R(M)) = A$ and A does not contain M , we have M is not a neutrosophic 5-valued refined *pre-open* set. Now, to find $Cl_R(Int_R(Cl_R(M)))$ we note that the only neutrosophic 5-valued refined closed sets in τ




FIGURE 1. Relations between different types of generalized neutrosophic n-valued refined open sets.

containing A are $1_X, A^c$ and B^c , so $Cl_R(Int_R(Cl_R(M))) = B^c$ which contains M , so M is a neutrosophic 5-valued refined β -open set but not *semi*-open nor *pre*-open.

The following diagram shows the relations between different types of generalized neutrosophic n -valued refined sets:

Theorem 2.19. *Let $\tau \in TOP_{(n,r,s)}(X)$ and $K \in \mathcal{R}_{(n,r,s)}(X)$. Then*

- (1) *If there is a neutrosophic n -valued refined open set U such that $K \sqsubseteq_R U \sqsubseteq_R Cl_R(K)$, then K is a neutrosophic n -valued refined pre-open set.*
- (2) *If there is a neutrosophic n -valued refined open set U such that $U \sqsubseteq_R K \sqsubseteq_R Cl_R(U)$, then K is a neutrosophic n -valued refined semi-open set.*

Proof. (1) $K \sqsubseteq_R U \sqsubseteq_R Int_R(Cl_R(U)) \sqsubseteq_R Int_R(Cl_R(Cl_R(K))) = Int_R(Cl_R(K))$.

(2) Since $Cl_R(Int_R(U)) = Cl_R(U)$ we have

$$Cl_R(Int_R(K)) \supseteq_R Cl_R(Int_R(U)) = Cl_R(U) \supseteq_R K.$$

□

Theorem 2.20. *Let $\tau \in TOP_{(n,r,s)}(X)$ and $K \in \mathcal{R}_{(n,r,s)}(X)$. Then the union of any collection of neutrosophic n -valued refined α -open, β -open, pre-open or semi-open sets is a neutrosophic n -valued refined α -open, β -open, pre-open or semi-open set respectively.*

Proof. We will prove it for neutrosophic n -valued refined β -open sets, and the remaining parts can be proved in the same manner. Let A_γ be a neutrosophic n -valued refined β -open set for every $\gamma \in \Delta$. Then $A_\gamma \sqsubseteq_R Cl_R(int_R(Cl_R(A_\gamma)))$ for every $\gamma \in \Delta$. Then from parts (7) and (14) of Theorem 2.11 we have:

$$\begin{aligned} Cl_R(int_R(Cl_R(\sqcup_{\gamma \in \Delta} A_\gamma))) &\sqsupseteq_R Cl_R(int_R(\sqcup_{\gamma \in \Delta} Cl_R(A_\gamma))) \sqsupseteq_R Cl_R(\sqcup_{\gamma \in \Delta} int_R(Cl_R(A_\gamma))) \sqsupseteq_R \\ \sqcup_{\gamma \in \Delta} Cl_R(int_R(Cl_R(A_\gamma))) &\sqsupseteq_R \sqcup_{\gamma \in \Delta} A_\gamma \quad \square \end{aligned}$$

Funding: This study is supported via funding from Prince sattam bin Abdulaziz University project number (PSAU/2023/R/1444).

Conflicts of Interest: The author declare that he has no conflict of interest.

REFERENCES

1. M. Abdel-Basset , A. Gamal, L. H. Son and F. Smarandache , *ABipolar Neutrosophic multi Criteria Decision Making Framework for Professional Selection*, Applied Sciences, 10(4),1202.
2. M. Abdel-Basset , R. Mohamed , A. E. N. H. Zaied , A. Gamal and F. Smarandache , *Solving the supply chain problem using the best-worst method based on a novel Plithogenic Model*, In Optimization Theory Based on Neutrosophic and Plithogenic Sets, Academic Press, 2020, 1-19.
3. M. Abdel-Basset and R. Mohamed , *A Novel Plithogenic TOPSIS-CRITIC Model for Sustainable Supply Chain Risk Management*, Journa of Cleaner Production, 2020,247, 119586.
4. M. Abdel-Basset , R. Mohamed , A. E. N. H. Zaied and F. Smarandache , *A Hybrid Plithogenic Decision-Making Approach with quality function deployment for selecting Supply Chain Sustainability metrics*, Symmetry, 2019, 11(7), 903.
5. Abdel-Basset, M., Ding, W., Mohamed, R. et al., *An integrated plithogenic MCDM approach for financial performance evaluation of manufacturing industries*, Risk Management,(22)2020, 192–218.
6. Abdel-Basset M., Mohamed R., Sallam K. and Elhoseny M., *A novel decision-making model for sustainable supply chain finance under uncertainty environment*, Journal of Cleaner Production,(269) 2020, 122324.
7. Abdel-Basst M., Mohamed R. and Elhoseny M., *A novel framework to evaluate innovation value proposition for smart product-service systems*, Environmental Technology and Innovation, (20) 2020, 101036.
8. M. ARAR , *A Note on Spaces with a Countable μ -Base*, Acta Math. Hungar., 144 (2) (2014), 494–498.
9. M. Arar and Saeid Jafari, *Neutrosophic μ -Topological Sapces*, Neutrosophic Sets and Systems, 38(2020), 51-66.
10. M. Arar, *About Neutrosophic Countably Compactness*, Neutrosophic Sets and Systems, 36(2020), 246-255.
11. M. M. ARAR , *On Countably μ -Paracompact Spaces*, Acta Math. Hungar., 149 (1) (2016), 50–57.
12. M. ARAR , *Strongly Generalized Neighborhood Systems*, Missouri Journal of Mathematical Sciences, 29(1). 2017, 43-49.
13. K. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, 20 (1986),87–96.

14. S. Broumi and F. Smarandache, *Extended Hausdorff Distance and Similarity Measures for Neutrosophic Refined Sets and Their Applications in Medical Diagnosis*, Journal of New Theory, 7 (2015), 64-78.
15. Said Broumi and Florentin Smarandache, *Extended Hausdorff Distance and Similarity Measures for Neutrosophic Refined Sets And Their Application in Medical Diagnosis*, Journal of New Theory, 7 (2015), 64–78.
16. Chinnadurai V and Sindhu M P, *A Novel Approach for Pairwise Separation Axioms on Bi-Soft Topology Using Neutrosophic Sets and An Output Validation in Real Life Application*, Neutrosophic Sets and Systems, 35 (2020), 435–463.
17. Masoud Ghods and Zahra Rostami, *Introduction to Topological Indices in Neutrosophic Graphs*, Neutrosophic Sets and Systems, 35 (2020), 68-77.
18. Madad Khan, Muhammad Zeeshan, Saima Anis, Abdul Sami Awan and Florentin Smarandache, *Neutrosophic Soft Fixed Points*, Neutrosophic Sets and Systems, 35 (2020), pp.531-546.
19. F. G. Lupiáñez, *On neutrosophic topology*, The International Journal of Systems and Cybernetics, 37(6) (2008), 797–800.
20. F. G. Lupiáñez, *Interval neutrosophic sets and topology*, The International
21. Mary Margaret A, Trinita Pricilla M and Shawkat Alkhazaleh, *Neutrosophic Vague Topological Spaces*, Neutrosophic Sets and Systems, 28(2019) pp.179-190.
22. Muhammad Riaz, Florentin Smarandache, Faruk Karaaslan, Masooma Raza Hashmi and Iqra Nawaz, *Neutrosophic Soft Rough Topology and its Applications to Multi-Criteria Decision-Making*, Neutrosophic Sets and Systems, 35 (2020) pp.198-219.
23. Rakhal Das and Binod Chandra Tripathy, *Neutrosophic Multiset Topological Space*, Neutrosophic Sets and Systems, vol. 35 (2020), pp.142-152.
24. A. Salama and S. AL-Blowi, *Neutrosophic Set and Neutrosophic Topological Spaces*, IOSR Journal of Mathematics, 3(4) (2012), 31–35.
25. A. E. Samuel and R. Narmadhagnanam, *Neutrosophic Refined Sets in Medical Diagnosis*, International Journal of Fuzz Mathematical Archive, 14(1), 2017, 117-123.
26. F. Smarandache, *Neutrosophic set - a generalization of the intuitionistic fuzzy set*, International Journal of Pure and Applied Mathematics, 24(3) (2005) 287–297.
27. F. Smarandache, *Neutrosophy and neutrosophic logic*, first international conference on neutrosophy, neutrosophic logic, set, probability, and statistics, University of New Mexico, Gallup, NM 87301, USA(2002).
28. F. Smarandache, *n-Valued Refined Neutrosophic Longic and Its Applications to Physics*, arXiv:1407.1041v1, 3 JULy, (2014).
29. S. Karatas and C. Kuru, *Neutrosophic Topology*, Neutrosophic Sets and Systems, University of New Mexico, 13 (2016), 90-95.
30. Shawkat Alkhazaleh, *Neutrosophic vague set theory*, Cretical Review, 10 (2015), 29-39.
31. Suman Das and Surapati Pramanik, *Generalized neutrosophic b-open sets in neutrosophic topological space*, Neutrosophic Sets and Systems, 35 (2020) 522-530.

Received: August 10, 2022. Accepted: January 06, 2023