

5-1-2023

New approach to bisemiring theory via the bipolar valued neutrosophic normal sets

M. Palanikumar

G. Selvi

Ganeshsree Selvachandran

Sher Lyn Tan

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Palanikumar, M.; G. Selvi; Ganeshsree Selvachandran; and Sher Lyn Tan. "New approach to bisemiring theory via the bipolar valued neutrosophic normal sets." *Neutrosophic Sets and Systems* 55, 1 (2023). https://digitalrepository.unm.edu/nss_journal/vol55/iss1/26

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



New approach to bisemiring theory via the bipolar valued neutrosophic normal sets

M.Palanikumar¹, G.Selvi², Ganeshsree Selvachandran³ and Sher Lyn Tan⁴

¹ Department of Mathematics, Saveetha School of Engineering, SIMATS, India; palanimaths86@gmail.com

² Department of Mathematics, Saveetha School of Engineering, SIMATS, India; mslalima11@gmail.com

³ Faculty of Business and Management, UCSI University, Jalan Menara Gading, 56000 Cheras, Kuala Lumpur, Malaysia; Ganeshsree@ucsiuniversity.edu.my; ganeshsree86@yahoo.com

⁴ Institute of Actuarial Science and Data Analytics, UCSI University, Jalan Menara Gading, 56000 Cheras, Kuala Lumpur, Malaysia; SherlynTan@ucsiuniversity.edu.my

Abstract. In this paper, we introduce the notion of bipolar-valued neutrosophic subbisemiring (BVNSBS), level sets of BVNSBS, and bipolar valued neutrosophic normal subbisemiring (BVNNSBS) of a bisemiring. The concept of BVNSBS is a new generalization of subbisemiring over bisemirings. We discussed the theory of (ξ, τ) -BVNSBS and (ξ, τ) -BVNNSBS over bisemirings and presented several illustrative examples to demonstrate the sufficiency and validity of the proposed theorems, lemmas, and propositions.

Keywords: Fuzzy set; Bipolar valued neutrosophic subbisemiring; Bipolar valued neutrosophic bisemiring; Homomorphism; Normal.

1. Introduction

Classical mathematics may not always be the solution for practical situations in economics, medical sciences, engineering, social sciences, and environmental sciences, which involves various uncertainties, imprecise and incomplete information. The limitation of classical mathematics that is unable to deal with uncertainties and fuzziness motivated the introduction of mathematical theory such as probability theory, fuzzy set theory [1], rough set theory [2], vague set theory [3], interval mathematics [4], and soft set theory [5]. However, these theories were insufficient and have limitations in dealing with uncertainties. Probability theory can only deal with stochastically stable problems, which may not apply to many problems in the field of economic, environmental, and social sciences. Interval mathematics takes calculation

M.Palanikumar, G.Selvi, Ganeshsree Selvachandran and Sher Lyn Tan, New approach to bisemiring theory via the bipolar valued neutrosophic normal sets

errors into account by constructing an interval estimate for the solution that is useful in many areas, but it is not appropriately adaptable for problems that arise from unreliable, inadequate, and change of information. On the other hand, the fuzzy set theory introduced by Zadeh [1] is most appropriate for dealing with uncertainties and vagueness. Membership of an element in a fuzzy set is a single value between the interval, but in real-life problems, the degree of non-membership may not always be equal to 1 minus the degree of membership as there may be some degree of hesitation. Works on fuzzy set theory are progressing rapidly and have resulted in the conception of many hybrid fuzzy models. In 1983, Atanasov [6] proposed intuitionistic fuzzy sets as a generalization of the notion of fuzzy set, which incorporated the degree of hesitation. Later, Zhang [7] introduced bipolar fuzzy sets in which the membership function is mapped to intervals, thereby allowing it to deal with complex problems in both positive and negative aspects. Later, Zhang [8] proposed that bipolar fuzzy logic should combine both fuzziness and polarity by introducing the (Yin) (Yang) bipolar fuzzy sets. Lee [9] introduced the operation in bipolar-valued fuzzy sets, whereas Lee [10] discovered that bipolar-valued fuzzy sets can represent the degree of satisfaction to counter property but fail to express uncertainties in assigning membership degree. These concepts have been widely applied to handle incomplete information arising from practical situations. However, these were still unable to address uncertainties such as indeterminate and inconsistent information.

In 1999, Smarandache [11] proposed the neutrosophic theory that deals with "the origin, nature, and scope of neutralities, as well as their interactions with different ideational spectra". The idea of neutrosophic logic is a logic that states that each proposition is estimated to have a degree of trust, degree of indeterminacy, and degree of falsity. Smarandache [12] further generalized the theory of intuitionistic fuzzy sets to the neutrosophic model, and introduced the truth, indeterminacy, and falsity components that represent the membership, indeterminacy, and non-membership values of a neutrosophic set, respectively. In contrast to intuitionistic fuzzy sets, neutrosophic sets used indeterminacy as a completely independent measure of the membership and non-membership information, and thus it can effectively describe uncertain and inconsistent information and overcome the limitation of the existing approaches in handling uncertain information.

The original neutrosophic theory was introduced from a philosophical standpoint. Hence, it may be difficult to be applied in practical problems. Subsequently, Wang et al. [13] generalized the neutrosophic set from a technical point of view and specified the set-theoretic operators on an instance of a neutrosophic set, called the single-valued neutrosophic set, which takes values from the subset of $[0, 1]$, thereby enabling it to be used feasibly for real-world problems. Over the years, subsequent developments and extensions of the neutrosophic set were proposed. Deli et al. [14] proposed bipolar neutrosophic sets as an extension of bipolar fuzzy sets [7]. Ye [15]

introduced the concept of simplified neutrosophic sets. Peng et al. [16] introduced multi-valued neutrosophic sets that allow the truth, indeterminacy, and falsity membership degrees to have a set of crisp values between zero and one, respectively. Das et al. [17] introduced the notion of neutrosophic fuzzy sets by combining fuzzy sets with neutrosophic fuzzy sets to overcome the difficulties in handling the non-standard interval of neutrosophic components.

On the other hand, the fuzzy set theory had been applied and contributed to the generalization of many fundamental concepts in algebra. Extensive research has been done on the fuzzy algebraic structure of semirings introduced by Vandiver [18], which is a generalization of a ring by relaxing the conditions on the additive structure requiring just a monoid rather than a group and have been proven useful for dealing with problems in various areas. The application of semirings had been studied extensively by Golan [19] and Glazek [20].

Ahsan, Saifullah and Farid Khan [21] initiated the study of fuzzy semirings, while Feng, Jun and Zhao [22], and Yousafzai et al. [23] studied semigroups and semirings using fuzzy set and soft sets, respectively. Furthermore, Mockor [24] introduced the notion of a semiring-valued fuzzy set for special commutative partially pre-ordered semiring and introduced F-transform and inverse F-transform for these fuzzy-type structures. Other than that, palanikumar et al. [25–30] studied the algebraic structure of various semirings that constitute a natural generalization of semirings.

Recently, many studies applied bipolar fuzzy information in various algebraic structures, for instance, semigroups [31–33] and BCK/BCI-algebras [34–37]. Zararsz et al. [38] discussed the notion of bipolar fuzzy metric spaces with application. Selvachandran and Salleh [39] introduced vague soft hyperrings and vague soft hyperideals. Jun, Kim and Lee [40] introduced bipolar fuzzy translation in BCK/BCI-algebra and investigated its properties, whereas Jun and Park [41] introduced bipolar fuzzy regularity, bipolar fuzzy regular subalgebra, bipolar fuzzy filter, and bipolar fuzzy closed quasi filter in BCH-algebras. Apart from that, Sen, Ghosh and Ghosh [42] extended the study of semirings and proposed the concept of bisemiring in 2004. Later, Hussain [43] defined the congruence relation between bisemiring and bisemiring homomorphisms, followed by the factor bisemiring. Hussain et al. [44] further generalized bisemiring to a new algebraic structure called n -semiring and congruence relations on homomorphisms and n -semirings.

To the best of our knowledge, studies on bisemiring theory using bipolar valued neutrosophic sets have not been studied extensively, and further generalization for bisemiring is still needed for various practical problems. In this paper, we introduce the notion of bipolar valued neutrosophic subbisemiring (BVNSBS), level sets of BVNSBS, and bipolar valued neutrosophic normal subbisemiring (BVNNSBS) of a bisemiring. The concept of BVNSBS is a new generalization of subbisemiring over bisemirings. We discussed the theory for (ξ, τ) -BVNSBS

and (ξ, τ) -BVNNSBS over bisemiring theory and presented several illustrative examples. The rest of the paper is organized as follows: Section 2 outlines the preliminary definitions and results, Section 3 introduces the notion of BVNSBS, Section 4 discusses the (ξ, τ) -BVNSBS and Section 5 discusses the (ξ, τ) -BVNNSBS.

2. Preliminaries

Definition 2.1. [9] Let U be the universe set. A bipolar valued fuzzy set ϑ in U is an object having the form $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$, where $\vartheta^- : U \rightarrow [-1, 0]$ and $\vartheta^+ : U \rightarrow [0, 1]$ are mappings. The positive membership degree $\vartheta^+(u)$ denoted the satisfaction degree of an element u to the property corresponding to a bipolar valued fuzzy set $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$, and the negative membership degree $\vartheta^-(u)$ denotes the satisfaction degree of u to some implicit counter-property of $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$. If $\vartheta^+(u) \neq 0$ and $\vartheta^-(u) = 0$, it is the situation that u is regarded as having only positive satisfaction for $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$. If $\vartheta^+(u) = 0$ and $\vartheta^-(u) \neq 0$, it is the situation that u does not satisfy the property of $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$ but somewhat satisfies the counter property of $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$. It is possible for an element u to be $\vartheta^+(u) \neq 0$ and $\vartheta^-(u) \neq 0$ when the membership function of the property overlaps that of its counter-property over some portion of the domain. For the sake of simplicity, we shall use the symbol $\vartheta = \langle U; \vartheta^-, \vartheta^+ \rangle$ for the bipolar valued fuzzy set $\vartheta = \{(u, \vartheta^+(u), \vartheta^-(u)) | u \in U\}$, and use the notion of bipolar fuzzy sets instead of the notion of bipolar valued fuzzy sets.

Definition 2.2. [11] A neutrosophic set K in a universe set U is an object having the structure $K = \{(m, \vartheta_K^T(m), \vartheta_K^I(m), \vartheta_K^F(m)) | m \in U\}$, where $\vartheta_K^T(m), \vartheta_K^I(m), \vartheta_K^F(m) : U \rightarrow [0, 1]$ represents the truth-membership function, the indeterminacy membership function and the falsity-membership function respectively. There is no restriction on the sum of $\vartheta_K^T, \vartheta_K^I, \vartheta_K^F$ and so $0 \leq \vartheta_K^T + \vartheta_K^I + \vartheta_K^F \leq 3$.

Definition 2.3. [11] Let $K = \{(m, \vartheta_K^T(m), \vartheta_K^I(m), \vartheta_K^F(m)) | m \in U\}$ and $L = \{(m, \vartheta_L^T(m), \vartheta_L^I(m), \vartheta_L^F(m)) | m \in U\}$ be any two neutrosophic sets of a set U . Then

$$K \cap L = \left\{ \langle m, \min\{\vartheta_K^T(m), \vartheta_L^T(m)\}, \min\{\vartheta_K^I(m), \vartheta_L^I(m)\}, \max\{\vartheta_K^F(m), \vartheta_L^F(m)\} \rangle \mid m \in U \right\},$$

$$K \cup L = \left\{ \langle m, \max\{\vartheta_K^T(m), \vartheta_L^T(m)\}, \max\{\vartheta_K^I(m), \vartheta_L^I(m)\}, \min\{\vartheta_K^F(m), \vartheta_L^F(m)\} \rangle \mid m \in U \right\}.$$

Definition 2.4. [11] For any neutrosophic set $K = \{(m, \vartheta_K^T(m), \vartheta_K^I(m), \vartheta_K^F(m)) | m \in U\}$ of a set U , we defined a (ξ, τ) -cut of as the crisp subset $\{\vartheta_K^T(m) \geq \xi, \vartheta_K^I(m) \geq \xi, \vartheta_K^F(m) \leq \tau | m \in U\}$ of U .

Definition 2.5. [11] Let K and L be any two neutrosophic set of U . Then

$$K \times L = \{\vartheta_{K \times L}^T(m, n), \vartheta_{K \times L}^I(m, n), \vartheta_{K \times L}^F(m, n) | \forall m, n \in U\},$$

where $\vartheta_{K \times L}^T(m, n) = \min\{\vartheta_K^T(m), \vartheta_L^T(n)\}$, $\vartheta_{K \times L}^I(m, n) = \frac{\vartheta_K^I(m) + \vartheta_L^I(n)}{2}$, $\vartheta_{K \times L}^F(m, n) = \max\{\vartheta_K^F(m), \vartheta_L^F(n)\}$.

M.Palanikumar, K.Arulmozhi, Ganeshsree Selvachandran and Sher Lyn Tan, New approach to bisemiring theory via the bipolar valued neutrosophic normal sets

Definition 2.6. [44] A fuzzy subset K of a bisemiring $(S, \uplus_1, \uplus_2, \uplus_3)$ is said to be a fuzzy subbisemiring of S if $\vartheta_K(m \uplus_1 n) \geq \min\{\vartheta_K(m), \vartheta_K(n)\}, \vartheta_K(m \uplus_2 n) \geq \min\{\vartheta_K(m), \vartheta_K(n)\}, \vartheta_K(m \uplus_3 n) \geq \min\{\vartheta_K(m), \vartheta_K(n)\}$, for all $m, n \in S$.

Definition 2.7. [44] Let $(S_1, +, \cdot, \times)$ and $(S_2, \boxplus, \circ, \otimes)$ be any two bisemirings. A function $\phi : S_1 \rightarrow S_2$ is said to be a homomorphism if $\phi(m + n) = \phi(m) \boxplus \phi(n), \phi(m \cdot n) = \phi(m) \circ \phi(n), \phi(m \times n) = \phi(m) \otimes \phi(n)$, for all $m, n \in S_1$.

3. Bipolar Valued Neutrosophic Subbisemiring (BVNSBS)

In what follows, let \mathcal{S} denote a bisemiring unless otherwise noted. In this section, we communicate the concept of bipolar valued neutrosophic subbisemiring, strongest neutrosophic relation on \mathcal{S} . Furthermore, we introduce the arbitrary intersection bipolar valued neutrosophic subbisemiring and list some properties.

Definition 3.1. A bipolar valued neutrosophic subset K of \mathcal{S} is said to be BVNSBS of \mathcal{S} if it satisfies the following conditions:

$$\left\{ \begin{array}{l} \left(\vartheta_K^{T+}(m \uplus_1 n) \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\}, \right. \\ \left. \vartheta_K^{T-}(m \uplus_1 n) \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} \right) \\ \left(\vartheta_K^{T+}(m \uplus_2 n) \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\}, \right. \\ \left. \vartheta_K^{T-}(m \uplus_2 n) \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} \right) \\ \left(\vartheta_K^{T+}(m \uplus_3 n) \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\}, \right. \\ \left. \vartheta_K^{T-}(m \uplus_3 n) \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} \right) \end{array} \right\} \left\{ \begin{array}{l} \left(\vartheta_K^{I+}(m \uplus_1 n) \geq \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \right. \\ \left. \vartheta_K^{I-}(m \uplus_1 n) \leq \frac{\vartheta_K^{I-}(m) - \vartheta_K^{I-}(n)}{2} \right) \\ \text{OR} \\ \left(\vartheta_K^{I+}(m \uplus_2 n) \geq \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \right. \\ \left. \vartheta_K^{I-}(m \uplus_2 n) \leq \frac{\vartheta_K^{I-}(m) - \vartheta_K^{I-}(n)}{2} \right) \\ \text{OR} \\ \left(\vartheta_K^{I+}(m \uplus_3 n) \geq \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \right. \\ \left. \vartheta_K^{I-}(m \uplus_3 n) \leq \frac{\vartheta_K^{I-}(m) - \vartheta_K^{I-}(n)}{2} \right) \end{array} \right\}$$

$$\left\{ \begin{array}{l} \left(\vartheta_K^{F+}(m \uplus_1 n) \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}, \right. \\ \left. \vartheta_K^{F-}(m \uplus_1 n) \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\} \right) \\ \left(\vartheta_K^{F+}(m \uplus_2 n) \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}, \right. \\ \left. \vartheta_K^{F-}(m \uplus_2 n) \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\} \right) \\ \left(\vartheta_K^{F+}(m \uplus_3 n) \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}, \right. \\ \left. \vartheta_K^{F-}(m \uplus_3 n) \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\} \right) \end{array} \right\}$$

for all $m, n \in \mathcal{S}$.

Example 3.2. Let $\mathcal{S} = \{l_1, l_2, l_3, l_4\}$ be the bisemiring with the following Cayley table:

\uplus_1	l_1	l_2	l_3	l_4	\uplus_2	l_1	l_2	l_3	l_4	\uplus_3	l_1	l_2	l_3	l_4
l_1	l_1	l_1	l_1	l_1	l_1	l_1	l_2	l_3	l_4	l_1	l_1	l_1	l_1	l_1
l_2	l_1	l_2	l_1	l_2	l_2	l_2	l_4	l_4		l_2	l_1	l_2	l_3	l_4
l_3	l_1	l_1	l_3	l_3	l_3	l_4	l_3	l_4		l_3	l_4	l_4	l_4	l_4
l_4	l_1	l_2	l_3	l_4	l_4	l_4	l_4	l_4		l_4	l_4	l_4	l_4	l_4

$(\vartheta_K^+(l), \vartheta_K^-(l))$	$l = l_1$	$l = l_2$	$l = l_3$	$l = l_4$
$(\vartheta_K^{T+}(l), \vartheta_K^{T-}(l))$	(0.55, -0.7)	(0.35, -0.6)	(0.15, -0.3)	(0.25, -0.4)
$(\vartheta_K^{I+}(l), \vartheta_K^{I-}(l))$	(0.65, -0.8)	(0.5, -0.5)	(0.3, -0.1)	(0.4, -0.2)
$(\vartheta_K^{F+}(l), \vartheta_K^{F-}(l))$	(0.25, -0.15)	(0.35, -0.25)	(0.65, -0.65)	(0.55, -0.45)

Clearly, K is an BVNSBS of \mathcal{S} .

Theorem 3.3. *The intersection of a family of BVNSBS^s of \mathcal{S} is a BVNSBS of \mathcal{S} .*

Proof. Let $\{O_i | i \in I\}$ be a family of BVNSBS^s of \mathcal{S} and $K = \bigcap_{i \in I} O_i$.

Let m and n in \mathcal{S} . Now,

$$\begin{aligned}
 \vartheta_K^{T+}(m \uplus_1 n) &= \inf_{i \in I} \vartheta_{O_i}^{T+}(m \uplus_1 n) \\
 &\geq \inf_{i \in I} \min\{\vartheta_{O_i}^{T+}(m), \vartheta_{O_i}^{T+}(n)\} \\
 &= \min\left\{\inf_{i \in I} \vartheta_{O_i}^{T+}(m), \inf_{i \in I} \vartheta_{O_i}^{T+}(n)\right\} \\
 &= \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\} \\
 \vartheta_K^{T-}(m \uplus_1 n) &= \sup_{i \in I} \vartheta_{O_i}^{T-}(m \uplus_1 n) \\
 &\leq \sup_{i \in I} \max\{\vartheta_{O_i}^{T-}(m), \vartheta_{O_i}^{T-}(n)\} \\
 &= \max\left\{\sup_{i \in I} \vartheta_{O_i}^{T-}(m), \sup_{i \in I} \vartheta_{O_i}^{T-}(n)\right\} \\
 &= \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\}.
 \end{aligned}$$

Now,

$$\begin{aligned}
 \vartheta_K^{I+}(m \uplus_1 n) &= \inf_{i \in I} \vartheta_{O_i}^{I+}(m \uplus_1 n) \\
 &\geq \inf_{i \in I} \frac{\vartheta_{O_i}^{I+}(m) + \vartheta_{O_i}^{I+}(n)}{2} \\
 &= \frac{\inf_{i \in I} \vartheta_{O_i}^{I+}(m) + \inf_{i \in I} \vartheta_{O_i}^{I+}(n)}{2} \\
 &= \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}
 \end{aligned}$$

$$\begin{aligned}
\vartheta_K^{I-}(m \uplus_1 n) &= \sup_{i \in I} \vartheta_{O_i}^{I-}(m \uplus_1 n) \\
&\leq \sup_{i \in I} \frac{\vartheta_{O_i}^{I-}(m) + \vartheta_{O_i}^{I-}(n)}{2} \\
&= \frac{\sup_{i \in I} \vartheta_{O_i}^{I-}(m) + \sup_{i \in I} \vartheta_{O_i}^{I-}(n)}{2} \\
&= \frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}.
\end{aligned}$$

Now,

$$\begin{aligned}
\vartheta_K^{F+}(m \uplus_1 n) &= \sup_{i \in I} \vartheta_{O_i}^{F+}(m \uplus_1 n) \\
&\leq \sup_{i \in I} \max\{\vartheta_{O_i}^{F+}(m), \vartheta_{O_i}^{F+}(n)\} \\
&= \max\left\{\sup_{i \in I} \vartheta_{O_i}^{F+}(m), \sup_{i \in I} \vartheta_{O_i}^{F+}(n)\right\} \\
&= \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}
\end{aligned}$$

$$\begin{aligned}
\vartheta_K^{F-}(m \uplus_1 n) &= \inf_{i \in I} \vartheta_{O_i}^{F-}(m \uplus_1 n) \\
&\geq \inf_{i \in I} \min\{\vartheta_{O_i}^{F-}(m), \vartheta_{O_i}^{F-}(n)\} \\
&= \min\left\{\inf_{i \in I} \vartheta_{O_i}^{F-}(m), \inf_{i \in I} \vartheta_{O_i}^{F-}(n)\right\} \\
&= \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\}.
\end{aligned}$$

Similarly, we can prove that other two operations. Hence K is an BVNSBS of \mathcal{S} .

Theorem 3.4. *If K and L are any two BVNSBS^s of \mathcal{S}_1 and \mathcal{S}_2 respectively, then $K \times L$ is a BVNSBS of $\mathcal{S}_1 \times \mathcal{S}_2$.*

Proof. Let K and L be two BVNSBS^s of \mathcal{S}_1 and \mathcal{S}_2 respectively. Let $m_1, m_2 \in \mathcal{S}_1$ and $n_1, n_2 \in \mathcal{S}_2$. Then (m_1, n_1) and (m_2, n_2) are in $\mathcal{S}_1 \times \mathcal{S}_2$. Now,

$$\begin{aligned}
\vartheta_{K \times L}^{T+}[(m_1, n_1) \uplus_1 (m_2, n_2)] &= \vartheta_{K \times L}^{T+}(m_1 \uplus_1 m_2, n_1 \uplus_1 n_2) \\
&= \min\{\vartheta_K^{T+}(m_1 \uplus_1 m_2), \vartheta_L^{T+}(n_1 \uplus_1 n_2)\} \\
&\geq \min\{\min\{\vartheta_K^{T+}(m_1), \vartheta_K^{T+}(m_2)\}, \min\{\vartheta_L^{T+}(n_1), \vartheta_L^{T+}(n_2)\}\} \\
&= \min\{\min\{\vartheta_K^{T+}(m_1), \vartheta_L^{T+}(n_1)\}, \min\{\vartheta_K^{T+}(m_2), \vartheta_L^{T+}(n_2)\}\} \\
&= \min\{\vartheta_{K \times L}^{T+}(m_1, n_1), \vartheta_{K \times L}^{T+}(m_2, n_2)\}.
\end{aligned}$$

Similarly, $\vartheta_{K \times L}^{T-}[(m_1, n_1) \uplus_1 (m_2, n_2)] \leq \max\{\vartheta_{K \times L}^{T-}(m_1, n_1), \vartheta_{K \times L}^{T-}(m_2, n_2)\}$.

Now,

$$\begin{aligned} \vartheta_{K \times L}^{I+}[(m_1, n_1) \uplus_1 (m_2, n_2)] &= \vartheta_{K \times L}^{I+}(m_1 \uplus_1 m_2, n_1 \uplus_1 n_2) \\ &= \frac{\vartheta_K^{I+}(m_1 \uplus_1 m_2) + \vartheta_L^{I+}(n_1 \uplus_1 n_2)}{2} \\ &\geq \frac{1}{2} \left[\frac{\vartheta_K^{I+}(m_1) + \vartheta_K^{I+}(m_2)}{2} + \frac{\vartheta_L^{I+}(n_1) + \vartheta_L^{I+}(n_2)}{2} \right] \\ &= \frac{1}{2} \left[\frac{\vartheta_K^{I+}(m_1) + \vartheta_L^{I+}(n_1)}{2} + \frac{\vartheta_K^{I+}(m_2) + \vartheta_L^{I+}(n_2)}{2} \right] \\ &= \frac{1}{2} [\vartheta_{K \times L}^{I+}(m_1, n_1) + \vartheta_{K \times L}^{I+}(m_2, n_2)]. \end{aligned}$$

Similarly, $\vartheta_{K \times L}^{I-}[(m_1, n_1) \uplus_1 (m_2, n_2)] \leq \frac{1}{2} [\vartheta_{K \times L}^{I-}(m_1, n_1) + \vartheta_{K \times L}^{I-}(m_2, n_2)]$.

Now,

$$\begin{aligned} \vartheta_{K \times L}^{F+}[(m_1, n_1) \uplus_1 (m_2, n_2)] &= \vartheta_{K \times L}^{F+}(m_1 \uplus_1 m_2, n_1 \uplus_1 n_2) \\ &= \max\{\vartheta_K^{F+}(m_1 \uplus_1 m_2), \vartheta_L^{F+}(n_1 \uplus_1 n_2)\} \\ &\leq \max\{\max\{\vartheta_K^{F+}(m_1), \vartheta_K^{F+}(m_2)\}, \max\{\vartheta_L^{F+}(n_1), \vartheta_L^{F+}(n_2)\}\} \\ &= \max\{\max\{\vartheta_K^{F+}(m_1), \vartheta_L^{F+}(n_1)\}, \max\{\vartheta_K^{F+}(m_2), \vartheta_L^{F+}(n_2)\}\} \\ &= \max\{\vartheta_{K \times L}^{F+}(m_1, n_1), \vartheta_{K \times L}^{F+}(m_2, n_2)\}. \end{aligned}$$

Similarly, $\vartheta_{K \times L}^{F-}[(m_1, n_1) \uplus_1 (m_2, n_2)] \geq \min\{\vartheta_{K \times L}^{F-}(m_1, n_1), \vartheta_{K \times L}^{F-}(m_2, n_2)\}$.

Similarly, we can prove other two operations. Hence, $K \times L$ is an BVNSBS of \mathcal{S} .

Corollary 3.5. *If K_1, K_2, \dots, K_n are the family of BVNSBS^s of $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_n$ respectively, then $K_1 \times K_2 \times \dots \times K_n$ is an BVNSBS of $\mathcal{S}_1 \times \mathcal{S}_2 \times \dots \times \mathcal{S}_n$.*

Definition 3.6. Let K be a bipolar valued neutrosophic subset in \mathcal{S} , the strongest neutrosophic relation on \mathcal{S} , that is a bipolar valued neutrosophic relation on K is O such that

$$\left\{ \begin{aligned} &\left(\begin{aligned} \vartheta_O^{T+}(m, n) &= \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\}, \\ \vartheta_O^{T-}(m, n) &= \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} \end{aligned} \right), & \left(\begin{aligned} \vartheta_O^{I+}(m, n) &= \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \\ \vartheta_O^{I-}(m, n) &= \frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2} \end{aligned} \right), \\ &\left(\begin{aligned} \vartheta_O^{F+}(m, n) &= \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}, \\ \vartheta_O^{F-}(m, n) &= \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\} \end{aligned} \right) \end{aligned} \right\}.$$

Theorem 3.7. *Let K be the BVNSBS of \mathcal{S} and O be the strongest bipolar valued neutrosophic relation of \mathcal{S} . Then K is an BVNSBS of \mathcal{S} if and only if O is an BVNSBS of $\mathcal{S} \times \mathcal{S}$.*

Proof. Let K be the BVNSBS of \mathcal{S} and O be the strongest bipolar valued neutrosophic relation of \mathcal{S} . Then for any $m = (m_1, m_2)$ and $n = (n_1, n_2)$ are in $\mathcal{S} \times \mathcal{S}$. Now,

$$\begin{aligned}\vartheta_O^{T+}(m \uplus_1 n) &= \vartheta_O^{T+}[(m_1, m_2) \uplus_1 (n_1, n_2)] \\ &= \vartheta_O^{T+}(m_1 \uplus_1 n_1, m_2 \uplus_1 n_2) \\ &= \min\{\vartheta_K^{T+}(m_1 \uplus_1 n_1), \vartheta_K^{T+}(m_2 \uplus_1 n_2)\} \\ &\geq \min\{\min\{\vartheta_K^{T+}(m_1), \vartheta_K^{T+}(n_1)\}, \min\{\vartheta_K^{T+}(m_2), \vartheta_K^{T+}(n_2)\}\} \\ &= \min\{\min\{\vartheta_K^{T+}(m_1), \vartheta_K^{T+}(m_2)\}, \min\{\vartheta_K^{T+}(n_1), \vartheta_K^{T+}(n_2)\}\} \\ &= \min\{\vartheta_O^{T+}(m_1, m_2), \vartheta_O^{T+}(n_1, n_2)\} \\ &= \min\{\vartheta_O^{T+}(m), \vartheta_O^{T+}(n)\}.\end{aligned}$$

Similarly, $\vartheta_O^{T-}(m \uplus_2 n) \leq \max\{\vartheta_O^{T-}(m), \vartheta_O^{T-}(n)\}$.

Now,

$$\begin{aligned}\vartheta_O^{I+}(m \uplus_1 n) &= \vartheta_O^{I+}[(m_1, m_2) \uplus_1 (n_1, n_2)] \\ &= \vartheta_O^{I+}(m_1 \uplus_1 n_1, m_2 \uplus_1 n_2) \\ &= \frac{\vartheta_K^{I+}(m_1 \uplus_1 n_1) + \vartheta_K^{I+}(m_2 \uplus_1 n_2)}{2} \\ &\geq \frac{1}{2} \left[\frac{\vartheta_K^{I+}(m_1) + \vartheta_K^{I+}(n_1)}{2} + \frac{\vartheta_K^{I+}(m_2) + \vartheta_K^{I+}(n_2)}{2} \right] \\ &= \frac{1}{2} \left[\frac{\vartheta_K^{I+}(m_1) + \vartheta_K^{I+}(m_2)}{2} + \frac{\vartheta_K^{I+}(n_1) + \vartheta_K^{I+}(n_2)}{2} \right] \\ &= \frac{\vartheta_O^{I+}(m_1, m_2) + \vartheta_O^{I+}(n_1, n_2)}{2} \\ &= \frac{\vartheta_O^{I+}(m) + \vartheta_O^{I+}(n)}{2}.\end{aligned}$$

Similarly, $\vartheta_O^{I-}(m \uplus_1 n) \leq \frac{\vartheta_O^{I-}(m) + \vartheta_O^{I-}(n)}{2}$.

Similarly, $\vartheta_O^{F+}(m \uplus_1 n) \leq \max\{\vartheta_O^{F+}(m), \vartheta_O^{F+}(n)\}$ and $\vartheta_O^{F-}(m \uplus_1 n) \geq \min\{\vartheta_O^{F-}(m), \vartheta_O^{F-}(n)\}$.

Similarly to prove other two operations. Hence O is an BVNSBS of $\mathcal{S} \times \mathcal{S}$.

Conversely assume that O is an BVNSBS of $\mathcal{S} \times \mathcal{S}$, then for any $m = (m_1, m_2)$ and $n = (n_1, n_2)$ are in $\mathcal{S} \times \mathcal{S}$. Now,

$$\begin{aligned}\min\{\vartheta_K^{T+}(m_1 \uplus_1 n_1), \vartheta_K^{T+}(m_2 \uplus_1 n_2)\} &= \vartheta_O^{T+}(m_1 \uplus_1 n_1, m_2 \uplus_1 n_2) \\ &= \vartheta_O^{T+}[(m_1, m_2) \uplus_1 (n_1, n_2)] \\ &= \vartheta_O^{T+}(m \uplus_1 n)\end{aligned}$$

$$\begin{aligned} &\geq \min\{\vartheta_O^{T+}(m), \vartheta_O^{T+}(n)\} \\ &= \min\{\vartheta_O^{T+}(m_1, m_2), \vartheta_O^{T+}(n_1, n_2)\} \\ &= \min\{\min\{\vartheta_K^{T+}(m_1), \vartheta_K^{T+}(m_2)\}, \min\{\vartheta_K^{T+}(n_1), \vartheta_K^{T+}(n_2)\}\}. \end{aligned}$$

If $\vartheta_K^{T+}(m_1 \uplus_1 n_1) \leq \vartheta_K^{T+}(m_2 \uplus_1 n_2)$, then $\vartheta_K^{T+}(m_1) \leq \vartheta_K^{T+}(m_2)$ and $\vartheta_K^{T+}(n_1) \leq \vartheta_K^{T+}(n_2)$.

We get $\vartheta_K^{T+}(m_1 \uplus_1 n_1) \geq \min\{\vartheta_K^{T+}(m_1), \vartheta_K^{T+}(n_1)\}$.

$$\begin{aligned} \max\{\vartheta_K^{T-}(m_1 \uplus_1 n_1), \vartheta_K^{T-}(m_2 \uplus_1 n_2)\} &= \vartheta_O^{T-}(m_1 \uplus_1 n_1, m_2 \uplus_1 n_2) \\ &= \vartheta_O^{T-}[(m_1, m_2) \uplus_1 (n_1, n_2)] \\ &= \vartheta_O^{T-}(m \uplus_1 n) \\ &\leq \max\{\vartheta_O^{T-}(m), \vartheta_O^{T-}(n)\} \\ &= \max\{\vartheta_O^{T-}(m_1, m_2), \vartheta_O^{T-}(n_1, n_2)\} \\ &= \max\{\max\{\vartheta_K^{T-}(m_1), \vartheta_K^{T-}(m_2)\}, \max\{\vartheta_K^{T-}(n_1), \vartheta_K^{T-}(n_2)\}\}. \end{aligned}$$

If $\vartheta_K^{T-}(m_1 \uplus_1 n_1) \geq \vartheta_K^{T-}(m_2 \uplus_1 n_2)$, then $\vartheta_K^{T-}(m_1) \geq \vartheta_K^{T-}(m_2)$ and $\vartheta_K^{T-}(n_1) \geq \vartheta_K^{T-}(n_2)$.

We get $\vartheta_K^{T-}(m_1 \uplus_1 n_1) \leq \max\{\vartheta_K^{T-}(m_1), \vartheta_K^{T-}(n_1)\}$ for all $m_1, n_1 \in \mathcal{S}$. Now,

$$\begin{aligned} \frac{1}{2} \left[\vartheta_K^{I+}(m_1 \uplus_1 n_1) + \vartheta_K^{I+}(m_2 \uplus_1 n_2) \right] &= \vartheta_O^{I+}(m_1 \uplus_1 n_1, m_2 \uplus_1 n_2) \\ &= \vartheta_O^{I+}[(m_1, m_2) \uplus_1 (n_1, n_2)] \\ &= \vartheta_O^{I+}(m \uplus_1 n) \\ &\geq \frac{\vartheta_O^{I+}(m) + \vartheta_O^{I+}(n)}{2} \\ &= \frac{\vartheta_O^{I+}(m_1, m_2) + \vartheta_O^{I+}(n_1, n_2)}{2} \\ &= \frac{1}{2} \left[\frac{\vartheta_K^{I+}(m_1) + \vartheta_K^{I+}(m_2)}{2} + \frac{\vartheta_K^{I+}(n_1) + \vartheta_K^{I+}(n_2)}{2} \right]. \end{aligned}$$

If $\vartheta_K^{I+}(m_1 \uplus_1 n_1) \leq \vartheta_K^{I+}(m_2 \uplus_1 n_2)$, then $\vartheta_K^{I+}(m_1) \leq \vartheta_K^{I+}(m_2)$ and $\vartheta_K^{I+}(n_1) \leq \vartheta_K^{I+}(n_2)$.

We get, $\vartheta_K^{I+}(m_1 \uplus_1 n_1) \geq \frac{\vartheta_K^{I+}(m_1) + \vartheta_K^{I+}(n_1)}{2}$.

Similarly, $\frac{1}{2} \left[\vartheta_K^{I-}(m_1 \uplus_1 n_1) + \vartheta_K^{I-}(m_2 \uplus_1 n_2) \right] \leq \frac{1}{2} \left[\frac{\vartheta_K^{I-}(m_1) + \vartheta_K^{I-}(m_2)}{2} + \frac{\vartheta_K^{I-}(n_1) + \vartheta_K^{I-}(n_2)}{2} \right]$.

If $\vartheta_K^{I-}(m_1 \uplus_1 n_1) \geq \vartheta_K^{I-}(m_2 \uplus_1 n_2)$, then $\vartheta_K^{I-}(m_1) \geq \vartheta_K^{I-}(m_2)$ and $\vartheta_K^{I-}(n_1) \geq \vartheta_K^{I-}(n_2)$.

We get, $\vartheta_K^{I-}(m_1 \uplus_1 n_1) \leq \frac{\vartheta_K^{I-}(m_1) + \vartheta_K^{I-}(n_1)}{2}$.

Similarly, $\max\{\vartheta_K^{F+}(m_1 \uplus_1 n_1), \vartheta_K^{F+}(m_2 \uplus_1 n_2)\} \leq \max\{\max\{\vartheta_K^{F+}(m_1), \vartheta_K^{F+}(m_2)\}, \max\{\vartheta_K^{F+}(n_1), \vartheta_K^{F+}(n_2)\}\}$.

If $\vartheta_K^{F+}(m_1 \uplus_1 n_1) \geq \vartheta_K^{F+}(m_2 \uplus_1 n_2)$, then $\vartheta_K^{F+}(m_1) \geq \vartheta_K^{F+}(m_2)$ and $\vartheta_K^{F+}(n_1) \geq \vartheta_K^{F+}(n_2)$.

We get, $\vartheta_K^{F+}(m_1 \uplus_1 n_1) \leq \max\{\vartheta_K^{F+}(m_1), \vartheta_K^{F+}(n_1)\}$.

Similarly, $\min\{\vartheta_K^{F-}(m_1 \uplus_1 n_1), \vartheta_K^{F-}(m_2 \uplus_1 n_2)\} \geq \min\{\min\{\vartheta_K^{F-}(m_1), \vartheta_K^{F-}(m_2)\}, \min\{\vartheta_K^{F-}(n_1), \vartheta_K^{F-}(n_2)\}\}$.

$\min\{\vartheta_K^{F-}(n_1), \vartheta_K^{F-}(n_2)\}$.

If $\vartheta_K^{F-}(m_1 \uplus_1 n_1) \leq \vartheta_K^{F-}(m_2 \uplus_1 n_2)$, then $\vartheta_K^{F-}(m_1) \leq \vartheta_K^{F-}(m_2)$ and $\vartheta_K^{F-}(n_1) \leq \vartheta_K^{F-}(n_2)$.

We get, $\vartheta_K^{F-}(m_1 \uplus_1 n_1) \geq \min\{\vartheta_K^{F-}(m_1), \vartheta_K^{F-}(n_1)\}$.

Similarly to prove other two operations. Hence K is an BVNSBS of \mathcal{S} .

Theorem 3.8. *Let K be bipolar valued neutrosophic subset in \mathcal{S} . Then $\vartheta = \{(\vartheta_K^{T+}, \vartheta_K^{T-}), (\vartheta_K^{I+}, \vartheta_K^{I-}), (\vartheta_K^{F+}, \vartheta_K^{F-})\}$ is an BVNSBS of \mathcal{S} if and only if all non empty level set $\vartheta^{(t,s)}$ is a subbisemiring of \mathcal{S} for $t, s \in [-1, 0] \times [0, 1]$.*

Proof. Assume that ϑ is an BVNSBS of \mathcal{S} . For each $t, s \in [-1, 0] \times [0, 1]$ and $a_1, a_2 \in \vartheta^{(t,s)}$. We have $\vartheta_K^{T+}(a_1) \geq t, \vartheta_K^{T+}(a_2) \geq t$ and $\vartheta_K^{I+}(a_1) \geq t, \vartheta_K^{I+}(a_2) \geq t$ and $\vartheta_K^{F+}(a_1) \leq s, \vartheta_K^{F+}(a_2) \leq s$. Now, $\vartheta_K^{T+}(a_1 \uplus_1 a_2) \geq \min\{\vartheta_K^{T+}(a_1), \vartheta_K^{T+}(a_2)\} \geq t$ and $\vartheta_K^{I+}(a_1 \uplus_1 a_2) \geq \frac{\vartheta_K^{I+}(a_1) + \vartheta_K^{I+}(a_2)}{2} \geq \frac{t+t}{2} = t$ and $\vartheta_K^{F+}(a_1 \uplus_1 a_2) \leq \max\{\vartheta_K^{F+}(a_1), \vartheta_K^{F+}(a_2)\} \leq s$. Since, $t, s \in [-1, 0] \times [0, 1]$, we have $\vartheta_K^{T-}(a_1) \leq t, \vartheta_K^{T-}(a_2) \leq t$ and $\vartheta_K^{I-}(a_1) \leq t, \vartheta_K^{I-}(a_2) \leq t$ and $\vartheta_K^{F-}(a_1) \geq s, \vartheta_K^{F-}(a_2) \geq s$. Now, $\vartheta_K^{T-}(a_1 \uplus_1 a_2) \leq \max\{\vartheta_K^{T-}(a_1), \vartheta_K^{T-}(a_2)\} \leq t$ and $\vartheta_K^{I-}(a_1 \uplus_1 a_2) \leq \frac{\vartheta_K^{I-}(a_1) + \vartheta_K^{I-}(a_2)}{2} \leq \frac{t+t}{2} = t$ and $\vartheta_K^{F-}(a_1 \uplus_1 a_2) \geq \min\{\vartheta_K^{F-}(a_1), \vartheta_K^{F-}(a_2)\} \geq s$. This implies that $a_1 \uplus_1 a_2 \in \vartheta^{(t,s)}$. Similarly, to prove other two operations. Hence, $\vartheta^{(t,s)}$ is a subbisemiring of \mathcal{S} for each $t, s \in [-1, 0] \times [0, 1]$.

Conversely, assume that $\vartheta^{(t,s)}$ is a subbisemiring of \mathcal{S} for each $t, s \in [-1, 0] \times [0, 1]$. Suppose if there exist $a_1, a_2 \in \mathcal{S}$ such that $\vartheta_K^{T+}(a_1 \uplus_1 a_2) < \min\{\vartheta_K^{T+}(a_1), \vartheta_K^{T+}(a_2)\}, \vartheta_K^{I+}(a_1 \uplus_1 a_2) < \frac{\vartheta_K^{I+}(a_1) + \vartheta_K^{I+}(a_2)}{2}$ and $\vartheta_K^{F+}(a_1 \uplus_1 a_2) > \max\{\vartheta_K^{F+}(a_1), \vartheta_K^{F+}(a_2)\}$. Select $t, s \in [0, 1]$ such that $\vartheta_K^{T+}(a_1 \uplus_1 a_2) < t \leq \min\{\vartheta_K^{T+}(a_1), \vartheta_K^{T+}(a_2)\}$ and $\vartheta_K^{I+}(a_1 \uplus_1 a_2) < t \leq \frac{\vartheta_K^{I+}(a_1) + \vartheta_K^{I+}(a_2)}{2}$ and $\vartheta_K^{F+}(a_1 \uplus_1 a_2) > s \geq \max\{\vartheta_K^{F+}(a_1), \vartheta_K^{F+}(a_2)\}$. Then $a_1, a_2 \in \vartheta^{(t,s)}$, but $a_1 \uplus_1 a_2 \notin \vartheta^{(t,s)}$. Suppose if there exist $a_1, a_2 \in \mathcal{S}$ such that $\vartheta_K^{T-}(a_1 \uplus_1 a_2) > \max\{\vartheta_K^{T-}(a_1), \vartheta_K^{T-}(a_2)\}, \vartheta_K^{I-}(a_1 \uplus_1 a_2) > \frac{\vartheta_K^{I-}(a_1) + \vartheta_K^{I-}(a_2)}{2}$ and $\vartheta_K^{F-}(a_1 \uplus_1 a_2) < \min\{\vartheta_K^{F-}(a_1), \vartheta_K^{F-}(a_2)\}$. Select $t, s \in [-1, 0]$ such that $\vartheta_K^{T-}(a_1 \uplus_1 a_2) > t \geq \max\{\vartheta_K^{T-}(a_1), \vartheta_K^{T-}(a_2)\}$ and $\vartheta_K^{I-}(a_1 \uplus_1 a_2) > t \geq \frac{\vartheta_K^{I-}(a_1) + \vartheta_K^{I-}(a_2)}{2}$ and $\vartheta_K^{F-}(a_1 \uplus_1 a_2) < s \leq \min\{\vartheta_K^{F-}(a_1), \vartheta_K^{F-}(a_2)\}$. Then $a_1, a_2 \in \vartheta^{(t,s)}$, but $a_1 \uplus_1 a_2 \notin \vartheta^{(t,s)}$. This contradicts to that $\vartheta^{(t,s)}$ is a subbisemiring of \mathcal{S} . Hence $\vartheta_K^{T+}(a_1 \uplus_1 a_2) \geq \min\{\vartheta_K^{T+}(a_1), \vartheta_K^{T+}(a_2)\}, \vartheta_K^{T-}(a_1 \uplus_1 a_2) \leq \max\{\vartheta_K^{T-}(a_1), \vartheta_K^{T-}(a_2)\}, \vartheta_K^{I+}(a_1 \uplus_1 a_2) \geq \frac{\vartheta_K^{I+}(a_1) + \vartheta_K^{I+}(a_2)}{2}, \vartheta_K^{I-}(a_1 \uplus_1 a_2) \leq \frac{\vartheta_K^{I-}(a_1) + \vartheta_K^{I-}(a_2)}{2}$ and $\vartheta_K^{F+}(a_1 \uplus_1 a_2) \leq \max\{\vartheta_K^{F+}(a_1), \vartheta_K^{F+}(a_2)\}, \vartheta_K^{F-}(a_1 \uplus_1 a_2) \geq \min\{\vartheta_K^{F-}(a_1), \vartheta_K^{F-}(a_2)\}$. Similarly to prove other two operations such as \uplus_2 and \uplus_3 . Hence $\tilde{\vartheta} = \{(\vartheta_K^{T+}, \vartheta_K^{T-}), (\vartheta_K^{I+}, \vartheta_K^{I-}), (\vartheta_K^{F+}, \vartheta_K^{F-})\}$ is an BVNSBS of \mathcal{S} .

Definition 3.9. Let K be any BVNSBS of \mathcal{S} and $a \in \mathcal{S}$. Then the pseudo bipolar valued neutrosophic coset $(aA)^z$ is defined by

$$\left\{ \left(\begin{aligned} ((a\vartheta_K^{T+})^z)(m) &= z(a)\vartheta_K^{T+}(m), \\ ((a\vartheta_K^{T-})^z)(m) &= z(a)\vartheta_K^{T-}(m) \end{aligned} \right), \left(\begin{aligned} ((a\vartheta_K^{I+})^z)(m) &= z(a)\vartheta_K^{I+}(m), \\ ((a\vartheta_K^{I-})^z)(m) &= z(a)\vartheta_K^{I-}(m) \end{aligned} \right), \right. \\ \left. \left(\begin{aligned} ((a\vartheta_K^{F+})^z)(m) &= z(a)\vartheta_K^{F+}(m), \\ ((a\vartheta_K^{F-})^z)(m) &= z(a)\vartheta_K^{F-}(m) \end{aligned} \right) \right\}.$$

for every $m \in \mathcal{S}$ and for some $z \in P$, where P is a any non-empty set.

Theorem 3.10. *Let K be any BVNSBS of \mathcal{S} , then the pseudo bipolar valued neutrosophic coset $(aA)^z$ is an BVNSBS of \mathcal{S} , for every $a \in \mathcal{S}$.*

Proof. Now, $((a\vartheta_K^{T+})^z)(m \uplus_1 n) = z(a) \vartheta_K^{T+}(m \uplus_1 n) \geq z(a) \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\} = \min\{z(a) \vartheta_K^{T+}(m), z(a) \vartheta_K^{T+}(n)\} = \min\{((a\vartheta_K^{T+})^z)(m), ((a\vartheta_K^{T+})^z)(n)\}$. Thus, $((a\vartheta_K^{T+})^z)(m \uplus_1 n) \geq \min\{((a\vartheta_K^{T+})^z)(m), ((a\vartheta_K^{T+})^z)(n)\}$. Now, $((a\vartheta_K^{I+})^z)(m \uplus_1 n) = z(a) \vartheta_K^{I+}(m \uplus_1 n) \geq z(a) \left[\frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2} \right] = \frac{z(a) \vartheta_K^{I+}(m) + z(a) \vartheta_K^{I+}(n)}{2} = \frac{((a\vartheta_K^{I+})^z)(m) + ((a\vartheta_K^{I+})^z)(n)}{2}$. Thus, $((a\vartheta_K^{I+})^z)(m \uplus_1 n) \geq \frac{((a\vartheta_K^{I+})^z)(m) + ((a\vartheta_K^{I+})^z)(n)}{2}$. Now, $((a\vartheta_K^{F+})^z)(m \uplus_1 n) = z(a) \vartheta_K^{F+}(m \uplus_1 n) \leq z(a) \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\} = \max\{z(a) \vartheta_K^{F+}(m), z(a) \vartheta_K^{F+}(n)\} = \max\{((a\vartheta_K^{F+})^z)(m), ((a\vartheta_K^{F+})^z)(n)\}$. Thus, $((a\vartheta_K^{F+})^z)(m \uplus_1 n) \leq \max\{((a\vartheta_K^{F+})^z)(m), ((a\vartheta_K^{F+})^z)(n)\}$. Also, $((a\vartheta_K^{T-})^z)(m \uplus_1 n) = z(a) \vartheta_K^{T-}(m \uplus_1 n) \leq z(a) \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} = \max\{z(a) \vartheta_K^{T-}(m), z(a) \vartheta_K^{T-}(n)\} = \max\{((a\vartheta_K^{T-})^z)(m), ((a\vartheta_K^{T-})^z)(n)\}$. Thus, $((a\vartheta_K^{T-})^z)(m \uplus_1 n) \leq \max\{((a\vartheta_K^{T-})^z)(m), ((a\vartheta_K^{T-})^z)(n)\}$. Now, $((a\vartheta_K^{I-})^z)(m \uplus_1 n) = z(a) \vartheta_K^{I-}(m \uplus_1 n) \leq z(a) \left[\frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2} \right] = \frac{z(a) \vartheta_K^{I-}(m) + z(a) \vartheta_K^{I-}(n)}{2} = \frac{((a\vartheta_K^{I-})^z)(m) + ((a\vartheta_K^{I-})^z)(n)}{2}$. Thus, $((a\vartheta_K^{I-})^z)(m \uplus_1 n) \leq \frac{((a\vartheta_K^{I-})^z)(m) + ((a\vartheta_K^{I-})^z)(n)}{2}$. Now, $((a\vartheta_K^{F-})^z)(m \uplus_1 n) = z(a) \vartheta_K^{F-}(m \uplus_1 n) \geq z(a) \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\} = \min\{z(a) \vartheta_K^{F-}(m), z(a) \vartheta_K^{F-}(n)\} = \min\{((a\vartheta_K^{F-})^z)(m), ((a\vartheta_K^{F-})^z)(n)\}$. Thus, $((a\vartheta_K^{F-})^z)(m \uplus_1 n) \geq \min\{((a\vartheta_K^{F-})^z)(m), ((a\vartheta_K^{F-})^z)(n)\}$. Similarly to prove other two operations such as \uplus_2 and \uplus_3 . Hence $(aA)^z$ is an BVNSBS of \mathcal{S} .

Definition 3.11. Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any function and K be any BVNSBS in \mathcal{S}_1 , O be any BVNSBS in $\Lambda(\mathcal{S}_1) = \mathcal{S}_2$. If $\vartheta_K = \{(\vartheta_K^{T+}, \vartheta_K^{T-}), (\vartheta_K^{I+}, \vartheta_K^{I-}), (\vartheta_K^{F+}, \vartheta_K^{F-})\}$ is a bipolar valued neutrosophic set in \mathcal{S}_1 , then ϑ_O is a bipolar valued neutrosophic set in \mathcal{S}_2 , defined by

$$\begin{aligned} \vartheta_O^{T+}(n) &= \begin{cases} \sup \vartheta_K^{T+}(m) & \text{if } m \in \Lambda^{-1}(n) \\ 0 & \text{otherwise} \end{cases} & ; \vartheta_O^{T-}(n) &= \begin{cases} \inf \vartheta_K^{T-}(m) & \text{if } m \in \Lambda^{-1}(n) \\ -1 & \text{otherwise} \end{cases} \\ \vartheta_O^{I+}(n) &= \begin{cases} \sup \vartheta_K^{I+}(m) & \text{if } m \in \Lambda^{-1}(n) \\ 0 & \text{otherwise} \end{cases} & ; \vartheta_O^{I-}(n) &= \begin{cases} \inf \vartheta_K^{I-}(m) & \text{if } m \in \Lambda^{-1}(n) \\ -1 & \text{otherwise} \end{cases} \\ \vartheta_O^{F+}(n) &= \begin{cases} \inf \vartheta_K^{F+}(m) & \text{if } m \in \Lambda^{-1}(n) \\ 1 & \text{otherwise} \end{cases} & ; \vartheta_O^{F-}(n) &= \begin{cases} \sup \vartheta_K^{F-}(m) & \text{if } m \in \Lambda^{-1}(n) \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

for all $m \in \mathcal{S}_1$ and $n \in \mathcal{S}_2$ is called the image of ϑ_K under Λ .

If $\vartheta_O = \{(\vartheta_O^{T+}, \vartheta_O^{T-}), (\vartheta_O^{I+}, \vartheta_O^{I-}), (\vartheta_O^{F+}, \vartheta_O^{F-})\}$ is a bipolar valued neutrosophic set in \mathcal{S}_2 , then neutrosophic set $\vartheta_K = \Lambda \circ \vartheta_O$ in \mathcal{S}_1 [ie, the bipolar valued neutrosophic set defined by $\vartheta_K(m) = \vartheta_O(\Lambda(m))$] is called the preimage of ϑ_O under Λ .

Theorem 3.12. *Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. The homomorphic image of BVNSBS of \mathcal{S}_1 is an BVNSBS of \mathcal{S}_2 .*

Proof. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any homomorphism. Then $\Lambda(m \vee_1 n) = \Lambda(m) \sqcup_1 \Lambda(n), \Lambda(m \vee_2 n) = \Lambda(m) \sqcup_2 \Lambda(n)$ and $\Lambda(m \vee_3 n) = \Lambda(m) \sqcup_3 \Lambda(n)$ for all $m, n \in \mathcal{S}_1$. Let $O = \Lambda(K)$, K is any BVNSBS of \mathcal{S}_1 . Let $\Lambda(m), \Lambda(n) \in \mathcal{S}_2$. Let $m \in \Lambda^{-1}(\Lambda(m))$ and $n \in \Lambda^{-1}(\Lambda(n))$ be such that $\vartheta_K^{T+}(m) = \sup_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{T+}(z), \vartheta_K^{T+}(n) = \sup_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{T+}(z)$ and $\vartheta_K^{T-}(m) = \inf_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{T-}(z), \vartheta_K^{T-}(n) = \inf_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{T-}(z)$. Now,

$$\begin{aligned} \vartheta_O^{T+}(\Lambda(m) \sqcup_1 \Lambda(n)) &= \sup_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{T+}(z') \\ &= \sup_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{T+}(z') \\ &= \vartheta_K^{T+}(m \vee_1 n) \\ &\geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\} \\ &= \min\{\vartheta_O^{T+} \Lambda(m), \vartheta_O^{T+} \Lambda(n)\}. \end{aligned}$$

Thus, $\vartheta_O^{T+}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \min\{\vartheta_O^{T+} \Lambda(m), \vartheta_O^{T+} \Lambda(n)\}$.

$$\begin{aligned} \vartheta_O^{T-}(\Lambda(m) \sqcup_1 \Lambda(n)) &= \inf_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{T-}(z') \\ &= \inf_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{T-}(z') \\ &= \vartheta_K^{T-}(m \vee_1 n) \\ &\leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} \\ &= \max\{\vartheta_O^{T-} \Lambda(m), \vartheta_O^{T-} \Lambda(n)\}. \end{aligned}$$

Thus, $\vartheta_O^{T-}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \max\{\vartheta_O^{T-} \Lambda(m), \vartheta_O^{T-} \Lambda(n)\}$.

Let $m \in \Lambda^{-1}(\Lambda(m))$ and $n \in \Lambda^{-1}(\Lambda(n))$ be such that $\vartheta_K^{I+}(m) = \sup_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{I+}(z),$
 $\vartheta_K^{I+}(n) = \sup_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{I+}(z), \vartheta_K^{I-}(m) = \inf_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{I-}(z), \vartheta_K^{I-}(n) = \inf_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{I-}(z)$.

Now,

$$\begin{aligned} \vartheta_O^{I+}(\Lambda(m) \sqcup_1 \Lambda(n)) &= \sup_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{I+}(z') \\ &= \sup_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{I+}(z') \\ &= \vartheta_K^{I+}(m \vee_1 n) \\ &\geq \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2} \\ &= \frac{\vartheta_O^{I+} \Lambda(m) + \vartheta_O^{I+} \Lambda(n)}{2}. \end{aligned}$$

Thus, $\vartheta_O^{I+}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \frac{\vartheta_O^{I+}\Lambda(m) + \vartheta_O^{I+}\Lambda(n)}{2}$.

Similarly, $\vartheta_O^{I-}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \frac{\vartheta_O^{I-}\Lambda(m) + \vartheta_O^{I-}\Lambda(n)}{2}$.

Let $\Lambda(m), \Lambda(n) \in \mathcal{S}_2$. Let $m \in \Lambda^{-1}(\Lambda(m))$ and $n \in \Lambda^{-1}(\Lambda(n))$ be such that

$$\vartheta_K^{F+}(m) = \inf_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{F+}(z), \vartheta_K^{F+}(n) = \inf_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{F+}(z), \vartheta_K^{F-}(m) = \sup_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{F-}(z)$$

and $\vartheta_K^{F-}(n) = \sup_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{F-}(z)$. Now,

$$\begin{aligned} \vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)) &= \inf_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{F+}(z') \\ &= \inf_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{F+}(z') \\ &= \vartheta_K^{F+}(m \vee_1 n) \\ &\leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\} \\ &= \max\{\vartheta_O^{F+}\Lambda(m), \vartheta_O^{F+}\Lambda(n)\}. \end{aligned}$$

Thus, $\vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \max\{\vartheta_O^{F+}\Lambda(m), \vartheta_O^{F+}\Lambda(n)\}$.

Similarly, $\vartheta_O^{F-}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \min\{\vartheta_O^{F-}\Lambda(m), \vartheta_O^{F-}\Lambda(n)\}$.

Similarly, to prove other two operations. Hence O is an BVNSBS of \mathcal{S}_2 .

Theorem 3.13. *Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. The homomorphic preimage of BVNSBS of \mathcal{S}_2 is an BVNSBS of \mathcal{S}_1 .*

Proof. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any homomorphism. Then $\Lambda(m \vee_1 n) = \Lambda(m) \sqcup_1 \Lambda(n)$, $\Lambda(m \vee_2 n) = \Lambda(m) \sqcup_2 \Lambda(n)$ and $\Lambda(m \vee_3 n) = \Lambda(m) \sqcup_3 \Lambda(n)$ for all $m, n \in \mathcal{S}_1$. Let $O = \Lambda(K)$, where O is any BVNSBS of \mathcal{S}_2 . Let $m, n \in \mathcal{S}_1$. Now, $\vartheta_K^{T+}(m \vee_1 n) = \vartheta_O^{T+}(\Lambda(m \vee_1 n)) = \vartheta_O^{T+}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \min\{\vartheta_O^{T+}\Lambda(m), \vartheta_O^{T+}\Lambda(n)\} = \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\}$. Thus, $\vartheta_K^{T+}(m \vee_1 n) \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\}$. Now, $\vartheta_K^{I+}(m \vee_1 n) = \vartheta_O^{I+}(\Lambda(m \vee_1 n)) = \vartheta_O^{I+}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \frac{\vartheta_O^{I+}\Lambda(m) + \vartheta_O^{I+}\Lambda(n)}{2} = \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}$. Thus, $\vartheta_K^{I+}(m \vee_1 n) \geq \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}$. Now, $\vartheta_K^{F+}(m \vee_1 n) = \vartheta_O^{F+}(\Lambda(m \vee_1 n)) = \vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \max\{\vartheta_O^{F+}\Lambda(m), \vartheta_O^{F+}\Lambda(n)\} = \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}$. Thus, $\vartheta_K^{F+}(m \vee_1 n) \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\}$. Also, $\vartheta_K^{T-}(m \vee_1 n) = \vartheta_O^{T-}(\Lambda(m \vee_1 n)) = \vartheta_O^{T-}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \max\{\vartheta_O^{T-}\Lambda(m), \vartheta_O^{T-}\Lambda(n)\} = \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\}$. Thus, $\vartheta_K^{T-}(m \vee_1 n) \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\}$. We have, $\vartheta_K^{I-}(m \vee_1 n) = \vartheta_O^{I-}(\Lambda(m \vee_1 n)) = \vartheta_O^{I-}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \frac{\vartheta_O^{I-}\Lambda(m) + \vartheta_O^{I-}\Lambda(n)}{2} = \frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}$. Thus, $\vartheta_K^{I-}(m \vee_1 n) \leq \frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}$. Now, $\vartheta_K^{F-}(m \vee_1 n) = \vartheta_O^{F-}(\Lambda(m \vee_1 n)) = \vartheta_O^{F-}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \min\{\vartheta_O^{F-}\Lambda(m), \vartheta_O^{F-}\Lambda(n)\} = \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\}$. Thus, $\vartheta_K^{F-}(m \vee_1 n) \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\}$. Similarly to prove other two operations, hence K is an BVNSBS of \mathcal{S}_1 .

Theorem 3.14. *Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. If $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ is a homomorphism, then $\Lambda(K_{(t,s)})$ is a level subbisemiring of BVNSBS O of \mathcal{S}_2 .*

Proof. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any homomorphism. Then $\Lambda(m \vee_1 n) = \Lambda(m) \sqcup_1 \Lambda(n)$, $\Lambda(m \vee_2 n) = \Lambda(m) \sqcup_2 \Lambda(n)$ and $\Lambda(m \vee_3 n) = \Lambda(m) \sqcup_3 \Lambda(n)$ for all $m, n \in \mathcal{S}_1$. Let $O = \Lambda(K)$, K is an BVNSBS of \mathcal{S}_1 . By Theorem 3.12, O is an BVNSBS of \mathcal{S}_2 . Let $K_{(t,s)}$ be any level subbisemiring of K . Suppose that $m, n \in K_{(t,s)}$. Then $\Lambda(m \vee_1 n), \Lambda(m \vee_2 n)$ and $\Lambda(m \vee_3 n) \in K_{(t,s)}$. Now, $\vartheta_O^{T+}(\Lambda(m)) = \vartheta_K^{T+}(m) \geq t, \vartheta_O^{T+}(\Lambda(n)) = \vartheta_K^{T+}(n) \geq t$. Thus, $\vartheta_O^{T+}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \vartheta_K^{T+}(m \vee_1 n) \geq t$. Now, $\vartheta_O^{I+}(\Lambda(m)) = \vartheta_K^{I+}(m) \geq t, \vartheta_O^{I+}(\Lambda(n)) = \vartheta_K^{I+}(n) \geq t$. Thus, $\vartheta_O^{I+}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \vartheta_K^{I+}(m \vee_1 n) \geq t$. Now, $\vartheta_O^{F+}(\Lambda(m)) = \vartheta_K^{F+}(m) \leq s, \vartheta_O^{F+}(\Lambda(n)) = \vartheta_K^{F+}(n) \leq s$. Thus, $\vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \vartheta_K^{F+}(m \vee_1 n) \leq s$, for all $\Lambda(m), \Lambda(n) \in \mathcal{S}_2$. Also, $\vartheta_O^{T-}(\Lambda(m)) = \vartheta_K^{T-}(m) \leq t, \vartheta_O^{T-}(\Lambda(n)) = \vartheta_K^{T-}(n) \leq t$. Thus, $\vartheta_O^{T-}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \vartheta_K^{T-}(m \vee_1 n) \leq t$. Now, $\vartheta_O^{I-}(\Lambda(m)) = \vartheta_K^{I-}(m) \leq t, \vartheta_O^{I-}(\Lambda(n)) = \vartheta_K^{I-}(n) \leq t$. Thus, $\vartheta_O^{I-}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \vartheta_K^{I-}(m \vee_1 n) \leq t$. Now, $\vartheta_O^{F-}(\Lambda(m)) = \vartheta_K^{F-}(m) \geq s, \vartheta_O^{F-}(\Lambda(n)) = \vartheta_K^{F-}(n) \geq s$. Thus, $\vartheta_O^{F-}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \vartheta_K^{F-}(m \vee_1 n) \geq s$, for all $\Lambda(m), \Lambda(n) \in \mathcal{S}_2$. Similarly to prove other operations, hence $\Lambda(K_{(t,s)})$ is a level subbisemiring of BVNSBS O of \mathcal{S}_2 .

Theorem 3.15. Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. If $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ is any homomorphism, then $K_{(t,s)}$ is a level subbisemiring of BVNSBS K of \mathcal{S}_1 .

Proof. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any homomorphism. Then $\Lambda(m \vee_1 n) = \Lambda(m) \sqcup_1 \Lambda(n)$, $\Lambda(m \vee_2 n) = \Lambda(m) \sqcup_2 \Lambda(n)$ and $\Lambda(m \vee_3 n) = \Lambda(m) \sqcup_3 \Lambda(n)$ for all $m, n \in \mathcal{S}_1$. Let $O = \Lambda(K)$, O is an BVNSBS of \mathcal{S}_2 . By Theorem 3.13, K is an BVNSBS of \mathcal{S}_1 . Let $\Lambda(K_{(t,s)})$ be a level subbisemiring of O . Suppose that $\Lambda(m), \Lambda(n) \in \Lambda(K_{(t,s)})$. Then $\Lambda(m \vee_1 n), \Lambda(m \vee_2 n)$ and $\Lambda(m \vee_3 n) \in \Lambda(K_{(t,s)})$. Now, $\vartheta_K^{T+}(m) = \vartheta_O^{T+}(\Lambda(m)) \geq t, \vartheta_K^{T+}(n) = \vartheta_O^{T+}(\Lambda(n)) \geq t$. Thus, $\vartheta_K^{T+}(m \vee_1 n) \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n)\} \geq t$. Now, $\vartheta_K^{I+}(m) = \vartheta_O^{I+}(\Lambda(m)) \geq t, \vartheta_K^{I+}(n) = \vartheta_O^{I+}(\Lambda(n)) \geq t$. Thus, $\vartheta_K^{I+}(m \vee_1 n) \geq \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2} \geq t$. Now, $\vartheta_K^{F+}(m) = \vartheta_O^{F+}(\Lambda(m)) \leq s, \vartheta_K^{F+}(n) = \vartheta_O^{F+}(\Lambda(n)) \leq s$. Thus, $\vartheta_K^{F+}(m \vee_1 n) = \vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)) \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n)\} \leq s$, for all $m, n \in \mathcal{S}_1$. Also, $\vartheta_K^{T-}(m) = \vartheta_O^{T-}(\Lambda(m)) \leq t, \vartheta_K^{T-}(n) = \vartheta_O^{T-}(\Lambda(n)) \leq t$. Thus, $\vartheta_K^{T-}(m \vee_1 n) \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n)\} \leq t$. Now, $\vartheta_K^{I-}(m) = \vartheta_O^{I-}(\Lambda(m)) \leq t, \vartheta_K^{I-}(n) = \vartheta_O^{I-}(\Lambda(n)) \leq t$. Thus, $\vartheta_K^{I-}(m \vee_1 n) \leq \frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2} \leq t$. Now, $\vartheta_K^{F-}(m) = \vartheta_O^{F-}(\Lambda(m)) \geq s, \vartheta_K^{F-}(n) = \vartheta_O^{F-}(\Lambda(n)) \geq s$. Thus, $\vartheta_K^{F-}(m \vee_1 n) = \vartheta_O^{F-}(\Lambda(m) \sqcup_1 \Lambda(n)) \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n)\} \geq s$, for all $m, n \in \mathcal{S}_1$. In the same way, prove the other two operations, hence $K_{(t,s)}$ is a level subbisemiring of BVNSBS K of \mathcal{S}_1 .

4. (ξ, τ) -Bipolar Valued Neutrosophic Subbisemiring

In this section, we discuss (ξ, τ) -bipolar valued neutrosophic subbisemiring. In what follows that, $(\xi^+, \tau^+) \in [0, 1]$ and $(\xi^-, \tau^-) \in [-1, 0]$ be such that $0 \leq \xi^+ < \tau^+ \leq 1$ and $-1 \leq \tau^- < \xi^- \leq 0$, both $(\xi, \tau) \in [0, 1]$ are arbitrary but fixed.

Definition 4.1. Let K be any bipolar valued neutrosophic subset of \mathcal{S} is called a (ξ, τ) -BVNSBS of \mathcal{S} if it satisfies the following conditions:

$$\left\{ \begin{array}{l} \left(\max\{\vartheta_K^{T+}(m \uplus_1 n), \xi^+\} \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+\}, \right. \\ \left. \min\{\vartheta_K^{T-}(m \uplus_1 n), \xi^-\} \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\} \right) \\ \left(\max\{\vartheta_K^{T+}(m \uplus_2 n), \xi^+\} \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+\}, \right. \\ \left. \min\{\vartheta_K^{T-}(m \uplus_2 n), \xi^-\} \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\} \right) \\ \left(\max\{\vartheta_K^{T+}(m \uplus_3 n), \xi^+\} \geq \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+\}, \right. \\ \left. \min\{\vartheta_K^{T-}(m \uplus_3 n), \xi^-\} \leq \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\} \right) \end{array} \right\} \left\{ \begin{array}{l} \left(\max\{\vartheta_K^{I+}(m \uplus_1 n), \xi^+\} \geq \min\left\{\frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \tau^+\right\} \right. \\ \left. \min\{\vartheta_K^{I-}(m \uplus_1 n), \xi^-\} \leq \max\left\{\frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}, \tau^-\right\} \right) \\ \text{OR} \\ \left(\max\{\vartheta_K^{I+}(m \uplus_2 n), \xi^+\} \geq \min\left\{\frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \tau^+\right\} \right. \\ \left. \min\{\vartheta_K^{I-}(m \uplus_2 n), \xi^-\} \leq \max\left\{\frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}, \tau^-\right\} \right) \\ \text{OR} \\ \left(\max\{\vartheta_K^{I+}(m \uplus_3 n), \xi^+\} \geq \min\left\{\frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \tau^+\right\} \right. \\ \left. \min\{\vartheta_K^{I-}(m \uplus_3 n), \xi^-\} \leq \max\left\{\frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}, \tau^-\right\} \right) \end{array} \right\} \\ \left\{ \begin{array}{l} \left(\min\{\vartheta_K^{F+}(m \uplus_1 n), \xi^+\} \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\}, \right. \\ \left. \max\{\vartheta_K^{F-}(m \uplus_1 n), \xi^-\} \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\} \right) \\ \left(\min\{\vartheta_K^{F+}(m \uplus_2 n), \xi^+\} \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\}, \right. \\ \left. \max\{\vartheta_K^{F-}(m \uplus_2 n), \xi^-\} \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\} \right) \\ \left(\min\{\vartheta_K^{F+}(m \uplus_3 n), \xi^+\} \leq \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\}, \right. \\ \left. \max\{\vartheta_K^{F-}(m \uplus_3 n), \xi^-\} \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\} \right) \end{array} \right\}$$

for all $m, n \in \mathcal{S}$.

Example 4.2. By the Example 3.2,

$(\vartheta_K^+(l), \vartheta_K^-(l))$	$l = l_1$	$l = l_2$	$l = l_3$	$l = l_4$
$(\vartheta_K^{T+}(l), \vartheta_K^{T-}(l))$	(0.85, -0.95)	(0.8, -0.75)	(0.7, -0.55)	(0.75, -0.65)
$(\vartheta_K^{I+}(l), \vartheta_K^{I-}(l))$	(0.95, -0.8)	(0.9, -0.7)	(0.8, -0.5)	(0.85, -0.55)
$(\vartheta_K^{F+}(l), \vartheta_K^{F-}(l))$	(0.65, -0.25)	(0.85, -0.35)	(0.95, -0.45)	(0.90, -0.40)

Clearly, K is a (0.60, 0.70)-BVNSBS of \mathcal{S} .

Theorem 4.3. The intersection of family of (ξ, τ) -BVNSBS^s of \mathcal{S} is a (ξ, τ) -BVNSBS of \mathcal{S} .

Proof. Let $\{O_i | i \in I\}$ be any family of (ξ, τ) -BVNSBS^s of \mathcal{S} and $K = \bigcap_{i \in I} O_i$.

Let m and n in \mathcal{S} . Now,

$$\begin{aligned} \max\{\vartheta_K^{T+}(m \uplus_1 n), \xi^+\} &= \inf_{i \in I} \max\{\vartheta_{O_i}^{T+}(m \uplus_1 n), \xi^+\} \\ &\geq \inf_{i \in I} \min\{\vartheta_{O_i}^{T+}(m), \vartheta_{O_i}^{T+}(n), \tau^+\} \\ &= \min \left\{ \inf_{i \in I} \vartheta_{O_i}^{T+}(m), \inf_{i \in I} \vartheta_{O_i}^{T+}(n), \tau^+ \right\} \\ &= \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+\} \end{aligned}$$

$$\begin{aligned}
\min\{\vartheta_K^{T-}(m \uplus_1 n), \xi^-\} &= \sup_{i \in I} \min\{\vartheta_{O_i}^{T-}(m \uplus_1 n), \xi^-\} \\
&\leq \sup_{i \in I} \max\{\vartheta_{O_i}^{T-}(m), \vartheta_{O_i}^{T-}(n), \tau^-\} \\
&= \max\left\{\sup_{i \in I} \vartheta_{O_i}^{T-}(m), \sup_{i \in I} \vartheta_{O_i}^{T-}(n), \tau^-\right\} \\
&= \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\}.
\end{aligned}$$

Now,

$$\begin{aligned}
\max\{\vartheta_K^{I+}(m \uplus_1 n), \xi^+\} &= \inf_{i \in I} \max\{\vartheta_{O_i}^{I+}(m \uplus_1 n), \xi^+\} \\
&\geq \inf_{i \in I} \min\left\{\frac{\vartheta_{O_i}^{I+}(m) + \vartheta_{O_i}^{I+}(n)}{2}, \tau^+\right\} \\
&= \min\left\{\frac{\inf_{i \in I} \vartheta_{O_i}^{I+}(m) + \inf_{i \in I} \vartheta_{O_i}^{I+}(n)}{2}, \tau^+\right\} \\
&= \min\left\{\frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \tau^+\right\}.
\end{aligned}$$

$$\begin{aligned}
\min\{\vartheta_K^{I-}(m \uplus_1 n), \xi^-\} &= \sup_{i \in I} \min\{\vartheta_{O_i}^{I-}(m \uplus_1 n), \xi^-\} \\
&\leq \sup_{i \in I} \max\left\{\frac{\vartheta_{O_i}^{I-}(m) + \vartheta_{O_i}^{I-}(n)}{2}, \tau^-\right\} \\
&= \max\left\{\frac{\sup_{i \in I} \vartheta_{O_i}^{I-}(m) + \sup_{i \in I} \vartheta_{O_i}^{I-}(n)}{2}, \tau^-\right\} \\
&= \max\left\{\frac{\vartheta_K^{I-}(m) + \vartheta_K^{I-}(n)}{2}, \tau^-\right\}.
\end{aligned}$$

Now,

$$\begin{aligned}
\min\{\vartheta_K^{F+}(m \uplus_1 n), \xi^+\} &= \sup_{i \in I} \min\{\vartheta_{O_i}^{F+}(m \uplus_1 n), \xi^+\} \\
&\leq \sup_{i \in I} \max\{\vartheta_{O_i}^{F+}(m), \vartheta_{O_i}^{F+}(n), \tau^+\} \\
&= \max\left\{\sup_{i \in I} \vartheta_{O_i}^{F+}(m), \sup_{i \in I} \vartheta_{O_i}^{F+}(n), \tau^+\right\} \\
&= \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\} \\
\max\{\vartheta_K^{F-}(m \uplus_1 n), \xi^-\} &= \inf_{i \in I} \max\{\vartheta_{O_i}^{F-}(m \uplus_1 n), \xi^-\} \\
&\geq \inf_{i \in I} \min\{\vartheta_{O_i}^{F-}(m), \vartheta_{O_i}^{F-}(n), \tau^-\} \\
&= \min\left\{\inf_{i \in I} \vartheta_{O_i}^{F-}(m), \inf_{i \in I} \vartheta_{O_i}^{F-}(n), \tau^-\right\} \\
&= \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\}.
\end{aligned}$$

Similarly to prove other operations. Hence, K is a (ξ, τ) -BVNSBS of \mathcal{S} .

Theorem 4.4. *If K and L are any two $(\xi, \tau) - BVNSBS^s$ of \mathcal{S}_1 and \mathcal{S}_2 respectively, then $K \times L$ is a $(\xi, \tau) - BVNSBS$ of $\mathcal{S}_1 \times \mathcal{S}_2$.*

Proof. Let K and L be two $(\xi, \tau) - BVNSBS^s$ of \mathcal{S}_1 and \mathcal{S}_2 respectively. Let $m_1, m_2 \in \mathcal{S}_1$ and $n_1, n_2 \in \mathcal{S}_2$. Then (m_1, n_1) and (m_2, n_2) are in $\mathcal{S}_1 \times \mathcal{S}_2$. Now

$$\begin{aligned} & \max \left\{ \vartheta_{K \times L}^{T+}[(m_1, n_1) \uplus_1 (m_2, n_2)], \xi^+ \right\} \\ &= \max \left\{ \vartheta_{K \times L}^{T+}(m_1 \uplus_1 m_2, n_1 \uplus_1 n_2), \xi^+ \right\} \\ &= \min \left\{ \max \{ \vartheta_K^{T+}(m_1 \uplus_1 m_2), \xi^+ \}, \max \{ \vartheta_L^{T+}(n_1 \uplus_1 n_2), \xi^+ \} \right\} \\ &\geq \min \left\{ \min \{ \vartheta_K^{T+}(m_1), \vartheta_K^{T+}(m_2), \tau^+ \}, \min \{ \vartheta_L^{T+}(n_1), \vartheta_L^{T+}(n_2), \tau^+ \} \right\} \\ &= \min \left\{ \{ \min \{ \vartheta_K^{T+}(m_1), \vartheta_L^{T+}(n_1) \}, \min \{ \vartheta_K^{T+}(m_2), \vartheta_L^{T+}(n_2) \} \}, \tau^+ \right\} \\ &= \min \left\{ \vartheta_{K \times L}^{T+}(m_1, n_1), \vartheta_{K \times L}^{T+}(m_2, n_2), \tau^+ \right\}. \end{aligned}$$

Also, $\min \left\{ \vartheta_{K \times L}^{T-}[(m_1, n_1) \uplus_1 (m_2, n_2)], \xi^- \right\} \leq \max \left\{ \vartheta_{K \times L}^{T-}(m_1, n_1), \vartheta_{K \times L}^{T-}(m_2, n_2), \tau^- \right\}$.

Now, $\max \left\{ \vartheta_{K \times L}^{I+}[(m_1, n_1) \uplus_1 (m_2, n_2)], \xi^+ \right\}$

$$\begin{aligned} &= \max \left\{ \vartheta_{K \times L}^{I+}(m_1 \uplus_1 m_2, n_1 \uplus_1 n_2), \xi^+ \right\} \\ &= \min \left\{ \frac{1}{2} \left[\max \{ \vartheta_K^{I+}(m_1 \uplus_1 m_2), \xi^+ \} + \max \{ \vartheta_L^{I+}(n_1 \uplus_1 n_2), \xi^+ \} \right] \right\} \\ &\geq \min \left\{ \frac{1}{2} \left[\min \left\{ \frac{\vartheta_K^{I+}(m_1) + \vartheta_K^{I+}(m_2)}{2}, \tau^+ \right\} + \min \left\{ \frac{\vartheta_L^{I+}(n_1) + \vartheta_L^{I+}(n_2)}{2}, \tau^+ \right\} \right] \right\} \\ &= \min \left\{ \frac{1}{2} \left[\frac{\vartheta_K^{I+}(m_1) + \vartheta_L^{I+}(n_1)}{2} + \frac{\vartheta_K^{I+}(m_2) + \vartheta_L^{I+}(n_2)}{2} \right], \tau^+ \right\} \\ &= \min \left\{ \frac{\vartheta_{K \times L}^{I+}(m_1, n_1) + \vartheta_{K \times L}^{I+}(m_2, n_2)}{2}, \tau^+ \right\}. \end{aligned}$$

Also, $\min \left\{ \vartheta_{K \times L}^{I-}[(m_1, n_1) \uplus_1 (m_2, n_2)], \xi^- \right\} \leq \max \left\{ \frac{\vartheta_{K \times L}^{I-}(m_1, n_1) + \vartheta_{K \times L}^{I-}(m_2, n_2)}{2}, \tau^- \right\}$.

Similarly, $\min \left\{ \vartheta_{K \times L}^{F+}[(m_1, n_1) \uplus_1 (m_2, n_2)], \xi^+ \right\}$

$$\begin{aligned} &= \min \left\{ \vartheta_{K \times L}^{F+}(m_1 \uplus_1 m_2, n_1 \uplus_1 n_2), \xi^+ \right\} \\ &= \max \left\{ \min \{ \vartheta_K^{F+}(m_1 \uplus_1 m_2), \xi^+ \}, \min \{ \vartheta_L^{F+}(n_1 \uplus_1 n_2), \xi^+ \} \right\} \\ &\leq \max \left\{ \max \{ \vartheta_K^{F+}(m_1), \vartheta_K^{F+}(m_2), \tau^+ \}, \max \{ \vartheta_L^{F+}(n_1), \vartheta_L^{F+}(n_2), \tau^+ \} \right\} \\ &= \max \left\{ \{ \max \{ \vartheta_K^{F+}(m_1), \vartheta_L^{F+}(n_1) \}, \max \{ \vartheta_K^{F+}(m_2), \vartheta_L^{F+}(n_2) \} \}, \tau^+ \right\} \\ &= \max \left\{ \vartheta_{K \times L}^{F+}(m_1, n_1), \vartheta_{K \times L}^{F+}(m_2, n_2), \tau^+ \right\}. \end{aligned}$$

Also, $\max \left\{ \vartheta_{K \times L}^{F-}[(m_1, n_1) \uplus_1 (m_2, n_2)], \xi^- \right\} \geq \min \left\{ \vartheta_{K \times L}^{F-}(m_1, n_1), \vartheta_{K \times L}^{F-}(m_2, n_2), \tau^- \right\}$.

In the same way, prove the other two operations. Hence $K \times L$ is a (ξ, τ) - BVNSBS of $\mathcal{S}_1 \times \mathcal{S}_2$.

Corollary 4.5. *If K_1, K_2, \dots, K_n are the family of (ξ, τ) - BVNSBS^s of $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_n$ respectively, then $K_1 \times K_2 \times \dots \times K_n$ is a (ξ, τ) - BVNSBS of $\mathcal{S}_1 \times \mathcal{S}_2 \times \dots \times \mathcal{S}_n$.*

Definition 4.6. Let K be any (ξ, τ) - bipolar valued neutrosophic subset in \mathcal{S} , the strongest (ξ, τ) - bipolar valued neutrosophic relation on \mathcal{S} , that is a (ξ, τ) - bipolar valued neutrosophic relation on K is O such that

$$\left\{ \begin{array}{l} \left(\begin{array}{l} \max\{\vartheta_O^{T+}(m, n), \xi^+\} = \min\{\vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+\}, \\ \min\{\vartheta_O^{T-}(m, n), \xi^-\} = \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\} \end{array} \right) \\ \left(\begin{array}{l} \max\{\vartheta_O^{I+}(m, n), \xi^+\} = \min\{\vartheta_K^{I+}(m), \vartheta_K^{I+}(n), \tau^+\}, \\ \min\{\vartheta_O^{I-}(m, n), \xi^-\} = \max\{\vartheta_K^{I-}(m), \vartheta_K^{I-}(n), \tau^-\} \end{array} \right) \\ \left(\begin{array}{l} \min\{\vartheta_O^{F+}(m, n), \xi^+\} = \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\}, \\ \max\{\vartheta_O^{F-}(m, n), \xi^-\} = \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\} \end{array} \right) \end{array} \right\}.$$

Theorem 4.7. *Let K be any (ξ, τ) – BVNSBS of \mathcal{S} and O be the strongest (ξ, τ) - bipolar valued neutrosophic relation of \mathcal{S} . Then K is a (ξ, τ) – BVNSBS of \mathcal{S} if and only if O is a (ξ, τ) – BVNSBS of $\mathcal{S} \times \mathcal{S}$.*

Theorem 4.8. *Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. The homomorphic image of (ξ, τ) – BVNSBS of \mathcal{S}_1 is a (ξ, τ) – BVNSBS of \mathcal{S}_2 .*

Proof. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any homomorphism. Then $\Lambda(m \vee_1 n) = \Lambda(m) \sqcup_1 \Lambda(n), \Lambda(m \vee_2 n) = \Lambda(m) \sqcup_2 \Lambda(n)$ and $\Lambda(m \vee_3 n) = \Lambda(m) \sqcup_3 \Lambda(n)$ for all $m, n \in \mathcal{S}_1$. Let $O = \Lambda(K)$, K is any (ξ, τ) -BVNSBS of \mathcal{S}_1 . Let $\Lambda(m), \Lambda(n) \in \mathcal{S}_2$. Let $m \in \Lambda^{-1}(\Lambda(m))$ and $n \in \Lambda^{-1}(\Lambda(n))$ be such that $\vartheta_K^{T+}(m) = \sup_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{T+}(z), \vartheta_K^{T+}(n) = \sup_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{T+}(z), \vartheta_K^{T-}(m) = \inf_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{T-}(z)$ and $\vartheta_K^{T-}(n) = \inf_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{T-}(z)$. Now,

$$\begin{aligned} \max \left[\vartheta_O^{T+}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^+ \right] &= \max \left[\sup_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{T+}(z'), \xi^+ \right] \\ &= \max \left[\sup_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{T+}(z'), \xi^+ \right] \\ &= \max \left[\vartheta_K^{T+}(m \vee_1 n), \xi^+ \right] \\ &\geq \min \left\{ \vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+ \right\} \\ &= \min \left\{ \vartheta_O^{T+} \Lambda(m), \vartheta_O^{T+} \Lambda(n), \tau^+ \right\}. \end{aligned}$$

Similarly, $\min [\vartheta_O^{T-}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^-] \leq \max \{ \vartheta_O^{T-} \Lambda(m), \vartheta_O^{T-} \Lambda(n), \tau^- \}$.

Let $\Lambda(m), \Lambda(n) \in \mathcal{S}_2$. Let $m \in \Lambda^{-1}(\Lambda(m))$ and $n \in \Lambda^{-1}(\Lambda(n))$ be such that $\vartheta_K^{I+}(m) = \sup_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{I+}(z)$ and $\vartheta_K^{I+}(n) = \sup_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{I+}(z)$, $\vartheta_K^{I-}(m) = \inf_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{I-}(z)$ and $\vartheta_K^{I-}(n) = \inf_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{I-}(z)$. Now,

$$\begin{aligned} \max [\vartheta_O^{I+}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^+] &= \max \left[\sup_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{I+}(z'), \xi^+ \right] \\ &= \max \left[\sup_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{I+}(z'), \xi^+ \right] \\ &= \max [\vartheta_K^{I+}(m \vee_1 n), \xi^+] \\ &\geq \min \left\{ \frac{\vartheta_K^{I+}(m) + \vartheta_K^{I+}(n)}{2}, \tau^+ \right\} \\ &= \min \left\{ \frac{\vartheta_O^{I+} \Lambda(m) + \vartheta_O^{I+} \Lambda(n)}{2}, \tau^+ \right\} \end{aligned}$$

Similarly, $\min [\vartheta_O^{I-}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^-] \leq \max \left\{ \frac{\vartheta_O^{I-} \Lambda(m) + \vartheta_O^{I-} \Lambda(n)}{2}, \tau^- \right\}$.

Let $m \in \Lambda^{-1}(\Lambda(m))$ and $n \in \Lambda^{-1}(\Lambda(n))$ be such that $\vartheta_K^{F+}(m) = \inf_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{F+}(z)$, $\vartheta_K^{F+}(n) = \inf_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{F+}(z)$, $\vartheta_K^{F-}(m) = \sup_{z \in \Lambda^{-1}(\Lambda(m))} \vartheta_K^{F-}(z)$ and $\vartheta_K^{F-}(n) = \sup_{z \in \Lambda^{-1}(\Lambda(n))} \vartheta_K^{F-}(z)$. Now,

$$\begin{aligned} \min [\vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^+] &= \min \left[\inf_{z' \in \Lambda^{-1}(\Lambda(m) \sqcup_1 \Lambda(n))} \vartheta_K^{F+}(z'), \xi^+ \right] \\ &= \min \left[\inf_{z' \in \Lambda^{-1}(\Lambda(m \vee_1 n))} \vartheta_K^{F+}(z'), \xi^+ \right] \\ &= \min [\vartheta_K^{F+}(m \vee_1 n), \xi^+] \\ &\leq \max \{ \vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+ \} \\ &= \max \{ \vartheta_O^{F+} \Lambda(m), \vartheta_O^{F+} \Lambda(n), \tau^+ \}. \end{aligned}$$

Similarly, $\max [\vartheta_O^{F-}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^-] \geq \min \{ \vartheta_O^{F-} \Lambda(m), \vartheta_O^{F-} \Lambda(n), \tau^- \}$. In the same way, prove the other two operations. Hence O is a (ξ, τ) -BVNSBS of \mathcal{S}_2 .

Theorem 4.9. Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings. The homomorphic preimage of (ξ, τ) -BVNSBS of \mathcal{S}_2 is a (ξ, τ) -BVNSBS of \mathcal{S}_1 .

Proof. Let $\Lambda : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be any homomorphism. Then $\Lambda(m \vee_1 n) = \Lambda(m) \sqcup_1 \Lambda(n)$, $\Lambda(m \vee_2 n) = \Lambda(m) \sqcup_2 \Lambda(n)$ and $\Lambda(m \vee_3 n) = \Lambda(m) \sqcup_3 \Lambda(n)$ for all $m, n \in \mathcal{S}_1$. Let $O = \Lambda(K)$, where O is any (ξ, τ) -BVNSBS of \mathcal{S}_2 . Let $m, n \in \mathcal{S}_1$. Then $\max \{ \vartheta_K^{T+}(m \vee_1 n), \xi^+ \} = \max \{ \vartheta_O^{T+}(\Lambda(m \vee_1 n)), \xi^+ \} = \max \{ \vartheta_O^{T+}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^+ \} \geq \min \{ \vartheta_O^{T+} \Lambda(m), \vartheta_O^{T+} \Lambda(n), \tau^+ \} = \min \{ \vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+ \}$. Thus, $\max \{ \vartheta_K^{T+}(m \vee_1 n), \xi^+ \} \geq \min \{ \vartheta_K^{T+}(m), \vartheta_K^{T+}(n), \tau^+ \}$. Also,

$\min\{\vartheta_K^{T-}(m \vee_1 n), \xi^-\} = \min\{\vartheta_O^{T-}(\Lambda(m \vee_1 n)), \xi^-\} = \min\{\vartheta_O^{T-}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^-\} \leq$
 $\max\{\vartheta_O^{T-}\Lambda(m), \vartheta_O^{T-}\Lambda(n), \tau^-\} = \max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\}$. Thus, $\min\{\vartheta_K^{T-}(m \vee_1 n), \xi^-\} \leq$
 $\max\{\vartheta_K^{T-}(m), \vartheta_K^{T-}(n), \tau^-\}$. Now, $\max\{\vartheta_K^{I+}(m \vee_1 n), \xi^+\} = \max\{\vartheta_O^{I+}(\Lambda(m \vee_1 n)), \xi^+\} =$
 $\max\{\vartheta_O^{I+}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^+\} \geq \min\{\vartheta_O^{I+}\Lambda(m), \vartheta_O^{I+}\Lambda(n), \tau^+\} = \min\{\vartheta_K^{I+}(m), \vartheta_K^{I+}(n), \tau^+\}$.
 Thus, $\max\{\vartheta_K^{I+}(m \vee_1 n), \xi^+\} \geq \min\{\vartheta_K^{I+}(m), \vartheta_K^{I+}(n), \tau^+\}$. Also, $\min\{\vartheta_K^{I-}(m \vee_1 n), \xi^-\} =$
 $\min\{\vartheta_O^{I-}(\Lambda(m \vee_1 n)), \xi^-\} = \min\{\vartheta_O^{I-}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^-\} \leq \max\{\vartheta_O^{I-}\Lambda(m), \vartheta_O^{I-}\Lambda(n), \tau^-\} =$
 $\max\{\vartheta_K^{I-}(m), \vartheta_K^{I-}(n), \tau^-\}$. Thus, $\min\{\vartheta_K^{I-}(m \vee_1 n), \xi^-\} \leq \max\{\vartheta_K^{I-}(m), \vartheta_K^{I-}(n), \tau^-\}$. Now,
 $\min\{\vartheta_K^{F+}(m \vee_1 n), \xi^+\} = \min\{\vartheta_O^{F+}(\Lambda(m \vee_1 n)), \xi^+\} = \min\{\vartheta_O^{F+}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^+\} \leq$
 $\max\{\vartheta_O^{F+}\Lambda(m), \vartheta_O^{F+}\Lambda(n), \tau^+\} = \max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\}$. Thus, $\min\{\vartheta_K^{F+}(m \vee_1 n), \xi^+\} \leq$
 $\max\{\vartheta_K^{F+}(m), \vartheta_K^{F+}(n), \tau^+\}$. Also, $\max\{\vartheta_K^{F-}(m \vee_1 n), \xi^-\} = \max\{\vartheta_O^{F-}(\Lambda(m \vee_1 n)), \xi^-\} =$
 $\max\{\vartheta_O^{F-}(\Lambda(m) \sqcup_1 \Lambda(n)), \xi^-\} \geq \min\{\vartheta_O^{F-}\Lambda(m), \vartheta_O^{F-}\Lambda(n), \tau^-\} = \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\}$.
 Thus, $\max\{\vartheta_K^{F-}(m \vee_1 n), \xi^-\} \geq \min\{\vartheta_K^{F-}(m), \vartheta_K^{F-}(n), \tau^-\}$. In the same way, prove the other
 two operations, hence K is a (ξ, τ) -BVNSBS of \mathcal{S}_1 .

5. (ξ, τ) -Bipolar Valued Neutrosophic Normal Subbisemiring

In this section, we interact the theory for (ξ, τ) -bipolar valued neutrosophic normal subbisemiring. Here *BVNNSBS* stands for bipolar valued neutrosophic normal subbisemiring.

Definition 5.1. Let K be any bipolar valued neutrosophic subset of \mathcal{S} is said to be a *BVNNSBS* of \mathcal{S} if it satisfies the following conditions:

$$\left\{ \begin{array}{l} \left(\begin{array}{l} \vartheta_K^{T+}(m \uplus_1 n) = \vartheta_K^{T+}(n \uplus_1 m), \\ \vartheta_K^{T-}(m \uplus_1 n) = \vartheta_K^{T-}(n \uplus_1 m) \end{array} \right) \\ \left(\begin{array}{l} \vartheta_K^{T+}(m \uplus_2 n) = \vartheta_K^{T+}(n \uplus_2 m), \\ \vartheta_K^{T-}(m \uplus_2 n) = \vartheta_K^{T-}(n \uplus_2 m) \end{array} \right) \\ \left(\begin{array}{l} \vartheta_K^{T+}(m \uplus_3 n) = \vartheta_K^{T+}(n \uplus_3 m), \\ \vartheta_K^{T-}(m \uplus_3 n) = \vartheta_K^{T-}(n \uplus_3 m) \end{array} \right) \end{array} \right\} \left\{ \begin{array}{l} \left(\begin{array}{l} \vartheta_K^{I+}(m \uplus_1 n) = \vartheta_K^{I+}(n \uplus_1 m), \\ \vartheta_K^{I-}(m \uplus_1 n) = \vartheta_K^{I-}(n \uplus_1 m) \end{array} \right) \\ \text{OR} \\ \left(\begin{array}{l} \vartheta_K^{I+}(m \uplus_2 n) = \vartheta_K^{I+}(n \uplus_2 m), \\ \vartheta_K^{I-}(m \uplus_2 n) = \vartheta_K^{I-}(n \uplus_2 m) \end{array} \right) \\ \text{OR} \\ \left(\begin{array}{l} \vartheta_K^{I+}(m \uplus_3 n) = \vartheta_K^{I+}(n \uplus_3 m), \\ \vartheta_K^{I-}(m \uplus_3 n) = \vartheta_K^{I-}(n \uplus_3 m) \end{array} \right) \end{array} \right\} \\ \\ \left\{ \begin{array}{l} \left(\begin{array}{l} \vartheta_K^{F+}(m \uplus_1 n) = \vartheta_K^{F+}(n \uplus_1 m), \\ \vartheta_K^{F-}(m \uplus_1 n) = \vartheta_K^{F-}(n \uplus_1 m) \end{array} \right) \\ \left(\begin{array}{l} \vartheta_K^{F+}(m \uplus_2 n) = \vartheta_K^{F+}(n \uplus_2 m), \\ \vartheta_K^{F-}(m \uplus_2 n) = \vartheta_K^{F-}(n \uplus_2 m) \end{array} \right) \\ \left(\begin{array}{l} \vartheta_K^{F+}(m \uplus_3 n) = \vartheta_K^{F+}(n \uplus_3 m), \\ \vartheta_K^{F-}(m \uplus_3 n) = \vartheta_K^{F-}(n \uplus_3 m) \end{array} \right) \end{array} \right\}$$

for all $m, n \in \mathcal{S}$.

Theorem 5.2. (a) *The intersection of a family of BVNNSBS^s of \mathcal{S} is a BVNNSBS of \mathcal{S} .*
 (b) *The intersection of a family of (ξ, τ) – BVNNSBS^s of \mathcal{S} is a (ξ, τ) – BVNNSBS of \mathcal{S} .*

Proof. Proof follows from Theorem 3.3 and Theorem 4.3.

Theorem 5.3. (a) *If K_1, K_2, \dots, K_n are the family of BVNNSBS^s of $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_n$ respectively, then $K_1 \times K_2 \times \dots \times K_n$ is a BVNNSBS of $\mathcal{S}_1 \times \mathcal{S}_2 \times \dots \times \mathcal{S}_n$.*
 (b) *If K_1, K_2, \dots, K_n are the family of (ξ, τ) – BVNNSBS^s of $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_n$ respectively, then $K_1 \times K_2 \times \dots \times K_n$ is a (ξ, τ) – BVNNSBS of $\mathcal{S}_1 \times \mathcal{S}_2 \times \dots \times \mathcal{S}_n$.*

Proof. Proof follows from Theorem 3.4 and Theorem 4.4.

Theorem 5.4. (a) *Let K be any BVNNSBS of \mathcal{S} and O be the strongest bipolar valued neutrosophic relation of \mathcal{S} . Then K is a BVNNSBS of \mathcal{S} if and only if O is a BVNNSBS of $\mathcal{S} \times \mathcal{S}$.*
 (b) *Let K be any (ξ, τ) – BVNNSBS of \mathcal{S} and O be the strongest (ξ, τ) bipolar valued neutrosophic relation of \mathcal{S} . Then K is a (ξ, τ) – BVNNSBS of \mathcal{S} if and only if O is a (ξ, τ) – BVNNSBS of $\mathcal{S} \times \mathcal{S}$.*

Proof. Proof follows from Theorem 3.7.

Theorem 5.5. *Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings.*
 (a) *The homomorphic image of any BVNNSBS of \mathcal{S}_1 is a BVNNSBS of \mathcal{S}_2 .*
 (b) *The homomorphic image of any (ξ, τ) – BVNNSBS of \mathcal{S}_1 is a (ξ, τ) – BVNNSBS of \mathcal{S}_2 .*

Proof. Proof follows from Theorem 3.12 and Theorem 4.8.

Theorem 5.6. *Let $(\mathcal{S}_1, \vee_1, \vee_2, \vee_3)$ and $(\mathcal{S}_2, \sqcup_1, \sqcup_2, \sqcup_3)$ be any two bisemirings.*
 (a) *The homomorphic preimage of any BVNNSBS of \mathcal{S}_2 is a BVNNSBS of \mathcal{S}_1 .*
 (b) *The homomorphic preimage of any (ξ, τ) – BVNNSBS of \mathcal{S}_2 is a (ξ, τ) – BVNNSBS of \mathcal{S}_1 .*

Proof. Proof follows from Theorem 3.13 and Theorem 4.9.

Acknowledgments: The authors would like to thank the Editors-in-Chief and the anonymous reviewers for their constructive suggestions and helpful comments that have led to the improvement in the quality of the paper.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Zadeh, L.A. Fuzzy sets. *Information and Control* 1965, 8(3), 338-353.
2. Pawlak, Z. Rough sets. *International Journal of Computer and Information Sciences*, 1982, 11(5), 341-356.
3. Gau, W.L.; Buehrer, D.J. Vague sets. *IEEE Transactions on Systems, Man, and Cybernetics*, 1993, 23(2), 610-614.
4. Gorzalczany, M.B. A method of inference in approximate reasoning based on interval-valued fuzzy sets. *Fuzzy Sets and Systems*, 1987, 21(1), 1-17.
5. Molodtsov, D. Soft set theory- first results. *Computers & Mathematics with Applications*, 1999, 37(4-5), 19-31.
6. Atanassov, K.T. Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 1986, 20(1), 87-96.
7. Zhang, W.R. Bipolar fuzzy sets and relations. A computational framework for cognitive modelling and multiagent decision analysis. *The Industrial Fuzzy Control and Intelligent, Proceedings of the First International Joint Conference of The North American Fuzzy Information Processing Society Biannual Conference*, San Antonio, TX, USA, 1994, 18-21.
8. Zhang, W.R. (Yin) (Yang) bipolar fuzzy sets. *IEEE World Congress on Computational Intelligence*, 1998 *IEEE International Conference on Fuzzy Systems Proceedings*, Anchorage, AK, USA, 4-9, May 1998, 1, 835-840.
9. Lee, K. Bipolar-valued fuzzy sets and their operations. *Proceedings of the International Conference on Intelligent Technologies*, Bangkok, Thailand, 2000.
10. Lee, K.M. Comparison of Interval-valued fuzzy sets, Intuitionistic fuzzy sets and bipolar-valued fuzzy sets. *Journal of Korean Institute of Intelligent Systems*, 2004, 14(2), 125-129.
11. Smarandache, F. A unifying field in logics, neutrosophy: Neutrosophic probability, set and logic; American Research Press, Detroit, USA, 1999.
12. Smarandache, F. Neutrosophic set a generalization of the intuitionistic fuzzy set. *International Journal of Pure and Applied Mathematics*, 2005, 24(3), 287.
13. Wang, H.; Smarandache, F.; Zhang, Y.; Sunderraman, R. Interval neutrosophic sets and logic theory and applications in computing. *Hexis*, Arizona, USA, 2005.
14. Deli, I., Ali, M.; Smarandache, F. Bipolar neutrosophic sets and their application based on multi-criteria decision-making problems. In *2015 International Conference on Advanced Mechatronic Systems*, Beijing, China, 22-24 August 2015.
15. Ye, J. A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets. *Journal of Intelligent & Fuzzy Systems* 2014, 26(5), 2459-2466.
16. Peng, J.; Wang, J.; Wu, X.; Wang, J.; Chen, X. Multi-valued neutrosophic sets and power aggregation operators with their applications in multi-criteria group decision-making problems. *International Journal of Computational Intelligence Systems*, 2015, 8(2), 345-363.
17. Das, S.; Roy, B.K.; Kar, M.B.; Kar, S.; Pamucar, D. Neutrosophic fuzzy set and its application in decision making. *Journal of Ambient Intelligence and Humanized Computing*, 2020, 11, 5017-5029.
18. Vandiver, H.S. Note on a simple type of algebra in which the cancellation law of addition does not hold. *Bulletin of the American Mathematical Society*, 1934, 40(12), 914-920.
19. Golan, J.S. *Semirings and their applications*; Springer: Dordrecht, Netherlands, 1999.
20. Glazek, K. *A guide to the literature on semirings and their applications in mathematics and information sciences*. Springer, Dordrecht, Netherlands, 2002.
21. Ahsan, J.; Saifullah, K.; Farid Khan, M. Fuzzy semirings. *Fuzzy Sets and Systems*, 1993, 60(3), 309-320.
22. Fang, F.; Jun, Y.B.; Zhao, X. Soft semirings. *Computers & Mathematics with Applications*, 2008, 56(10), 2621-2628.

23. Yousafzai, F.; Khalaf, M.M.; Khan, M.U.I.; Borumand Saeid, A.; Iqbal, Q. Some studies in fuzzy non-associative semigroups. *Journal of Intelligent & Fuzzy Systems*, 2017, 32(3), 1917-1930.
24. Mocko, J. Semiring-valued fuzzy sets and F-transform. *Mathematics*, 2021, 9(23), 1-24.
25. Palanikumar, M.; Arulmozhi, K.; On vague subbisemirings of bisemirings, *Bulletin of the International Mathematical Virtual Institute*, 2022, 11(3), 419-428.
26. Palanikumar, M.; Arulmozhi, K. (α, β) -neutrosophic subbisemiring of bisemiring, *Neutrosophic Sets and Systems*, 2022, 48, 368-385.
27. Palanikumar, M.; Arulmozhi, K. On intuitionistic fuzzy normal subbisemiring of bisemiring, 2021, *Nonlinear Studies*, 2021, 28(3), 717-721.
28. Palanikumar, M.; Arulmozhi, K. On various tri-ideals in ternary semirings. *Bulletin of the International Mathematical Virtual Institute*, 2021, 11(1), 79-90.
29. Palanikumar, M.; Iampan, A.; Lejo J. Manavalan M-Bi-base generator of ordered gamma-semigroups, *ICIC Express Letters Part-B*, 2022 13(8), 795-802.
30. Palanikumar, M.; Arulmozhi, K. On New Ways of various ideals in ternary semigroups, *Matrix Science Mathematic*, 2020, 4(1), 06-09.
31. Palanikumar, M.; Arulmozhi, K.; Jana, C. Multiple attribute decision-making approach for Pythagorean neutrosophic normal interval-valued aggregation operators, *Comp. Appl. Math.* 2022, 41(90), 1-27.
32. Palanikumar, M.; Iampan, A.; Spherical fermatean interval valued fuzzy soft set based on multi criteria group decision making, *International Journal of Innovative Computing, Information and Control*, 2022, 18(2), 607-619.
33. Palanikumar, M.; Iampan, A. Novel approach to decision making based on type-II generalized fermatean bipolar fuzzy soft sets, *International Journal of Innovative Computing, Information and Control*, 2022, 18(3), 769-781.
34. Jun, Y.B.; Song, S.Z. Subalgebras and closed ideals of BCH-algebras based on bipolar-valued fuzzy sets. *Scientiae Mathematicae Japonicae Online*, 2008, 427-437.
35. Lee, K.J. Bipolar fuzzy subalgebras and bipolar fuzzy ideals of BCK/BCI-algebras. *Bulletin of the Malaysian Mathematical Sciences Society*, 2009, 32(3), 361-373.
36. Muhiuddin, G.; Harizavi, H.; Jun, Y.B. Bipolar-valued fuzzy soft hyper BCK ideals in hyper BCK algebras. *Discrete Mathematics, Algorithms and Applications*, 2020, 12(2), 2050018.
37. Jana, C.; Senapati, T.; Shum, P.C.; Pal, M. Bipolar fuzzy soft subalgebras and ideals of BCK/BCI-algebras based on bipolar fuzzy points. *Journal of Intelligent & Fuzzy Systems*, 2019, 37(2), 2785-2795.
38. Zararsz, Z.; Riaz, M. Bipolar fuzzy metric spaces with application. *Comp. Appl. Math.* 2022, 41(49).
39. Selvachandran, G.; Salleh, A.R. Algebraic Hyperstructures of vague soft sets associated with hyperrings and hyperideals. *The Scientific World Journal*, 2015, 2015, 780121.
40. Jun, Y.B.; Kim, H.S.; Lee, K.J. Bipolar fuzzy translations in BCK/BCI-algebras. *Journal of the Chungcheong Mathematical Society*, 2009, 22(3), 399-408.
41. Jun, Y.B.; Park, C.H. Filters of BCH-algebras based on bipolar-valued fuzzy sets. *International Mathematical Forum*, 2009, 4(13), 631-643.
42. Sen, M.K.; Ghosh, S.; Ghosh, S. An introduction to bisemirings. *Southeast Asian Bulletin of Mathematics*, 2004, 28(3), 547-559.
43. Hussain, F. Factor bisemirings. *Italian Journal of Pure and Applied Mathematics*, 2015, 34, 81-88.
44. Hussain, F.; Hashim, R.M.; Khan, A.; Naeem, M. Generalization of bisemirings. *International Journal of Computer Science and Information Security*, 2016, 14(9), 275-289.

Received: August 05, 2022. Accepted: January 10, 2023

M.Palanikumar, K.Arulmozhi, Ganeshsree Selvachandran and Sher Lyn Tan, New approach to bisemiring theory via the bipolar valued neutrosophic normal sets