

7-29-2023

On Symbolic Plithogenic Algebraic Structures and Hyper Structures

A. A.A. Agboola

M. A. Ibrahim

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Agboola, A. A.A. and M. A. Ibrahim. "On Symbolic Plithogenic Algebraic Structures and Hyper Structures." *Neutrosophic Sets and Systems* 56, 1 (2023). https://digitalrepository.unm.edu/nss_journal/vol56/iss1/17

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



On Symbolic Plithogenic Algebraic Structures and Hyper Structures

A.A.A. Agboola¹ and M.A. Ibrahim²

¹Department of Mathematics, Federal University of Agriculture, PMB 2240, Abeokuta, Nigeria;
agboolaaaa@funaab.edu.ng

²Department of Mathematics, Auburn University, Auburn, AL 36849, USA;
mai0015@auburn.edu

Correspondence: agboolaaaa@funaab.edu.ng

Abstract. The objective of this paper is to study symbolic Plithogenic algebraic structures and hyper structures. In particular, we study symbolic Plithogenic group, symbolic Plithogenic ring, symbolic Plithogenic hypergroup and symbolic Plithogenic canonical hypergroup.

Keywords: Plithogeny; Plithogenic; Plithogenic Set; Fuzzy Set; Intuitionistic Fuzzy Set; Neutrosophy; Neutrosophic Set; Plithogenic Group; Plithogenic Ring; Plithogenic Hypergroup; Plithogenic canonical Hypergroup.

1. Introduction

The concepts of Plithogeny, Plithogenic logic/set, Plithogenic probability and Plithogenic statistics were introduced by Smarandache in [26]. Plithogenic set/logic is an extension of the classical logic/set, fuzzy logic/set of Zadeh [37], intuitionistic fuzzy logic/set of Atanassov [11], neutrosophic logic/set of Smarandache [30] and quadruple neutrosophic logic/set of Smarandache [29]. Smarandache in [23], [25] and [28] introduced and studied symbolic Plithogenic algebraic structures and hyper structures. In [22], Merkepsi and Abobala studied symbolic 2-Plithogenic rings, in [10], Al-Basheer et al. studied symbolic 3-Plithogenic rings and in [17], Gayen et al. studied Plithogenic Hypersoft Subgroup. Also in [32], Taffach and Hatip studied Symbolic

2-Plithogenic Number Theory And Algebraic Equations, in [33], Taffach and Othman studied Symbolic 2-Plithogenic Modules over Symbolic 2-Plithogenic Rings and in [34], Taffach studied Symbolic 2-Plithogenic Vector Spaces. In [18], [24] and [27], applications of Plithogenic set/logic were presented. In the present paper, we study symbolic Plithogenic algebraic structures and hyper structures. In particular, we study symbolic Plithogenic group, symbolic Plithogenic ring, symbolic Plithogenic hypergroup and symbolic Plithogenic canonical hypergroup and present their basic properties.

2. Symbolic Plithogenic Set

A symbolic Plithogenic set SPX is defined by

$$SPX = \{(a, a_1P_1, a_2P_2, a_3P_3, \dots, a_nP_n) : a, a_i \in \mathbb{R} \text{ or } \mathbb{C} \text{ or any AlgebraicStructure}\} \quad (1)$$

where P_i 's are the Plithogenic parameters/variables. a is called the non-Plithogenic part of SPX , a_iP_i is called the Plithogenic part of SPX and a_i 's are called the coefficients of P_i s where $i = 1, 2, 3, \dots, n$. For a positive integer k , P_i has the following properties :

$$P_i^k = P_i, \quad \forall i \text{ and } k \geq 2, \quad (2)$$

$$kP_i = P_i + P_i + P_i + \dots + P_i \quad [k \text{ summand}] \quad \forall i, \quad (3)$$

$$0P_i = 0 \quad \forall i, \quad (4)$$

$$P_i^{-1} = \frac{1}{P_i} \text{ does not exist } \quad \forall i. \quad (5)$$

when $n = 1$, equation (1) reduces to

$$SPX = \{(a, a_1P_1) : a, a_1 \in \mathbb{R} \text{ or } \mathbb{C} \text{ or any AlgebraicStructure}\} \quad (6)$$

and SPX becomes the usual Neutrosophic set with $P_1 = I$.

When $n = 3$, equation (1) reduces to

$$SPX = \{(a, a_1P_1, a_2P_2, a_3P_3) : a, x_i \in \mathbb{R} \text{ or } \mathbb{C} \text{ or any AlgebraicStructure}\} \quad (7)$$

and SPX becomes the usual Neutrosophic Quadruple set with $P_1 = T$, $P_2 = I$ and $P_3 = F$.

When $n = 2$, equation (1) reduces to

$$SPX = \{(a, a_1P_1, a_2P_2) : a, a_i \in \mathbb{R} \text{ or } \mathbb{C} \text{ or any AlgebraicStructure}\} \quad (8)$$

which is called symbolic 2-Plithogenic set.

3. Symbolic Plithogenic Algebraic Structure

All the symbolic Plithogenic sets to be considered in this section and the section after will be symbolic 2-Plithogenic sets of the form given by equation (8) and we are going to assume throughout the prevalence order $P_1 > P_2$ so that

$$P_1P_1 = P_{\min\{1,1\}} = P_1, \quad (9)$$

$$P_2P_2 = P_{\min\{2,2\}} = P_2, \quad (10)$$

$$P_1P_2 = P_2P_1 = P_{\min\{1,2\}} = P_1. \quad (11)$$

Definition 3.1. Let $+$, $-$ and \cdot be the usual arithmetic operations of addition, subtraction and multiplication of numbers respectively and let k be a nonzero scalar. If $x = (a, a_1P_1, a_2P_2)$ and $y = (b, b_1P_1, b_2P_2)$ are arbitrary elements of the symbolic Plithogenic set SPX where $a, b, a_i, b_i \in \mathbb{R}$ or \mathbb{C} , then:

$$x \pm y = (a \pm b, (a_1 \pm b_1)P_1, (a_2 \pm b_2)P_2), \quad (12)$$

$$kx = (ka, ka_1P_1, ka_2P_2), \quad (13)$$

$$x \cdot y = (ab, (ab_1 + a_1b + a_1b_1 + a_1b_2 + a_2b_1)P_1, (ab_2 + a_2b + a_2b_2)P_2). \quad (14)$$

When $k = 0$, then we have

$$0x = (0a, 0a_1P_1, 0a_2P_2) = (0, 0P_1, 0P_2) = (0, 0, 0). \quad (15)$$

Notation 3.2. In what follows next, we will use the symbols $SP\mathbb{N}$, $SP\mathbb{Z}$, $SP\mathbb{Q}$, $SP\mathbb{R}$ and $SP\mathbb{C}$ to denote the Plithogenic sets of natural, integer, rational, real and complex numbers respectively.

Example 3.3. $(SP\mathbb{Q}, \cdot)$, $(SP\mathbb{R}, \cdot)$ and $(SP\mathbb{C}, \cdot)$ are symbolic Plithogenic groups.

Definition 3.4. Let $(X, *)$ be any algebraic structure and let SPX be the corresponding symbolic Plithogenic set. The couple $(SPX, *)$ is called a symbolic Plithogenic algebraic structure. SPX will be named according to the name of the underlying algebraic structure X . For instance if X is a group, SPX will be called a symbolic Plithogenic group, if X is a ring, SPX will be called a symbolic Plithogenic ring, if X is a hypergroup, SPX will be called a symbolic Plithogenic hypergroup and so on .

Theorem 3.5. Let $(G, *)$ be a group and let SPG be the corresponding symbolic Plithogenic group. Then:

- (i) $G \subset SPG$.
- (ii) $(SPG, *)$ is a semigroup.
- (iii) $(SPG, *)$ is not a group.

Proof. (i) This follows from the definition of SPG .

(ii) Let $x = (a, a_1P_1, a_2P_2)$, $y = (b, b_1P_1, b_2P_2)$ and $z = (c, c_1P_1, c_2P_2)$ be arbitrary elements of SPG . Then:

$$\begin{aligned} x * y &= (ab, (ab_1 + a_1b + a_1b_1 + a_1b_2 + a_2b_1)P_1, (ab_2 + a_2b + a_2b_2)P_2) \in SPG. \text{ Now,} \\ (x * y) * z &= (abc, (abc_1 + ab_1c + a_1bc + a_1b_1c + a_1b_2c + a_2b_1c + ab_1c_1 + a_1bc_1 + a_1b_1c_1 \\ &\quad + a_1b_2c_1 + a_2b_1c_1 + ab_1c_2 + a_1bc_2 + a_1b_1c_2 + a_1b_2c_2 + a_2b_1c_2 + ab_2c_1 \\ &\quad + a_2bc_1 + a_2b_2c_1)P_1, (abc_2 + ab_2c + a_2bc + a_2b_2c + ab_2c_2 + a_2bc_2 + a_2b_2c_2)P_2) \\ x * (y * z) &= (abc, (abc_1 + ab_1c + ab_1c_1 + ab_1c_2 + ab_2c_1 + a_1bc + a_1bc_1 + a_1b_1c + a_1b_1c_1 \\ &\quad + a_1b_1c_2 + a_1b_2c_1 + a_1bc_2 + a_1b_2c + a_1b_2c_2 + a_2bc_1 + a_2b_1c + a_2b_1c_1 \\ &\quad + a_2b_1c_2 + a_2b_2c_1)P_1, (abc_2 + ab_2c + a_2bc + a_2b_2c + ab_2c_2 + a_2bc_2 + a_2b_2c_2)P_2) \\ &= x * (y * z). \end{aligned}$$

This shows that $(SPG, *)$ is a semigroup.

(iii) Since P_1^{-1} and P_2^{-1} do not exist, it follows that we cannot find x^{-1} , $\forall x \in SPG$. Hence, $(SPG, *)$ is not a group. \square

Remark 3.6. If $(G, +)$ is a group, then the symbolic Plithogenic group $(SPG, +)$ is a group.

Example 3.7. $(SP\mathbb{Z}, +)$, $(SP\mathbb{Q}, +)$, $(SP\mathbb{R}, +)$ and $(SP\mathbb{C}, +)$ are abelian groups.

Theorem 3.8. Every symbolic Plithogenic group (SPG, \cdot) has at least 2 nontrivial idempotent elements.

Proof. Since $P_1P_1 = P_1, P_2P_2 = P_2$ in SPG , the required result follows. \square

Theorem 3.9. Let $(G, *)$ be a finite group of order n . Then $(SPG, *)$ is a finite symbolic Plithogenic group of order n^3 .

Example 3.10. Let \mathbb{Z}_2 be the group of integers modulo 2. Then

$$SP\mathbb{Z}_2 = \{(0, 0, 0), (1, 0, 0), (0, P_1, 0), (0, 0, P_2), (0, P_1, P_2), (1, P_1, 0), (1, 0, P_2), (1, P_1, P_2)\}$$

is a symbolic Plithogenic group of integers modulo 2. The elements $(0, P_1, 0)$, $(0, 0, P_2)$ and $(1, P_1, P_2)$ of $SP\mathbb{Z}_2$ are nontrivial idempotent elements.

Definition 3.11. Let $\phi : SPG \rightarrow SPH$ be a mapping from the symbolic Plithogenic group $(SPG, *)$ to the symbolic Plithogenic group (SPH, \star) . ϕ is called a symbolic Plithogenic group homomorphism if the following conditions hold:

A.A.A. Agboola and M.A. Ibrahim, On Symbolic Plithogenic Algebraic Structures and Hyper Structures

- (i) $\phi(x * y) = \phi(x) \star \phi(y), \forall x, y \in SPG,$
- (ii) $\phi(P_i) = P_i, i = 1, 2.$

The kernel of ϕ denoted by $\text{Ker}\phi$ is defined by

$$\text{Ker}\phi = \{x \in SPG : \phi(x) = \text{identity element of } SPH\}.$$

Example 3.12. Let $(G, +)$ be a group and let $\phi : SPG \times SPG \rightarrow SPG$ be a mapping defined by

$$\phi(a, b) = a \quad \forall (a, b) \in SPG \times SPG.$$

Then ϕ is a symbolic Plithogenic group homomorphism.

If $G = \mathbb{Z}_2$, then

$$\begin{aligned} \text{Ker}\phi = & \{((0, 0, 0), (0, 0, 0)), ((0, 0, 0), (1, 0, 0)), ((0, 0, 0), (0, P_1, 0)), ((0, 0, 0), (0, 0, P_2)), \\ & ((0, 0, 0), (0, P_1, P_2)), ((0, 0, 0), (1, P_1, 0)), ((0, 0, 0), (1, 0, P_2)), ((0, 0, 0), (1, P_1, P_2))\} \end{aligned}$$

which is a subgroup of $SP\mathbb{Z}_2 \times SP\mathbb{Z}_2$.

Example 3.13. Let $G = \mathbb{Z}$, let SPG be the corresponding symbolic Plithogenic group of integers and let $G(I)$ be the neutrosophic group of integers. If $\phi : SPG \rightarrow G(I)$ is a mapping defined by

$$\phi(x) = (a, (b + c)I), \forall x = (a, bP_1, cP_2) \in SPG,$$

then ϕ is a group homomorphism and $\text{Ker}\phi = \{(0, kP_1, -kP_2) : k \in \mathbb{Z}\}$ which is a subgroup of SPG .

Definition 3.14. Let $(R, +, \cdot)$ be any ring. The triple $(SPR, +, \cdot)$ is called a symbolic Plithogenic ring. If R is commutative with unity, so also is SPR .

Theorem 3.15. Let $(R, +, \cdot)$ be any ring. Then $(SPR, +, \cdot)$ is a ring.

Proof. Using Definition 3.1, it can easily be shown that $(SPR, +)$ is an abelian group and (SPR, \cdot) is a semigroup. Also, for arbitrary $x, y, z \in SPR$, it can be shown that $x(y + z) = xy + xz$ and $(y + z)x = yx + zx$. Hence, $(SPR, +, \cdot)$ is a ring. \square

Theorem 3.16. Every symbolic Plithogenic ring $(SPR, +, \cdot)$ has at least 2 nontrivial idempotent elements.

Theorem 3.17. Let $(R, +, \cdot)$ be a finite ring of order n . Then $(SPR, +, \cdot)$ is a finite symbolic Plithogenic ring of order n^3 .

Example 3.18. Let \mathbb{Z}_2 be the ring of integers modulo 2. Then

$$SP\mathbb{Z}_2 = \{(0, 0, 0), (1, 0, 0), (0, P_1, 0), (0, 0, P_2), (0, P_1, P_2), (1, P_1, 0), (1, 0, P_2), (1, P_1, P_2)\}$$

is a symbolic Plithogenic ring of integers modulo 2.

Lemma 3.19. Let $(SPR, +, \cdot)$ be a symbolic Plithogenic ring and let $x = (a, a_1P_1, a_2P_2)$ and $y = (b, b_1P_1, b_2P_2)$ be any two nonzero elements of SPR .

- (a) x is idempotent if and only if all the following hold:
 - (i) a is idempotent,
 - (ii) $a + a_2$ is idempotent and
 - (iii) $a + a_1 + a_2$ is idempotent.
- (b) x and y are zero divisors if and only if all the following hold:
 - (i) a and b are zero divisors,
 - (ii) $a + a_2$ and $b + b_2$ are zero divisors and
 - (iii) $a + a_1 + a_2$ and $b + b_1 + b_2$ are zero divisors.

Example 3.20. Let $SP\mathbb{Z}_6$ be the symbolic Plithogenic ring of integers modulo 6. Then

- (i) $(1, 3P_1, 3P_2), (1, 5P_1, 3P_2), (3, 5P_1, P_2)$ and $(4, P_1, 5P_2)$ are idempotent elements.
- (ii) $(2, P_1, P_2)$ and $(3, 5P_1, P_2)$ are zero divisors.

Definition 3.21. Let $\phi : SPR \rightarrow SPS$ be a mapping from the symbolic Plithogenic ring $(SPR, +, \cdot)$ to the symbolic Plithogenic ring $(SPS, +, \cdot)$. ϕ is called a symbolic Plithogenic ring homomorphism if the following conditions hold:

- (i) $\phi(x + y) = \phi(x) + \phi(y), \forall x, y \in SPR,$
- (ii) $\phi(xy) = \phi(x)\phi(y), \forall x, y \in SPR,$
- (iii) $\phi(P_i) = P_i, i = 1, 2.$

The kernel of ϕ denoted by $\text{Ker}\phi$ is defined by

$$\text{Ker}\phi = \{x \in SPR : \phi(x) = \text{identity element of } SPS\}.$$

Example 3.22. Let $(R, +, \cdot)$ be a ring and let $\phi : SPR \times SPR \rightarrow SPR$ be a mapping defined by

$$\phi(a, b) = b \quad \forall (a, b) \in SPR \times SPR.$$

Then ϕ is a symbolic Plithogenic ring homomorphism.

If $R = \mathbb{Z}_2$, then

$$\begin{aligned} \text{Ker}\phi = & \{((0, 0, 0), (0, 0, 0)), ((1, 0, 0), (0, 0, 0)), ((0, P_1, 0), (0, 0, 0)), ((0, 0, P_2), (0, 0, 0)), \\ & ((0, P_1, P_2), (0, 0, 0)), ((1, P_1, 0), (0, 0, 0)), ((1, 0, P_2), (0, 0, 0)), ((1, P_1, P_2), (0, 0, 0))\} \end{aligned}$$

which is a subring of $SP\mathbb{Z}_2 \times SP\mathbb{Z}_2$.

Theorem 3.23. Let $\psi : R \rightarrow S$ be a ring homomorphism and let $\phi : SPR \rightarrow SPS$ be a mapping from a symbolic Plithogenic ring SPR into a symbolic Plithogenic ring SPS defined by

$$\phi(x) = (\psi(a), \psi(b)P_1, \psi(c)P_2), \quad \forall x = (a, bP_1, cP_2) \in SPR.$$

Then ϕ is a ring homomorphism.

Proof. Let $x = (a, bP_1, cP_2)$ and $y = (d, eP_1, fP_2)$ be two arbitrary elements in SPR . Then

$$\begin{aligned} x + y &= (a + d, (b + e)P_1, (c + f)P_2), \\ xy &= (ad, (ae + bd + be + bf + ce)P_1, (af + cd + cf)P_2), \\ \phi(x) &= (\psi(a), \psi(b)P_1, \psi(c)P_2), \\ \phi(y) &= (\psi(d), \psi(e)P_1, \psi(f)P_2), \\ \therefore \phi(x + y) &= (\psi(a + d), \psi(b + e)P_1, \psi(c + f)P_2), \\ &= (\psi(a) + \psi(d), \psi(b)P_1 + \psi(e)P_1, \psi(c)P_2 + \psi(f)P_2) \\ &= (\psi(a), \psi(b)P_1, \psi(c)P_2) + (\psi(d), \psi(e)P_1, \psi(f)P_2), \\ &= \phi(x) + \phi(y), \\ \phi(xy) &= (\psi(ad), \psi(ae + bd + be + bf + ce)P_1, \psi(af + cd + cf)P_2), \\ &= (\psi(a)\psi(d), (\psi(a)\psi(e) + \psi(b)\psi(d) + \psi(b)\psi(e) + \psi(b)\psi(f) + \psi(c)\psi(e))P_1, \\ &\quad (\psi(a)\psi(f) + \psi(c)\psi(d) + \psi(c)\psi(f))P_2), \\ &= [(\psi(a), \psi(b)P_1, \psi(c)P_2)][(\psi(d), \psi(e)P_1, \psi(f)P_2)], \\ &= \phi(x)\phi(y). \end{aligned}$$

Accordingly, ϕ is a ring homomorphism. \square

Example 3.24. Let $R = \mathbb{Z}_6$, $S = \mathbb{Z}_2$ and let $\psi : \mathbb{Z}_6 \rightarrow \mathbb{Z}_2$ be a ring homomorphism defined by $\psi(\bar{x}_6) = \bar{x}_2$. Let $\phi : SP\mathbb{Z}_6 \rightarrow SP\mathbb{Z}_2$ be a symbolic Plithogenic ring homomorphism defined by

$$\phi((a, bP_1, cP_2)) = (\psi(a), \psi(b)P_1, \psi(c)P_2), \forall (a, bP_1, cP_2) \in SP\mathbb{Z}_6.$$

Then, $\text{Ker}\psi = \{0, 2, 4\}$ and $\text{Ker}\phi = \{(i, jP_1, kP_2) : i, j, k = 0, 2, 4\}$.

4. Symbolic Plithogenic Algebraic Hyper Structure

Definition 4.1. Let H be a nonempty set and $* : H \times H \rightarrow \mathbb{P}^*(H)$ be a hyperoperation. The couple $(H, *)$ is called a hypergroupoid.

For any two nonempty subsets A and B of H and $x \in H$, we define

$$\begin{aligned} A * B &= \bigcup_{a \in A, b \in B} a * b, \\ A * x &= A * \{x\} \text{ and} \\ x * B &= \{x\} * B. \end{aligned}$$

A hypergroupoid $(H, *)$ is called a semihypergroup if $\forall a, b, c \in H$ we have $(a * b) * c = a * (b * c)$, which means that

$$\bigcup_{u \in a * b} u * c = \bigcup_{v \in b * c} a * v.$$

A hypergroupoid $(H, *)$ is called a quasihypergroup if $\forall a \in H$ we have $a * H = H * a = H$. This condition is also called the reproduction axiom.

If a hypergroupoid $(H, *)$ is both a semihypergroup and a quasihypergroup, then it is called a hypergroup.

- Example 4.2.**
- (i) Let H be a nonempty set and let $x * y = H, \forall x, y \in H$. Then $(H, *)$ is a hypergroup called a total hypergroup.
 - (ii) Let $(H, .)$ be a group and let P be a nonempty subset of H . If $x * y = xPy, \forall x, y \in H$, then, $(H, *)$ is a hypergroup called a P -hypergroup.
 - (iii) Let $(H, .)$ be a group. If $x * y = \langle x, y \rangle, \forall x, y \in H$, where $\langle x, y \rangle$ is the subgroup generated by x and y , then $(H, *)$ is a hypergroup.

Definition 4.3. Let $(H, *)$ and (K, \circ) be two hypergroups. A mapping $\phi : H \rightarrow K$, is called:

- (i) an inclusion homomorphism if $\phi(x * y) \subseteq \phi(x) \circ \phi(y), \forall x, y \in H$;
- (ii) a good homomorphism if $\phi(x * y) = \phi(x) \circ \phi(y), \forall x, y \in H$.

Definition 4.4. Let H be a nonempty set and let $+$ be a hyperoperation on H . The couple $(H, +)$ is called a canonical hypergroup if the following conditions hold:

- (i) $x + y = y + x, \forall x, y \in H$,
- (ii) $x + (y + z) = (x + y) + z, \forall x, y, z \in H$,
- (iii) there exists a neutral element $0 \in H$ such that $x + 0 = \{x\} = 0 + x, \forall x \in H$,
- (iv) for every $x \in H$, there exists a unique element $-x \in H$ such that $0 \in x + (-x) \cap (-x) + x$,
- (v) $z \in x + y$ implies $y \in -x + z$ and $x \in z - y, \forall x, y, z \in H$.

Example 4.5. Let $H = \{0, a, b, c\}$ be a set and let $+$ be a hyperoperation on H defined in the Cayley table below.

$+$	0	a	b	c
0	0	a	b	c
a	a	$\{0, b\}$	$\{a, c\}$	b
b	b	$\{a, c\}$	$\{0, b\}$	a
c	c	b	a	0

Then $(H, +)$ is a canonical hypergroup.

Definition 4.6. Let $(H, +)$ and $(K, +)$ be two canonical hypergroups. A mapping $\phi : H \rightarrow K$ is called:

- (a) a homomorphism if:
 - (i) $\phi(x + y) \subseteq \phi(x) + \phi(y), \forall x, y \in H$ and
 - (ii) $\phi(0) = 0$.
- (b) a good or strong homomorphism if:
 - (i) $\phi(x + y) = \phi(x) + \phi(y), \forall x, y \in H$ and
 - (ii) $\phi(0) = 0$.

The kernel of ϕ denoted by $\text{Ker}\phi$ is the set $\{x \in H : \phi(x) = 0\}$.

Definition 4.7. Let $(H, *)$ be any hypergroup. The couple $(SPH, *)$ is called a symbolic Plithogenic hypergroup. If $x = (a, a_1P_1, a_2P_2)$ and $y = (b, b_1P_1, b_2P_2)$ are any two elements of SPH , the composition of x and y in SPH denoted by $x * y$ is defined as

$$x * y = \{(c, c_1P_1, c_2P_2) : c \in a * b, c_1 \in (a * b_1 \cup a_1 * b \cup a_1 * b_1 \cup a_1 * b_2 \cup a_2 * b_1)P_1, c_2 \in (a * b_2 \cup a_2 * b \cup a_2 * b_2)P_2\} \tag{16}$$

- Example 4.8.**
- (i) Let $(H, *)$ be a total hypergroup. Then $(SPH, *)$ is a symbolic Plithogenic total hypergroup.
 - (ii) Let $(H, *)$ be a P-hypergroup. Then $(SPH, *)$ is a symbolic Plithogenic P-hypergroup.

Theorem 4.9. Let $(H, *)$ be a hypergroup and let $(SPH, *)$ be the corresponding symbolic Plithogenic hypergroup. Then:

- (i) $(SPH, *)$ is a semigroup.
- (ii) $(SPH, *)$ generally is not a hypergroup.

Proof. Let $x = (a, a_1P_1, a_2P_2), y = (b, b_1P_1, b_2P_2)$ and $z = (c, c_1P_1, c_2P_2)$ be arbitrary elements of SPH .

(i)

$$\begin{aligned} x * y &= (a, a_1P_1, a_2P_2) * (b, b_1P_1, b_2P_2) \\ &= (a * b, (a * b_1 \cup a_1 * b \cup a_1 * b_1 \cup a_1 * b_2 \cup a_2 * b_1)P_1), \\ &\quad (a * b_2 \cup a_2 * b \cup a_2 * b_2)P_2) \\ &\subset SPH. \end{aligned}$$

This shows that $(SPH, *)$ is a groupoid. Next,

$$\begin{aligned} (x * y) * z &= [(a, a_1P_1, a_2P_2) * (b, b_1P_1, b_2P_2)] * (c, c_1P_1, c_2P_2) \\ &= [(a * b, (a * b_1 \cup a_1 * b \cup a_1 * b_1 \cup a_1 * b_2 \cup a_2 * b_1)P_1), \\ &\quad (a * b_2 \cup a_2 * b \cup a_2 * b_2)P_2)] * (c, c_1P_1, c_2P_2) \\ &= (a * b * c, (a * b * c_1 \cup a * b_1 * c \cup a_1 * b * c \cup a_1 * b_1 * c \cup a_1 * b_2 * c \\ &\quad \cup a_2 * b_1 * c \cup a * b_1 * c_1 \cup a_1 * b * c_1 \cup a_1 * b_1 * c_1 \cup a_1 * b_2 * c_1 \cup a_2 * b_1 * c_1 \\ &\quad \cup a * b_1 * c_2 \cup a_1 * b * c_2 \cup a_1 * b_1 * c_2 \cup a_1 * b_2 * c_2 \cup a_2 * b_1 * c_2 \cup a * b_2 * c_1 \\ &\quad \cup a_2 * b * c_1 \cup a_2 * b_2 * c_1)P_1, (a * b * c_2 \cup a * b_2 * c \cup a_2 * b * c \cup a_2 * b_2 * c \\ &\quad \cup a * b_2 * c_2 \cup a_2 * b * c_2 \cup a_2 * b_2 * c_2)P_2) \\ x * (y * z) &= (a * b * c, (a * b * c_1 \cup a * b_1 * c \cup a * b_1 * c_1 \cup a * b_1 * c_2 \cup a * b_2 * c_1 \cup a_1 * b * c \\ &\quad \cup a_1 * b * c_1 \cup a_1 * b_1 * c \cup a_1 * b_1 * c_1 \cup a_1 * b_1 * c_2 \cup a_1 * b_2 * c_1 \cup a_1 * b * c_2 \\ &\quad \cup a_1 * b_2 * c \cup a_1 * b_2 * c_2 \cup a_2 * b * c_1 \cup a_2 * b_1 * c \cup a_2 * b_1 * c_1 \cup a_2 * b_1 * c_2 \\ &\quad \cup a_2 * b_2 * c_1)P_1, (a * b * c_2 \cup a * b_2 * c \cup a_2 * b * c \cup a_2 * b_2 * c \cup a * b_2 * c_2 \\ &\quad \cup a_2 * b * c_2 \cup a_2 * b_2 * c_2)P_2) \\ &= x * (y * z). \end{aligned}$$

Accordingly, $(SPH, *)$ is a semigroup.

(ii) For all $x = (a, a_1P_1, a_2P_2)$ in SPH , it can be shown that $x * SPH \neq SPH \neq SPH * x$. This shows that reproduction axiom failed to hold in SPH . Hence, $(SPH, *)$ is not a hypergroup. \square

Definition 4.10. Let $(SPH, *)$ and (SPK, \circ) be any two symbolic Plithogenic hypergroups and let $\phi : SPH \rightarrow SPK$ be a mapping from SPH into SPK .

(a) ϕ is called a symbolic Plithogenic hypergroup homomorphism if the following conditions hold:

- (i) $\phi(x * y) \subseteq \phi(x) \circ \phi(y), \forall x, y \in SPH.$
- (ii) $\phi(P_i) = P_i,$ for $i = 1, 2.$

(b) ϕ is called a symbolic Plithogenic good hypergroup homomorphism if the following conditions hold:

- (i) $\phi(x * y) = \phi(x) \circ \phi(y), \forall x, y \in SPH.$
- (ii) $\phi(P_i) = P_i,$ for $i = 1, 2.$

Theorem 4.11. *Let $\psi : (H, *) \rightarrow (K, \circ)$ be a good hypergroup homomorphism from a hypergroup $(H, *)$ into a hypergroup (K, \circ) and let $\phi : SPH \rightarrow SPK$ be a mapping from a symbolic Plithogenic hypergroup SPH into a symbolic Plithogenic hypergroup SPK defined by*

$$\phi(x) = (\psi(a), \psi(b)P_1, \psi(c)P_2), \quad \forall x = (a, bP_1, cP_2) \in SPH.$$

Then ϕ is a good hypergroup homomorphism.

Proof. Let $x = (a, bP_1, cP_2)$ and $y = (d, eP_1, fP_2)$ be two arbitrary elements in $SPR.$ Then

$$\begin{aligned} x * y &= (a * d, (a * e \cup b * d \cup b * e \cup b * f \cup c * e)P_1, (a * f \cup c * d \cup c * f)P_2), \\ \phi(x) &= (\psi(a), \psi(b)P_1, \psi(c)P_2), \\ \phi(y) &= (\psi(d), \psi(e)P_1, \psi(f)P_2), \\ \therefore \phi(x * y) &= (\psi(a * d), \psi((a * e \cup b * d \cup b * e \cup b * f \cup c * e)P_1, \psi(a * f \cup c * d \cup c * f)P_2)), \\ &= (\psi(a) \circ \psi(d), (\psi(a) \circ \psi(e) \cup \psi(b) \circ \psi(d) \cup \psi(b) \circ \psi(e) \cup \psi(b) \circ \psi(f) \cup \psi(c) \circ \psi(e))P_1, \\ &\quad (\psi(a) \circ \psi(f) \cup \psi(c) \circ \psi(d) \cup \psi(c) \circ \psi(f))P_2), \\ &= [(\psi(a), \psi(b)P_1, \psi(c)P_2)] \circ [(\psi(d), \psi(e)P_1, \psi(f)P_2)], \\ &= \phi(x) \circ \phi(y). \end{aligned}$$

Accordingly, ϕ is a good hypergroup homomorphism. \square

Definition 4.12. Let $(C, +)$ be any canonical hypergroup. The couple $(SPC, +)$ is called a symbolic Plithogenic canonical hypergroup. If $x = (a, a_1P_1, a_2P_2)$ and $y = (b, b_1P_1, b_2P_2)$ are any two elements of $SPC,$ the composition of x and y in SPC denoted by $x + y$ is defined as

$$x + y = \{(c, c_1P_1, c_2P_2) : c \in a + b, c_1 \in a_1 + b_1, c_2 \in a_2 + b_2\}. \tag{17}$$

Theorem 4.13. *Let $(SPC, +)$ be a symbolic Plithogenic canonical hypergroup. Then $(SPC, +)$ is a canonical hypergroup.*

Proof. Let $x = (a, a_1P_1, a_2P_2)$, $y = (b, b_1P_1, b_2P_2)$ and $z = (c, c_1P_1, c_2P_2)$ be arbitrary elements of SPC . Then

$$\begin{aligned} x + y &= (a, a_1P_1, a_2P_2) + (b, b_1P_1, b_2P_2) \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + b, u_1 \in a_1 + b_1, u_2 \in a_2 + b_2\} \\ &= \{(u, u_1P_1, u_2P_2) : u \in b + a, u_1 \in b_1 + a_1, u_2 \in b_2 + a_2\} \\ &= y + x. \end{aligned}$$

Next,

$$\begin{aligned} (x + y) + z &= ((a, a_1P_1, a_2P_2) + (b, b_1P_1, b_2P_2)) + (c, c_1P_1, c_2P_2) \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + b, u_1 \in a_1 + b_1, u_2 \in a_2 + b_2\} + (c, c_1P_1, c_2P_2) \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + b + c, u_1 \in a_1 + b_1 + c_1, u_2 \in a_2 + b_2 + c_2\} \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + (b + c), u_1 \in a_1 + (b_1 + c_1), u_2 \in a_2 + (b_2 + c_2)\} \\ &= (a, a_1P_1, a_2P_2) + ((b, b_1P_1, b_2P_2) + (c, c_1P_1, c_2P_2)) \\ &= x + (y + z). \end{aligned}$$

Since SPC is a symbolic Plithogenic canonical hypergroup, it follows that $(0, 0P_1, 0P_2) = (0, 0, 0) \in SPC$ so that

$$\begin{aligned} x + (0, 0, 0) &= (a, a_1P_1, a_2P_2) + (0, 0, 0) \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + 0, u_1 \in a_1 + 0, u_2 \in a_2 + 0\} \\ &= \{(u, u_1P_1, u_2P_2) : u \in \{a\}, u_1 \in \{a_1\}, u_2 \in \{a_2\}\} \\ &= \{(a, a_1P_1, a_2P_2)\} \\ &= \{x\} \text{ and similarly,} \\ (0, 0, 0) + x &= \{x\}. \end{aligned}$$

Also,

$$\begin{aligned} x + (-x) \cap (-x) + x &= [(a, a_1P_1, a_2P_2) + (-a, -a_1P_1, -a_2P_2)] \cap [(-a, -a_1P_1, -a_2P_2) \\ &\quad + (a, a_1P_1, a_2P_2)] \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + (-a), u_1 \in a_1 + (-a_1), u_2 \in a_2 + (-a_2)\} \\ &\quad \cap \{(v, v_1P_1, v_2P_2) : v \in (-a) + a, v_1 \in (-a_1) + (a_1), v_2 \in (-a_2) + (a_2)\} \\ &= \{(u, u_1P_1, u_2P_2) : u \in \{0\}, u_1 \in \{0\}, u_2 \in \{0\}\} \\ &\quad \cap \{(v, v_1P_1, v_2P_2) : v \in \{0\}, v_1 \in \{0\}, v_2 \in \{0\}\} \\ \therefore (0, 0, 0) &\in x + (-x) \cap (-x) + x \end{aligned}$$

which shows that $-x$ is the unique inverse of x , $\forall x \in SPC$.

Lastly, suppose that $z \in x + y$, then

$$\begin{aligned} (c, c_1P_1, c_2P_2) &\in (a, a_1P_1, a_2P_2) + (b, b_1P_1, b_2P_2) \\ &= \{(u, u_1P_1, u_2P_2) : u \in a + b, u_1 \in a_1 + b_1, u_2 \in a_2 + b_2\} \\ &= \{(u, u_1P_1, u_2P_2) : b \in -a + u, b_1 \in -a_1 + u_1, b_2 \in -a_2 + u_2\} \\ &= \{(b, b_1P_1, b_2P_2) : b \in -a + u, b_1 \in -a_1 + u_1, b_2 \in -a_2 + u_2\} \\ \therefore (b, b_1P_1, b_2P_2) &\in -(a, a_1P_1, a_2P_2) + (c, c_1P_1, c_2P_2) \\ \text{that is } y &\in -x + z \text{ and similarly,} \\ z &\in x + y \Rightarrow x \in z - y. \end{aligned}$$

Accordingly, $(SPC, +)$ is a canonical hypergroup. \square

Example 4.14. Let $\psi, \psi_1, \psi_2 : C_1 \rightarrow C_2$ be good canonical hypergroup homomorphisms and let SPC_1 and SPC_2 be two symbolic Plithogenic canonical hypergroups. If $\phi : SPC_1 \rightarrow SPC_2$ is a mapping defined by

$$\phi(x) = (\psi(a), \psi_1(a_1)P_1, \psi_2(a_2)P_2), \forall x = (a, a_1P_1, a_2P_2) \in SPC_1,$$

then ϕ is a good canonical hypergroup homomorphism and

$$\begin{aligned} \text{Ker}\phi &= \{(a, a_1P_1, a_2P_2) \in SPC_1 : \phi((a, a_1P_1, a_2P_2)) = (0, 0P_1, 0P_2)\} \\ &= \{(a, a_1P_1, a_2P_2) \in SPC_1 : (\psi(a), \psi_1(a_1)P_1, \psi_2(a_2)P_2) = (0, 0P_1, 0P_2)\} \\ &= \{(a, a_1P_1, a_2P_2) \in SPC_1 : \psi(a) = 0, \psi_1(a_1) = 0, \psi_2(a_2) = 0\} \\ &= \{(a, a_1P_1, a_2P_2) \in SPC_1 : a \in \text{Ker}(\psi), a_1 \in \text{Ker}(\psi_1), a_2 \in \text{Ker}(\psi_2)\} \\ &= \{(\text{Ker}\psi, \text{Ker}\psi_1P_1, \text{Ker}\psi_2P_2)\}. \end{aligned}$$

5. Conclusion

We have in this paper studied symbolic Plithogenic algebraic structures and hyper structures. In particular, we studied symbolic Plithogenic group, symbolic Plithogenic ring, symbolic Plithogenic hypergroup and symbolic Plithogenic canonical hypergroup, and we presented their basic properties.

Funding: This research received no external funding.

Acknowledgments: The authors acknowledge useful comments and suggestions of Professor Florentin Smarandache towards the improvement of the paper.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Adeleke E.O., Agboola A.A.A and Smarandache F., Refined Neutrosophic Rings I, International Journal of Neutrosophic Science (IJNS), 2020, vol. 2 (2), pp. 77-81.
(DOI:10.5281/zenodo.3728222)
2. Adeleke E.O., Agboola A.A.A. and Smarandache F., Refined Neutrosophic Rings II, International Journal of Neutrosophic Science (IJNS), 2020, vol. 2 (2), pp. 89-94.
(DOI:10.5281/zenodo.3728235)
3. Agboola A.A.A., On Refined Neutrosophic Algebraic Structures, Neutrosophic Sets and Systems (NSS), 2015, vol.10, pp. 99-101.
4. Agboola A.A.A., Akinola A.D. and Oyebola O.Y., Neutrosophic Rings I, Int. J. of Math. Comb., 2011, vol. 4, pp. 1-14.
5. Agboola A.A.A., Adeleke E.O. and Akinleye S.A., Neutrosophic Rings II, Int. J. of Math. Comb., 2012, vol. 2, pp. 1-8.
6. Agboola A.A.A., Akwu A.O. and Oyebo Y.T., Neutrosophic Groups and Neutrosopic Subgroups, Int. J. of Math. Comb., 2012, vol. 3, pp. 1-9.
7. Agboola A.A.A. and Davvaz B., On Neutrosophic Canonical Hyperrings and Neutrosophic Hyperrings, Neutrosophic Sets and Systems (NSS), 2014, vol. 2, pp. 34-41.
8. Agboola A.A.A., Davvaz B. and Smarandache F., Neutrosophic Quadruple Hyperstructures, Annals of Fuzzy Mathematics and Informatics, 2017, vol. 14, pp. 29-42.
9. Akinleye S.A., Smarandache F. and Agboola A.A.A., On Neutrosophic Quadruple Algebraic Structures, Neutrosophic Sets and Systems (NSS), 2016, vol. 12, pp. 122-126.
10. Al-Basheer O., Hajjari A. and Dalla R. On The Symbolic 3-Plithogenic Rings and Their Algebraic Properties, Neutrosophic Sets and Systems (NSS), 2023, vol 54, pp. 57-67.
11. Atanassov K., Intuitionistic fuzzy sets, Fuzzy Sets and Systems (FSS), 1986, vol. 20, pp. 87-96.
12. Corsini P., Prolegomena of Hypergroup Theory, Second edition, Avaiain editore, 1993.
13. Corsini P. and Leoreanu V., Applications of Hyperstructure Theory, Advances in Mathematics, Kluwer Academic Publishers, Dordrecht, 2003.
14. Davvaz B., Isomorphism theorems of hyperrings, Indian J. Pure Appl. Math., vol. 35(3), pp. 321-333, 2004.
15. Davvaz B. and Leoreanu-Fotea V., Hyperring Theory and Applications, International Academic Press, USA, 2007.
16. De Salvo M., Hyperrings and hyperfields, Annales Scientifiques de l'Universite de Clermont-Ferrand II, 1984, vol. 22, pp. 89-107.
17. Gayen S., Smarandache F., Jha S., Singh M.K., Broumi S. and Kumar R., Introduction to Plithogenic Hypersoft Subgroup, Neutrosophic Sets and Systems (NSS), 2020, vol 33, pp. 208-233.
18. Gomathy S., Nagarajan D. and Lathamaheswari M. Plithogenic sets and their applications in decision making, Neutrosophic Sets and Systems (NSS), 2020, vol 38, pp. 453-469.
19. Ibrahim M.A., Agboola A.A.A., Adeleke E.O. and Akinleye S.A., Introduction to Neutrosophic Subtraction Algebra and Neutrosophic Subtraction Semigroup, International Journal of Neutrosophic Science (IJNS), 2020, vol. 2 (1), pp. 47-62.
(DOI:10.5281/zenodo.3724603)

20. Krasner K., A class of hyperrings and hyperfields, *Int. J. Math. and Math. Sci.*, 1983, vol. 6 (2), pp. 307-311.
21. Marty F., Sur une generalization de la notion de groupe, 8th Congress Math. Scandinaves, Stockholm, Sweden, 1934, pp. 45-49.
22. Merkepci H and Abobala M. On the Symbolic 2-Plithogenic Rings, *International Journal of Neutrosophic Science (IJNS)* 2023, Vol. 20 (03), pp. 115-122.
DOI: <http://doi.org/10.54216/IJNS.200311>
23. Smarandache F., Introduction to the symbolic Plithogenic Algebraic Structures (revisited), *Neutrosophic Sets and Systems (NSS)*, 2023, vol 53 (1), pp. 653-665.
24. Smarandache F. and Abdel-Basset M., *Optimization Theory based on Neutrosophic and Plithogenic Sets*, Academic Press, 2020.
25. Smarandache F., An Overview of Plithogenic set and Symbolic Plithogenic Algebraic Structures, *Journal of Fuzzy Extension and Applications (JFEA)* 2023, vol. 4 (1), pp. 48-55.
DOI: <http://doi.org/10.22105/jfea.2023.382196.1252>
26. Smarandache F., *Plithogeny, Plithogenic Set, Logic, Probability, and Statistics*, 141 pages, Pons Editions, Brussels, Belgium, 2017. arXiv.org (Cornell University, Computer Science - Artificial Intelligence, 03Bxx:
27. Smarandache F., *Physical Plithogenic Set*, 71st Annual Gaseous Electronic Conference, Session LW1, Oregon Convention Center Room, Portland, Oregon, USA, November 5-9, 2018.
28. Smarandache F., *Plithogenic Algebraic Structures*, Chapter in "Nidus idearum Scilogs, V: joining the dots" (third version) Pons Publishing Brussels, 2019, pp. 123-125.
29. Smarandache F., *Neutrosophic Quadruple Numbers, Refined Neutrosophic Quadruple Numbers, Absorbance Law, and the Multiplication of Neutrosophic Quadruple Numbers*. In *Symbolic Neutrosophic Theory*, Chapter 7, Europa Nova, Brussels, Belgium, 2015, pp. 186-193.
30. Smarandache F., *Neutrosophy, Neutrosophic Probability, Set, and Logic*, ProQuest Information & Learning, Ann Arbor, Michigan, USA, 105 p., 1998.
<http://fs.unm.edu/eBook-Neutrosophic6.pdf> (edition online).
31. F. Smarandache, (T,I,F)- Neutrosophic Structures, *Neutrosophic Sets and Systems (NSS)*, 2015, vol. 8, pp. 3-10.
32. Taffach N.M. and Hatip A., A Brief Review on the Symbolic 2-Plithogenic Number Theory And Algebraic Equations, *Galoitica Journal of Mathematical Structures And Applications (GJMSA)* 2023, Vol. 5 (01), pp. 36-44.
DOI: <http://doi.org/10.54216/GJMSA.050103>
33. Taffach N.M. and Othman K.B., An Introduction to Symbolic 2-Plithogenic Modules over Symbolic 2-Plithogenic Rings, *Neutrosophic Sets and Systems (NSS)*, 2023, vol 54, pp. 33-44.
34. Taffach N.M., An Introduction to Symbolic 2-Plithogenic Vector Spaces Generated from Fusion of Symbolic Plithogenic Sets and Vector Spaces, *Neutrosophic Sets and Systems (NSS)*, 2023, vol 54, pp. 45-57.
35. Vasantha Kandasamy W.B. and Smarandache F., *Neutrosophic Rings*, Hexis, Phoenix, Arizona, 2006.
<http://fs.unm.edu/NeutrosophicRings.pdf>

36. Vougiouklis T., The fundamental relation in hyperrings. The general hyperfield, Proc. Fourth Int. Congress on Algebraic Hyperstructures and Applications (AHA 1990), World Scientific, 1991, pp. 203-211.
37. Zadeh L.A., Fuzzy sets, Information and Control, 1965, vol. 8, pp. 338-353.

Received: March 25, 2023. Accepted: July 18, 2023