

Neutrosophic Sets and Systems

Volume 59 *Neutrosophic Sets and Systems*,
Vol. 59, 2023 - Special Issue on Symbolic
Plithogenic Algebraic Structures

Article 5

10-28-2023

An Introduction to The Dual Symbolic 3-Plithogenic And 4-Plithogenic Numbers

Khadija Ben Othman

Maretta Sarkis

Djamal Lhiani

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Othman, Khadija Ben; Maretta Sarkis; and Djamal Lhiani. "An Introduction to The Dual Symbolic 3-Plithogenic And 4-Plithogenic Numbers." *Neutrosophic Sets and Systems* 59, 1 (2023).
https://digitalrepository.unm.edu/nss_journal/vol59/iss1/5

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



An Introduction to The Dual Symbolic 3-Plithogenic And 4-Plithogenic Numbers

¹Khadija Ben Othman, ²Maretta Sarkis, ³ Djamal Lhiani

¹. Umm Al-Qura Univerity, Mekka, Saudi Arabia

²Abu Dhabi University, Abu Dhabi, United Arab Emirates

³University of Blida 1, Department of Mathematics, Algeria

Co-khagijabenothman33@gmail.com

Abstract:

The objective of this paper is to use dual numbers with symbolic 3-plithogenic and 4-plithogenic numbers in one numerical system called dual symbolic 3-plithogenic/4-plithogenic numbers.

Also, the elementary algebraic properties of the suggested systems will be discussed in terms of theorems and related examples that explain the validity of these algebraic number systems.

Keywords: Symbolic 3-plithogenic number, dual number, dual symbolic 3-plithogenic number, Symbolic 4-plithogenic number, dual symbolic 4-plithogenic number.

Introduction and preliminaries.

Dual numbers are considered as a generalization of real numbers, where they are defined as follows:

$D = \{a + bt; t^2 = 0, a, b \in R\}$ [1]. Dual numbers make together a commutative ring with many interesting properties.

Addition on D is defined as follows:

$$(a_0 + b_0t) + (a_1 + b_1t) = (a_0 + a_1) + (b_0 + b_1)t$$

Multiplication on D is defined as follows:

$$(a_0 + b_0t) \cdot (a_1 + b_1t) = (a_0a_1) + (a_0b_1 + b_0a_1)t$$

In [2-4], smarandache presented symbolic n-plithogenic sets, then they were used in generalizing many famous algebraic structures such as rings, matrices, and other structures [6-11].

We refer to many similar numerical systems that generalize real number, such as neutrosophic numbers, split-complex number, and weak fuzzy numbers [12-18]. These generalized numbers were applicable in cryptography and matrix theory [19-24].

Through this paper, we use symbolic 3-plithogenic real numbers and symbolic 4-plithogenic real numbers to build a new generalization of real numbers, and we present some of its elementary algebraic properties.

Main concepts.

Definition.

The set of symbolic 3-plithogenic dual numbers is defined as follows:

$$3 - SP_D = \{(x_0 + x_1t) + (y_0 + y_1t)P_1 + (z_0 + z_1t)P_2 + (s_0 + s_1t)P_3; x_i, y_i, z_i, s_i \in R, t^2 = 0\}.$$

Definition.

Addition of $3 - SP_D$ is defined:

$$[(m_0 + m_1t) + (k_0 + k_1t)P_1 + (s_0 + s_1t)P_2 + (r_0 + r_1t)P_3] + [(n_0 + n_1t) + (l_0 + l_1t)P_1 + (q_0 + q_1t)P_2 + (g_0 + g_1t)P_3] = (m_0 + n_0) + (m_1 + n_1)t + [(k_0 + l_0) + (k_1 + l_1)t]P_1 + [(s_0 + q_0) + (s_1 + q_1)t]P_2 + [(r_0 + g_0) + (r_1 + g_1)t]P_3.$$

$(3 - SP_D, +)$ is an abelian group.

Remark.

A symbolic 3-plithogenic dual number $X = (x_0 + x_1t) + (y_0 + y_1t)P_1 + (z_0 + z_1t)P_2 + (s_0 + s_1t)P_3$

can be written:

$$X = (x_0 + y_0P_1 + z_0P_2 + s_0P_3) + t(x_1 + y_1P_1 + z_1P_2 + s_1P_3).$$

Definition.

Let

$$X = (x_0 + x_1P_1 + x_2P_2 + x_3P_3) + t(\acute{x}_0 + \acute{x}_1P_1 + \acute{x}_2P_2 + \acute{x}_3P_3) = M_1 + M_2t,$$

$$Y = (y_0 + y_1P_1 + y_2P_2 + y_3P_3) + t(\acute{y}_0 + \acute{y}_1P_1 + \acute{y}_2P_2 + \acute{y}_3P_3) = N_1 + N_2t \in 3 - SP_D,$$

then:

Multiplication on $3 - SP_D$ is defined as follows:

$$X.Y = M_1N_1 + t(M_1N_2 + N_1M_2)$$

Example.

Consider $X = (1 + P_1 + P_2 + P_3) + t(2 - P_3), Y = P_1 + t(1 - P_3)$, we have:

$$X + Y = (1 + 2P_1 + P_2 + P_3) + t(3 - 2P_3)$$

$$X.Y = (1 + P_1 + P_2 + P_3)P_1 + t[(1 + P_1 + P_2 + P_3)(1 - P_3) + (2 - P_3)P_1] =$$

$$(2P_1 + P_2 + P_3) + t[(1 - P_3 + P_2 - P_3 + P_3 - P_3 + P_1 - P_3) + 2P_1 - P_3] =$$

$$(2P_1 + P_2 + P_3) + t(1 + P_1 + P_2 - 3P_3).$$

Remark.

$(3 - SP_D, +, \cdot)$ Is a commutative ring.

Invertibility:

Theorem.

Let $X = (m_0 + m_1P_1 + m_2P_2 + m_3P_3) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3) \in 3 - SP_D$, then

X is invertible if and only if $m_0 \neq 0, m_0 + m_1 \neq 0, m_0 + m_1 + m_2 \neq 0, m_0 + m_1 +$

$m_2 + m_3 \neq 0$ and:

$$\begin{aligned} X^{-1} = \frac{1}{X} = & \left[\frac{1}{m_0} + \left(\frac{1}{m_0 + m_1} - \frac{1}{m_0} \right) P_1 + \left(\frac{1}{m_0 + m_1 + m_2} - \frac{1}{m_0 + m_1} \right) P_2 \right. \\ & \left. + \left(\frac{1}{m_0 + m_1 + m_2 + m_3} - \frac{1}{m_0 + m_1 + m_2} \right) P_3 \right] \\ & - t \left[\frac{n_0}{(m_0)^2} + \left(\frac{n_0 + n_1}{(m_0 + m_1)^2} - \frac{n_0}{(m_0)^2} \right) P_1 \right. \\ & \left. + \left(\frac{n_0 + n_1 + n_2}{(m_0 + m_1 + m_2)^2} - \frac{n_0 + n_1}{(m_0 + m_1)^2} \right) P_2 \right. \\ & \left. + \left(\frac{n_0 + n_1 + n_2 + n_3}{(m_0 + m_1 + m_2 + m_3)^2} - \frac{n_0 + n_1 + n_2}{(m_0 + m_1 + m_2)^2} \right) P_3 \right] \end{aligned}$$

Proof.

X is invertible if and only if $\frac{1}{X}$ is defined as follows:

$$\begin{aligned} \frac{1}{X} &= \frac{1}{(m_0 + m_1P_1 + m_2P_2 + m_3P_3) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3)} \\ &= \frac{(m_0 + m_1P_1 + m_2P_2 + m_3P_3) - t(n_0 + n_1P_1 + n_2P_2 + n_3P_3)}{[(m_0 + m_1P_1 + m_2P_2 + m_3P_3) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3)][(m_0 + m_1P_1 + m_2P_2 + m_3P_3) - t(n_0 + n_1P_1 + n_2P_2 + n_3P_3)]} \\ &= \frac{(m_0 + m_1P_1 + m_2P_2 + m_3P_3) - t(n_0 + n_1P_1 + n_2P_2 + n_3P_3)}{(m_0 + m_1P_1 + m_2P_2 + m_3P_3)^2} \end{aligned}$$

So that $m_0 + m_1P_1 + m_2P_2 + m_3P_3$ is invertible in $3 - SP_R$.

This is equivalent to $m_0 \neq 0, m_0 + m_1 \neq 0, m_0 + m_1 + m_2 \neq 0, m_0 + m_1 + m_2 + m_3 \neq 0$.

On the other hand, $\frac{1}{X} = \frac{1}{m_0 + m_1P_1 + m_2P_2 + m_3P_3} - t \frac{(n_0 + n_1P_1 + n_2P_2 + n_3P_3)^2}{(m_0 + m_1P_1 + m_2P_2 + m_3P_3)^2}$

Put $Y = \left[\frac{1}{m_0} + \left(\frac{1}{m_0 + m_1} - \frac{1}{m_0} \right) P_1 + \left(\frac{1}{m_0 + m_1 + m_2} - \frac{1}{m_0 + m_1} \right) P_2 + \left(\frac{1}{m_0 + m_1 + m_2 + m_3} - \frac{1}{m_0 + m_1 + m_2} \right) P_3 \right] - t \left[\frac{n_0}{(m_0)^2} + \left(\frac{n_0 + n_1}{(m_0 + m_1)^2} - \frac{n_0}{(m_0)^2} \right) P_1 + \left(\frac{n_0 + n_1 + n_2}{(m_0 + m_1 + m_2)^2} - \frac{n_0 + n_1}{(m_0 + m_1)^2} \right) P_2 + \left(\frac{n_0 + n_1 + n_2 + n_3}{(m_0 + m_1 + m_2 + m_3)^2} - \frac{n_0 + n_1 + n_2}{(m_0 + m_1 + m_2)^2} \right) P_3 \right]$.

Compute the result of XY to get:

$$XY = 1$$

So that, $X^{-1} = \frac{1}{X} = Y$

Example.

Take $X = (1 + P_3) + t(2 + P_3) \in 3 - SP_D$:

$$X^{-1} = \frac{1}{1} + \left(\frac{1}{2} - \frac{1}{1} \right) P_2 - t \left[\frac{2}{1} + \left(\frac{2}{1} - \frac{2}{1} \right) P_1 + \left(\frac{2}{1} - \frac{2}{1} \right) P_2 + \left(\frac{3}{4} - \frac{2}{1} \right) P_3 \right] = 1 - \frac{1}{2} P_3 - t \left(2 - \frac{5}{4} P_3 \right).$$

Natural power.

Theorem.

Let $X = (m_0 + m_1P_1 + m_2P_2 + m_3P_3) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3) \in 3 - SP_D$, then:

$$X^n = (m_0)^n + ((m_0 + m_1)^n - (m_0)^n)P_1 + ((m_0 + m_1 + m_2)^n - (m_0 + m_1)^n)P_2 + ((m_0 + m_1 + m_2 + m_3)^n - (m_0 + m_1 + m_2)^n)P_3 + n(n_0 + n_1P_1 + n_2P_2 + n_3P_3)[(m_0)^{n-1} + ((m_0 + m_1)^{n-1} - (m_0)^{n-1})P_1 + ((m_0 + m_1 + m_2)^{n-1} - (m_0 + m_1)^{n-1})P_2 + ((m_0 + m_1 + m_2 + m_3)^{n-1} - (m_0 + m_1 + m_2)^{n-1})P_3] \text{ for } n \in \mathbb{N}.$$

Proof.

Let $X = A + Bt; A, B \in 3 - SP_D$, then:

$A^n = A^n + nA^{n-1}Bt$, we get:

$$A^n = (m_0)^n + ((m_0 + m_1)^n - (m_0)^n)P_1 + ((m_0 + m_1 + m_2)^n - (m_0 + m_1)^n)P_2 + ((m_0 + m_1 + m_2 + m_3)^n - (m_0 + m_1 + m_2)^n)P_3, \text{ then the proof holds.}$$

Example.

Take $X = (1 + P_3) + t(2 - P_3) \in 3 - SP_D$

$$X^3 = 1 + (1 - 1)P_1 + (1 - 1)P_2 + (8 - 1)P_3 + 3t(2 - P_3)[1 + (1 - 1)P_1 + (1 - 1)P_2 + (4 - 1)P_3] = 1 + 7P_3 + 3t[(2 - P_3)(1 + 3P_3)] = 1 + 7P_3 + t(6 + 6P_3).$$

Idempotency.

Definition.

Let $X \in 3 - SP_D$, then X is called idempotent if and only if $X^2 = X$.

Theorem.

Let $X = (m_0 + m_1P_1 + m_2P_2 + m_3P_3) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3) \in 3 - SP_D$, then X is called idempotent if and only if:

1. $m_0 + m_1P_1 + m_2P_2 + m_3P_3$ is idempotent.
2. $(n_0 + n_1P_1 + n_2P_2 + n_3P_3)[2m_0 - 1 + 2m_1P_1 + 2m_2P_2 + 2m_3P_3] = 0$

Proof.

$X = M + Nt$ is idempotent if and only if:

$$X^2 = X \Rightarrow \begin{cases} M^2 = M \\ 2MN = N \Rightarrow N(2M - 1) = 0 \end{cases}$$

For $M = m_0 + m_1P_1 + m_2P_2 + m_3P_3, N = n_0 + n_1P_1 + n_2P_2 + n_3P_3 \in 3 - SP_R$.

This implies the proof.

Definition.

The set of symbolic 4-plithogenic dual numbers is defined as follows:

$$4 - SP_D = \{(x_0 + x_1t) + (y_0 + y_1t)P_1 + (z_0 + z_1t)P_2 + (s_0 + s_1t)P_3 + (l_0 + l_1t)P_4; x_i, y_i, z_i, s_i, l_i \in R, t^2 = 0\}.$$

Definition.

Addition of $4 - SP_D$ is defined:

$$[(m_0 + m_1t) + (k_0 + k_1t)P_1 + (s_0 + s_1t)P_2 + (r_0 + r_1t)P_3 + (d_0 + d_1t)P_4] + [(n_0 + n_1t) + (l_0 + l_1t)P_1 + (q_0 + q_1t)P_2 + (g_0 + g_1t)P_3 + (c_0 + c_1t)P_4] =$$

$$(m_0 + n_0) + (m_1 + n_1)t + [(k_0 + l_0) + (k_1 + l_1)t]P_1 + [(s_0 + q_0) + (s_1 + q_1)t]P_2 + [(r_0 + g_0) + (r_1 + g_1)t]P_3 + [(d_0 + c_0) + (d_1 + c_1)t]P_4.$$

$(4 - SP_D, +)$ is an abelian group.

Remark.

A symbolic 4-plithogenic dual number $X = (x_0 + x_1t) + (y_0 + y_1t)P_1 + (z_0 + z_1t)P_2 + (s_0 + s_1t)P_3 + (d_0 + d_1t)P_4$

can be written:

$$X = (x_0 + y_0P_1 + z_0P_2 + s_0P_3 + d_0P_4) + t(x_1 + y_1P_1 + z_1P_2 + s_1P_3 + d_1P_4).$$

Definition.

Let

$$X = (x_0 + x_1P_1 + x_2P_2 + x_3P_3 + x_4P_4) + t(\acute{x}_0 + \acute{x}_1P_1 + \acute{x}_2P_2 + \acute{x}_3P_3 + \acute{x}_4P_4) = M_1 + M_2t,$$

$$Y = (y_0 + y_1P_1 + y_2P_2 + y_3P_3 + y_4P_4) + t(\acute{y}_0 + \acute{y}_1P_1 + \acute{y}_2P_2 + \acute{y}_3P_3 + \acute{y}_4P_4) = N_1 + N_2t \in 4 - SP_D,$$

then:

Multiplication on $4 - SP_D$ is defined as follows:

$$X.Y = M_1N_1 + t(M_1N_2 + N_1M_2)$$

Example.

Consider $X = (1 + P_4) + t(2 - P_3), Y = P_1 + t(1 - P_4)$, we have:

$$X + Y = (1 + P_1 + P_4) + t(3 - P_3 - P_4)$$

$$X.Y = (1 + P_4)P_1 + t[(1 + P_4)(1 - P_4) + (2 - P_3)P_1] = (P_1 + P_4) + t[(1 - P_4) + 2P_1 - P_3] = (P_1 + P_4) + t(1 + 2P_1 - P_3 - P_4).$$

Remark.

$(4 - SP_D, +, \cdot)$ Is a commutative ring.

Invertibility:

Theorem.

Let

$$X = (m_0 + m_1P_1 + m_2P_2 + m_3P_3 + m_4P_4) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3 + n_4P_4) \in 4 - SP_D,$$

then X is invertible if and only if $m_0 \neq 0, m_0 + m_1 \neq 0, m_0 + m_1 + m_2 \neq 0, m_0 + m_1 + m_2 + m_3 \neq 0, m_0 + m_1 + m_2 + m_3 + m_4 \neq 0$ and:

$$X^{-1} = \frac{1}{X} = \left[\frac{1}{m_0} + \left(\frac{1}{m_0+m_1} - \frac{1}{m_0} \right) P_1 + \left(\frac{1}{m_0+m_1+m_2} - \frac{1}{m_0+m_1} \right) P_2 + \left(\frac{1}{m_0+m_1+m_2+m_3} - \frac{1}{m_0+m_1+m_2} \right) P_3 + \left(\frac{1}{m_0+m_1+m_2+m_3+m_4} - \frac{1}{m_0+m_1+m_2+m_3} \right) P_4 \right] - t \left[\frac{n_0}{(m_0)^2} + \left(\frac{n_0+n_1}{(m_0+m_1)^2} - \frac{n_0}{(m_0)^2} \right) P_1 + \left(\frac{n_0+n_1+n_2}{(m_0+m_1+m_2)^2} - \frac{n_0+n_1}{(m_0+m_1)^2} \right) P_2 + \left(\frac{n_0+n_1+n_2+n_3}{(m_0+m_1+m_2+m_3)^2} - \frac{n_0+n_1+n_2}{(m_0+m_1+m_2)^2} \right) P_3 + \left(\frac{n_0+n_1+n_2+n_3+n_4}{(m_0+m_1+m_2+m_3+m_4)^2} - \frac{n_0+n_1+n_2+n_3}{(m_0+m_1+m_2+m_3)^2} \right) P_4 \right].$$

Proof.

X is invertible if and only if $\frac{1}{X}$ is defined as follows:

$$\frac{1}{X} = \frac{1}{(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)+t(n_0+n_1P_1+n_2P_2+n_3P_3+n_4P_4)} = \frac{(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)-t(n_0+n_1P_1+n_2P_2+n_3P_3+n_4P_4)}{[(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)+t(n_0+n_1P_1+n_2P_2+n_3P_3+n_4P_4)][(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)-t(n_0+n_1P_1+n_2P_2+n_3P_3+n_4P_4)]} = \frac{(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)-t(n_0+n_1P_1+n_2P_2+n_3P_3+n_4P_4)}{(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)^2},$$

So that $m_0 + m_1P_1 + m_2P_2 + m_3P_3 + m_4P_4$ is invertible in $4 - SP_R$.

This is equivalent to $m_0 \neq 0, m_0 + m_1 \neq 0, m_0 + m_1 + m_2 \neq 0, m_0 + m_1 + m_2 + m_3 \neq 0, m_0 + m_1 + m_2 + m_3 + m_4 \neq 0$.

On the other hand,

$$\frac{1}{X} = \frac{1}{m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4} - t \frac{(n_0+n_1P_1+n_2P_2+n_3P_3+n_4P_4)^2}{(m_0+m_1P_1+m_2P_2+m_3P_3+m_4P_4)^2}$$

Put

$$Y = \left[\frac{1}{m_0} + \left(\frac{1}{m_0+m_1} - \frac{1}{m_0} \right) P_1 + \left(\frac{1}{m_0+m_1+m_2} - \frac{1}{m_0+m_1} \right) P_2 + \left(\frac{1}{m_0+m_1+m_2+m_3} - \frac{1}{m_0+m_1+m_2} \right) P_3 + \left(\frac{1}{m_0+m_1+m_2+m_3+m_4} - \frac{1}{m_0+m_1+m_2+m_3} \right) P_4 \right] - t \left[\frac{n_0}{(m_0)^2} + \left(\frac{n_0+n_1}{(m_0+m_1)^2} - \frac{n_0}{(m_0)^2} \right) P_1 + \left(\frac{n_0+n_1+n_2}{(m_0+m_1+m_2)^2} - \frac{n_0+n_1}{(m_0+m_1)^2} \right) P_2 + \left(\frac{n_0+n_1+n_2+n_3}{(m_0+m_1+m_2+m_3)^2} - \frac{n_0+n_1+n_2}{(m_0+m_1+m_2)^2} \right) P_3 + \left(\frac{n_0+n_1+n_2+n_3+n_4}{(m_0+m_1+m_2+m_3+m_4)^2} - \frac{n_0+n_1+n_2+n_3}{(m_0+m_1+m_2+m_3)^2} \right) P_4 \right].$$

Compute the result of XY to get:

$$XY = 1$$

$$\text{So that, } X^{-1} = \frac{1}{X} = Y$$

Natural power.

Theorem.

Let

$$X = (m_0 + m_1P_1 + m_2P_2 + m_3P_3 + m_4P_4) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3 + n_4P_4) \in 4 - SP_D,$$

then:

$$X^n = (m_0)^n + ((m_0 + m_1)^n - (m_0)^n)P_1 + ((m_0 + m_1 + m_2)^n - (m_0 + m_1)^n)P_2 + ((m_0 + m_1 + m_2 + m_3)^n - (m_0 + m_1 + m_2)^n)P_3 + ((m_0 + m_1 + m_2 + m_3 + m_4)^n - (m_0 + m_1 + m_2 + m_3)^n)P_4 + n(n_0 + n_1P_1 + n_2P_2 + n_3P_3 + n_4P_4)[(m_0)^{n-1} + ((m_0 + m_1)^{n-1} - (m_0)^{n-1})P_1 + ((m_0 + m_1 + m_2)^{n-1} - (m_0 + m_1)^{n-1})P_2 + ((m_0 + m_1 + m_2 + m_3)^{n-1} - (m_0 + m_1 + m_2)^{n-1})P_3 + ((m_0 + m_1 + m_2 + m_3 + m_4)^{n-1} - (m_0 + m_1 + m_2 + m_3)^{n-1})P_4] \text{ for } n \in N.$$

Proof.

Let $X = A + Bt; A, B \in 4 - SP_D$, then:

$$A^n = A^n + nA^{n-1}Bt, \text{ we get:}$$

$$A^n = (m_0)^n + ((m_0 + m_1)^n - (m_0)^n)P_1 + ((m_0 + m_1 + m_2)^n - (m_0 + m_1)^n)P_2 + ((m_0 + m_1 + m_2 + m_3)^n - (m_0 + m_1 + m_2)^n)P_3 + ((m_0 + m_1 + m_2 + m_3 + m_4)^n - (m_0 + m_1 + m_2 + m_3)^n)P_4, \text{ then the proof holds.}$$

Idempotency.

Definition.

Let $X \in 4 - SP_D$, then X is called idempotent if and only if $X^2 = X$.

Theorem.

Let $X = (m_0 + m_1P_1 + m_2P_2 + m_3P_3 + m_4P_4) + t(n_0 + n_1P_1 + n_2P_2 + n_3P_3 + n_4P_4) \in 4 - SP_D$, then X is called idempotent if and only if:

1. $m_0 + m_1P_1 + m_2P_2 + m_3P_3 + m_4P_4$ is idempotent.
2. $(n_0 + n_1P_1 + n_2P_2 + n_3P_3 + n_4P_4)[2m_0 - 1 + 2m_1P_1 + 2m_2P_2 + 2m_3P_3 + 2m_4P_4] = 0$

Proof.

$X = M + Nt$ is idempotent if and only if:

$$X^2 = X \Rightarrow \begin{cases} M^2 = M \\ 2MN = N \Rightarrow N(2M - 1) = 0 \end{cases}$$

For $M = m_0 + m_1P_1 + m_2P_2 + m_3P_3 + m_4P_4, N = n_0 + n_1P_1 + n_2P_2 + n_3P_3 + n_4P_4 \in 4 - SP_R$.

This implies the proof.

Conclusion

In this paper, we have studied for the first time the combination of symbolic 3-plithogenic numbers and 4-plithogenic numbers with dual numbers. The novel algebraic structures generated by them are called dual symbolic 3-plithogenic numbers and dual symbolic 4-plithogenic numbers.

We have determined the invertibility condition and the formula of the inverse for dual symbolic 3-plithogenic and 4-plithogenic numbers.

References

1. Akar, M.; Yuce, S.; Sahin, S. On The Dual Hyperbolic Numbers and The Complex Hyperbolic Numbers. *Jcscm* **2018**, *8*, DOI: 10.20967/jcscm.2018.01.001.
2. Smarandache, F., " Introduction to the Symbolic Plithogenic Algebraic Structures (revisited)", *Neutrosophic Sets and Systems*, vol. 53, 2023.
3. Florentin Smarandache, Plithogenic Algebraic Structures. Chapter in "Nidus idearum Scilogs, V: joining the dots" (third version), Pons Publishing Brussels, pp. 123-125, 2019.
4. Florentin Smarandache: Plithogenic Set, an Extension of Crisp, Fuzzy, Intuitionistic Fuzzy, and Neutrosophic Sets – Revisited, *Neutrosophic Sets and Systems*, vol. 21, 2018, pp. 153-166.
5. Ali, R., and Hasan, Z., "An Introduction To The Symbolic 3-Plithogenic Vector Spaces", *Galoitica Journal Of Mathematical Structures and Applications*, vol. 6, 2023.
6. Rawashdeh, A., "An Introduction To The Symbolic 3-plithogenic Number Theory", *Neoma Journal Of Mathematics and Computer Science*, 2023.

7. Ben Othman, K., "On Some Algorithms For Solving Symbolic 3-Plithogenic Equations", Neoma Journal Of Mathematics and Computer Science, 2023.
8. Ali, R., and Hasan, Z., " An Introduction To The Symbolic 3-Plithogenic Modules ", Galoitica Journal Of Mathematical Structures and Applications, vol. 6, 2023.
9. M. B. Zeina, N. Altounji, M. Abobala, and Y. Karmouta, "Introduction to Symbolic 2-Plithogenic Probability Theory," *Galoitica: Journal of Mathematical Structures and Applications*, vol. 7, no. 1, 2023.
10. Nader Mahmoud Taffach , Ahmed Hatip.," A Brief Review on The Symbolic 2-Plithogenic Number Theory and Algebraic Equations ", Galoitica Journal Of Mathematical Structures and Applications, Vol.5, 2023.
11. Merkepçi, H., and Abobala, M., " On The Symbolic 2-Plithogenic Rings", International Journal of Neutrosophic Science, 2023.
12. Sarkis, M., " On The Solutions Of Fermat's Diophantine Equation In 3-refined Neutrosophic Ring of Integers", Neoma Journal of Mathematics and Computer Science, 2023.
13. Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", *Mathematical Problems in Engineering*, Hindawi, 2021
14. Abobala, M., "A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", *International Journal of Mathematics and Mathematical Sciences*, hindawi, 2021
15. Khaldi, A., " A Study On Split-Complex Vector Spaces", Neoma Journal Of Mathematics and Computer Science, 2023.
16. Ahmad, K., " On Some Split-Complex Diophantine Equations", Neoma Journal Of Mathematics and Computer Science, 2023.

17. Ali, R., " On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces", Neoma Journal Of Mathematics and Computer Science, 2023.
18. Hatip, A., "An Introduction To Weak Fuzzy Complex Numbers ", Galoitica Journal Of Mathematical Structures and Applications, Vol.3, 2023.
19. Merkepci, M., Abobala, M., and Allouf, A., " The Applications of Fusion Neutrosophic Number Theory in Public Key Cryptography and the Improvement of RSA Algorithm ", Fusion: Practice and Applications, 2023.
20. Merkepci, M.; Sarkis, M. An Application of Pythagorean Circles in Cryptography and Some Ideas for Future Non Classical Systems. Galoitica Journal of Mathematical Structures and Applications **2022**.
21. Merkepci, M., and Abobala, M., " Security Model for Encrypting Uncertain Rational Data Units Based on Refined Neutrosophic Integers Fusion and El Gamal Algorithm ", Fusion: Practice and Applications, 2023.
22. Alhasan, Y., Alfahal, A., Abdulfatah, R., Ali, R., and Aljibawi, M., " On A Novel Security Algorithm For The Encryption Of 3×3 Fuzzy Matrices With Rational Entries Based On The Symbolic 2-Plithogenic Integers And El-Gamal Algorithm", International Journal of Neutrosophic Science, 2023.
23. Alfahal, A., Alhasan, Y., Abdulfatah, R., Mehmood, A., and Kadhim, T., " On Symbolic 2-Plithogenic Real Matrices And Their Algebraic Properties ", International Journal of Neutrosophic Science, 2023.
24. Abobala, M., and Allouf, A., " On A Novel Security Scheme for The Encryption and Decryption Of 2×2 Fuzzy Matrices with Rational Entries Based on The Algebra of Neutrosophic Integers and El-Gamal Crypto-System", Neutrosophic Sets and Systems, vol.54, 2023.

25. Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Concept Of Symbolic 7-Plithogenic Real Matrices", *Pure Mathematics For Theoretical Computer Science, Vol.1, 2023.*
26. Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Symbolic 8-Plithogenic Matrices", *Pure Mathematics For Theoretical Computer Science, Vol.1, 2023.*
27. Ben Othman, K., Albasheer, O., Nadweh, R., Von Shtawzen, O., and Ali, R., "On The Symbolic 6-Plithogenic and 7-Plithogenic Rings", *Galoitica Journal of Mathematical Structures and Applications, Vol.8, 2023.*

Received 18/6/2023, Accepted 10/10/2023