

Neutrosophic Sets and Systems

Volume 59 *Neutrosophic Sets and Systems*,
Vol. 59, 2023 - Special Issue on Symbolic
Plithogenic Algebraic Structures

Article 17

10-28-2023

The Computing of Pythagoras Triples in Symbolic 2-Plithogenic Rings

Abuobida Mohammed A. Alfahal

Yaser Ahmad Alhasan

Raja Abdullah Abdulfatah

Ahmad Abd Al-Aziz

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Alfahal, Abuobida Mohammed A.; Yaser Ahmad Alhasan; Raja Abdullah Abdulfatah; and Ahmad Abd Al-Aziz. "The Computing of Pythagoras Triples in Symbolic 2-Plithogenic Rings." *Neutrosophic Sets and Systems* 59, 1 (2023). https://digitalrepository.unm.edu/nss_journal/vol59/iss1/17

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact disc@unm.edu.



The Computing of Pythagoras Triples in Symbolic 2-Plithogenic Rings

Abuobida Mohammed A. Alfahal^{1,*}, Yaser Ahmad Alhasan², Raja Abdullah Abdulfatah³, Ahmad Abd Al-Aziz⁴

¹Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

²Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

³Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

⁴Cairo University, Department of Mathematics, Cairo, Egypt

Emails: ¹a.alfahal@psau.edu.sa; ²y.alhasan@psau.edu.sa; ³r.abdulfatah@psau.edu.sa ;

Corresponding author: a.alfahal@psau.edu.sa

Abstract:

This paper is dedicated to finding a general algorithm for generating different solutions for Pythagoras' non-linear Diophantine equation in four variables $x^2 + y^2 = z^2$ in symbolic 2-plithogenic rings, which are known as Pythagoras triples.

Also, we present some examples of those triples in some finite symbolic 2-plithogenic rings.

Keywords: symbolic 2-plithogenic ring, Pythagoras triples, Diophantine equations

Introduction and Preliminaries

Symbolic n-plithogenic algebraic structures are a new generalization of classical algebraic structures, as they have serious algebraic properties to study.

In the previous literature, we can clearly note several algebraic studies that were interested in discovering the properties of these algebraic structures, for example, we can find some applications of plithogenic structures in probability, ring theory, linear spaces, matrices, and equations [1-10].

Researchers have studied Pythagorean triples in the ring of ordinary algebraic numbers [11-14]. Several efficient algorithms for calculating these quadruples have been presented, as solutions to the corresponding Diophantine equation.

This has motivated us to study Pythagoras triples in the symbolic 2-plithogenic commutative case, where we find a general algorithm for generating different solutions for Pythagoras non-linear Diophantine equation in four variables $x^2 + y^2 = z^2$ in symbolic 2-plithogenic rings.

Definition.

The symbolic 2-plithogenic ring of real numbers is defined as follows:

$$2 - SP_R = \{t_0 + t_1P_1 + t_2P_2; t_i \in R, P_1 \times P_2 = P_2 \times P_1 = P_2, P_1^2 = P_2^2 = P_2\}$$

The addition operation on $2 - SP_R$ is defined as follows:

$$(t_0 + t_1P_1 + t_2P_2) + (t'_0 + t'_1P_1 + t'_2P_2) = (t_0 + t'_0) + (t_1 + t'_1)P_1 + (t_2 + t'_2)P_2$$

The multiplication on $2 - SP_R$ is defined as follows:

$$\begin{aligned} (t_0 + t_1P_1 + t_2P_2)(t'_0 + t'_1P_1 + t'_2P_2) \\ = t_0t'_0 + (t_0t'_1 + t_1t'_0 + t_1t'_1)P_1 + (t_0t'_2 + t_1t'_2 + t_2t'_2 + t_2t'_0 + t_2t'_1)P_2 \end{aligned}$$

Main Discussion

Definition.

Let R be a ring, then (t, s, k) is called a Pythagoras triple if and only if $t^2 + s^2 = k^2; t, s, k \in R..$

Theorem.

Let $T = t_0 + t_1P_1 + t_2P_2, S = s_0 + s_1P_1 + s_2P_2, K = k_0 + k_1P_1 + k_2P_2$ are three arbitrary symbolic 3-plithogenic elements $T, S, K \in 2 - SP_R$, then (T, S, K) are Pythagoras triple in $2 - SP_R$ if and only if:

$$\begin{cases} (t_0, s_0, k_0), (t_0 + t_1, s_0 + s_1, k_0 + k_1) \text{ are pythagoras triples in } R \\ (t_0 + t_1 + t_2, s_0 + s_1 + s_2, k_0 + k_1 + k_2) \text{ is pythagoras triple in } R \end{cases}$$

Proof.

According to [], we have:

$$T^2 = t_0^2 + [(t_0 + t_1)^2 - t_0^2]P_1 + [(t_0 + t_1 + t_2)^2 - (t_0 + t_1)^2]P_2$$

$$S^2 = s_0^2 + [(s_0 + s_1)^2 - s_0^2]P_1 + [(s_0 + s_1 + s_2)^2 - (s_0 + s_1)^2]P_2$$

$$T^2 = k_0^2 + [(k_0 + k_1)^2 - k_0^2]P_1 + [(k_0 + k_1 + k_2)^2 - (k_0 + k_1)^2]P_2$$

The equation $T^2 + S^2 = K^2$ is equivalent to:

$$t_0^2 + s_0^2 = k_0^2 \text{ (equation 1),}$$

$$(t_0 + t_1)^2 + (s_0 + s_1)^2 = (k_0 + k_1)^2 \text{ (equation 2),}$$

$$(t_0 + t_1 + t_2)^2 + (s_0 + s_1 + s_2)^2 = (k_0 + k_1 + k_2)^2 \text{ (equation 3),}$$

Equation (1) implies that (t_0, s_0, k_0) is a Pythagoras triple in R .

Equation (2) implies that $(t_0 + t_1, s_0 + s_1, k_0 + k_1)$ is a Pythagoras triple in R .

Equation (3) implies that $(t_0 + t_1 + t_2, s_0 + s_1 + s_2, k_0 + k_1 + k_2)$ is a Pythagoras triple in R .

Thus the proof is complete.

Theorem.

Let $(t_0, s_0, k_0), (t_1, s_1, k_1), (t_2, s_2, k_2)$ be three Pythagoras triples in the ring R , then (T, S, K) Pythagoras triple in $3 - SP_R$, where:

$$T = t_0 + [t_1 - t_0]P_1 + [t_2 - t_1]P_2.$$

$$S = s_0 + [s_1 - s_0]P_1 + [s_2 - s_1]P_2.$$

$$K = k_0 + [k_1 - k_0]P_1 + [k_2 - k_1]P_2.$$

Proof.

We have: $t_0 + (t_1 - t_0) = t_1, t_0 + (t_1 - t_0) + (t_2 - t_1) = t_2.$

$s_0 + (s_1 - s_0) = s_1, s_0 + (s_1 - s_0) + (s_2 - s_1) = s_2,$

$k_0 + (k_1 - k_0) = k_1, k_0 + (k_1 - k_0) + (k_2 - k_1) = k_2.$

This implies that (T, S, K) Pythagoras triple in $2 - SP_R$ according to the theorem.

Examples.

We have:

$$\left\{ \begin{array}{l} (t_0, s_0, k_0) = (3, 4, 5) \\ (t_1, s_1, k_1) = (6, 8, 10) \\ (t_2, s_2, k_2) = (4, 3, 5) \end{array} \right.$$

Are three Pythagoras triples in Z .

The corresponding symbolic 2-plithogenic Pythagoras triple is (T, S, K) , where:

$$T = 3 + [6 - 3]P_1 + [4 - 6]P_2 = 3 + 3P_1 - 2P_2$$

$$S = 4 + [8 - 4]P_1 + [3 - 8]P_2 = 4 + 4P_1 - 5P_2$$

$$K = 5 + [10 - 5]P_1 + [5 - 10]P_2 = 5 + 5P_1 - 5P_2$$

Example.

Find all Pythagoras triples in $2 - SP_{Z_2}$, where Z_2 is the ring of integers module 2.

First, we find all Pythagoras triples in Z .

$$L_1 = (0,0,0), L_2 = (1,0,1), L_3 = (0,1,1), L_4 = (1,1,0)$$

Remark that for every permutation of the set $\{L_1, L_2, L_3, L_4\}$, we get a different symbolic 2-plithogenic Pythagoras triple.

We discuss all possible cases:

Permutation (1).

$$\begin{cases} Y_1 = P_1 - P_2 = P_1 + P_2 \\ \quad \dot{Y}_1 = P_2 \\ \quad Y_1'' = P_1 = P_1 \end{cases}$$

Permutation (2).

$$\begin{cases} Y_2 = P_2 \\ \dot{Y}_2 = P_1 - P_2 = P_1 + P_2 \\ \quad Y_2'' = P_1 = P_1 \end{cases}$$

Permutation (3).

$$\begin{cases} Y_3 = P_1 \\ \dot{Y}_3 = P_1 + P_2 \\ \quad Y_3'' = P_2 \end{cases}$$

Permutation (4).

$$\begin{cases} Y_4 = P_1 + P_2 \\ \quad \dot{Y}_4 = P_1 \\ \quad Y_4'' = P_2 \end{cases}$$

Permutation (5).

$$\begin{cases} Y_5 = P_1 \\ \quad \dot{Y}_5 = P_2 \\ Y_5'' = P_1 + P_2 \end{cases}$$

Permutation (6).

$$\begin{cases} Y_6 = P_2 \\ \quad \dot{Y}_6 = P_1 \\ Y_6'' = P_1 + P_2 \end{cases}$$

Permutation (7).

$$\begin{cases} Y_7 = 1 + P_2 \\ \check{Y}_7 = P_1 \\ Y_7'' = 1 + P_1 + P_2 \end{cases}$$

Permutation (8).

$$\begin{cases} Y_8 = P_2 \\ \check{Y}_8 = 1 + P_1 + P_2 \\ Y_8'' = 1 + P_1 + P_2 \end{cases}$$

Permutation (9).

$$\begin{cases} Y_9 = 1 + P_1 \\ \check{Y}_9 = P_1 + P_2 \\ Y_9'' = 1 + P_2 \end{cases}$$

Permutation (10).

$$: \begin{cases} Y_{10} = P_1 + P_2 \\ \check{Y}_{10} = 1 + P_1 \\ Y_{10}'' = 1 + P_2 \end{cases}$$

Permutation (11).

$$\begin{cases} Y_{11} = 1 + P_2 \\ \check{Y}_{11} = 1 + P_1 \\ Y_{11}'' = P_1 + P_2 \end{cases}$$

Permutation (12).

$$\begin{cases} Y_{12} = 1 + P_1 + P_2 \\ \check{Y}_{12} = 1 + P_1 \\ Y_{12}'' = P_2 \end{cases}$$

Permutation (13).

$$\begin{cases} Y_{13} = 1 + P_1 + P_2 \\ \check{Y}_{13} = 1 + P_2 \\ Y_{13}'' = P_1 \end{cases}$$

Permutation (14).

$$\begin{cases} Y_{14} = 1 + P_1 \\ \check{Y}_{14} = 1 + P_2 \\ Y_{14}'' = P_1 + P_2 \end{cases}$$

Permutation (15).

$$\begin{cases} Y_{15} = 1 + P_1 \\ \check{Y}_{15} = 1 + P_1 + P_2 \\ Y_{15}'' = P_2 \end{cases}$$

Permutation (16).

$$\begin{cases} Y_{16} = 1 + P_2 \\ Y'_{16} = 1 + P_1 + P_2 \\ Y''_{16} = P_1 \end{cases}$$

Permutation (17).

$$\begin{cases} Y_{17} = 1 + P_2 \\ Y'_{17} = P_1 + P_2 \\ Y''_{17} = 1 + P_1 \end{cases}$$

Permutation (18).

$$\begin{cases} Y_{18} = 1 + P_2 \\ Y'_{18} = P_1 \\ Y''_{18} = 1 + P_1 + P_2 \end{cases}$$

Permutation (19).

$$: \begin{cases} Y_{19} = P_1 \\ Y'_{19} = 1 + P_1 + P_2 \\ Y''_{19} = 1 + P_2 \end{cases}$$

Permutation (20).

$$\begin{cases} Y_{20} = P_1 \\ Y'_{20} = 1 + P_1 + P_2 \\ Y''_{20} = 1 + P_2 \end{cases}$$

Permutation (21).

$$: \begin{cases} Y_{21} = P_1 + P_2 \\ Y'_{21} = 1 + P_2 \\ Y''_{21} = 1 + P_1 \end{cases}$$

Permutation (22).

$$\begin{cases} Y_{22} = 1 + P_1 \\ Y'_{22} = P_1 + P_2 \\ Y''_{22} = 1 + P_2 \end{cases}$$

Permutation (23).

$$\begin{cases} Y_{23} = P_1 \\ Y'_{23} = 1 + P_2 \\ Y''_{23} = 1 + P_1 + P_2 \end{cases}$$

Permutation (24).

$$\begin{cases} Y_{24} = P_1 + P_2 \\ Y'_{24} = 1 + P_1 \\ Y''_{24} = 1 + P_2 \end{cases}$$

Also, other quadraples $(L_i, L_j, L_k, L_s); 1 \leq i, j, k, s \leq 4$ give Pythagoras triples with i, j, k, s are not distinct at all.

We continuo our discussions.

Permutation (25).

$$\begin{cases} Y_{25} = (0,0,0) \\ Y'_{25} = (0,0,0) \\ Y''_{25} = (0,0,0) \end{cases}$$

Permutation (26).

$$\begin{cases} Y_{26} = P_1 + P_2 \\ Y'_{26} = 0 \\ Y''_{26} = P_1 + P_2 \end{cases}$$

Permutation (27).

$$\begin{cases} Y_{27} = 0 \\ Y'_{27} = P_1 + P_2 \\ Y''_{27} = P_1 + P_2 \end{cases}$$

Permutation (28).

$$\begin{cases} Y_{28} = P_1 + P_2 \\ Y'_{28} = P_1 + P_2 \\ Y''_{28} = 0 \end{cases}$$

Permutation (29).

$$\begin{cases} Y_{29} = P_2 \\ Y'_{29} = 0 \\ Y''_{29} = P_2 \end{cases}$$

Permutation (30).

$$\begin{cases} Y_{30} = 0 \\ Y'_{30} = P_2 \\ Y''_{30} = P_2 \end{cases}$$

Permutation (31).

$$\begin{cases} Y_{31} = P_2 \\ Y'_{31} = P_2 \\ Y''_{31} = 0 \end{cases}$$

Permutation (32).

$$\begin{cases} Y_{32} = 1 \\ Y'_{32} = 0 \\ Y''_{32} = 1 \end{cases}$$

Permutation (33).

$$: \begin{cases} Y_{33} = 1 + P_2 \\ \check{Y}_{33} = 0 \\ Y_{33}'' = 1 + P_2 \end{cases}$$

Permutation (34).

$$\begin{cases} Y_{34} = 1 + P_2 \\ \check{Y}_{34} = P_2 \\ Y_{34}'' = 1 \end{cases}$$

Permutation (35).

$$\begin{cases} Y_{35} = 1 \\ \check{Y}_{35} = P_2 \\ Y_{35}'' = 1 + P_2 \end{cases}$$

Permutation (36).

$$\begin{cases} Y_{36} = 1 + P_1 \\ \check{Y}_{36} = 0 \\ Y_{36}'' = 1 + P_1 \end{cases}$$

Permutation (37).

$$\begin{cases} Y_{37} = 1 \\ \check{Y}_{37} = P_1 + P_2 \\ Y_{37}'' = 1 + P_2 + P_1 \end{cases}$$

Permutation (38).

$$\begin{cases} Y_{38} = 1 + P_2 \\ \check{Y}_{38} = P_1 + P_2 \\ Y_{38}'' = 1 + P_1 \end{cases}$$

Permutation (39).

$$\begin{cases} Y_{39} = 0 \\ \check{Y}_{39} = 1 \\ Y_{39}'' = 1 \end{cases}$$

Permutation (40).

$$\begin{cases} Y_{40} = P_2 \\ \check{Y}_{40} = 1 + P_2 \\ Y_{40}'' = 1 \end{cases}$$

Permutation (41).

$$\begin{cases} Y_{41} = P_2 \\ \check{Y}_{41} = 1 \\ Y_{41}'' = 1 + P_2 \end{cases}$$

Permutation (42).

$$\begin{cases} Y_{42} = 0 \\ \check{Y}_{42} = 1 + P_1 + P_2 \\ Y_{42}'' = 1 + P_1 + P_2 \end{cases}$$

Permutation (43).

$$\begin{cases} Y_{43} = P_1 + P_2 \\ \check{Y}_{43} = 1 + P_1 + P_2 \\ Y_{43}'' = 1 \end{cases}$$

Permutation (44).

$$\begin{cases} Y_{44} = P_1 + P_2 \\ \check{Y}_{44} = 1 \\ Y_{44}'' = 1 + P_1 + P_2 \end{cases}$$

Permutation (45).

$$\begin{cases} Y_{45} = 1 \\ \check{Y}_{45} = 1 \\ Y_{45}'' = 0 \end{cases}$$

Permutation (46).

$$\begin{cases} Y_{46} = 1 + P_2 \\ \check{Y}_{46} = 1 + P_2 \\ Y_{46}'' = 0 \end{cases}$$

Permutation (47).

$$\begin{cases} Y_{47} = 1 \\ \check{Y}_{47} = 1 + P_2 \\ Y_{47}'' = P_2 \end{cases}$$

Permutation (48).

$$\begin{cases} Y_{48} = 1 + P_2 \\ \check{Y}_{48} = 1 \\ Y_{48}'' = P_2 \end{cases}$$

Permutation (49).

$$\begin{cases} Y_{49} = 1 + P_1 + P_2 \\ \check{Y}_{49} = 1 + P_1 + P_2 \\ Y_{49}'' = 0 \end{cases}$$

Permutation (50).

$$\begin{cases} Y_{50} = 1 \\ \check{Y}_{50} = 1 + P_1 + P_2 \\ Y_{50}'' = P_1 + P_2 \end{cases}$$

Permutation (51).

$$\begin{cases} Y_{51} = 1 + P_1 + P_2 \\ Y_{51}' = 1 \\ Y_{51}'' = P_1 + P_2 \end{cases}$$

By continuing this argument, we can get all Pythagoras triples in $2 - SP_{Z_2}$

Conclusion.

In this paper, we have studied Pythagoras triples in symbolic 2-plithogenic commutative rings, where necessary and sufficient conditions for a symbolic 2-plithogenic triple (x, y, z) to be a Pythagoras triple.

Also, we have presented some related examples that explain how to find 2-plithogenic triples from classical triples.

Acknowledgments " This study is supported via funding from Prince sattam bin Abdulaziz University project number (PSAU/2023/R/1445) ".

References

- [1] M. B. Zeina, N. Altounji, M. Abobala, and Y. Karmouta, "Introduction to Symbolic 2-Plithogenic Probability Theory," *Galoitica: Journal of Mathematical Structures and Applications*, vol. 7, no. 1, 2023.
- [2] Merkepci, H., and Abobala, M., " On The Symbolic 2-Plithogenic Rings", *International Journal of Neutrosophic Science*, 2023.
- [3] Nader Mahmoud Taffach , Ahmed Hatip., "A Review on Symbolic 2-Plithogenic Algebraic Structures " *Galoitica Journal Of Mathematical Structures and Applications*, Vol.5, 2023.
- [4] Rawashdeh, A., "An Introduction To The Symbolic 3-plithogenic Number Theory", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [5] Ben Othman, K., "On Some Algorithms For Solving Symbolic 3-Plithogenic Equations", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [6] Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Concept Of Symbolic 7-Plithogenic Real Matrices", *Pure Mathematics For Theoretical Computer Science*, Vol.1, 2023.

- [7] Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Symbolic 8-Plithogenic Matrices", Pure Mathematics For Theoretical Computer Science, Vol.1, 2023.
- [8] Mohamed, M., "Modified Approach for Optimization of Real-Life Transportation Environment: Suggested Modifications", American Journal of Business and Operations research, 2021.
- [9] Nabeeh, N., Alshaimaa, A., and Tantawy, A., "A Neutrosophic Proposed Model For Evaluation Blockchain Technology in Secure Enterprise Distributed Applications", Journal of Cybersecurity and Information Management, 2023.
- [10] Abualkishik, A., Almajed, R., Thompson, W., "Improving The Performance of Fog-assisted Internet of Things Networks Using Bipolar Trapezoidal Neutrosophic Sets", Journal of Wireless and Ad Hoc Communication, 2023.
- [11] R. A. Trivedi, S. A. Bhanotar, Pythagorean Triplets - Views, Analysis and Classification, IIOSR J. Math., 11 (2015).
- [12] J. Rukavicka, Dickson's method for generating Pythagorean triples revisited, Eur. J. Pure Appl. Math., 6 (2013).
- [13] T. Roy and F. J. Sonia, A Direct Method To Generate Pythagorean Triples And Its Generalization To Pythagorean Quadruples And n-tuples, , Jadavpur University, India (2010)
- [14] Paul Oliverio, " Self-Generating Pythagorean Quadruples And TV-Tuples", Los Angeles, (1993)

Received 3/6/2023, Accepted 18/9/2023