

11-18-2009

Unraveling the intricacies of spatial organization of the ErbB receptors and downstream signaling pathways

Michelle Costa

Follow this and additional works at: https://digitalrepository.unm.edu/cbe_etds

Recommended Citation

Costa, Michelle. "Unraveling the intricacies of spatial organization of the ErbB receptors and downstream signaling pathways." (2009). https://digitalrepository.unm.edu/cbe_etds/31

This Dissertation is brought to you for free and open access by the Engineering ETDs at UNM Digital Repository. It has been accepted for inclusion in Chemical and Biological Engineering ETDs by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.

Michelle N. Costa


Candidate

Chemical and Nuclear Engineering

Department


This dissertation is approved, and it is acceptable in quality and form for publication:

Approved by the Dissertation Committee:



, Chairperson

Pedro T. Embri



**UNRAVELING THE INTRICACIES OF SPATIAL
ORGANIZATION OF THE ERBB RECEPTORS AND
DOWNSTREAM SIGNALING PATHWAYS**

BY

MICHELLE N. COSTA

B.S., Chemical Engineering, University of New Mexico, 2006

DISSERTATION

Submitted in Partial Fulfillment of the
Requirements for the Degree of

**Doctor of Philosophy
Engineering**

The University of New Mexico
Albuquerque, New Mexico

August 2009

© 2009, Michelle N. Costa

Dedication

To the memory of

Thelma I. Salley

And

Joe Costa Jr

Acknowledgments

This work would not be possible without the love and support of my family. I would like to thank my parents Dan and Linda Costa whose love is unconditional and are always by my side through thick and thin. I would like to thank my brother Ryan Costa for his love and encouragement. I would like to thank my grandparents Bobby and Thelma Salley who always loved and believed in me. I would like to thank my fiancé Luis Archuleta who helped me in so many ways in graduate school. I would like to thank him for his love and support.

I would like to thank my advisors Jeremy Edwards, Chi-Chi May, Dion Vlachos, and Bridget Wilson. I would like to thank my committee members Jeremy Edwards, Chi-Chi May, Pedro Embid, and Dimiter Petsev. I would like to thank all of the wonderful people who supported me along the way Norah Torrez-Martinez, Purushart Prakash, and Chi-Chi May. I would like to thank my funding INCBN IGERT. This work was supported in part by the University of New Mexico Center for Advanced Research Computing and utilized the *ristra* and *nano* supercomputers. I would like to thank Florencia Sanchez for helping prepare this dissertation.

**UNRAVELING THE INTRICACIES OF SPATIAL
ORGANIZATION OF THE ERBB RECEPTORS AND
DOWNSTREAM SIGNALING PATHWAYS**

BY

MICHELLE N. COSTA

ABSTRACT OF DISSERTATION

Submitted in Partial Fulfillment of the
Requirements for the Degree of

**DOCTOR OF PHILOSOPHY
ENGINEERING**

The University of New Mexico
Albuquerque, New Mexico

August 2009

Unraveling the Intricacies of Spatial Organization of the ErbB Receptors and Downstream Signaling Pathways

Michelle N. Costa

B.S Chemical Engineering University of New Mexico-Albuquerque, 2006

Ph.D. Engineering University of New Mexico, 2009

Abstract

Faced with the complexity of diseases such as cancer which has 10^{12} mutations, altering gene expression, and disrupting regulatory networks, there has been a paradigm shift in the biological sciences and what has emerged is a much more quantitative field of biology. Mathematical modeling can aid in biological discovery with the development of predictive models that provide future direction for experimentalist. In this work, I have contributed to the development of novel computational approaches which explore mechanisms of receptor aggregation and predict the effects of downstream signaling. The coupled spatial non-spatial simulation algorithm, CSNSA is a tool that I took part in developing, which implements a spatial kinetic Monte Carlo for capturing receptor interactions on the cell membrane with Gillespie's stochastic simulation algorithm, SSA, for temporal cytosolic interactions. Using this framework we determine that receptor clustering significantly enhances downstream signaling.

In the next study the goal was to understand mechanisms of clustering. Cytoskeletal interactions with mobile proteins are known to hinder diffusion. Using a Monte Carlo approach we simulate these interactions, determining at what cytoskeletal distribution and receptor concentration optimal clustering occurs and when it is inhibited. We investigate oligomerization induced trapping to determine mechanisms of clustering, and our results show that the cytoskeletal interactions lead to receptor clustering. After exploring the mechanisms of clustering we determine how receptor aggregation effects downstream signaling. We further proceed by implementing the adaptively coarse grained Monte Carlo, ACGMC to determine if “receptor-sharing” occurs when receptors are clustered. In our proposed “receptor-sharing” mechanism a cytosolic species binds with a receptor then disassociates and rebinds a neighboring receptor. We tested our hypothesis using a novel computational approach, the ACGMC, an algorithm which enables the spatial temporal evolution of the system in three dimensions by using a coarse graining approach. In this framework we are modeling EGFR reaction-diffusion events on the plasma membrane while capturing the spatial-temporal dynamics of proteins in the cytosol. From this framework we observe “receptor-sharing” which may be an important mechanism in the regulation and overall efficiency of signal transduction. In summary, I have helped to develop predictive computational tools that take systems biology in a new direction.

Table of Contents

Dedication	iv
Acknowledgements.....	v
Abstract.....	vii
Table of Figures	xv
Table of Tables	xvii
Chapter 1. Spatial Stochastic Modeling: A new way of dissecting biology	1
Abstract	3
Introduction	4
Quantitative Spatial Experimental Approaches in Biology	7
Fluorescence Recovery After Photobleaching (FRAP)	7
Single Particle Tracking (SPT)	8
Forster Resonance Energy Transfer (FRET)	10
Electron Microscopy (EM).....	10
Transmission Electron Microscopy (TEM)	11
Scanning Electron Microscopy (SEM)	12
Discoveries from Quantified Experiments.....	12
Paradigm Shift in Membrane Biology	12
Lipid Rafts.....	13
Cytoskeletal Interactions: Picket-Fence Model.....	15
Moving Ahead in Membrane Biology	16
Modeling Approaches	17
Cornerstones of Modeling Techniques	20

Choosing an Appropriate Method.....	24
Spatial-Temporal Stochastic Methods.....	26
Sorting Out Microdomains: An example using the SKMC.....	32
A New Direction for Systems Biology.....	35

Chapter 2. Coupled Stochastic Spatial and Non-Spatial simulations of ErbB1

Signaling Pathways Demonstrate the Importance of Spatial Organization in Signal

Transduction	37
Abstract	38
Background	38
Methodology/Principal Findings.....	38
Conclusions/Significance	38
Introduction	40
Results	43
Establishing Parameters for the Spatial Model	43
Validating the CSNSA hybrid approach	46
Predicting the Impact of Receptor Density vs. Clustering.....	49
Discussion	51
Methods	53
Coupled Spatial, Non-spatial Simulation Algorithm (CSNSA)	53
Spatial Kinetic Monte Carlo (SKMC)	59
Stochastic Simulation Algorithm (SSA)	61
Interfacial Reactions.....	61
Sensitivity Analysis.....	63
Acknowledgements	64

Chapter 3. Picket Fence Densities Determine Enhanced vs. Inhibitive Role in	
Receptor Aggregation	65
Abstract	66
Introduction	67
Materials and Methods	69
Spatial Kinetic Monte Carlo (SKMC)	69
Picket Fences.....	74
Quantification of Microdomains	78
Results	79
Clustering vs. Picket Fence Density.....	83
Clustering vs. Receptor Concentration	85
Mechanism of Clustering: Oligomerization Induced Trapping	89
Predimerization vs. Dimerization.....	91
Discussion.....	93
Acknowledgements.....	95

Chapter 4. Adaptively Coarse Grained Monte Carlo Method for Capturing the	
“Receptor–Sharing” Mechanism	96
Abstract	97
Introduction	98
Results	101
Cell Signaling Events are Dependent on Spatial Localization.....	101
Evidence of a Concentration Gradient	103
The “Receptor-Sharing” Mechanism	105
Quantifying the “Receptor-Sharing” Mechanism	108
Inhibiting the “Receptor-Sharing” Mechanism.....	112
Discussion	113
Materials and Methods	115
Adaptively Coarse-Grained Monte Carlo (ACGMC)	
Adaptively Coarse-Graining & Diffusion Transition Rates.....	119
Validation of Approach.....	121

Chapter 5. Future Work	139
Introduction	141
Computational Improvements	141
Computational Predictions	146
Understanding the impact of IFN γ R clustering on immune response pathways	146
A Monte Carlo Based Approach for Determining Optimal Drug Efficacy in Different Cytoskeletal Distributions	149
Microdomains and the Underlying Cytoskeleton Alter the Efficiency of “Receptor-Sharing”	151
Conclusion	154
 Appendix	156
Appendix A. The effective kinetic rate constant for diffusion-reaction problems derived by Lauffenburger and Linderman	157
Appendix B. Understanding the impact of IFN γ R clustering on immune response pathways.....	158
Appendix C. A Monte Carlo Based Approach for Determining Optimal Drug Efficacy in Different Cytoskeletal Distributions	160
Appendix D. Microdomains and the Underlying Cytoskeleton Alter the Efficiency of “Receptor-Sharing”	194
Appendix E. Abbreviations Used.....	210
 References	213

Table of Figures

Figure 1	21
Figure 2	24
Figure 3	34
Figure 4	45
Figure 5	47
Figure 6	48
Figure 7	50
Figure 8	57
Figure 9	58
Figure 10	62
Figure 11	75
Figure 12	77
Figure 13	80
Figure 14	82
Figure 15	84
Figure 16	87
Figure 17	88
Figure 18	90
Figure 19	92
Figure 20	102
Figure 21	104
Figure 22	106

Figure 23 107

Figure 24 109

Figure 25 111

Figure 26 118

Table of Tables

Table 1	44
Table 2	56
Table 3	71
Table 4	73
Table 5	123

Chapter1. Spatial Stochastic Modeling: A new way of dissecting biology

Michelle Costa¹, Bridget Wilson^{2,3}, Dionisios G. Vlachos⁴, & Jeremy S. Edwards^{1,3,4,*}

¹ Department of Chemical and Nuclear Engineering, University of New Mexico,
Albuquerque, NM 87131

² Department of Pathology, University of New Mexico Health Sciences Center
Albuquerque, NM 87131

³ Cancer Research and Treatment Center, University of New Mexico Health Sciences
Center, Albuquerque, NM 87131

⁴ Department of Chemical Engineering, University of Delaware, Newark, DE 19716

⁵ Molecular Genetics and Microbiology, University of New Mexico Health Sciences
Center, Albuquerque, NM 87131

Abstract

The field of membrane biology has undergone a paradigm shift and in the process shaken the foundation of mathematical models built under the well mixed assumption. Advances in technology have led us to a new frontier in the biological sciences with techniques such as SPT, FRET, and EM at our fingertips the possibilities are endless. In order to complement experimental innovation, novel computational platforms must be developed which accurately describe discrete, stochastic, spatial heterogeneity observed in biological systems. The traditional method for modeling signal transduction pathways is using ordinary differential equations (ODEs). ODEs are deterministic, continuum based models that inherently assume spatial homogeneity. However, it is becoming clear that spatial modeling tools will be needed to fully understand the complexities of signaling pathways. In this review we will begin by discussing advancements made in biology through the development of mathematical models and the experiments that gave them their birth. We will go beyond these initial experiments and discuss the new cutting edge experimental techniques which brought about an evolution to the membrane biology community. We will then discuss computational methods; their advantageous and drawbacks. We then propose novel computational platforms which must evolve in order to complement experiments and bring predictability back to systems biology.

Keywords: Mathematical modeling, signal transduction, plasma membrane, ODE, PDE, spatial modeling, Spatial Monte Carlo

Introduction

The physical sciences have always been interwoven with mathematics, through a foundation built equally from theory and experimentation. The life sciences, in contrast, have been a much more observatory science, relying more on description rather than theory and quantification. However, being confronted with complex diseases has altered this approach; cancer, for example, can have as many as 10^{12} mutations (Tomlinson, Novelli et al. 1996; Tomlinson, Sasieni et al. 2002; d'Onofrio and Tomlinson 2007), altering gene expression, and disrupting regulatory networks. This complexity has driven a paradigm shift in biology, complementing the traditional reductionist approach with a systems level approach which builds upon theory. A novel science, quantitative biology or systems biology, has emerged as a result; quantitative biology or systems biology derives principles from fields such as mathematics, physics, chemistry and engineering in order to develop predictive models of biological systems.

The need to develop novel mathematics (Faeder, Blinov et al. 2005; Mayawala, Vlachos et al. 2005; Chatterjee and Vlachos 2006; Mayawala, Vlachos et al. 2006; Borisov, Chistopolsky et al. 2008; Collins, Chatterjee et al. 2008; Hsieh, Yang et al. 2008) and acquire quantified experimental data (Janes, Albeck et al. 2003; Conzelmann, Saez-Rodriguez et al. 2006; Kumar, Zaman et al. 2006; Aksamitiene, Hoek et al. 2007; Wilson, Pfeiffer et al. 2007; Andrews, Lidke et al. 2008) has come full circle with experimentation being done to compliment theoretical models (Janes, Albeck et al. 2005; Birtwistle, Hatakeyama et al. 2007; Hsieh, Yang et al. 2008; Kumar, Afeyan et al. 2008) and models guiding further hypotheses. Developing a

mathematical model of a cell signaling pathway involves detailed experimentation. For example, cells are stimulated, lysed at various time points, and target proteins are immunoprecipitated. This data provides initial protein concentration as well as dynamic data which are crucial to building a mathematical model of the system. The model in return, plays a predictive role, guiding experimentation, by testing hypothesis which may be infeasible experimentally due to poor cell viability, lack of current technologies, cost, and time.

Over the past decade many mathematical models have been developed, to sort out complex cell signaling networks; an example of this is taken from the mitogen activated protein kinase, MAPK pathway. Intensively studied, the MAPK pathway regulates cellular functions such as proliferation, differentiation, apoptosis, adhesion and migration. The first mathematical model describing this pathway (Huang 1996) was an ordinary differential equation (ODE) model containing 35 reactions. It provided insight into the sensitivity of this system. In 1999, using an ODE approach, Kholodenko et al. published an EGFR signaling model. Although it did not exceed the Huang model in number of reactions, it contained signaling events upstream at the cell membrane where signaling cascades are initiated. In 2002, Schoeberl et al. developed a much more extensive ODE model, containing 125 reactions with 94 species, which spans receptor activation at the cell membrane to ERK phosphorylation and includes receptor degradation and internalization. In 2005, Sasagawa et al. published a much more complex ODE model that linked two receptors to activation of the MAPK pathway. A year later Kiyatkin et al. published a model incorporating additional pathways, such as the GAB1 and Src; it was yet

another model exploding in complexity but built on the well-mixed assumption inherent to ODEs.

What has become clear from the work of these original ODE models is that cell signaling is very complex. Although mathematical models have grown exponentially, ODE have several limitations. Proteins are distinct entities, making signal propagation a discrete process, which is not captured by continuum ODE's. Experimental evidence shows that signaling receptors are not well mixed on the plasma membrane but rather confined to enrichment and depletion zones. Therefore, overall reaction rates for species on the plasma membrane are limited by diffusion which is not taken into account with ODEs. The limitations of ODE models are calling for a new theoretical framework, a spatial-temporal approach which will more accurately describe the biology. In this review we describe a new generation of experimentation and its impact on biology. Then we discuss the evolution of mathematical models, from the ODE to spatial Monte Carlo methods, focusing on what theoretical frameworks must be developed to complement novel experimentation.

Quantitative Spatial Experimental Approaches in Biology

Systems biology relies on the ability to make quantitative measurements. Quantitative measurements of biological systems have been a challenge, and in this section we will summarize the critical experimental techniques for studying the spatial-temporal dynamics of cell membranes: fluorescence recovery after photobleaching (FRAP), single particle tracking (SPT), Forster resonance energy transfer (FRET), and electron microscopy (EM) focusing on how they facilitate systems biology. Then we will talk about the paradigm shift which has occurred in membrane biology as a result of these techniques; highlighting the discoveries of microdomains such as lipid rafts and picket fences.

Fluorescence Recovery After Photobleaching (FRAP)

One of the first methods developed to study the dynamics of proteins in living cells, FRAP has resurfaced due to techniques such as confocal-microscope-based photo-bleaching, and the discovery of GFP (Patterson and Lippincott-Schwartz 2002; Lippincott-Schwartz, Altan-Bonnet et al. 2003). FRAP relies on the fact that extended excitation of a fluorescent protein or molecule leads to photobleaching. Therefore, when a laser is directed to a small region of the cell membrane, photobleaching of this region occurs. The surrounding molecules, not subjugated to these repetitive cycles, will diffuse into the photobleached area. This recovery of fluorescent signal is recorded over time with a low-intensity laser light, making it possible to obtain information about the kinetic parameters, such as diffusion coefficient, mobile fraction, and binding dissociation rate constants (Lippincott-Schwartz, Altan-Bonnet et al. 2003).

FRAP is a powerful technique and has provided estimates of protein diffusion coefficients. However, one of the drawbacks of FRAP is its poor spatial resolution, which makes it difficult to interpret the data. Due to the complexities of the plasma membrane, a variety of mechanisms may determine diffusivity such as obstruction, transient binding, confinement, and hydrodynamic interactions (Saxton and Jacobson 1997). A method to overcome these shortcomings is SPT, which will be reviewed in the following section.

Single Particle Tracking (SPT)

A methodology for deciphering the dynamics of membrane proteins, SPT, enables individual trajectories of molecules and multimolecular complexes to be resolved (Saxton and Jacobson 1997; Kusumi, Ike et al. 2005; Bates, Wiseman et al. 2006). SPT involves labeling proteins with a probe (bead, fluorophore, gold particle, or q-dot), imaging and then tracking the centroid of the imaged probe over time (Bates, Wiseman et al. 2006).

Observables, such as molecular trajectories, translational diffusion in different cell areas, and access to different modes of motion can be determined using SPT (Marguet, Lenne et al. 2006). Given an individual trajectory, it is possible to determine the mode of motion which could be immobile, directed, confined, tethered, normal diffusion, and anomalous diffusion (Saxton and Jacobson 1997). The mode of motion is classified by looking at the time dependence of the mean-square displacement (MSD), (Saxton and Jacobson 1997; Kusumi, Nakada et al. 2005; Bates, Wiseman et al. 2006). The capability to determine the mode of motion makes SPT a very powerful tool, providing insight into the dynamics of signaling receptors.

There are various methods for imaging individual proteins, all of which rely on attaching a moiety to the protein of interest. For example, colloidal gold can be used as a probe due to its strong light scattering ability. Once contrast enhancement and background subtraction are performed, the label is darker than its surroundings (Saxton and Jacobson 1997). However, one of the major concerns with gold particles (size ~30-40 nm diameter) is that may decrease the diffusion rate of the protein (Bates, Wiseman et al. 2006). Multivalency is another issue, which may cause clustering and underestimation of the diffusion coefficient (Kusumi, Nakada et al. 2005). Additionally, researchers have used Fluorescence (both traditional fluors and Q-Dots) SPT. Organic fluorophores are another popular probe because they are less susceptible to cross-linking. One of their limitations, however, is that they are susceptible to photobleaching thus limiting their use to very short observation times (Bates, Wiseman et al. 2006). The semiconductor quantum dot has overcome these limitations, making it the preferred probe. With its novel properties such as improved photostability, optical tenability, and multicolor light emission it has taken live cell imaging and in vivo animal models to a new realm of optical resolution (Giepmans, Deerinck et al. 2005; Giepmans, Adams et al. 2006; Smith, Ruan et al. 2006).

Forster Resonance Energy Transfer (FRET)

FRET is an incredibly powerful technology going beyond traditional immunoassays to reveal direct protein-protein interactions. FRET works when a donor in close proximity ($<10\text{nm}$) and with preferred orientation brings about energy transfer which in turn induces emission from the acceptor. FRET donors and acceptors are chosen on the premise that there is an ($> 30\%$) overlap (Sekar and Periasamy 2003) between the emission spectrum of the donor and excitation spectrum of the acceptor. Microscopy techniques capture FRET, when the donor channel signal is quenched and the acceptor channel signal is sensitized. The transfer of energy between an acceptor and a donor can be measured by microscopy and thus spatial organization of two molecules relative to one another can be measured.

Electron Microscopy (EM)

Great advances have been made in biology due to imaging; electron microscopy stands at the cornerstone of these advancements. This technique provides a much greater magnification (10^4) than the traditional light microscopes due to the electron's small de Broglie wavelength. Different types of microscopy have been developed we will be discussing those of most biological importance.

Transmission Electron Microscopy (TEM)

TEM utilizes a high voltage electron beam which is focused on an ultra thin Specimen. As the beam passes through, an image is created which is then magnified and focused(Egerton 2005). An electron gun cases a tungsten filament cathode, the electron source; once a voltage of 40 to 400 keV is applied to the cathode, an electron beam propagates towards the sample. This technique applies a magnification between 1,000X to 1,000,000X, but it is limited by the density of the specimen due to scattering.

Viewing a single atom was beyond the reach of the traditional TEM in the 1970's, when Crewe et al. developed the scanning transmission electron microscope (STEM) capable of viewing bright spots that were arguably individual atoms(Crewe and Wall 1970; Crewe, Wall et al. 1970). STEM builds on the foundation of TEM, by focusing the electron beam on a narrow strip which is scanned over the raster. The data collected can be analyzed with methods such as energy dispersive X-ray (EDX) spectroscopy, electron energy loss spectroscopy (EELS) and annular dark-field imaging (ADF) which allow direct correlation of image. STEM allows, a high contrast image of biological samples without staining due to its use of dark-field microscopy. STEM has been applied to biological problems to resolve and distinguish structure on the molecular level.

Scanning Electron Microscopy (SEM)

SEM is another electron microscopy technique that scans a focused electron beam across the area of the specimen. Scanning produces a mixed array of signal such as cathodoluminescence, back scattered electrons, characteristic x-rays, secondary electrons specimen current, and transmitted electrons which once analyzed give a large depth of field yielding a characteristic three-dimensional appearance.

Discoveries from Quantified Experiments

The technologies presented in the previous section have moved us into the next frontier of biological discoveries. Studying cellular behavior in vivo for example, has lead to novel insights; from the cell membrane with diffusivity, stoichiometry, and protein translation being deciphered, to the cytosol/nucleus with transcription factor binding, to transport through the nuclear pore complex, motor proteins on linear tracks, to reveal new spatial-temporal dynamics (Joo, Balci et al. 2008). In this section we focus on membrane biology, which has been transformed by the previously discussed experimental techniques (Kusumi, Nakada et al. 2005).

Paradigm Shift in Membrane Biology

The original fluid mosaic model portrays protein movement as Brownian motion in a sea of lipids. Although this model is the foundation of the membrane biology literature, it has been plagued by two inconsistencies. The first discrepancy being that the diffusion coefficient for both proteins and lipids is a factor of 5 to 50 times smaller in the plasma membrane than within artificial membranes. The second discrepancy was that oligomers or molecular complexes exhibit a much lower diffusion coefficient (factor of 40) in the plasma membrane in comparison with

artificial membranes. These inadequacies of the original fluid mosaic model were indicative of an alternative model and have led researchers on a 30 year journey to uncover the true nature of the plasma membrane.

Lipid Rafts

The first indication of discrete microdomains within the cell membrane came a year after Singer and Nicholson (1972)(Singer and Nicolson 1972) proposed the fluid mosaic model, when a study by Yu et al. (Yu, Fischman et al. 1973) demonstrated the existence of detergent-resistant sphingolipid-rich domains in erythrocyte's plasma membrane. This was followed up in the 1980's with studies by van Meer et al.(van Meer, Poorthuis et al. 1980; van Meer, Stelzer et al. 1987) in erythrocytes; they showed asymmetry in the distribution of phospholipids throughout the plasma membrane. In the early 90's Lisanti et al. (Lisanti, Caras et al. 1989; Lisanti, Caras et al. 1991; Lisanti and Rodriguez-Boulan 1991; Hannan, Lisanti et al. 1993) used fluorescence microscopy to link sphingolipids and GPI (glycosylphosphatidylinositol)-anchored proteins. In addition, an isolation method, detergent extraction was developed (Brown and Rose 1992) to separate these sphingolipid-rich microdomains.

First observed in the 70's, lipid microdomains have been given various names such as detergent-resistant membranes (DRMs), and detergent-insoluble glycolipid-enriched membrane domains (DIGs) (Jacobson and Dietrich 1999), but the name which has stuck with these complex structures is 'lipid rafts'. The idea of lipid rafts has stirred both excitement as well as controversy in the biological community; the evidence of their existence is only suggestive and their biological role is divisive.

Some researchers speculate that lipid rafts could be ‘vanishingly small’ and may hold no biological significance (Edidin 2001; Kovbasnjuk, Edidin et al. 2001), while others hypothesize that lipid rafts may be hubs of signal transduction (Kabouridis and Jury 2008). While caveats remain in the experimental evidence both in support as well as opposition, data suggest a role for lipid rafts in both health: regulation of signaling pathways (Maffucci, Brancaccio et al. 2003), transport of substrates (Saltiel and Kahn 2001), and uptake of LCFAs (long-chain fatty acids) into adipose tissue (Pohl, Ring et al. 2004; Vial and Evans 2005), as well as disease: cardiovascular disease (Zuo, Ushio-Fukai et al. 2005; Maguy, Hebert et al. 2006) (angiotensin II receptor, activated G-proteins, adrenergic, and, cholinergic colocalization in lipid rafts), carcinogenesis (Nagy, Vereb et al. 2002; Mocanu, Fazekas et al. 2005; Yang, Raymond-Stintz et al. 2007) (delayed catalyses of ceramide synthesis decrease apoptosis), immune response (Oliver, Pfeiffer et al. 2004; Wilson, Steinberg et al. 2004) (alteration in lipid raft composition affect localization of immunogenic receptors), and neurological diseases (Park, Hwang et al. 2003; Abad-Rodriguez, Ledesma et al. 2004) (accumulation of Ab (amyloid b-peptide) in lipid rafts); for an interesting review see (Michel and Bakovic 2007).

Cytoskeletal Interactions: Picket-Fence Model

Fujiwara et al. developed powerful microscopy techniques and tools with the goal of visualizing membrane microdomains or lipid rafts. However, they made an unexpected observation. When a lipid probe, DOPE, was measured at high time resolution, it underwent short-term confined diffusion followed by a long-term “hop diffusion”(Fujiwara, Ritchie et al. 2002). These results shed light on the thirty-year-old enigma; decreased diffusion on the plasma membrane is the result of a highly compartmentalized plasma membrane in which the diffusivity of proteins is greatly hindered.

Clues for compartmentalization due to cytoskeletal interactions were evident before the single-particle tracking era, when FRAP experiments observed an increase in diffusivity for blebbed membranes after partial depolymerization of actin filaments (Tank, Wu et al. 1982; Wu, Tank et al. 1982; Paller 1994). Investigating spectrin-deficient mutant mouse erythrocytes, Sheetz et al. (Sheetz, Schindler et al. 1980)observed increased diffusivity (10-fold) in comparison to the normal cell type. In an effort to explain these findings Sheetz et al.(Sheetz 1983) proposed the membrane skeleton “picket fence” model, which describes the cell membrane as a landscape of boundaries on the cytoplasmic surface (due to interactions with the membrane cytoskeleton) and protrusions (“pickets”) expanding into the extracellular surface.

The picket fence model was expanded by the work of Fujiwara et al.(Fujiwara, Ritchie et al. 2002) and Murase et al.(Murase, Fujiwara et al. 2004), by examining the involvement of the membrane skeleton, extracellular matrices,

extracellular domains of membrane proteins, and cholesterol-rich raft domains on phospholipid diffusion. They noted altered diffusion when the membrane skeleton was disrupted. Their results seemed counterintuitive, namely that a phospholipid on the extracellular leaflet of the plasma membrane would be affected by the membrane skeleton which hinders diffusion on the cytoplasmic leaflet. They reconciled this observation by expanding the picket fence model, suggesting that transmembrane proteins are confined by the membrane skeleton and thus hinder phospholipid diffusion on the extracellular face. These transmembrane proteins were termed “pickets” and adding on to the cytoskeletal membrane model now referenced as the “picket-fence” model.

The “picket-fence” model not only resolved the diffusion coefficient inconsistency of the fluid mosaic model, but it also explained oligomerization induced trapping, the phenomenon of decreased diffusion in oligomers observed on the membrane. Many signal transduction pathways are initiated by monomers forming dimers or higher order oligomers. When oligomers are formed the diffusivity decreases due to the additional hindrance endured by a larger molecular complex transversing through the membrane cytoskeleton mesh work. The mechanism of oligomerization induced trapping can enhance the local receptor concentration within a corral and form a “hub” for signal transduction.

Moving Ahead in Membrane Biology

In the midst of a sea of lipids, a landscape of complexity arises, ranging from the coalescences of phospholipids into “lipid rafts” to the hindering interactions caused by the cytoskeleton. This thirty year journey has enlighten the membrane

biology community and provides a new direction for systems biology as a whole. What has been established is that the plasma membrane is a highly compartmentalized surface, which affects the diffusivity of signaling proteins in lipids in the membrane, and hence the initiation and activation of signal transduction pathways. These developments highlight a need for computational algorithms that take into account the biological complexity occurring at the cell membrane.

In this next section we will discuss the following mathematical approaches: temporal deterministic, temporal stochastic, spatial-temporal deterministic and spatial-temporal stochastic and for what systems they are best suited. We will then go into greater detail with the spatial stochastic approach given that it most accurately describes the unique conditions on the plasma membrane. We will then provide an example using the spatial kinetic Monte Carlo to model the initial events of a signaling pathway on the plasma membrane.

Modeling Approaches

Mathematical modeling is an important tool in systems biology. When developing a mathematical model of a biological system it is extremely important that the modeling assumptions are consistent with the system of interest. For example, an inherent assumption in ODE models is that the system is well-mixed, or in other words there are no concentration gradients or spatial heterogeneities in the system. Additionally, ODE models also assume that the species being modeled have large populations acting as a continuum, whereas, small populations require model approaches that handle discrete entities and the inherent stochasticity.

Limits placed on bimolecular reactions span the scale of diffusional to reactional. On the far right are reaction limited conditions inherent for well mixed systems where diffusional effects are negligible. On the far left are diffusion limited conditions with the innate assumption that collision equates with reaction. Such a range of limitations has succumbed to theory born in generality but molded by the assumptions describing the system. Take for example, Smoluchowski seminal work, (Smoluchowski 1917) derived under diffusion limitations; it predicts the evolution of a system of clusters which coalesce forming larger clusters. The Smoluchowski coagulation equation is shown below

$$\frac{d}{dt}n(k,t) = \frac{1}{2} \sum_{j=1}^{k-1} K(j,k-j)n(j,t)n(k-j,t) - n(k,t) \sum_{j=1}^{\infty} K(j,k)n(j,t)$$

where $n(k,t)$ is the concentration of clusters of size k at time t and K is the coagulation kernel. This equation only takes into account coalescence, and speaking in biomolecular reactions terms signifies the far left, collision is reaction. Within the gap of diffusion-limited and reaction-limited conditions falls mathematics which considers both diffusion and reaction. These modeling techniques will be discussed in greater detail in the next section.

Variation of system dynamics could be described either as continuous and predictive, or discrete and random. Mathematics encapsulates variation in dynamics ranging from deterministic approaches with exact solutions, to stochastic approaches with probabilistic solutions. Take for example the stochastic simulation algorithm (SSA), developed by Dan Gillespie in 1977 (Gillespie 1977). The SSA is a formulation of the exact numerical simulation for the underlying master equation. This stochastic temporal method describes a chemical reaction as the “reaction

probability per unit time” (Gillespie 1977) instead of the continuum, deterministic definition of “reaction rate”. The unique feature of the SSA is that the reaction transition rates are computed in terms of total number of molecules rather than numbers of molecules per unit volume as is the standard approach with ODEs. Using total number of molecules, reactants are modeled as discrete entities undergoing reaction events.

Each reaction has a propensity such that there exists a constant c_1 which is related to the deterministic chemical rate constant k_1 by $c_1 = \frac{k_1}{V}$. Multiplying c_1 by the number of reactants gives a transition rate for that reaction; summing transition rates yields a total transition rate. A probability for a event is calculated by taking the transition rate for an individual reaction and dividing it by the total transition rate. Generating a pseudo random number that falls within this calculated probability enables selection of an event and gives the SSA stochasticity.

Cornerstones of Modeling Techniques

Two themes emerge from the previous discussion biomolecular reactions in the presence or absence of diffusional limitations and stochasticity vs deterministcity. Figure 1. shows mathematical techniques that vary on the y-axis from stochastic to deterministic and on the x-axis from methods that take into account diffusion and those that neglect diffusion. This graph defines four approaches temporal deterministic, temporal stochastic, spatial-temporal deterministic, and spatial-temporal stochastic. We have placed mathematical techniques into each category; some techniques span the gray area between two approaches. We will discuss these mathematic platforms beginning with ordinary differential equations, ODEs categorized as temporal deterministic.

One of the most popular techniques in systems biology, ordinary differential equations (ODEs) are used to model the temporal evolution of protein concentrations. ODEs are widely applicable at high population levels in a homogenous system, conditions commonly encountered in the cytosol and extracellular space (Figure 2). The limitations of ODEs come from their intrinsic assumptions: spatial homogeneity, deterministic behavior, and the continuum. An example of the inappropriate use of ODEs is on the plasma membrane, where spatial heterogeneity presides, resulting in a disconnect between biology and algorithm. In addition to the shortcomings in the acquired data, membrane microdomains tend to discretize species populations, hence, leading to a breakdown in the continuum and inaccuracies in the deterministic solution. The loss of information and inaccuracies within the results jeopardizes the algorithm's ability to be predictive.




Figure 1. Modeling Methods Suited for Different Aspects of Cell Biology. This diagram of the cell shows complementary modeling techniques for a given domain of the cell.

The SSA is a method suitable for homogenous system with low species populations that exhibit inherent stochastic noise. It is best suited in the nucleus, which exhibits spatial-homogeneity and small numbers of transcription factors producing innate stochastic responses.

The spatial-temporal toolbox of systems biology represents the other half of modeling approaches (Figure 1). The spatial-temporal deterministic approaches (e.g., partial differential equations, PDEs) are primarily used in systems with a single source and sink. The applications of PDE based models range from the diffusional aspects of synaptic transmission (Tai, Bond et al. 2003; Song, Zhang et al. 2004; Zhang, Suen et al. 2005), cytosol nuclear translocation (Haugh and Lauffenburger 1997; Brown and Kholodenko 1999; Kholodenko 2003; Kholodenko 2006), and receptor-ligand dynamics on the plasma membrane (Goldstein and Dembo 1995; Haugh 2002; Monine, Berezhkovskii et al. 2005). The former methods seem to be appropriately applied, utilizing the strengths of the method to gain important spatial-temporal information. The latter application could be questionable due to the characteristics of PDEs, not taking into account a sink-sink overlap. Being that the plasma membrane is a kaleidoscope of microdomains (Sheetz, Schindler et al. 1980; Kusumi, Nakada et al. 2005) this assumption is only valid at low receptor densities; conditions which cause the continuum to fail.

The remaining niches in the systems biology toolbox, spatial-temporal stochastic methods, are best suited for the plasma membrane, organelle sites, and biochemical cascades. Spatial-stochastic algorithms are not fruitful at high species populations and unnecessary in homogenous solutions. One of the major drawbacks

of these spatial stochastic approaches is the required computational rigor. Spatial stochastic methods, however, provide a powerful computational platform and are ideal for studying the dynamics on the plasma membrane. They will therefore be the focus of our discussion.




Figure 2. Choosing Appropriate Modeling Technique. We categorized mathematical techniques on the x-axis from spatial-temporal (diffusion limited circumstances) to temporal methods (well mixed solutions) as the ratio of the effective reaction rate per diffusion rate increases. Along the y-axis stochastic solutions to deterministic solutions are shown as a function of number of molecules.

Choosing an Appropriate Method

The art of computational biology comes from understanding the system. Figure 2 shows mathematical models applied to proper biological processes First deciding whether to use a temporal vs. spatial-temporal approach requires calculating the effective kinetic rate constant for diffusion-reaction problems. The following equation was derived by Lauffenburger and Linderman (Lauffenburger and Linderman 1993):

$$k_{PDE} = \frac{2\pi D_{AB} k_{Areal}}{2\pi D_{AB} + k_{Areal} \ln\left(\frac{b}{s}\right)} = \frac{4\pi D_{AB} k_{Areal}}{4\pi D_{AB} + k_{Areal} \ln\left(\frac{1}{\pi \rho_A s^2}\right)} \quad (1)$$

where $D_{AB} = D_A + D_B$, D_A and D_B are the diffusivities of A and B, ρ_A is the density of molecules of A (number of molecules per unit area), $b = \sqrt{1/\pi \rho_A}$ is one-half of the mean displacement between molecules of A distributed in a certain area, k_{Areal} is the intrinsic reaction rate constant in units of (receptors /area)⁻¹ s⁻¹, and s is the encounter radius. The relationship between k_{Areal} and k is given as, $k_{Areal} = ks^2$. Using Eq 1. and combining it with the expression for resistance in a series model

$$k_{PDE} = \left(\frac{1}{k_{Areal}} + \frac{1}{k_{Diffusion}} \right)^{-1} \quad (2),$$

the diffusion-limited reaction rate constant, $k_{Diffusion}$ is determined to be

$$k_{Diffusion} = \frac{4\pi D_{AB}}{\ln\left(\frac{1}{\pi \rho_A s^2}\right)} \quad (3).$$

Using $k_{Diffusion}$, k_{Areal} one can determine if in the diffusion limited case $k_{Diffusion} \ll k_{Areal}$, or

in the reaction limited case $k_{Diffusion} \gg k_{Areal}$. Using the ratio of $\frac{k_{Areal}}{k_{Diffusion}}$ we obtain a

dimensionless parameter which provides insight into the correct modeling technique (spatial-temporal/temporal). As shown in Figure 1. starting with a small value for

$\frac{k_{Areal}}{k_{Diffusion}}$ a spatial temporal model is most desired; however increasing $\frac{k_{Areal}}{k_{Diffusion}}$ validates

the temporal approach. Within the gray area between spatial-temporal and temporal models fall techniques such as compartmentalized ODEs that model neighboring organelles as compartments with their own set of initial conditions.

A similar calculation must be preformed when determining whether to use a stochastic or a deterministic approach. This decision must be based on the number of molecules; at high concentrations stochastic variations maybe negligible and better approximated with a continuum model. At low concentrations discrete behavior becomes apparent and undoubtedly leads to intrinsic noisy systems. Figure 1 depicts this concept starting with a stochastic method (SSA/MC) at low molecular numbers and switching to a continuum method (ODE/PDE) at high molecular concentrations.

Spatial-Temporal Stochastic Methods

Since the focus of this review is on mathematical modeling of the plasma membrane we will spend some time now discussing the method most appropriate for the plasma membrane: spatial-temporal stochastic algorithms. These methods are often implemented either off-lattice or on-lattice. The former being difficult to derive and computationally intensive, and the latter providing ease of implementation and computational efficiency. Two types of off-lattice approaches will be discussed the first agent based off-lattice method derives probabilities from a distance factor and the second Metropolis Monte Carlo (MC) method calculates transition rates by free

energy minimization. The agent based off-lattice approach will be briefly described followed by the Metropolis MC which will be summarized in a historical setting leading into the on-lattice spatial kinetic MC.

Agent based off-lattice approaches calculate probabilities for reaction by taking into account the distance between two reactants. Calculating reaction probabilities based on separation distance is by no means trivial. Rigorous theoretical derivations(Sung, Shin et al. 1997; Kim and Shin 1999) have resulted in promising numerical algorithms (Popov, Agmon et al. 2004; Park, Shin et al. 2005; Hsieh, Yang et al. 2008). However many of these derivations rely on assumptions that are contrary to know biological systems of the plasma membrane, such as uniform concentration of reactant B , immobile reactant A , as well as collision which equates with reaction.

The Metropolis MC algorithm was developed in 1953 (Metropolis, Rosenbluth et al. 1953) being the first method to numerically solve the underlying master equation. Describing a physical system with a Hamiltonian, Metropolis et al. constructed the probability for various states of the system weighted in terms of a Markov process by the novelty of a pseudo-random number. A configurational space which includes thermal vibrations and microscopic process, termed as “rare events” was sampled.

At the time this method proved “suitable for fast computing machines” and appropriate for describing equilibrium conditions. However the receptor interactions on the plasma membrane do not occur under equilibrium conditions but rather are dynamic involving billions of diffusion and millions of reaction events. The time scales between thermal vibrations and diffusion (let alone the diffusion which is

necessary for reaction to occur) are large and define the system as “stiff”. This means that most computational time will be spent performing fast vibrations instead of slow diffusion events, or “rare events” that are necessary for the system to evolve. This “stiffness” translates to small time increments making the original off-lattice MC impractical for modeling biological events occurring on the plasma membrane.

With the many drawbacks faced when implementing off-lattice methods, spatial stochastic modeling in biological systems would prove infeasible. However from the popularity of the Metropolis MC, came the desire to apply the MC to study dynamical behavior and thus lead to algorithms such as the dynamic MC (DMC) and spatial kinetic MC (SKMC). Dynamic simulations are now possible using the SKMC because it integrates out the effects of thermal vibrations on the microscopic rates. In these simulations slow events such as reaction and diffusion occur on the spatial coordinates of a lattice and are fired, neglecting the vibrational time. Probabilities for reaction and diffusion can be easily computed by calculating a transition rate for either reaction or diffusion to a neighboring lattice site.

An example of the “null-event” SKMC algorithm is implemented by calculating probabilities for reaction and diffusion events. This is done by normalizing transition rates with a microscopic rate Γ_{\max} that makes probabilities less than 1 and gives a null bin at the i th site. The normalized rate Γ_{\max} is the maximum sum of all microscopic rates at a site,

$$\Gamma_{\max} = \max_{i=1,\dots,N_L} \sum_{j=1}^{N_p} \Gamma_j \mathcal{E}_{ij}$$

in which N_p is the number of processes, N_L is the number of lattice sites, Γ_j is the transition rate for that process, and ε_{ij} is the participation matrix being 1 if the i th site gives rise to the j th process or 0 if it does not. Using the normalization rate Γ_{\max} probabilities p^x for successful events (reaction or diffusion) are computed as

$$p^x = \frac{\Gamma^x}{\Gamma_{\max}} .$$

Since the probabilities do not sum to one for a molecule at the i th site in

the j th process there is a probability for a null event or a null bin for each process.

The null bin penalizes sites with lower transition probabilities giving rise to more null events while higher transition probabilities result in less null events. Although the idea of null events suggest additional computational cost an excellent review by Chatterjee et al.(Chatterjee and Vlachos 2007) describes null event algorithms and shows how time advancement is independent of the chosen process.

The SKMC is dynamical due to time increments which were not present in the original Metropolis MC. The way in which time is updated varies with the MC method, although consideration should be given to events which occur most frequently in order to have a greater time resolution (Chatterjee and Vlachos 2007). Diffusion for example is a frequent process that can be used to update time

$$\Delta t = \frac{1}{\sum_{i=1}^{\text{No.of sites}} \left(\sigma_i \sum_{j \in B_i} \Gamma_{i \rightarrow j}^d (1 - \sigma_j) \right)} .$$

The term in the denominator describes all available sites a diffusion event can occur with a transition rate $\Gamma_{i \rightarrow j}^d$.

Overcoming the Limitations of SKMC Algorithms

Lattice based MCs usually implement periodic boundary conditions, PBCs and are computationally feasible at roughly 10^4 - 10^6 lattice sites (from 100×100 to 1000×1000 nm² with an inner lattice spacing of 1 nm). Large time scales represent another limitation of MC methods. Several novel algorithms have been developed to overcome these limitations: coarse graining techniques and τ -leaping methods. Coarse-graining MC (CGMC) techniques were developed to address the computational limitations occurring in large length scales. Grouping microscopic lattice points into coarse cells, low activity regimes are less computationally expensive, enabling the availability of more resources for high activity regimes or areas of particular interest. Although many of these CGMC techniques have been applied, their implementation of a uniform coarse cell size makes them inefficient for high activity regimes. A multiresolution framework incorporating a dynamic CGMC framework was developed by Chatterjee et al. (Chatterjee, Katsoulakis et al. 2005; Chatterjee and Vlachos 2006; Chatterjee and Vlachos 2007) called an adaptively coarse-grained Monte Carlo (ACGMC). The ACGMC enables large scale simulation with sharp concentration gradients, proving an adequate method for length disparities.

τ -leaping methods were developed in an effort to overcome disparity in time scales and to increase time increments. The first of these methods was introduced by Gillespie and later followed up by others. The τ -leap method computes a coarse-time increment, τ , being larger than the microscopic time and ‘fires’ all reactions multiple times and updates species populations appropriately within the time increment τ . The number of times reactions are ‘fired’ is determined randomly from a Poisson distribution. Although errors in the form of negative concentrations

are observed with the unbounded Poisson distribution in which the number of firings is larger than the populations of species, methods using a binomial distribution have proved more accurate for biological systems.

Although MCs are limited computationally they remain the most accurate at capturing biological phenomena, taking into account the inherent stochastic, discrete nature of biological systems as well as important spatial implications; essential aspects for capturing reaction and diffusion events on the plasma membrane. In this next section we will provide an example using an SKMC to gain biological insight into receptor dynamics on the plasma membrane.

Sorting Out Microdomains: An example using the SKMC

In an earlier section we discussed lipid rafts and picket fences on the plasma membrane in this example we illustrate the predicative capabilities of the SKMC at dissecting the effects of these microdomains. We define both a lipid raft and a picket fence on the lattice, and elucidate the effects of these microdomains in receptor aggregation. Lipid rafts range in size from 20 to 350 nm diameter (Jacobson and Dietrich 1999; Edidin 2001; Edidin 2001; Jacobson, Mouritsen et al. 2007) and proteins exhibit a decreased diffusivity on the order of 5 to 50 times on the plasma membrane (Kusumi, Ike et al. 2005; Kusumi, Nakada et al. 2005). We define a lipid raft on the lattice by specifying a region of 400 nm^2 and decreasing receptor diffusion by a factor of 50 in this area, as shown in Figure 3A. Similarly we define a picket fence as shown in Figure 3B. at the perimeter of 400 nm^2 region and allow breaks in the fence to occur every 230 nm at a time of 10 milliseconds.

To determine the effects that the picket fence and lipid raft have on receptor colocalization the Hopkins statistic test was performed as well as the p-value test (indicated as passed by the pink background). Initially 100 receptors were uniformly distributed on the lattice this is evident from the Hopkins test which shows our data following a normal distribution. As time increases to 0.08 seconds and 0.5 seconds we observe a right shift of our data (lipid raft Figure 3A) in the Hopkins test indicative of the clustered state. These results were further confirmed by the passing of the Chi-squared test. Observing the state of the system at similar time points reveals no apparent clustering in the picket fence microdomain. Additional simulations could be performed varying the diffusivity in the lipid raft, the size of the

lipid raft, varying the time between breaks and the number of breaks in the picket fence, and the area enclosed by the picket fence to further elucidate the effects of microdomains on receptor aggregate




Figure 3. Microdomains vs. Receptor Clustering. Schematic illustrates the lattice (left) and Hopkins test with pink boxes indicating passing of the Chi squared test. Two microdomains were tested lipid raft (left) and picket fence (right). The rows represent different time points (top-to-bottom) time is 0s, 1s, & 2s.

A New Direction for Systems Biology

This example illustrates how mathematical models can be predictive, guide experimentations, and interpret results. Using the SKMC approach we outlined here how one could interpret results from SPT experiments, predict the spatial-temporal evolution of the system based on conditions of the plasma membrane with results that would further guide experimentation.

In this review we have presented modern experimental techniques along with biological discoveries on the plasma membrane which challenge the predictability of mathematical models. The motivation for new computational approaches is driven by two factors: first the prospects of state of the art experimental technologies outdate previous mathematical methods, and two biological discoveries demonstrate an inhomogeneous, stochastic nature whose complexities can no longer be ignored.

Experimental technologies have evolved well beyond the standard western blot, immunoprecipitation experiments that were originally used to construct mathematical models. At the time, ODEs were complementary to western blot techniques by being able to predict the temporal evolution of a system. However the capacity of SPT and FRET to provide resolution of single protein diffusion and interaction in vivo and in real time can not be complemented with an ODE but rather must be complemented with an equally influential theoretical technique.

In addition to being outdated with experimental technologies the paradigm shift on the plasma membrane exemplifies the true complexity of cell biology which is not captured with ODEs. The plasma membrane is a highly compartmentalized structure,

a stage for discrete receptor diffusion and reaction events' resulting in heterogeneity, which in order to be accurately modeled needs a spatial-temporal stochastic approach. The plasma membrane was our focal point, however as more experimental studies are conducted we may discover intricacies in other biological systems as well providing additional motivation for these algorithms.

Although ODEs are not sufficient on the plasma membrane and do not complement high-tech experimental technologies, they are a tool of systems biology. When used appropriately ODEs are an insightful, predictive method for understanding biology. The challenge to systems biology comes not in the discontinuity of ODE methods but in terms of developing the future mathematical tools to guide and complement experimentation.

Chapter 2. Coupled Stochastic Spatial and Non-Spatial simulations of ErbB1 Signaling Pathways Demonstrate the Importance of Spatial Organization in Signal Transduction

Michelle N. Costa¹, Krishnan Radhakrishnan^{2,3}, Bridget S. Wilson^{2,3}, Dionisios G. Vlachos⁴, and Jeremy S. Edwards^{1,3,5,*}

¹ Department of Chemical and Nuclear Engineering, University of New Mexico, Albuquerque, NM 87131

² Department of Pathology, University of New Mexico Health Sciences Center Albuquerque, NM 87131

³ Cancer Research and Treatment Center, University of New Mexico Health Sciences Center, Albuquerque, NM 87131

⁴ Department of Chemical Engineering, University of Delaware, Newark, DE 19716

⁵ Molecular Genetics and Microbiology, University of New Mexico Health Sciences Center, Albuquerque, NM 87131

*Correspondence to Jeremy S. Edwards (jsedwards@salud.unm.edu)

Abstract

Background: The ErbB family of receptors activates intracellular signaling pathways that control cellular proliferation, growth, differentiation and apoptosis. Given these central roles, it is not surprising that overexpression of the ErbB receptors is often associated with carcinogenesis. Therefore, extensive laboratory studies have been devoted to understanding the signaling events associated with ErbB activation. Systems biology approaches have contributed significantly to current understanding of ErbB signaling networks.

Methodology/Principal Findings. Although computational models have grown in complexity over the years, little work has been done to consider the spatial-temporal dynamics of receptor interactions and to evaluate how spatial organization of membrane receptors influences signaling transduction. Herein, we explore the impact of spatial organization of the epidermal growth factor receptor (ErbB1/EGFR) on the initiation of downstream signaling. We describe the development of an algorithm that couples a spatial stochastic model of membrane receptors with a nonspatial stochastic model of the reactions and interactions in the cytosol. This novel algorithm provides a computationally efficient method to evaluate the effects of spatial heterogeneity on the coupling of receptors to cytosolic signaling partners.

Conclusions/Significance: Mathematical models of signal transduction rarely consider the contributions of spatial organization due to high computational costs. A hybrid stochastic approach simplifies analyses of the spatio-temporal aspects of cell signaling and, as an example, demonstrates that receptor clustering contributes significantly to the efficiency of signal propagation from ligand-engaged growth factor receptors.

Keywords: ErbB Signaling, EGFR, spatial modeling, Spatial Monte Carlo

Introduction

The ErbB family of receptors, under normal physiological conditions, regulates key cellular processes such as growth, proliferation and differentiation (Yarden and Sliwkowski 2001; Linggi and Carpenter 2006; Lajoie, Partridge et al. 2007).

Overexpression of these receptors deregulates normal cellular function and is a contributing factor to tumorigenesis (Britten 2004). There are four members of the ErbB family (ErbB1, ErbB2, ErbB3 and ErbB4) and each family member has its own unique ligand specificity (Hynes and Lane 2005), kinase activity (Linggi and Carpenter 2006) and spatial organization on the membrane (Lajoie, Partridge et al. 2007; Yang, Raymond-Stintz et al. 2007). In our current study, we have focused solely on the epidermal growth factor receptor (typically abbreviated ErbB1 or EGFR) and the ErbB1 activation of ERK, which is a mitogen activated protein kinase (Santos, Verveer et al. 2007). Ligand binding to ErbB1 stabilizes a conformation of the extracellular domain that allows receptor dimerization (Blinov, Yang et al. 2006). Dimerized receptors are active tyrosine kinases, capable of transphosphorylation (Blinov, Yang et al. 2006). Phosphorylation of receptor cytoplasmic tails results in recruitment of SH2-containing adaptor and signaling proteins, such as Grb2, Sos, and Shc, that form a signaling scaffold to activate ERK (Blinov, Faeder et al. 2006).

Due to the importance of the ErbB1-activated ERK pathway, several ordinary differential equation (ODE) models have been developed to gain insight into this pathway (Kholodenko, Demin et al. 1999; Schoeberl, Eichler-Jonsson et al. 2002; Hendriks, Opresko et al. 2003; Sasagawa, Ozaki et al. 2005). While ODE models have provided insight into the dynamics of this pathway, these models assume that the cell is a

homogeneous well-mixed system. In other words, the ODE models neglect spatial localization and organization, such as membrane receptor clustering (Yarden and Sliwkowski 2001; Orton, Sturm et al. 2005). Over the past decade, ODE models of ErbB1-induced ERK pathway system have evolved in complexity, becoming both larger and having more experimentally constrained parameters (Kiyatkin, Aksamitiene et al. 2006). The first ErbB1/EGFR model was introduced in 1996 and had 35 reactions (Huang and Ferrell 1996), whereas the most complete models available contain hundreds of reactions (Sasagawa, Ozaki et al. 2005; Kiyatkin, Aksamitiene et al. 2006).

The question remains whether these well-mixed deterministic models are capable of quantitatively describing the temporal dynamics of signaling, since there is significant evidence that cell membrane organization promotes the formation of localized “signaling platforms” (Jiang and Sorkin 2002; Borisov, Markevich et al. 2005; Bluthgen, Bruggeman et al. 2006; Kholodenko and Sauro 2008). Major advances in our understanding of the membrane have led to a revision of the original Fluid Mosaic model (Singer and Nicholson, 1972), to a more ordered structure with distinct membrane microdomains of lipids and proteins (Schroeder, Gallegos et al. 2001; Gallegos, Storey et al. 2006; Lillemeier, Pfeiffer et al. 2006). Advanced microscopy techniques have demonstrated that membrane properties, such as transient confinement zones and corrals, may restrict and govern the spatial-temporal dynamics of lipids and membrane proteins (Fujiwara, Ritchie et al. 2002; Ritchie and Kusumi 2003; Murase, Fujiwara et al. 2004; Koyama-Honda, Ritchie et al. 2005; Orr, Hu et al. 2005; Andrews, Lidke et al. 2008). The challenge is to develop computational approaches that can account for membrane spatial heterogeneity and evaluate the impact on signal propagation.

Spatial modeling has been implemented in many scientific disciplines, including physics, material sciences, chemistry, engineering and biological systems. Modeling approaches vary, including partial differential equations (Mac Gabhann and Popel 2005), agent-based modeling (Hsieh, Yang et al. 2008) and spatial Monte Carlo (MC) methods (Woolf and Linderman 2004; Chatterjee and Vlachos 2005; Chatterjee and Vlachos 2007). Spatial MC platforms are particularly powerful numerical simulation tools for studying the dynamics of membrane components (Woolf and Linderman ; Woolf and Linderman 2003; Brinkerhoff, Woolf et al. 2004; Mayawala, Vlachos et al. 2005). The application of spatial MC methods has been implemented by our group (Mayawala, Vlachos et al. 2005) to study ErbB reaction/diffusion and herein to study the effect of spatial heterogeneity on signal propagation. We report the development of a new computational framework that merges a spatial kinetic Monte Carlo (SKMC) algorithm for modeling reaction and diffusion events on the membrane with a stochastic simulator algorithm (SSA) (Gillespie 1977) for modeling cytosolic reactions. This new algorithm, the Coupled Spatial, Non-spatial Simulation Algorithm (CSNSA), has enabled us to determine the effects that receptor clustering has on the initiation of signaling.

Results

Establishing Parameters for the Spatial Model

One goal of our study was to evaluate whether simulation results from a spatial stochastic model would differ significantly from deterministic solutions that assume all components are well-mixed. As a starting point, we began with the original ODE model developed by Kholodenko and colleagues (Kholodenko, Demin et al. 1999). We noted, however, that the ODE model produced results that deviated from the same group's experimental data (Kholodenko, Demin et al. 1999). We performed a sensitivity analysis using the PottersWheel MatLab toolbox (Maiwald and Timmer 2008) to identify the most important enzymatic reaction parameters in the system. Based upon this analysis, we determined that incorporation of receptor degradation mechanisms results in a better fit to the experimental data (Figure 4A). Additional reactions added during our model development are illustrated in blue within Figure 4B and the entire set of reaction parameters are summarized in Table 1. Our model modifications are consistent with other models that include negative feedback reactions (Schoeberl, Eichler-Jonsson et al. 2002; Hendriks, Opresko et al. 2003; Sasagawa, Ozaki et al. 2005). In addition, it is noteworthy that the new parameters fit using the ODE model were not explicitly dependent on receptor diffusion. Appendix A. describes our analytical approach to demonstrate the validity of this fit, based upon very small error introduced by omission of diffusive properties of degradative machinery.

Table 1

Initial concentrations in nM units are Ru (varied), EGF = 20.42Vol_{Extracellular} (Vol_{Extracellular} is the volume of the cell (diameter of 20 μ m) multiplied by the ratio of the volume of incubation medium per cell over the cytoplasmic water volume ~ 33.3), PLC γ = 105, Grb2 = 85, and Sos = 34. First and second-order rate constants are in units of s⁻¹ and nM⁻¹ s⁻¹ and the Michaelis-Menten constants Km and Vmax are in units of nM and nM s⁻¹. Reactions are categorized as membrane reactions (handled by the SKMC), interfacial reactions (cytosolic species associating or dissociating with receptor) handled by the SKMC, and cytosolic reactions (handled by the SSA).

Reactions	Rate Constants	
Membrane Reactions		
1. EGF + Ru \leftrightarrow Rb	Kf = 0.003	Kb = 0.06
2. Rb + Rb \leftrightarrow RbRb	Kf = 0.01	Kb = 0.1
3. RbRb \leftrightarrow R	Kf = 1	Kb = 0.01
4. R \rightarrow RbRb	Vmax = 268	Km = 56.2
5. R-Sh \leftrightarrow R-pSh	Kf = 6	Kb = 0.06
6. R-PLC γ \leftrightarrow R-pPLC γ	Kf = 1	Kb = 0.05
Interfacial Reactions		
1. R + Shc \leftrightarrow R-Sh	Kf = 0.09	Kb = 0.6
2. R-pSh \leftrightarrow R + pShc	Kf = 0.3	Kb = 9x10 ⁴
3. R-pSh + Grb2 \leftrightarrow R-pSh-G	Kf = 0.003	Kb = 0.1
4. R-pSh-G \leftrightarrow R + pSh-G	Kf = 0.3	Kb = 9x10 ⁴
5. R-pSh-G + Sos \leftrightarrow R-pSh-G-S	Kf = 0.01	Kb = 2.14x10 ⁻²
6. R-pSh-G-S \leftrightarrow R + pSh-G-S	Kf = 0.12	Kb = 2.4x10 ⁴
7. R-pSh + G-S \leftrightarrow R-pSh-G-S	Kf = 0.009	Kb = 4.29x10 ⁻²
8. R + Grb \leftrightarrow R-G	Kf = 0.003	Kb = 0.05
9. R-G + Sos \leftrightarrow R-G-S	Kf = 0.01	Kb = 0.06
10. R-G-S \leftrightarrow R + G-S	Kf = 0.03	Kb = 4.5x10 ³
11. R + PLC γ \leftrightarrow R-PLC γ	Kf = 0.06	Kb = 0.2
12. R-pPLC γ \leftrightarrow R + pPLC γ	Kf = 0.3	Kb = 0.006
13. R-pShGS \rightarrow R-pShGS + E	Kf = 8	
14. R-GS \rightarrow R-GS + E	Kf = 48	
15. R + E \rightarrow Deg + E	Vmax = 4.7	Km = 82
16. R-pShGS + E \rightarrow Deg + E + pShGS	Vmax = 7560	Km = 78
17. R-GS + E \rightarrow Deg + E + GS	Vmax = 5520	Km = 7560
Cytosolic Reactions		
1. G-S \leftrightarrow Grb2 + Sos	Kf = 1.5x10 ³	Kb = 10 ⁴
2. pShc \rightarrow Shc	Vmax = 2.4	Km = 14.2
3. pShc + Grb2 \leftrightarrow pSh-G	Kf = 0.003	Km = 0.1
4. pSh-G + Sos \leftrightarrow pSh-G-S	Kf = 0.03	Kb = 0.064
5. pSh-G-S \leftrightarrow pSh + G-S	Kf = 0.1	Kb = 0.021
6. pPLCg \rightarrow PLCg	Vmax = 2	Km = 13
7. pPLCg \leftrightarrow pPLCg-I	Kf = 1	Kb = 0.003




Figure 4. Parameter optimization and summary of reaction network. A) Optimization of modeling parameters based upon sensitivity analysis and ODE solution. Green line: Kinetics of Shc phosphorylation in EGF-stimulated hepatocytes (20 nM EGF) as determined by Kholodenko et al. (Kholodenko, Demin et al. 1999). Red line: results obtained using the ODE model of (Kholodenko, Demin et al. 1999). Blue line: improved fit of ODE solution to experimental data after incorporation of receptor degradation reactions. B) Summary of reaction network in the ODE and CSNSA models. Note that, in the spatial CSNSA model, stars mark membrane reactions handled by the spatial stochastic Monte Carlo algorithm. All remaining reactions are governed by the Gillespie algorithm. Additional reactions that were added to the original ODE model from Kholodenko et al. (Kholodenko, Demin et al. 1999) are shown in blue. Numbering of reactions is arbitrary.

Validating the CSNSA hybrid approach

The novelty of the *CSNSA* approach lies in its computationally efficient framework that considers receptor diffusion and reaction in the 2-dimensional confines of the plasma membrane, while cytosolic reactions occur stochastically under well-mixed conditions. The simulated space is illustrated in Figure 5, with a full description of the *CSNSA* algorithm in the Methods section below. As an initial test, results were compared with the ODE solution (as described in Figure 4 and compared to the experiment results in (Kholodenko, Demin et al. 1999). The simulation space was populated with a receptor density of 141 receptors per μm^2 , each diffusing at $1 \times 10^{-14} \text{ m}^2\text{s}^{-1}$ (Kusumi, Ike et al. 2005), and a initial random distribution. In both ODE and *CSNSA* models, reactions were initiated by addition of EGF ligand (20 nM). Results show that, when receptors are randomly distributed, the two approaches give similar results for the rate and extent of ErbB1 phosphorylation (Figure 6A) and for the recruitment of PLC γ (Figure 6B). The *CSNSA* model predicts a slightly lower peak value and less sustained recruitment of Shc (Figure 6C), when compared to the ODE solution. These results emphasize that the accuracy of the *CSNSA* hybrid stochastic model is comparable to deterministic solutions in the absence of local concentration gradients or membrane inhomogeneities.




Figure 5. Illustration of the simulated space of the cell, consisting of two distinct domains: the cell membrane and the cytosol. The CSNSA model incorporates a Monte Carlo approach to simulate receptor diffusions and interactions on the cell membrane and couples to a spatial stochastic algorithm (Gillespie) for all cytosol interactions.




Figure 6. Comparison of the CSNSA and ODE solutions for receptor phosphorylation, PLC γ and SHC recruitment following EGF stimulation. Simulated kinetics of ErbB1 phosphorylation (A), PLC γ recruitment (B) and Shc phosphorylation after EGF (20 nM) using the ODE model (dashed lines) or the CSNSA model (solid black line). Results (A,B) from both simulation methods compare well with experimental data (solid circles) reported by Kholodenko et al. (Kholodenko, Demin et al. 1999)

Predicting the Impact of Receptor Density vs. Clustering

We next used the CSNSA to determine the effects of receptor spatial distribution and density on downstream signaling. We defined three different conditions, as shown in the schematic of Figure 7. In the first condition (magenta), the simulation space contained modest density of dispersed receptors (106 receptors per μm^2). In the second condition (dark blue), the simulation space contained a high density of well dispersed receptors (705 receptors per μm^2). The final simulation condition (cyan) began with a dense cluster of receptors, which was initially confined to a central region of 705 receptors per μm^2 and then permitted to diffuse over time to encompass the entire simulation space for a final density of 106 receptors per μm^2 . For each regime we examined how initial receptor density and clustering conditions influence coupling to four of ErbB1's signaling partners. The temporal profiles of the cytosolic species Grb2, Sos, and pShc and membrane-bound PLC γ are shown in Figure 7B-E.

All temporal profiles of the CSNSA were compared with their ODE solutions (shown in purple and red). The most notable differences came from the clustered regime (cyan), which had the same receptor concentration of 106 receptors per μm^2 as the non-clustered regime (magenta) but was initially confined to a smaller region. The clustered regime showed a marked increase in the amplitude of signal propagation in comparison to the ODE solution. The data demonstrates that spatial models are needed to accurately predict the consequence of membrane heterogeneity on signal propagation and set the stage for more refined considerations of signaling platforms.




Figure 7. The spatial model predicts that receptor clustering enhances signaling efficiency by creating locally high receptor densities. A) Schematic illustration of three simulation cases: dispersed (left), high-receptor density (middle), and highly clustered (right). See legend for key to colored lines in each plot. Results predict the kinetics of Grb2 activation (B), PLC γ phosphorylation (C), Shc phosphorylation (D) and Sos activation (E). Active Grb2 is equivalent to: RGrb2 + RGrb2Sos + RpShcGrb2 + RpShcGrb2Sos + Grb2Sos + pShcGrb2 + pShcGrb2Sos; Total phosphorylated PLC γ = RpPLC γ + pPLC γ + pPLC γ I; total phosphorylated Shc = RpShc + RpShcGrb2 + RpShcGrb2Sos + pShc + pShcGrb2 + pShcGrb2Sos; total Sos = RGrb2Sos + RpShcGrb2Sos + Grb2Sos + pShcGrb2Sos.

Discussion

In this work, we describe a new, efficient computation framework for evaluating the contributions of spatial organization to important cellular processes. Although applied here to study ErbB1 signal initiation at the plasma membrane, the algorithm should be readily adaptable to other receptor systems, organelle sites and biochemical cascades. We show that, when considering well-mixed systems, solutions obtained using the CSNSA hybrid model and the more traditional ODE solutions are comparable. However, given the growing evidence for membrane compartmentalization at both the plasma membrane and internal organelles (Smith, Simon et al. 1995; Yang, Simon et al. 2001; Yang, Raymond-Stintz et al. 2007), we propose that the spatial stochastic model will more accurately predict the outcomes of events that take place between membrane proteins and lipids and their cytosolic binding partners.

As an example, we used CSNSA to demonstrate that receptor clustering creates a more efficient signaling environment. The existence of receptor clusters is well established (Nagy, Vereb et al. 2002; Lillemeier, Pfeiffer et al. 2006; Wilson, Pfeiffer et al. 2007), but the significance of this membrane organization has been approached in only a few recent publications (Mayawala, Vlachos et al. 2006; Hsieh, Yang et al. 2008). Our previous work concluded that ligand-independent ErbB1 dimerization is likely to be dependent on two factors: density and the probability of receptor “fluxing” from a closed (dimerization-incompetent) to an open (dimerization-competent) conformation (Ozcan, Klein et al. 2006; Hsieh, Yang et al. 2008). Because clustering creates locally high receptor concentrations, it enhances the probability for collision between receptors that are transiently in the conformationally “open” state (Hsieh, Yang et al. 2008). Here, we

show that ErbB1 clustering also enhances the signaling output of receptors, based upon the more efficient recruitment of PLC γ , Grb2, Sos and Shc.

The importance of spatial effects is emerging as an important topic in systems biology, with technologies such as single particle tracking and electron microscopy demonstrating unique spatial domains (Fujiwara, Ritchie et al. 2002; Hansen, Prior et al. 2003; Ritchie, Iino et al. 2003; Ritchie, Shan et al. 2005; Suzuki, Ritchie et al. 2005; Birtwistle, Hatakeyama et al. 2007; Andrews, Lidke et al. 2008). In this work, we apply a novel algorithm to show a direct link between spatial heterogeneity and downstream signaling. We propose that future studies of receptor signaling should seek to gain a fundamental understanding of the spatial interactions and spatial organization of the receptors and to apply these concepts to predictions of signaling output. ErbB receptor clustered domains have been observed in many cancers using different microscopy techniques (Nagy, Vereb et al. 2002; Yang, Raymond-Stintz et al. 2007). Understanding this bigger picture of spatial-temporal protein interactions will drive forth knowledge of cell signaling events and offer the potential to lead towards better drug treatment options.

Methods

Coupled Spatial, Non-spatial Simulation Algorithm (CSNSA)

The Coupled Spatial Non-spatial Simulation Algorithm, CSNSA, is a hybrid model that considers the diffusive behavior and organization of receptors and other membrane components within a 2-D framework, bordered by a well-mixed cytosol. A spatial kinetic Monte Carlo algorithm is employed to capture the spatial-temporal dynamics of receptors on the cell membrane (Mayawala, Vlachos et al. 2005); this is a null-event algorithm that allows ease of implementation and variation of the underlying model. For computational simplicity, the cytosol is treated as a well-mixed solution and modeled with the stochastic simulation algorithm of Gillespie (Gillespie 1977). This assumption is reasonable in the cytosol, given that the diffusivity of proteins in the cytosol ($1 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$) (Morimatsu, Takagi et al. 2007) is four orders of magnitude larger than that in the plasma membrane ($1 \times 10^{-14} \text{ m}^2 \text{ s}^{-1}$) (Kusumi, Ike et al. 2005).

The two algorithms are coupled using the CSNSA, which employs a novel algorithm that selects and executes reactions that allow the molecular species to evolve in space and time. The coupling method takes into account the stochastic nature of biological systems. The first step of the CSNSA is to select a spatial domain (cell membrane or cytosol) and thus the corresponding algorithm for the next event. The selection is made by computing the probabilities of a membrane (SKMC) event or a cytosolic (SSA) event, which are calculated as:

$$P_{SKMC} = \frac{\Gamma_{tot,SKMC}}{\Gamma_{tot}} \quad \text{and}$$

$$P_{SSA} = \frac{\Gamma_{tot,SSA}}{\Gamma_{tot}}$$

where Γ_{tot} is defined as,

$$\Gamma_{tot} = \Gamma_{tot,SKMC} + \Gamma_{tot,SSA}.$$

The total transition rate for the SKMC, $\Gamma_{tot,SKMC}$, is the sum of all transition rates for all SKMC events, or more specifically the transition rate for diffusion ($\Gamma_{tot,Diff}$) and the sum of the reaction events ($\Gamma_{tot,k}$) for all N_{Rxn} reaction types (Table II),

$$\Gamma_{tot,SKMC} = \Gamma_{tot,Diff} + \sum_{k=1}^{N_{Rxn}} \Gamma_{tot,k}, \text{ where } \Gamma_{tot,k} \text{ is the total transition rate for each reaction type}$$

defined over all lattice sites N_L , $\Gamma_{tot,k} = \sum_{i=1}^{N_L} \Gamma_{i,k}$. $\Gamma_{tot,Diff}$ is defined as the sum of the

transition diffusion rate Γ_{Diff} over all lattice sites N_L , $\Gamma_{tot,Diff} = \sum_{i=1}^{N_L} \Gamma_{i,Diff}$. Thus, $\Gamma_{tot,SKMC}$ is

defined as:

$$\Gamma_{tot,SKMC} = \sum_{i=1}^{N_L} \Gamma_{i,Diff} + \sum_{k=1}^{N_{Rxn}} \sum_{i=1}^{N_L} \Gamma_{i,k}.$$

The SSA only accounts for stochastic variations in species populations and does not consider the spatial organization in the cytosol, and therefore does not contain a diffusion

term. The $\Gamma_{tot,SSA}$ is defined as the sum of Γ_k over all reaction types, $\Gamma_{tot,SSA} = \sum_{k=1}^{N_{Rxn}} \Gamma_k$.

The combined MC method operates like a single MC method by considering the superposition of all processes. Time is updated in a “combined” manner from Γ_{tot} with

an average time step as, $\Delta t = \frac{1}{\Gamma_{tot}}$. Given that the two algorithms are different (null-

event vs. rejection free), the CSNSA is a kind of a hybrid method. In order to properly match time scales, upon selection of a spatial event, the SKMC model is continuously executed until a successful event is selected, as shown in Figure 10, based on probability theory described in (Chatterjee and Vlachos 2007). The complete algorithm, which is shown in Figure 8, was implemented in Fortran 90. Due to the stochasticity of the algorithm, 10 simulations with different seeds for the random number generator were used for statistics. The *CSNSA* was benchmarked by comparison of an ODE model in a reaction-limited system, where the diffusion was made fast compared to the reaction rates (Figure 7). The typical CPU time for 50 receptors/lattice is ~15 min, for 125 receptors/lattice is ~2880 min, and for 500 receptors/lattice is ~14400 min on an Intel® Xeon™ CPU 3.2 GHz processor with 8.00 GB of Ram.

Table 2. Membrane Microscopic Events and Transition rates

Γ is defined on a square lattice with lattice species M, monomers, D, dimers, and pD, phosphorylated dimmers. Sx are species either within the cytosol SC or in the extracellular space SL. Details are provided in the text.

Microscopic Event	Transition Rate
Diffusion	$\Gamma_{i \rightarrow j}^D = \frac{1}{4} \Gamma^D \sigma_i (1 - \sigma_j) \quad j \in B_i$ <p>σ_i is the occupancy (discrete) that is 1, if site i is filled, and 0, if site i is empty (a single index indicating the site is used to simplify notation).</p> <p>$\Gamma^D = \frac{D}{a^2}$, where a is the microscopic lattice pixel dimension taken equal to the encounter radius, and D is the diffusivity of a receptor</p> <p>B_i denotes the set of sites to which diffusion from site i can occur which includes all 4 first-nearest neighboring sites</p>
Reactions	
Ligand Association Reaction ($S_L + M \rightarrow M^*$)	$\Gamma_i^R = k [S_L] \sigma_i$ <p>k is the macroscopic reaction rate constant with units as $[s^{-1}]$</p>
Ligand Disassociation Reaction ($M^* \rightarrow S_L + M$)	$\Gamma_i^R = k \sigma_i$ <p>k is the macroscopic reaction rate constant with units as $[s^{-1}]$</p>
Dimerization Reaction ($M^* + M^* \rightarrow D$)	$\Gamma_i^R = \frac{k}{2} \sigma_i \sigma_j$ <p>k is the macroscopic reaction rate constant with units as $[(\text{receptors/sites})^{-1} s^{-1}]$</p>
Decomposition Reaction ($D \rightarrow M^* + M^*$)	$\Gamma_i^R = k \sigma_i$ <p>k is the macroscopic reaction rate constant with units as $[s^{-1}]$</p>
Phosphorylation/Dephosphorylation Reaction ($D \leftrightarrow pD$)	$\Gamma_i^R = k \sigma_i$ <p>k is the macroscopic reaction rate constant with units as $[s^{-1}]$</p>
Cytosolic Association Reaction	$\Gamma_i^R = k [S_C] \sigma_i$ <p>k is the macroscopic reaction rate constant with units as $[s^{-1}]$</p>
Cytosolic Disassociation Reaction	$\Gamma_i^R = k \sigma_i$ <p>k is the macroscopic reaction rate constant with units as $[s^{-1}]$</p>




Figure 8. Schematic of *CSNSA*. Coupled Spatial Nonspatial Simulation Algorithm, *CSNSA*, combines the spatial stochastic algorithm (Gillespie 1977) depicted in the right branch, with the spatial kinetic Monte Carlo algorithm (Mayawala, Vlachos et al. 2005) in the left branch. Upon selection of a branch, a successful event has been executed, species populations are updated, transition rates and probabilities are recomputed, and time advances. The *CSNSA* is described in greater detail within the text.




Figure 9. Schematic of the SSA algorithm, as coupled to the hybrid algorithm. This algorithm is used for all cytosolic interactions. Being a rejection free algorithm, a successful event (reaction) is chosen and executed in each iteration. Our algorithm differs from the original Gillespie algorithm (26) in the time updating.

Spatial Kinetic Monte Carlo (SKMC)

Once an algorithm is selected and executed, transition probabilities are computed again at each time step. Computing $\Gamma_{tot,SKMC}$ involves computing the Γ values for the SKMC over the entire lattice. This computation is the most CPU intensive step in the simulation algorithm. We, therefore, used an optimized computation method. In order to maximize efficiency, a local region that is affected by the previous reaction event is defined (Mayawala, Vlachos et al. 2005), and the Γ for each lattice site is computed for this region both before and after the event has been executed. This eliminates scanning the entire lattice before and after an event is implemented, and the new $\Gamma_{tot,SKMC}$ is calculated by:

$$\Gamma_{tot,SKMC} = \Gamma_{tot,SKMC}^{old} - \Gamma_{local}^{old} + \Gamma_{local}^{new}$$

where, $\Gamma_{tot,SKMC}^{old}$ is the total transition probability computed initially or at a previous successful MC event, Γ_{local}^{old} is the sum of transition probabilities of all sites affected by an executed event based on the old configuration, and Γ_{local}^{new} is the sum of transition probabilities of all sites affected by an executed event based on the new configuration.

The SKMC algorithm is a modified null-event lattice MC method; for further details see Mayawala et al. (Mayawala, Vlachos et al. 2005). All reactions that are on the lattice or reacting with a species on the lattice are handled by the SKMC. Hereafter and in Figure 5, * denotes membrane reactions and ` denotes interfacial reactions. These reactions include ligand association and dissociation, receptor dimerization and decomposition, receptor phosphorylation and dephosphorylation, and phosphorylated receptor associating with and disassociating from cytosolic species. When an interfacial

reaction occurs, a molecule of the cytosolic species is subtracted from the cytosolic population and the membrane species is converted to a new species at the same location on the lattice.

The spatial domain is a two-dimensional lattice with periodic boundary conditions. The initial condition of the lattice is dependent on user specifications and can either be randomly populated or clustered in pre-defined domains. The algorithm is implemented by selecting an occupied lattice site, choosing a successful (reaction or diffusion) or unsuccessful (null) event based on the probabilities, and if a successful event was chosen, executing the event.

An event is selected by computing the probability distribution for all events, defined as $P_i^x = \frac{\Gamma_i^x}{\Gamma_{\max}}$, for lattice site i and event x . Table II shows the events executed by this algorithm and the equations for computing Γ^x for each event. Γ_{\max} is defined as

$$\Gamma_{\max} = 4 \left(\frac{\Gamma^d}{4} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right) + \max \left\{ \sum_{\text{all backward reaction events}} \Gamma^r \right\}$$

where the multiple of four accounts for events occurring in each of the four directions on the square lattice.

The spatial algorithm is coupled with the Stochastic Simulation Algorithm (SSA); therefore, unlike the original SKMC algorithm (Mayawala, Vlachos et al. 2005), the *CSNSA* is recursive in that it continuously selects an event until a successful event is chosen and executed as shown in Figure 8; therefore time is not updated if an unsuccessful event is selected.

Stochastic Simulation Algorithm (SSA)

The non-spatial SSA developed by Gillespie (Gillespie 1977) was used to model protein association reactions in the cytosol (Figure 9). The algorithm begins with initializing species populations and time; then propensities for all reactions are computed, and an event is randomly selected and the time is updated. This is a rejection free method; therefore, a reaction event is chosen and time is updated by an increment whose average is $\Delta t = \frac{1}{\Gamma_{tot}}$.

Interfacial Reactions

Interfacial reactions occur when a cytosolic species binds to or detaches from a receptor on the square lattice. In the former case, a molecule from the cytosolic species is subtracted from the cytosol population and a new product is produced at the site that was previously occupied by the reacting receptor. In the latter case, the converse procedure occurs. An example is shown in Table I(Interfacial Reaction #1) in which cytosolic species Shc binds to receptor R, occupying site k producing product R-Shc at site k.

The rate constants for cytosolic reactions are calculated by computing the cytosolic volume to be $V_{cyt} = 1/3 \pi L^2$ (units of μm^3), where r is the radius of the cell, and L is the lattice dimension. Next we compute the number of molecules per μm^3 , N_{sp} . By multiplying the product of V_{cyt} and N_{sp} with the rate constant (given in terms of molecules⁻¹ s⁻¹ for bimolecular reactions or s⁻¹ for unimolecular reactions), we obtain a transition rate with units of molecules s⁻¹.




Figure 10. The spatial kinetic Monte Carlo algorithm, as implemented in the *CSNS4*. This algorithm differs from the original algorithm of Mayawala et al (Mayawala, Vlachos et al. 2006) in the time update, which occurs recursively until a successful event is selected. Time is not updated when a null event occurs. A detailed description is provided in the text.

Sensitivity Analysis

To elucidate a mechanism that agrees with the experimental results (Kholodenko, Demin et al. 1999) and explains the biological nature of our system, we modified the reaction scheme developed by Kholodenko et al. (Kholodenko, Demin et al. 1999). A sensitivity analysis was performed on the reaction mechanism, using the decoupled direct method and the backward differentiation formula method, as implemented in the NASA Glenn chemical kinetics and sensitivity analysis code LSENS (Radhakrishnan 1991; Radhakrishnan 1999). In addition to the species concentrations, these methods automatically follow the temporal evolution of the first-order sensitivity coefficients $d\mathbf{C}/d\eta_j$. The vector \mathbf{C} contains the concentrations of all biochemical species and η_j is a parameter of interest, such as an initial concentration or a rate constant. The parameters of the new system were refined, and fits were performed for the new reactions shown in blue in Figure 4 and for the Michaelis-Menten reactions using PottersWheel, which were determined to be sensitive using the LSENS. The time scale of reactions in this model neglects the long term receptor production rate.

Acknowledgements

The image in Figure 5 was provided by Ulises Martinez.

Chapter 3. Picket Fence Densities Determine Enhanced vs. Inhibitive Role in Receptor Aggregation.

Michelle Costa¹, Dionisios G. Vlachos³ & Jeremy S. Edwards^{1,2,4,*}

¹ Department of Chemical and Nuclear Engineering, University of New Mexico,
Albuquerque, NM 87131

² Cancer Research and Treatment Center, University of New Mexico Health Sciences
Center, Albuquerque, NM 87131

³ Department of Chemical Engineering, University of Delaware, Newark, DE 19716

⁴ Molecular Genetics and Microbiology, University of New Mexico Health Sciences
Center, Albuquerque, NM 87131

Correspondence to Jeremy S. Edwards (jsedwards@salud.unm.edu)

Abstract

Experimental evidence suggests the cell membrane is a highly order structure of the cell which is compartmentalized by the underlying membrane cytoskeleton, MSK. The interaction between the cell membrane and the cytoskeleton led to the “picket-fence” model which was proposed to understand certain aspects of membrane compartmentalization. The picket fence model assumes that the membrane cytoskeleton sterically hinders and confines the motion of receptors and lipids in the membrane. However, the impact of receptor confinement on receptor clustering and aggregation, as well as downstream signaling remains controversial. Some evidence suggests that the MSK enhances dimerization, while other evidence links the MSK to a decrease in dimerization or signal activation. Although both scenarios may be in opposition, each may hold truth. Our hypothesis is that the rate of receptor dimerization is a function of the picket fence density and receptor concentration. Although it has been difficult to measure experimentally the relation between “picket-fence” density and receptor aggregation has been of great interest in recent years. Herein, we take a computational approach which enables us to test our hypothesis. Our results suggest that the peak in receptor clustering depends on the picket fence density as well as receptor concentration.

Keywords: EGFR, spatial modeling, Clustering. MSK, Picket Fences

Introduction

Membrane biology has gone down many new and exciting avenues since the original fluid mosaic model (Singer and Nicolson 1972) was proposed by Singer and Nicolson over 30 years ago (Vereb, Szollosi et al. 2003; Wisniewska, Draus et al. 2003;

Kusumi, Nakada et al. 2005). Great advances in our understanding of the membrane can be primarily attributed to new technology; i.e. high resolution microscopy captures structure on the nanometer scale, (Danuser and Waterman-Storer 2003; Morone, Fujiwara et al. 2006; Morone, Nakada et al. 2008) while protein tracking experiments (Ritchie, Iino et al. 2003; Murase, Fujiwara et al. 2004; Koyama-Honda, Ritchie et al. 2005; Suzuki, Ritchie et al. 2005; Andrews, Lidke et al. 2008) have revealed interesting spatial-temporal dynamics of membrane bound receptors. These new technologies are providing insights and pointing to inadequacies in the original fluid mosaic model.

There are two critical discrepancies between the experimental data and the fluidic mosaic model that demand further investigation (Kusumi, Nakada et al. 2005). First, the diffusion coefficients for both proteins and lipids in the plasma membrane were found to be smaller than those in artificial membranes (Saffman and Delbruck 1975; Cherry, Godfrey et al. 1982; Peters and Cherry 1982). Secondly, a dramatic drop of diffusion rates was observed for protein oligomers or molecular complexes (Nelson, Horvat et al. 1999; Roess, Horvat et al. 2000; Hegener, Prenner et al. 2004). These discrepancies clearly indicate that our understanding of membrane biology is incomplete and have led to a plethora of experimental studies giving a deeper understanding of the plasma membrane and the membrane skeleton (MSK) (Dietrich, Bagatolli et al. 2001; Fujiwara, Ritchie et al. 2002; Wilson, Pfeiffer et al. 2007; Morone, Nakada et al. 2008).

It has been hypothesized in the literature that the cell membrane is compartmentalized into microdomains, such as protein islands (Wilson, Pfeiffer et al. 2007) and lipid rafts (Nagy, Vereb et al. 2002). The “picket-fence” model is a non-mutually exclusive model that has been proposed as a mechanism for microdomains

formation(Ritchie, Iino et al. 2003; Ritchie and Kusumi 2004). In the picket-fence model the MSK acts as the fence by corralling transmembrane proteins while integral proteins serve as the “pickets” hindering receptor as well as lipid mobility (Nakada, Ritchie et al. 2003; Murase, Fujiwara et al. 2004). The “picket-fence” model accurately explains changes in diffusion rates; with lipids undergoing short term confined diffusion (with diffusion rates consistent with data on artificial membrane) followed by hop diffusion in between compartments (Ritchie, Shan et al. 2005; Suzuki, Ritchie et al. 2005). Restricted motion by the “picket-fence” also potentially explains differences in receptor diffusion between the artificial and the plasma membrane (Woelf and Linderman 2003; Brinkerhoff, Woelf et al. 2004; Hegener, Prenner et al. 2004).

Herein, we have utilized spatial stochastic simulations to test how picket fence density affects receptor clustering and hence signaling. While many *in silico* methods have been developed to study signaling which includes receptor interactions (i.e. dimerization), spatial information is often and receptors are assumed to be well mixed (Kholodenko, Demin et al. 1999; Sasagawa, Ozaki et al. 2005). Here we utilize the spatial kinetic Monte Carlo, SKMC to investigate the effect of the MSK on receptor clustering.

Materials and Methods

Spatial Kinetic Monte Carlo (SKMC)

Simulations were performed using the SKMC algorithm which is a modified null-event lattice based MC which was developed by our group (Mayawala, Vlachos et al. 2005; Mayawala, Vlachos et al. 2005). The algorithm is implemented by selecting an occupied lattice site, choosing a successful (reaction or diffusion) or unsuccessful (null) event based on probabilities, and if a successful event is chosen, executing the event. Transition rates are computed for both reaction and diffusion; they are presented in Table I. The model which was used as well as the reaction parameters is shown in Table II. The transition rate for diffusion is defined as,

$$\Gamma_{i \rightarrow j}^d = \frac{1}{4} \Gamma^d \sigma_i (1 - \sigma_j) j \in B_i \quad (1)$$

where $\Gamma^d = \frac{4D}{a^2}$ is four times its diffusion coefficient, D , divided by its inner

lattice distance, a . B_i signifies the set of sites in which diffusion from site σ_i can occur.

Diffusion takes place in four directions, each direction with its occupancy function σ_j which is discrete, equaling 1 if the site is filled, or 0, if the site j is empty. This means

that Eq. 1 can be either 0 or $\frac{1}{4} \Gamma^d$ depending on the occupancy of the neighboring site.

The probability distribution for both reaction and diffusion is $P_i^X = \frac{\Gamma_i^X}{\Gamma_{\max}}$ defined

for lattice site i and event x . Γ_{\max} is a normalization constant defined as

$$\Gamma_{\max} = 4 \left(\frac{\Gamma^d}{4} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right) + \max \left\{ \sum_{\text{all backward reaction events}} \Gamma^r \right\}$$

where the multiple of four accounts for events occurring in four directions on the square lattice; for further details see (Mayawala, Vlachos et al. 2005).

The spatial domain is a two-dimensional lattice with periodic boundary conditions. The initial condition of the lattice is dependent on user specifications and can either be randomly populated or clustered in pre-defined domains. Simulations were performed 10 times for statistical significance.

Table 3

Γ is defined on a square lattice with lattice species M, monomers, D, dimers, and pD, phosphorylated dimmers. Sx are species either within the cytosol SC or in the extracellular space SL. Details

Ligand Association Reaction	$\Gamma_i^R = k [S_L] \sigma_i$
Microscopic Event ($S_L + M \rightarrow M^*$)	Transition Rate k is the macroscopic reaction rate constant
Diffusion	with units as $\frac{1}{s}$ $\Gamma_i^D = \frac{1}{4} \sum_{j \in B_i} \Gamma^D \sigma_i (1 - \sigma_j)$
Ligand Disassociation Reaction ($M^* \rightarrow S_L + M$)	$\Gamma_i^R = k \sigma_i$ k is the macroscopic reaction rate constant with units as $[s^{-1}]$
Dimerization Reaction ($M^* + M^* \rightarrow D$)	$\Gamma_i^R = \frac{k}{2} \sigma_i \sigma_j$ $\Gamma^D = \frac{D}{a^2}$, where a is the microscopic lattice pixel dimension k is the macroscopic reaction rate constant with units as $[(\text{receptors/sites})^{-1} s^{-1}]$ taken equal to the encounter radius,
Decomposition Reaction ($D \rightarrow M^* + M^*$)	$\Gamma_i^R = k \sigma_i$ D is the diffusivity of a receptor k is the macroscopic reaction rate constant with units as $[s^{-1}]$ B_i denotes the set of sites to which
Phosphorylation/Dephosphorylation Reaction ($D \leftrightarrow pD$)	$\Gamma_i^R = k \sigma_i$ k is the macroscopic reaction rate constant with units as $[s^{-1}]$ first-nearest neighboring sites
Reactions	

Table 4

Initial concentrations in nM units are Ru (varied), $EGF = 20.42 \text{Vol}_{\text{Extracellular}}$ ($\text{Vol}_{\text{Extracellular}}$ is the volume of the cell (diameter of 20 μm) multiplied by the ratio of the volume of incubation medium per cell over the cytoplasmic water volume ~ 33.3). Reaction #5 was included in the predimerization simulations

Reactions	Rate Constants	
1. $EGF + Ru \leftrightarrow Rb$	$K_f = 0.003$	$K_b = 0.06$
2. $Rb + Rb \leftrightarrow RbRb$	$K_f = 0.01$	$K_b = 0.1$
3. $RbRb \leftrightarrow R$	$K_f = 1$	$K_b = 0.01$
4. $R \rightarrow RbRb$	$V_{\text{max}} = 268$	$K_m = 56.2$
5. $Ru + Ru \leftrightarrow RuRu^*$	$K_f = 0.01$	$K_b = 0.4$
*Predimerization		

Picket Fences

In order to model cytoskeletal interactions with the cell membrane boundaries, “picket fences” were placed on the lattice; previous work has investigated the use of a lattice to perform these simulations (Niehaus, Vlachos et al. 2008). Three different picket fence densities as shown in Figure 11, 25 corrals/lattice, 100 corrals/lattice, and 400 corrals/lattice were tested. The “picket-fences” occupy lattice sites and therefore prevent reaction and diffusion events in the direction of the boundary. Take for example, a receptor neighbored by a “picket fence” on the j th lattice site, it has an occupancy function, $\sigma_j = 1$ of one, yielding a zero transition probability, $\Gamma_{i \rightarrow j}^d = 0$ in the direction of the boundary. Similarly, a receptor separated from its partner by a “picket fence” can not dimerize being that the neighboring lattice site is occupied by the boundary.

Single particle tracking reveals short term receptor confinement followed by long term “hop” diffusion. Cytoskeletal rearrangements and actin depolarization are responsible for “hop” diffusion (Murase, Fujiwara et al. 2004; Kusumi, Nakada et al. 2005). In order to simulate “hop” diffusion breaks were randomly inserted into the boundaries an average of time step τ_{PF} of 10 milliseconds per $0.04 \mu\text{m}^2$ of picket fences as was observed experimentally (Fujiwara, Ritchie et al. 2002). Placing breaks in the “picket fence” enables receptors to diffuse out of their corrals. After an iteration of the SKMC breaks are closed and τ_{PF} is set to zero.

This is consistent with the rate kinetics of actin dimerization (Adams, Matov et al. 2004; Vallotton, Gupton et al. 2004; Danuser and Waterman-Storer 2006; Deshpande, McMeeking et al. 2006; Andrews, Lidke et al. 2008) which is on the order of the algorithm’s time step, Δt .




Figure 11. Picket Fence Distribution. The picket fence densities of $25 \text{ corrals}/\mu\text{m}^2$, $100 \text{ corrals}/\mu\text{m}^2$ and $400 \text{ corrals}/\mu\text{m}^2$ were tested.

We wanted to confirm that receptors have an equal probability of escape from each picket fence density. In order to do this we determined the ratio of the number of escapes to the number of collisions. These results are plotted in Figure 12, showing equal escape probability.




Figure 12 Number of Collisions and Escapes for each Picket Fence Density. The number of collisions (dashed line) and number of escapes (solid line) is shown as a function of time (s) for all picket fence densities 400 corrals/ μm^2 (red lines), 100 corrals/ μm^2 (green lines), and 25 corrals/ μm^2 (blue lines).

Quantification of Microdomains

In order to quantify microdomains in each simulation, the Fuzzy c-means (FCM) was performed to determine the number of clusters and the size of clusters. The FCM is a data clustering technique wherein each data point belongs to a cluster to some degree which is specified by its membership grade (Bezdek). An initial cluster number was specified in order to determine aggregation due to each corral and was then further reduced using the FCM. If the number of corrals was greater than the number of

molecules $x_{corrals} \geq x_{molecules}$ $N_{Clus} = \frac{x_{molecules}}{2}$, the cluster number was defined as the number of molecules divided by two; signifying that two or more receptors define a cluster.

However, if the number of corrals is less than the number of molecules,

$x_{corrals} \leq x_{molecules}$ $N_{Clus} = x_{corrals}$, the cluster number was defined as the number of corrals, specifying clustering due to each corral. The FCM algorithm performs an analysis unaware of the “picket fences”, such that different clusters occupy the same corral or a single cluster spans multiple corrals. Assuming that the mechanism of clustering is “picket fences”, we combined or separated clusters dependent on which corral they occupied. For example, clusters occupying the same corral were combined and clusters with receptor members occupying more than one corral were separated or eliminated. Performing this analysis we determined the number of clusters occupying a corral as well as the average cluster size.

Results

Clustering vs. Picket Fence Density

The first question we wanted to address is how does the density of picket fences effect receptor clustering. We tested this by looking at three different densities of picket fences: the first was a low picket fence density of 25 corrals/ μm^2 followed by a density of 100 corrals/ μm^2 and a hight density of 400 corrals/ μm^2 . Receptors were randomly placed on the lattice as shown in Figure 13(first row) for all picket fence densities. Random distributions were confirmed with the Hopkins test (Jain, Hamper et al. 1988), showing that the data follows a Gaussian distribution, and the Chi-squared Goodness-of-fit test (Snedecor and Cochran 1989) which validates the null hypothesis. Receptors were allowed to diffuse and at 1s (second row) there is a slight right-shift of the data in the Hopkins test. At 2s receptors cluster, indicated by the dramatic right-shift of the data in the Hopkins test. The Chi-squared values indicated a non-random, or clustered state.




Figure 13. Picket Fence Density vs. Receptor Clustering. Schematic illustrates the lattice (left) and Hopkins test with the chi-square parameters for that lattice. Three different picket fence densities representing the columns (left-to-right) 25 corral/μm², 100 corral/μm², and 400 corral/μm². The rows represent different time points (top-to-bottom) time is 0s, 1s, & 2s.

Comparing the three picket fence densities at 2s (third row) we see that both the Hopkins as well as the Goodness-of-fit test confirm the greatest amount of clustering at the picket fence density of 100 corrals/ μm^2 . While the densest picket fence density demonstrated the least amount of clustering as indicated by Hopkins and the Goodness-of-fit test; followed by least dense density.

In order to understand this result, we performed the fuzzy c-means clustering (FCM) algorithm (Bezdek) on the data. The FCM function computes the membership of receptors to a given cluster (Figure 14). The results were then analyzed and clusters were rejected if they border corrals, or contained less than two receptors in a corral. Performing this analysis we were able to determine the number of clusters and the cluster size (receptors per cluster). Ten simulations were performed and results were averaged for statistical significance. The results for each picket fence density are shown in (Figure 13). In this simulation we obtained an average cluster size of 5 receptors per cluster and 19 clusters (each occupying a separate corral) per lattice for the 25 corrals/ μm^2 densities. When the corral density was increased to 100 corrals/ μm^2 the average cluster size decreased to 3 receptors per cluster and the number of clusters increase to 28 clusters per lattice. The densest distribution of 400 corrals/ μm^2 showed a decrease in the number of clusters 17 clusters per lattice as well as cluster size of 2 receptors per cluster.




Figure 14. Cluster Analysis. Each picket fence density at a time of 2s was analyzed using fuzzy c-means clustering. Green boxes indicate the clusters which were due to a picket fence compartment. The diagram illustrates the member of a particular cluster. From this analysis the number of clusters as well as the average size of clusters was determined.

Clustering vs. Receptor Concentration

Hypothesizing that receptor concentration may affect our results we preformed simulations for receptor concentrations of 30, 50, 200, and 300 (number of receptors per lattice). The results show (Figure 15) that at a low receptor concentration (30-50 receptors per lattice) the greatest amount of clustering occurs at the least dense picket fence density of 25 corrals/ μm^2 where 8~13 clusters are observed. At a receptor concentration between 50 to 100 (receptors per lattice) we observe a switch within the data, with the most clustering at the 100 corrals/ μm^2 density. Although for higher receptor concentrations the greatest amount of clustering remains at 100 corrals/ μm^2 , there is an increase in the slope between 25 corrals/ μm^2 and 100 corrals/ μm^2 . Simulations at greater receptor concentrations were not preformed, due to computational limitations, but we predict a switch with the greatest amount of clustering occurring at the densest 400 corrals/ μm^2 density in which the cluster size would approximate the number of corrals at this density.




Figure 15. Number of Clusters and Cluster Size vs. Picket Fence Density. The number of clusters (blue line) and size of clusters (green line) or number of receptors per cluster is plotted as a function of picket fence density.

Picket Fence Densities Rescale MSD

To test a possible mechanism for clustering, we investigated how dimerization may lead to clustering via oligomerization induced trapping (Ritchie, Iino et al. 2003; Suzuki, Ritchie et al. 2005). We looked at the mean square displacement, MSD, of both monomer and dimer for each of the picket fence densities (rows) as a function of time, Figure 16 **Error! Reference source not found.** (left column). In the MSD plots we observed hindered diffusion for all picket fence densities Figure 16. At a density of 400 corrals/ μm^2 the monomer escapes at $\sim 0.3\text{s}$ (noted by the shift in MSD) and then is confined until $\sim 1.25\text{s}$; the dimer remains confined to a MSD of $\sim 0.002\text{ }\mu\text{m}^2$ within the 2s simulation. Looking at the single particle tracking trajectories (Figure 16 right most panels) the monomer travels across a much larger area than when it is bound to a partner. The MSD trajectories of the 100 corrals/ μm^2 density show both monomer and dimer escaping at $\sim 1.1\text{s}$; the dimer is more confined moving in MSD area of $\sim 0.013\text{ }\mu\text{m}^2$ while the monomer's area is $\sim 0.024\text{ }\mu\text{m}^2$. The single particle trajectories showed less difference between a monomer and dimer in the area traveled when compared with the trajectories of the 400 corrals/ μm^2 . The MSD plots for the 25 corrals/ μm^2 show escapes for monomer and dimer at $\sim 0.5\text{s}$. The dimer escapes and then seems to be confined from 1.1s till 1.6s and then escapes again, while the monomer "hops" to another corral at 1.1s where it is confined to $0.024\text{ }\mu\text{m}^2$ area and then escapes again at $\sim 1.5\text{s}$. The single particle trajectories showed the monomer to have traveled a greater area, but compared with the 400 corrals/ μm^2 the differences are small. The results indicated that dimerized receptors

cover less area, which supports oligomerization induced trapping; these results are more pronounced in the densest density of 400 corrals/ μm^2 .

Next we investigated the MSD trajectories as a function of the number of collisions (Figure 17). Rescaling the time in the MSD plots to number of collisions showed that escapes were only present in the 25 corrals/ μm^2 density. Rescaling the 100 and 400 corrals/ μm^2 density to 3×10^6 number of collisions, we see an escape for the monomer (400 corrals/ μm^2) at $\sim 1.3 \times 10^6$ number of collisions, however, the dimer remains confined. At 100 corrals/ μm^2 density both monomer and dimer escape after $\sim 1.55 \times 10^6$ number of collisions. The relation of the different densities shows how time, collisions, and escapes are being scaled. The 25 corrals/ μm^2 density shows a larger area traveled and greater number of escapes occurring in a smaller time and number of collision frame. These results illustrates time scaling of receptor signaling events based on the density of the membrane cytoskeleton.




Figure 16. Mean Squared Displacement as a Function of Time for Monomers and Dimers. The mean squared displacement is plotted as a function of time for both monomer and dimer for each of the picket fence densities. The right image shows single particle tracking on the lattice for both monomer and dimer.




Figure 17. Mean Squared Displacement as a Function of Number of Collisions for Monomers and Dimers. The mean squared displacement is plotted as a function of number of collisions for both monomer and dimer for each of the picket fence densities. The densities of 400 corals/ μm^2 , 100 corals/ μm^2 , 25 corals/ μm^2 mean square displacements are plotted in lower left column. The right column shows 400 corals/ μm^2 , and 100 corals/ μm^2 rescaled.

Oligomerization Induced Trapping

We wanted to determine if clustering is a result of oligomerization induced trapping. To test this we compare all three picket fence densities (25, 100, and 400 corrals/ μm^2) in the presence (Figure 18 left column) and absence (Figure 18 right column) of ligand stimulus. Adding ligand stabilizes EGFR in an open confirmation, enabling receptors to form dimers. In these simulations receptors were not allowed to predimerize. When performing the analysis a dimer was counted as a single species, as are monomers. Performing the Hopkins statistic test and the Chi-squared test showed clustering when ligand was added, indicating that clustering occurs as a result of dimerization.





Figure 18. Oligomerization Induced Trapping. Right column shows system in the absence of ligand. Left column shows system with ligand. Rows are for 25, 100, and 400 corral/μm² densities. All lattices are shown at 2s.

Predimerization vs. Dimerization

We wanted to test the effects of clustering for the case of predimerization vs. dimerization. An additional reaction was added to the system to account for dimerization in the absence of ligand,



with $k_f = 0.01 \text{ (nM s)}^{-1}$ and $k_b = 0.4 \text{ s}^{-1}$ (Shankaran, Wiley et al. 2006). Simulations were carried out as before with receptors initially distributed randomly and then being allowed to diffuse as time progresses. The $100 \text{ fence}/\mu\text{m}^2$ picket fence density was used for both cases and simulations were carried out to 2s which is when dimerization reaches steady state.

The results are shown in Figure 19, where we see the initial effects of clustering for the predimerization simulation at a time of 1s (Figure 19, right column middle). This was confirmed through the Hopkins statistic which showed a shift in our data, as well as the Chi-squared value which was 10.23, greater than Chi-squared value of 9.49, disproving the null hypothesis. However as time increased to 2s the amount of clustering proved to be statistically similar, leading to the conclusion that pre-dimerization initially increase clustering.




Figure 19. Dimerization vs. Predimerization. The picket fence density of $100 \text{ corrals}/\mu\text{m}^2$ is shown for both dimerization (left column) and predimerization (right column) at times 0s, 1s, and 2s (rows). The Hopkins test as well as the Chi-squared values are shown to the left of each lattice.

Discussion

The role that MSK microdomains, such as picket fences, have on receptor aggregation as well as downstream signaling has been controversial (Allen, Halverson-Tamboli et al. 2007). Some evidence indicates that picket fences may have an inhibitory role in cell signaling (Tank, Wu et al. 1982; Berk and Hochmuth 1992; Ganguly, Pucadyil et al. 2008), while other evidence points to an increase in receptor clustering which enhances downstream signaling (Douglass and Vale 2005; Heneberg, Lebduska et al. 2006; Chichili and Rodgers 2007). Through this work, we have shown that both are valid and dependent on the density of picket fences as well as receptor concentration. At low receptor concentrations (30 to 50 receptors per lattice) increasing picket fence density has an inhibitory effect on clustering; whereas at normal to high receptor concentrations, the greatest amount of clustering was observed at a picket fence density of 100 corrals/ μm^2 . Increasing from 200 to 300 (receptors per lattice), an increase in number of clusters occurs at the picket fence density of 400 corrals/ μm^2 . We hypothesize that at greater receptor concentrations there would be a shift in the number of clustering towards increasing picket fence densities.

The ability of the cell to synchronize cytoskeletal interactions in conjunction with signaling events has been shown experimentally (Nakada, Ritchie et al. 2003; Sheetz, Sable et al. 2006). Coordinating microdomain densities to regulate cell signaling could prove to be an important mechanism exploited during oncogenesis. Our data (Figure 13, Figure 16, & Figure 17) shows a time delay in clustering, which could activate some signaling pathways while suppressing other pathways. This time delay is dependent of picket fence density, at 400 corrals/ μm^2 , we see a shift in the data (Figure 13, left column

center) which indicates a more clustered state. This concept may be of importance to the activation of ERK, which can lead to either differentiation or proliferation dependent on its transient vs. sustained signal (Thrane, Schwarze et al. 2001; Sasagawa, Ozaki et al. 2005). Our results (Figure 19) showed clustering at 1s in the case of predimerization, whereas clustering was not observed until 2s for dimerization. These results support oligomerization induced trapping as a mechanism for clustering. Such a mechanism is further supported by looking at the MSD plots and the SPT simulations for dimers in comparison to their monomer components.

In summary our results show how microdomains on the plasma membrane can both inhibit and enhance clustering. When receptor aggregation is enhanced, oncogenic phenotypes, such as self-sufficiency of growth factors and an amplification of proliferative pathways, contribute to the diseased state. Often times oncogenic events are well coordinated and a mechanism of turning on and off signaling pathways via rearrangement of MSK could facilitate the cancer cell.

Acknowledgements

The research was supported by INCBN IGERT.

Chapter 4. Adaptively Coarse Grained Monte Carlo Method for Capturing the “Receptor–Sharing” Mechanism

Michelle N. Costa¹, Dionisios G. Vlachos & Jeremy S. Edwards^{1,2,4,*}

¹ Department of Chemical and Nuclear Engineering, University of New Mexico,
Albuquerque, NM 87131

² Cancer Research and Treatment Center, University of New Mexico Health Sciences
Center, Albuquerque, NM 87131

³ Department of Chemical Engineering, University of Delaware, Newark, DE 19716

⁴ Molecular Genetics and Microbiology, University of New Mexico Health Sciences
Center, Albuquerque, NM 87131

Correspondence to Jeremy S. Edwards (jsedwards@salud.unm.edu)

Abstract

The concept of the cell being nothing more than a bag of enzymes has come to pass, in its place the understanding that spatial organization dictates the efficiency of signal transduction pathways. It has been understood for sometime that receptor colocalization is an important part of signal transduction. When receptors form aggregates signal is processed in an effective way, reducing the limitations imposed by diffusion. In this work we investigated the extent to which receptor colocalization enhances downstream signaling. We observed a significant increase in association rates in comparison to dimerization rates when receptors are clustered. This result leads to our hypothesis that a “receptor-sharing” mechanism exists and contributes to the overall efficiency of signal transduction. The “receptor-sharing” mechanism occurs when cytosolic species binds with a receptor then disassociates and rebinds a neighboring receptor. We tested our hypothesis using a novel computational approach, the ACGMC, an algorithm which enables the spatial temporal evolution of the system in three dimensions by using a coarse graining approach. In this framework we are modeling EGFR reaction-diffusion events on the plasma membrane while capturing the spatial-temporal dynamics of proteins in the cytosol. From this framework we observe “receptor-sharing” which may be an important mechanism in the regulation and overall efficiency of signal transduction.

Keywords: receptor-sharing, adaptively coarse grained Monte Carlo, ErbB Signaling, EGFR, spatial modeling,

Introduction

To respond to the complexities of their external environment and internal queues, living cells have evolved complex signal transduction pathways, which even surpass the complexity of modern processing units (*1-3*). Most signal transduction pathways originate at the plasma membrane with receptor-receptor ligand interactions, followed by receptor-receptor interactions establishing signaling scaffolds and platforms (*4-7*). The rate limiting step in a signal transduction pathway is often receptor-receptor interaction (*8-12*), i.e. dimerization which is diffusion limited and essential to the propagation of signal. The highly controlled process of dimerization is often disrupted in tumor cells which, overexpress receptors (*13, 14*).

The prototypical signal transduction pathway (i.e. the ErbB network) begins with a receptor, (i.e. tyrosine kinase receptor) binding to its ligand when present (*15*). A ligand bound receptor has a greater affinity (eg. conformational orientation) to then form a dimer with neighboring receptors (*8, 16, 17*). Upon dimerization, receptor activation is an autocatalytic process occurring on the time scale of 1 nM s^{-1} (*18*), for example, the C-terminal transphosphorylates very rapidly due to the close proximity of the tyrosine kinase tails. The phosphorylated receptor leads to the recruitment of cytosolic species: adaptor proteins (Grb2, Sos, Shc)(*19-22*), cytosolic membrane tethered species (PLC α , Ras, Rap)(*20, 23-26*), signaling scaffolds (KSR)(*27-29*), scaffold binding proteins (Raf, MEK) (*30-33*) and transcription factors (ERK)(*29, 33-36*). The signaling cascade establishes itself at the plasma membrane, making it the “hub” of signal transduction (*6, 25, 29, 37-39*).

Spatial organization (i.e. receptor clustering) is an important part of signal transduction, which is often neglected in systems biology when modeling with ODEs (9, 18, 35). Exemplifying the essentiality of clustering, are $\text{INF}\gamma$ receptors which are expressed at low levels (40) ranging from 10^2 to 10^3 receptors per cell) on T-cells and macrophages that have a diameter $\sim 20\mu\text{m}$ (41). The $\text{INF}\gamma$ receptors are experimentally found, using techniques such as electron microscopy and immunoprecipitation, aggregated in caveolar domains; explaining how receptors expressed at low levels propagate signal (40, 42, 43). Another example illustrative of the need for receptors to colocalize, comes from the ErbB family of receptors which are observed in electron-microscopy images co-clustering (13, 44). It is known the formation of homo and heterodimers in the ErbB family leads to the activation of different signaling cascades (33, 45). The formation of ErbB1-ErbB1 homo, ErbB1-ErbB2 hetero dimer leads to PLC β , Grb2, and Shc activation while ErbB1-ErbB3 and ErbB1-ErbB2 initiates binding of multiple PI3K (45). Clustering facilitates dimerization and thus the activation of signal transduction pathways (10, 46-48).

A quantitative understanding of the receptor spatial organization and the downstream signaling proteins is still lacking. We previously showed through a novel algorithm, the *CSNSA*, how receptor clustering enhances downstream signaling. In a study investigating aspects of clustering (49-51) synthetic bivalent ligands for Fc ϵ RI were constructed from DNA fragments, and it was determined that the effective initiation of Fc ϵ RI requires receptor aggregation (51). However there has not been a mechanism to describe the effects of receptor aggregation on downstream signaling. Herein, we provide simulation results in support of the hypothesis that a “*receptor-sharing*”

mechanism exists such that a cytosolic species disassociating from its receptor can “*hop on*” or “*share*” a neighboring receptor when receptors are in the clustered state. Although such a mechanism is difficult to test experimentally, we have implemented a novel computational approach, the adaptively coarse-grained Monte Carlo, (ACGMC) (52-54) to test our hypothesis. In our mathematical model we are simulating the spatial-temporal interactions of EGFR on the plasma membrane while simultaneously capturing the spatial-temporal dynamics within the cytosol using a coarse-graining technique. To our knowledge this is the first coupled spatial-temporal membrane cytosol MC simulation.

Results

Cell Signaling Events are Dependent on Spatial Localization

Two spatial distributions of receptors were tested, a uniform distribution and a clustered distribution in which receptors cluster in a “lipid raft” with a size of 200 nm. Plotting dimerization vs. time (Figure 20A) we observe an increase in dimerization events for the clustered distribution when compared with the uniform distribution. Dimerization was enhanced when receptors were in the clustered distribution, to further investigate downstream signaling events we plot cytosolic binding events over time. and observe a transient increase in binding events. Figure 20B shows a much more dramatic difference between the clustered and uniformly distributed receptor conditions in terms of the number of binding events when compared with the number of dimerization events (Figure 20AB). This observation was suggestive of a mechanism that would enhance signal in the clustered state.




Figure 20. Reaction events for clustered vs. uniformly distributed receptor state. A) EGFR Dimerization (events) as a function of time. B) EGFR Association (events) for receptor binding cytosolic species as a function of time. Teal line: receptors clustered in a lipid raft of 200 nm. Magenta Line: receptors uniformly distributed.

Evidence of a Concentration Gradient

Observing an enhancement of cytosolic association events we hypothesized that a concentration gradient exists under the plasma membrane. To test this we looked at the total concentration of cytosolic species within the cytosol over time, where the total cytosolic species, CS is defined as

$CS = Grb2 + Grb2Sos + Shc + pShc + pShcGrb2 + pShcGrb2Sos$. As cytosolic species bind with phosphorylated receptors on the plasma membrane the concentration of these species in the cytosol decreases, as is observed in Figure 21. Comparing receptors which are in a clustered state with those that are uniformly distributed we observe additional concavity (Figure 21) in the cytosolic profile of the clustered state. The solutions deviate significantly between ~3s and ~16s but as time progress the solutions appear to reach a steady state, with small differences in the solutions.




Figure 21. Concentration of cytosolic species over time. The temporal profile of the concentration of cytosolic species in units of nM. Green line: receptors clustered in a lipid raft of 200 nm. Magenta Line: receptors uniformly distributed.

The “Receptor-Sharing” Mechanism

In previous work we showed how receptor aggregation enhances downstream signaling (ref), however our results assume the cytosolic species are well-mixed. We now know that cytosolic species are not well-mixed and their spatial organization also plays a role in cell signaling. We hypothesize that receptor aggregation increases downstream signaling via a “receptor-sharing” mechanism. Figure 22 demonstrates the “receptor-sharing” mechanism; a cytosolic species (CS1) bound with a receptor (R1-CS1) in the clustered state disassociates and “hops” on a neighboring receptor (R2-CS1). In order to observe this mechanism we identified single particle tracking trajectories shown in Figure 23. Cytosolic trajectories are shown in green while cytosolic bound receptor #1 (R1-CS1) trajectories are in red and cytosolic bound receptor #2 (R2-CS1) trajectories are in blue. Figure 23A&B shows a cytosolic species bind to receptor, the receptor-bound species diffuses then disassociating rebinds a neighboring receptor a distance of 20 nm from the original receptor. The “receptor-sharing” event occurred within 80ms, the cytosolic species traveled 10 nm in the z-direction rebinding a receptor 20nm away from the original receptor that it had previously bound. An example of a rebinding event that is clearly not via the receptor sharing mechanism is shown in Figure 23C&D. In this trajectory the cytosolic species binds receptor, diffuses with receptor, disassociates, and enters the cytosol where it diffuses in the z-direction essentially entering a well mixed pool. After 30s the cytosolic species diffuses to the plasma membrane and rebinds receptor, a random event not captured by the “receptor-sharing” mechanism.




Figure 22. Schematic of "receptor-sharing". The cytosolic species (CS1) binds with receptor #1(R1) then undergoes a "receptor-sharing" event by disassociating with R1 and binding the neighboring receptor #2 (R2).




Figure 23. Single particle tracking of receptor-sharing and non receptor-sharing events. Three-dimensional SPT of cytosolic species (CS1), diffusing in the cytosol (green), diffusing on the plasma membrane while bound to receptor 1, R1-CS1 (red) or bound to receptor 2, R2-CS1 (blue). A) “Receptor-sharing” event circled in yellow as viewed in x-z plane B) as viewed in x-y-z plane. C) Non “receptor-sharing” event as viewed in x-z plane D) as viewed in x-y-z plane.

Quantifying the “Receptor-Sharing” Mechanism

In order to quantify the receptor sharing mechanism we calculated the time of receptor sharing, t_{RS} , being defined as

$$t_{RS} = \frac{\langle r \rangle^2}{6D_C}$$

in which $\langle r \rangle^2$ is the mean square displacement in a microdomain of size $0.3 \mu\text{m}$ (55) and D_C is the cytosolic diffusivity coefficient of $1 \mu\text{m}^2/\text{s}$ (56, 57). Calculating the time of receptor sharing we are able to quantify a receptor sharing event defined as a receptor binding event occurring within 0.05, seconds $t_{RS} \leq 0.05\text{s}$ after it has disassociated from a neighboring receptor.

Turning up cytosolic diffusion in essence eliminates “receptor-sharing”. We preformed simulations by increasing cytosolic diffusion two orders of magnitude, $100 \mu\text{m}^2/\text{s}$. Figure 24 shows plots of the number of binding events as a function of time between binding events for normal cytosolic diffusion and high cytosolic diffusion. We observe a statistical difference in the data at 0.05s, with the number of receptor-sharing events being approximately ~ 5 for normal cytosolic diffusion and ~ 1 for high cytosolic diffusion. Performing a two-tailed t-test at 0.05s gives $10.25 > 6.5$ with a 0.005 P-value, indicating the significance between the two distributions.




Figure 24. Number of binding events vs. time between binding events. Blue line: Cytosolic species with normal diffusion at $1\mu\text{m}^2/\text{s}$. Cyan line: with high cytosolic at $100\mu\text{m}^2/\text{s}$.

Inhibiting the “Receptor-Sharing” Mechanism

The mechanism of “receptor-sharing” may have important therapeutic applications. We added tyrosine kinase inhibitors, Iressa (Gefitinib) and Tarceva (Erlotinib) to our system. These inhibitors block signal transduction by competitively binding with the receptor’s phosphorylation sites on the tyrosine kinase tail (concentration and binding kinetics are provided in the legend of **Figure 25**). We wanted to determine the effectiveness of TKI’s at reducing the “receptor-sharing” mechanism. In **Figure 25**, we looked at the number of association events as a function of time for Iressa and Tarceva. The effects of Iressa appear to be more potent at reducing both association events and “receptor-sharing” events.




Figure 25. Inhibition of the "receptor-sharing" mechanism. A) Association (events) vs. time (s) B) Number of binding events vs. time between binding events for green line: EGFR clustered in a lipid raft of 200 nm without drug, red line: treated with Iressa (Gefitinib), and blue line: treated with Tarceva (Erlotinib). An inhibitor concentration of 33 nM was used. Tarceva binding kinetics $k_f = 3 \text{ nM}^{-1} \text{ s}^{-1}$ (74) and $k_b = 1 \text{ s}^{-1}$, Iressa binding kinetics $k_f = 0.7 \text{ nM}^{-1} \text{ s}^{-1}$ and $k_b = 1 \text{ s}^{-1}$ (75)

Discussion

It has been established that receptor aggregation enhances the efficiency of signal transduction(58-60). With the underlying mechanism being that a local concentration gradient reduces diffusional limitations, thus increasing the number of dimers. Our results (Figure 20A) confirmed this statement, showing a slight increase in dimerization for receptors in the clustered state. However, when we looked at the number of association events the results were much more dramatic (Figure 20B), showing a significant increase in the number of association events for clustered receptors. These results were suggestive of an additional mechanism.

We hypothesized that a “receptor-sharing” mechanism exists and contributes to the overall efficiency of signal transduction. In our proposed “receptor-sharing” mechanism (Figure 22), a cytosolic species binds to a receptor, and then undergoes a “receptor-sharing” event by disassociating from the previously bound receptor and rebinding a neighboring receptor in close proximity. To confirm that the “receptor-sharing” mechanism exists we performed single particle tracking of cytosolic species. We observed a rebinding event which occurred via the “receptor-sharing” mechanism, disassociating and rebinding within 0.05s. In support of this mechanism, one would expect a concentration gradient to be established under the plasma membrane. This was confirmed by our results Figure 21, which showed additional concavity in the cytosolic concentration profile when receptor aggregation was present.

In order to quantify the “receptor-sharing” mechanism we established a time limit between disassociating and rebinding which would serve as a cut-off for classifying receptor binding events. The time limit was defined as:

$$t_{RS} = \frac{\langle r \rangle^2}{6D_C}$$

in which $\langle r \rangle^2$ is the mean square displacement, D_C is the cytosolic diffusivity coefficient, and 6 represents diffusion in three dimensions. The cytosolic diffusivity coefficient is two orders of magnitude higher in the cytosol than on the plasma membrane (56, 57). With measured receptor diffusivity coefficient being of the order $10^{-2} \mu\text{m}^2/\text{s}$ (56) we used a cytosolic diffusivity coefficient of $1 \mu\text{m}^2/\text{s}$. We then defined an area, $0.03 \mu\text{m}^2$, where “receptor-sharing” occurs. This area was based on the size of receptor aggregates ($0.1\text{-}0.3 \mu\text{m}$) (13, 44), microdomains (e.g. lipid rafts $0.02\text{-}0.5 \mu\text{m}$) (55), and membrane cytoskeletal corrals ($0.03\text{-}0.3 \mu\text{m}$) (61, 62). The “receptor-sharing” event is thus quantified as disassociation event-rebinding event occurring within 0.05s.

In this work we tested TKIs to determine their effects on “receptor-sharing”. Our results showed a decrease in association events and in “receptor-sharing” events when TKIs were added to the system. Given that “receptor-sharing” involves promiscuous signaling proteins interacting with receptors, and propagating signal more efficiently, this mechanism may be an important consideration for drug developers. Drug therapeutics which decrease “receptor-sharing” may be more effective at inhibiting signal propagation. The efficiency of “receptor-sharing” (unpublished data) is dependent on the binding kinetics, and thus therapeutics could potentially be tailored to reduce this mechanism. The “receptor-sharing” mechanism also hints at the selection of drug targets which may instead include promiscuous adaptor proteins.

It has long been established that signal transduction involves the coordination of proteins in time and space (47, 63-67). Experimental studies have shown disruption of

membrane recruitment reduces the amount of signal transduction (66, 68, 69). The plasma membrane, the nucleus of signal transduction, connects the extracellular with the intracellular via spatial organization of tethered membrane bound proteins, adaptor proteins, and signaling scaffolds (15, 70-72). The “receptor-sharing” mechanism may contribute significantly to this establishment. It may also prove to be a key regulator of signal transduction.

The predictive abilities of mathematical models enable us to test hypotheses which would be unattainable experimentally. In this work using a novel algorithm ACGMC, we tested our hypothesis that a “receptor-sharing” mechanism exists and contributes to the efficiency of signal transduction. Testing such a mechanism experimentally is not feasible using today’s technology. With limitations such as cytotoxicity to probes, poor conjugation, endosomal uptake, and resolution limitations, the importance of mathematical approaches becomes evident. This work demonstrates the predictive capacity of mathematical modeling to go beyond the present technology and address important biological questions.

Materials and Methods

Adaptively Coarse-Grained Monte Carlo (ACGMC)

The Adaptively Coarse-Grained Monte Carlo (ACGMC) algorithm extends for three-dimensional spatial modeling building on our previous algorithm, the SKMC, by including nine additional lattices eight of which extend into the cytosol and a lattice at the cytosol nucleus boundary. This framework allows us to study the formation of signaling scaffolds and to observe concentration gradients which result from these scaffolds.

The ACGMC, like the SKMC, begins by selecting a spatial location which could be either the cell membrane (1st lattice) or cytosol (2nd-9th lattices) or the nucleus (10th lattice). The selection is made by computing the probabilities for a spatial event (lattices 1-10).

$$\begin{aligned} P_{Lat\#1} &= \frac{\Gamma_{tot,Lat\#1}}{\Gamma_{tot}}, \\ P_{Lat\#2} &= \frac{\Gamma_{tot,Lat\#2}}{\Gamma_{tot}}, \text{and} \\ \dots P_{Lat\#10} &= \frac{\Gamma_{tot,Lat\#10}}{\Gamma_{tot}} \quad (1), \end{aligned}$$

where Γ_{tot} is defined as,

$$\Gamma_{tot} = \sum_{i=1}^{10} \Gamma_{tot,Lat\#i} \quad (2).$$

The probability distribution for the lattices, $\Gamma_{tot,Lat\#i}$, is defined as the sum of all transition rates for all SKMC events; however, the events differ in dimensionality and

type as shown in Table I. For example, on the cell membrane (1st lattice) receptors diffuse in 2D but react with cytosolic species (2nd lattice) on the lattice below such that reaction occurs in 2.5D. In the cytosol (3rd-9th lattice) species are able to diffuse and react in 3D. Cytosolic species in the boundary lattice which borders the cell membrane (2nd lattice) are able to react with neighboring receptors on the cell membrane but are not allowed to diffuse onto the membrane. At the boundary nucleus (10th lattice) species diffuse and react in 2.5D.

In the cytosol the following reaction occurs: $M_1 + M_2 \xrightleftharpoons[kb]{kf} M_{12}$, such that the product M_{12} occupies the single site k. This reaction is valid due to the large site separation distance of 10 nm in comparison to the small molecular masses of cytosolic species Shc, Grb2, and Sos 62 kDa 25 kDa, 11 kDa (20, 22, 73).

Once a lattice is chosen an event is selected, either reaction or diffusion for the SKMC algorithm as shown in Figure 26. Events are chosen, as in our previous algorithm(48), with the exception of dimensionality which changes the way Γ_{\max} and Γ_i^X are computed. The transition probabilities for cytosolic reactions in 3D are

$$M_1 + M_2 \xrightleftharpoons[kb]{kf} M_{12}, \quad \Gamma_i^r = \frac{k}{6} \sigma_i \sigma_j \quad (3)$$

in which reacting species (M_1 and M_2) occupy adjacent sites i and j and k has units of (molecules/site)⁻¹sec⁻¹. Diffusion in 3D is taken into account similarly

$$\Gamma_{i \rightarrow j}^d = \frac{1}{6} \Gamma^d \sigma_i (1 - \sigma_j) \quad j \in B_i \quad (4)$$

where B_i denotes the set of sites to which diffusion from site i can occur.

Table I lists all events, both reaction and diffusion, which make up Γ_{\max} . Γ_{\max} is defined as before but multiplying by a factor of 6 for all 3D event, by a factor of 4 for all 2D events, and a factor of 5 for all 2.5D events such that

$$\Gamma_{\max} = 4 \left(\frac{\Gamma^d}{4} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right)_{2D \text{ Events}} + 5 \left(\frac{\Gamma^d}{5} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right)_{2.5D \text{ Events}} + 6 \left(\frac{\Gamma^d}{6} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right)_{3D \text{ Events}} + \max \left\{ \sum_{\text{all backward reaction events}} \Gamma^r \right\} \quad (5)$$

The ACGMC operates like a single MC in which time is updated in a “combined” manner from Γ_{tot} with an average time step as, $\Delta t = \frac{1}{\Gamma_{tot}}$. Execution times vary depending on a chosen algorithm. The complete algorithm, which is shown in Figure 26, was implemented in Fortran 90. Due to the stochasticity of the algorithm, 10 simulations with different seeds for the random number generator were used for statistics.




Figure 26. Adaptively Coarse Grained Monte Carlo (ACGMC). Schematic of algorithmic details.

Adaptively Coarse-Graining & Diffusion Transition Rates

In ACGMC, we introduce coarse-graining by grouping microscopic sites into coarse cells, as shown in Figure 27. In our modeling schematic lattices one thru seven have microscopic spacing while lattices eight thru ten are coarse grained in the z-axis. Each coarse lattice L_C has m_x , m_y , and m_z coarse cells along x, y, z, such that the total number of cells is $m = m_x m_y m_z$. Each coarse cell C_j ($1 \leq j \leq m$) has q_j microscopic sites, in which $q_j = q_j^x q_j^y q_j^z$ is the number of microscopic sites in C_j along x, y and z axis. Diffusion between coarse-grained cells was rigorously derived by Chatterjee et al. (52),

$$\bar{C}_m(j \rightarrow i) = \frac{\Gamma_d}{q_j^z (q_j^z + q_i^z)} \eta_j (1 - \bar{\eta}_i) \quad (6)$$

in which, Γ_d is the diffusion transition rate, η_j , is the coarse-grained occupancy function at C_j is defined as

$$\eta_j = \sum_{p \in C_j} \sigma_p \quad (7).$$

While the occupancy of a coarse cell is in terms of the coarse cell coverage,

$$\bar{\eta}_j = \frac{1}{q_j} \sum_{p \in C_j} \sigma_p \quad (8)$$

which for the microscopic case reduces down to either zero for unoccupied or one for occupied.




Figure 27. Modeling Schematic. 1st lattice is the plasma membrane, lattices 2-10 are within the cytosol, lattice 10 is the cytosol-nucleus boundary. Lattices 1-7 have microscopic spacing of 10 nm, while lattices 8-10 are coarse grained with spacing of 0.5 μ m, and 1 μ m.

Validation of Approach

In order to validate our approach we initially placed all molecules within lattices one thru nine and allowed them to diffuse a distance of $1\ \mu\text{m}$ into lattice ten over time. Turning off reaction we have reflective boundary conditions at the cytosol membrane boundary (lattice two) and at the cytosol nucleus boundary (lattice 10) and periodic boundary conditions in the plane parallel to the plasma membrane. Using a diffusivity coefficient of $10\ \mu\text{m}^2\ \text{s}^{-1}$ and a distance $1\ \mu\text{m}$ and implementing reflective boundary conditions we compared the ACGMC with the PDE solution. Starting with a cytosolic species concentration $105\ \text{nM}$ we observe similar predications between the ACGMC and PDE profiles (Figure 28).




Figure 28. Validation of ACGMC. Concentration profile of cytosolic species (nM) at diffusivity of $10\mu\text{m}^2/\text{s}$ and a distance of $1\mu\text{m}$. magenta line: ACGMC blue line: PDE.

Table 5. Membrane & Cytocolic Microscopic Events and Transition

Microscopic event	Transition Rate	Lattices
Diffusion	$\Gamma_{i \rightarrow j}^D = \frac{1}{L_d} \Gamma^d \sigma_i (1 - \sigma_j) \quad j \in B_i$ <ul style="list-style-type: none"> - σ_i is the occupancy (discrete) that is 1, if site i is filled, and 0, if site i is empty (a single index indicating the site is used to simplify notation). - $\Gamma^d = D/a^2$, where a is the microscopic lattice pixel dimension taken equal to the encounter radius, and is the diffusivity of a receptor - L_d is the dimensions in which a particle can move in - B_i denotes the set of sites to which diffusion from site i can occur and is dependent on selected lattice for nearest neighboring sites 	$L_d = 4$ for Lattice #1 (cell membrane) $L_d = 5$ for Lattice #2 & #10 (boundary lattices) $L_d = 6$ for Lattices #3-#9 (cytosolic lattices)
Ligand Association Reaction ($S_i + M \xrightarrow{k} M^*$)	$\Gamma_i^R = k [S_i] \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #1 (cell membrane)
Ligand Dissociation Reaction ($M^* \xrightarrow{k} S_i + M$)	$\Gamma_i^R = k \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #1 (cell membrane)
Dimerization Reaction ($M^* + M^* \xrightarrow{k} D$)	$\Gamma_i^R = \frac{k}{2} \sigma_i \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [(receptors/site)$^{-1}$ s^{-1}]. 	Lattice #1 (cell membrane)
Decomposition Reaction ($D \xrightarrow{k} M^* + M^*$)	$\Gamma_i^R = k \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #1 (cell membrane)
Phosphorylation/Dephosphorylation Reaction ($D \xrightleftharpoons[k]{k} PD$)	$\Gamma_i^R = k \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #1 (cell membrane)
Cytosolic Association Reaction ($M_1 + M_2 \xrightarrow{k} M_{12}$)	$\Gamma_i^R = k \sigma_i \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #1 & #2 (cell membrane-cytosolic)
Cytosolic Dissociation Reaction ($M_{12} \xrightarrow{k} M_1 + M_2$)	$\Gamma_i^R = k \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #1 & #2 (cell membrane-cytosolic)
Cytosolic 1 st Order Reaction ($M_C \xrightarrow{k} M_C^*$)	$\Gamma_i^R = k \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #2-#10 (cytosolic-cytosolic)
Cytosolic Dimerization Reaction ($M_1 + M_2 \xrightarrow{k} M_{12}$)	$\Gamma_i^R = k \sigma_i \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #2-#10 (cytosolic-cytosolic)
Cytosolic Decomposition Reaction ($M_{12} \xrightarrow{k} M_1 + M_2$)	$\Gamma_i^R = k \sigma_i$ <ul style="list-style-type: none"> - k is the macroscopic reaction rate constant with units as [s^{-1}] 	Lattice #2-#10 (cytosolic-cytosolic)
Cytosolic Flux Event	$\Gamma_{i,Lat}^J = D(C_{Lat} - C_{SSA}) \sigma_{SSA} (1 - \sigma_j) \quad j \in B_i$ <ul style="list-style-type: none"> - C_{Lat} is the concentration of a given species within the spatial domain with units of [molecules/μm^3] - C_{SSA} is the concentration of a given species within the well-mixed domain with units of [molecules/μm^3] - r_{SSA} is the radius of the cell minus the cytosolic spatial domain - σ_j is the occupancy (discrete) that is 1, if site j is filled, and 0, if site j is empty (for lattice #10). - D diffusivity coefficient - B_i denotes the set of sites to which species can be populated on Lattice #10 	Lattice #10-SSA (cytosolic lattice-cytosolic well mixed)

Chapter 5. Future Work

Introduction

The Human Genome Project was one of the first modern biological endeavors which altered the way in which biology was practiced. Producing vast amounts of data the necessity became analysis and the need to develop computational tools. Data analysis of the human genome revealed higher order complexity and connectivity providing a new systematic prospective to biology. Unlike the reductionistic approach in which components are studied in isolation, the systematic approach investigates the role of a component within the system in order to understand the dynamics and stability of that system. Mathematical modeling has become essential to the systematic approach, with the ability to make predictions of the evolving system in time and space.

In this work I have showed mathematical models complementing experimentation with the ability to make novel predictions which may be unattainable experimentally. I envision this work proceeding in two directions: the first to make algorithmic improvements and increase computational efficiency, and the second to make future biological predictions of cell signaling processes. I will discuss each of these directions along with current work.

Computational Improvements

The algorithms which were developed implement a hybrid null-event spatial kinetic Monte Carlo, SKMC method. The algorithm calculates probabilities for every event, by calculating the maximum transition rate, Γ_{\max} which is defined as

$$\Gamma_{\max} = 4 \left(\frac{\Gamma^d}{4} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right) + \max \left\{ \sum_{\text{all backward reaction events}} \Gamma^r \right\}$$

where four takes into diffusion in two-dimensions on the plasma membrane. The probability for a given event is then defined as $P_i^X = \frac{\Gamma_i^X}{\Gamma_{\max}}$ the transition rate Γ_i^X for the given event divided by the the maximum transition rate. The algorithm will then select an occupied lattice site and update diffusion and reaction probabilities based on the state of the system. The maximum transition rate is greater than reaction and diffusion transition rates which gives rise to a null event probability P_i^{null} . Using this method events with low probabilities are penalized with additional null events. Incorporating null events means loss of computational time in selecting an occupied lattice site, determining probabilities based on neighboring species, and selecting an event.

Although the null-event algorithm has been used throughout this work computational efficiency could be improved by making this a rejection free algorithm. The stochastic simulation algorithm, SSA is an example of a rejection free method in

which probabilities are determined by computing a total transition rate, $\Gamma_{\text{total}} = \sum_{X=1}^{N_J} \sum_{i=1}^{N_L} \Gamma_i^X$

for all lattice sites, N_L and all events, N_J . Probabilities are then defined as $P_i^X = \frac{\Gamma_i^X}{\Gamma_{\text{total}}}$.

In order to eliminate the null bin, the total transition rate would be dependent on the state of the system.

Parrallization would be the next step to improving the overall algorithm efficiency. The *CSNSA* code could be parrallized in such a way that the lattice would be broken up into quadrants. Information would be passed between the quadrants using MPI. The most logical way to parrallize the *ACGMC* algorithm would be to break up the lattices and submit them to different processors. The current *ACGMC* algorithm operates in a similar

fashion by sending information about the current lattices as well as the surrounding lattices to the MC subroutine. Modifications would be made by implementing MPI to pass information between lattices. Parallelization I would expect to result in the greatest computational speedup.

Computational Predictions

Understanding the impact of IFN γ R clustering on immune response pathways

Introduction

Within a sea of lipids, a highly compartmentalized landscape exists, limiting the diffusion of signaling receptors and altering the spatial-temporal activation of signaling pathways. In this work we have unraveled the intricacies of spatial organization of the ErbB receptors and downstream signaling pathways, showing that clustering enhances signal transduction via “receptor-sharing” mechanism and that the MSK is a mechanism of receptor clustering. However, in order to come full circle, an understanding of how signal transduction regulates protein expression levels that as a result, restructures the plasma membrane, needs to be further elucidated.

Background

Caveolin-1, a 22-24 kDa integral protein, is the backbone to caveolar microdomains, and its expression level determines the number, size, and flexibility of caveolar domains. Caveolin-1 is regulated by many signal transduction pathways; PKC ϵ and androgen receptor signaling pathways lead to its upregulation whereas MAPK-Ras, c-Myc, INF γ R leads to its down regulation. A431 cells with decreased caveolin-1 expression have diverse membrane morphologies that alter the spatial organization of signaling receptors.

Electron microscopy images show interferon- γ receptor, INF γ R, colocalized within caveola microdomains. INF γ R, being responsible for the activation of macrophages during the early steps of innate immunity, induces direct antimicrobial mechanisms and up-regulates antigen processing and presentation pathways. Interferon- γ activates these immune responses via the Janus kinase (JAK) and signal transducer and activator of transcription (STAT) pathway. The initial signaling events of INF γ R are often the rate-limiting step and dependent on the spatial distribution of the INF γ R.

Experimental evidence shows that the relationship between INF γ and caveolar domains goes beyond spatial organization on the plasma membrane, gene networks of caveolin-1, and INF γ seem to be entwined. Stimulating macrophages with INF γ had an inhibitory effect on caveolin-1, the marker protein of caveola. In addition, when transfecting cell lines HT20 and DLD1 with caveolin-1 cDNA, there is downregulation in iNos, a metabolic product of INF γ immune response.

Motivation

A complex regulatory network exists between caveolin-1, $\text{INF}\gamma$ and iNos. This is a model system to explore two relationships: colocalization of $\text{INF}\gamma\text{R}$ on downstream signaling pathways, and protein expression levels on the structure of caveolar microdomains. In order to elucidate these relationships, a novel computational approach must be developed which would combine a spatial stochastic algorithm (CSNSA) with a deterministic circuit-based platform (BioXyce). This would enable spatial-temporal receptor dynamics to be coupled with regulatory and metabolic networks.

Approach

The computational challenges of this work are coupling deterministic and stochastic approaches which require the appropriate closure. A stochastic closure similar to that used in Katsoulakis et al. would be implemented. The hybrid algorithm would use the CSNSA to capture receptor diffusion reaction events on the plasma membrane, and the BioXyce circuit simulator would be implemented in the cytosol, nucleus, and mitochondria to simulate regulatory, metabolic, and signal transduction processes.

The coupled system would operate like a single MC with a stochastic variable time step Δt_s . Probabilities would be computed for the MC, based on transition rates for reaction diffusion events (MC). The MC algorithm would be iterated until a successful event is executed and time is updated $\Delta t_s = \frac{1}{\Gamma_{tot}}$ in which Γ_{tot} is the total of the transition rates. The MC would pause and the ODE would be executed for the time step, Δt_s . The procedure would be iterated until the final time is reached. Simulations would compare the hybrid algorithm with the CSNSA to verify appropriate levels of noise and temporal profiles of species.

Simulations

To elucidate the relationship between caveolin-1 and $\text{INF}\gamma\text{R}$, we will perform simulations at different concentrations of caveolin-1 and test the downstream signal propagation. We will then look at the upregulation and downregulation of gene networks based on different levels of signal. Gene networks will determine the protein expression levels of caveolin-1 and thus alter the size, density, and number of caveolar microdomains. Simulations will be preformed, looking at different combinations of regulatory networks and observing the diffusivity of $\text{INF}\gamma\text{R}$ through the caveolar microdomains. This work will be presented at the Q-bio conference, Santa Fe, for further details refer to Appendix B. It will be written up in a manuscript.

A Monte Carlo Based Approach for Determining Optimal Drug Efficacy in Different Cytoskeletal Distributions

Introduction

In the previous work we investigated various cytoskeletal densities in order to determine the effects on receptor clustering. Our results showed different receptor dynamics being dependent on cytoskeletal density. Experimental techniques show the MSK to vary in general morphology and distribution in different cell types. Based upon these fundamental differences in the cytoskeletal distributions, drug therapies can be tailored to specific tissue types. Although drugs have been designed for maximum potency in the past, emerging evidence suggests that the drug's disassociation rate may be altered to maximize signal as well as reduce desensitization. We hypothesized that optimal drug parameters are dependent on MSK density. Using a computational approach, we have tested different dissociation constants in various MSK densities, with the goal of finding an optimal drug design criteria that is specific for different cell types.

Background

Recent work using single particle tracking has revealed that the density of the MSK varies with cell type. Measurements of time proteins spent in confinement yield compartment sizes ranging from 32 nm for CHO-B1 cells to 230 nm for normal rat kidney fibroblast (NRK) cells. Similarly, three-dimensional reconstructed images of the plasma membrane in NRK cells and FRSK cells revealed differences in MSK density. The variation in MSK density appears to be of physiological significance, as observed in hippocampal neurons. These specialized cells contain a diffusional barrier highly concentrated with the MSK and transmembrane proteins located between the

somatodendritic and axonal domains, limiting the diffusion of even phospholipids. Such diffusive barriers have been observed in tight junctions, the neck of the bud in budding yeast, and between compartments in sperm. Although few studies have investigated the variation in MSK density and its role in cellular function, one could hypothesize that the diversity of MSK densities facilitates cell type specific physiology.

Motivation

With such variation in the MSK, we propose the concept of tailoring therapeutics with cell specificity in mind, an idea which to our knowledge has not been suggested before. Our hypothesis is that the membrane cytoskeletal distributions have diverse drug-binding parameters, k_{off} which correspond to optimal inhibition. The motivation for our hypothesis comes from previous work done by Woolf et al., performing Monte Carlo (MC) simulations a minimum rate of G-protein-coupled receptor (GPCR) phosphorylation was observed at an intermediate koff value. Their results were explained in terms of the spatial-temporal dynamics of receptors at different k_{off} parameters.

Approach & Results

In this work we have implemented a lattice-based spatial kinetic Monte Carlo (SKMC) algorithm which incorporates “picket-fence” boundaries enabling us to test different membrane cytoskeletal densities. Adding a drug to our system that inhibits epidermal growth factor receptor (ErbB or EGFR) homo and hetero dimerization, we are able to determine the optimal binding parameter k_{off} at a given cytoskeletal density. Our results show optimal inhibition at koff values of 100 s^{-1} , 8500 s^{-1} in the case without picket fences and at 100 s^{-1} , and 6000 s^{-1} at a picket fence distribution of $25 \text{ corrals}/\mu\text{m}^2$.

Future work

Future work will explain our results by computing the mean square displacement, MSD for all receptors bound to drug for a given k_{off} value. The results will show the MSD traveled by receptors bound to drug for a specific koff parameter, enabling us to compare

and contrast MSD as a function of time for different k_{off} parameters. This work is being prepared in a manuscript (see Appendix C.).

Microdomains and the Underlying Cytoskeleton Alter the Efficiency of “Receptor-Sharing”

Introduction

In our previous work we quantified the “receptor-sharing” mechanism; however, understanding the “receptor-sharing” mechanism in terms of microdomains and that the underlying cytoskeleton is of biological importance. In this work we implemented the ACGMC algorithm with microdomains and at different picket fence densities.

Background

Signal transduction, the cell’s communication network, comes to a crossroads on the highly compartmentalized plasma membrane. Often oversimplified in mathematical models, the plasma membrane is a facet of heterogeneity that alters receptor organization from transient confinement zones or signal transduction “hot spots” to depletion zones. The spatial heterogeneity of receptors comes as a result of sphingolipid-rich microdomains ‘lipid rafts’, caveolar formation, clathrin cages, and the hindering interactions with the membrane cytoskeleton. Diffusional limitations are the result, herein altering the initiation and activation of signal transduction pathways. Localized concentration gradients become a consequence of the membrane architecture and enhance the efficiency of signal propagation.

Motivation

Signal transduction and plasma membrane microdomains have typically been studied in isolation. Although growing evidence shows their intimate nature, with coalescing lipids and transient meshes altering signal transduction which come full circle, activating transcription factors that regulate protein expression levels on the cell membrane. The motivation for this work is to understand signal transduction in the context of membrane microdomains. Using the adaptively coarse-grained Monte Carlo (ACGMC), we observe signal transduction in terms of our previously proposed “receptor-sharing” mechanism under various microdomain conditions.

Future Work

We performed simulations which compare receptor sharing in different plasma membrane conditions such as lipid rafts, and with the underlying membrane cytoskeleton. Simulations have been run to test receptor sharing within lipid rafts at different parameters such as lipid raft size and receptor diffusivity through a lipid raft. Similarly, we have tested receptor sharing at three different MSK densities at varying receptor concentrations. Future work will include running additional simulations at biologically relevant diffusivities in lipid raft domains. The data will be further interrupted by quantification of the “receptor-sharing” mechanism and be presented in the most appropriate figures. This work will be written up in a manuscript; preliminary results are presented in Appendix D.

Monte Carlo Simulations Reveal Formation and Activation of the MAPK Pathway

Introduction

In our previous work we have looked at several mechanisms that are applicable for other signal transduction networks. In this work we investigate the mitogen-activated protein kinase, MAPK pathway to determine mechanisms of efficiency and adaptability. We utilize the adaptively coarse- grained Monte Carl (ACGMC) algorithm to observe the spatial temporal dynamics of scaffold formation.

Background

Signal transduction pathways process internal queues, relaying information that brings about a physiological response. Within this finely tuned system are mechanisms for efficiency and adaptability; to illustrate this are signaling scaffolds. The function of signaling scaffolds is similar to a circuit board in that both upstream and downstream signaling proteins can be spatially arranged and compartmentalized signaling nodes. Scaffolds are often necessary to catalyze enzymatic reactions, phosphorylate or dephosphorylate kinases, and to facilitate both positive and negative feedback loops.

Among the most extensively studied signaling scaffold systems is the mitogen-activated protein kinase, MAPK pathway which in mammalian cells regulates many key cellular processes such as proliferation, survival, motility, and differentiation. The initial signaling events in the MAPK pathway involve receptor ligation, dimerization, and phosphorylation which initiate the recruitment of adaptor signaling proteins such as Grb2 (growth factor receptor-binding protein 2), Shc (Src homology 2 and collagen domain protein), and SOS (son of the sevenless). The interactions of these adaptor proteins mimic the functionality of signaling scaffolds. Grb2 through its SH2 domain binds phosphorylated tyrosine residues of either EGFR or Shc and also has the capability of binding the guanine nucleotide exchange factor (GEF) SOS through its N-terminal SH3 domain. SOS exchanging a guanine transforms membrane bound (Ras-GDP) to its active form (Ras-GTP), thus inducing the MAPK pathway.

The aggregation of EGFR trickles down to the signaling cascade, effecting the spatial distribution of signaling scaffolds. A receptor bound scaffold in close proximity with another receptor can disassociate and then reattach to the proximal receptor,

enabling “receptor sharing” to occur between proteins in the signaling cascade. The effect of spatial distribution on signaling scaffolds has not been well elucidated. In this study we aim to uncover the downstream effects of receptor aggregation on the formation of signaling scaffolds.

Future Work

Implementing the Sasagawa et al. model of the MAPK pathway in the ACGMC, we will ask what cytosolic species are better at “receptor-sharing.” We will test binding rate affinities to observe optimal “receptor-sharing.” We will look at phosphatases and their receptor-sharing abilities and how they can dramatically alter signal transduction. We will cluster receptors to see if this leads to a colocalization of the Ras membrane-tethered protein. We will observe the formation of signaling scaffolds to observe its spatial temporal dynamics.

Conclusions

In the post Human Genome era, a new vision of personalized medicine is emerging which will be obtained through decades of multidisciplinary research that undoubtedly will build its foundation from predictions of mathematical models. Personalized medicine could entail going to a doctor having your human genome sequenced, analyzed, and a mathematical model generated which would describe cell signaling processes, gene and metabolic networks. The mathematical model would be personalized built from a patient’s genome and updated overtime. The personalized model could determine optimal drug treatment, and therefore provide patient specific care.

Although such a vision is beyond the horizon it will be achieved by multidisciplinary approaches relying on both experimentation and theory. It will incorporate the understanding of spatial-temporal dynamics of signaling proteins and its connection with the compartmentalized plasma membrane. This work will serve as a minute stepping stone in the direction of personalized medicine.

Appendix

Appendix A. The effective kinetic rate constant for diffusion-reaction problems derived by Lauffenburger and Linderman is

$$k_{PDE} = \frac{2\pi D_{AB} k_{Areal}}{2\pi D_{AB} + k_{Areal} \ln\left(\frac{b}{s}\right)} = \frac{4\pi D_{AB} k_{Areal}}{4\pi D_{AB} + k_{Areal} \ln\left(\frac{1}{\pi \rho_A s^2}\right)} \quad (1)$$

where $D_{AB} = D_A + D_B$, D_A and D_B are the diffusivities of A and B, ρ_A is the density of molecules of A (number of molecules per unit area), $b = \sqrt{1/\pi \rho_A}$ is one-half of the mean displacement between molecules of A distributed in a certain area, k_{Areal} is the intrinsic reaction rate constant in units of (receptors /area)⁻¹ s⁻¹, and s is the encounter radius. The relationship between k_{Areal} and k is given as, $k_{Areal} = ks^2$.

Using Eq 1. and combining it with the expression for resistance in a series model

$$k_{PDE} = \left(\frac{1}{k_{Areal}} + \frac{1}{k_{Diffusion}} \right)^{-1} \quad (2),$$

the diffusion-limited reaction rate constant, $k_{Diffusion}$ is determined to be

$$k_{Diffusion} = \frac{4\pi D_{AB}}{\ln\left(\frac{1}{\pi \rho_A s^2}\right)} \quad (3).$$

for diffusion limited cases, $k_{Diffusion} \ll k_{Areal}$, $k_{PDE} \approx k_{Areal}$ as was the case for the fitted parameters in this work. In diffusion limited cases, k_{PDE} is the fitted parameter and can be related back to the ODE parameter using Eq. 1. Using this approach, small differences were found in enzymatic reactions leading to the conclusion that omission of diffusion-limited contributions contributed negligible effects on the model.

Appendix B. Understanding the impact of IFN γ R clustering on immune response pathways

Michelle N. Costa¹, Elebeoba E. May²

Short Abstract — Interferon- γ plays an important role in macrophage activation during the early steps of innate immunity. Propagation of immune response via IFN γ is dependent on the spatial localization of the IFN γ R. EM images have found IFN γ R to be colocalized in caveolar membrane domains; whether this enhances or restricts signal remains to be elucidated. However, in an interesting twist, experimental evidence points to IFN γ as a negative regulator of caveolin-1. In order to understand the spatial-temporal dynamics of IFN γ R membrane localization and further investigate the impact of IFN γ R activation on gene and metabolic pathways that regulate caveolin-1 production we developed a simulation-based model using a coupled CSNSA-BioXyce platform that combines a spatial Monte Carlo method (CSNSA) with a circuit-based intracellular network simulator (BioXyce). In this work we explore the impact of receptor spatial organization on immune effector mechanisms and to complete the circle, the impact of IFN γ mediated effectors on spatial organization.

Keywords — Spatial organization, gene networks, IFN γ R, caveolin-1, spatial modeling

Purpose

Interferon- γ induces direct antimicrobial mechanisms and up-regulates antigen processing and presentation pathways (9). Interferon- γ activates these immune responses via the Janus kinase (JAK) and signal transducer and activator of transcription (STAT) pathway. The initial signaling events of IFN γ receptor (IFN γ R) are often the rate limiting step and dependent on the spatial distribution of the IFN γ receptors (3).

Electron microscopy using immunogold labeled particles has revealed aggregation of IFN γ R in membrane domains (1). With low numbers of receptors (10^2 to 10^3 receptors per cell (1)) spanning large distances (20um T cell and macrophage (3)) aggregation and colocalization are necessary mechanisms in the signal transduction pathway. Although the cell membrane is a vastly complex structure filled with heterogeneous microdomains IFN γ R has been observed to colocalize in caveolar membrane domains (1, 2, 4, 5).

Experimental evidence shows the link between IFN γ and caveolar domains goes beyond spatial organization on the plasma membrane, gene networks of caveolin-1 and IFN γ seem to be entwined. Stimulating macrophages with IFN γ had an inhibitory effect on caveolin-1, the marker protein of caveola (7). In addition when transfecting cell lines HT20 and DLD1 with caveolin-1 cDNA there is downregulation in iNOS (6), a metabolic product of IFN γ immune response. A complex regulatory network exists between caveolin-1, IFN γ and iNOS.

This distinct network has yet to be fully elucidated. Starting with a model of the IFN γ immune response (8) we have added gene networks of caveolin-1 as well as gene and metabolic networks of iNOS. In this study our aim is to understand the effects that spatial clustering has on IFN γ R downstream signaling using the coupled spatial non-spatial simulation algorithm (CSNSA) to simulate the IFN γ /IFN γ R mediated activation of the JAK-STAT signal transduction cascade. We then investigate the relationship between iNOS and caveolin-1 using a modeling framework which combines the CSNSA with BioXyce, where BioXyce is used to simulate the STAT mediated intracellular reactions that lead to the production of IFN γ activated genes and the metabolic reactions that lead to the production of immune effector molecules. We discuss the challenges and benefits of the coupled platform in providing a multiscale understanding of host immune response mechanisms.

References

- [1] Takaoka, A., et al. (2000) *Science* **288**, 2357-60.
- [2] Sadr, R., et al. (2001) *Cytokine* **14**, 19-26.
- [3] Wada, H., et al. (2008) *Nature* **452**, 768-72.
- [4] Lambert, et al. (2000) *Cytokine* **12**, 715-9.
- [5] Sadr, R., Lortat-Jacob, H. & Morel, G. (2000) *Cytokine* **12**, 711-4.
- [6] Sanchez, F. A., et al. (2008) *Am J Physiol Heart Circ Physiol*
- [7] Felley-Bosco, E. et al. (2000) *Proc Natl Acad Sci USA* **97**, 1433-9
- [8] Yamada, S., Shiono, S. Joo, A., and Yoshimura, A. (2003) *FEBS Letters* **534** 190-196
- [9] Schroder, K., Hertzog, P.J., Ravasi, T., Hume, D. (2004) *Jrnl Leukocyte Bio./* **75**, 163-18

¹Department of Chemical and Nuclear Engineering, University of New Mexico, Albuquerque, NM 87131.

E-mail: mcosta@unm.edu

²Complex Systems and Discrete Mathematics Department, Sandia National Laboratories.

E-mail: cemay@sandia.gov

Appendix C. A Monte Carlo Based Approach for Determining Optimal Drug Efficacy in Different Cytoskeletal Distributions

Michelle N. Costa^{4*}, Monica R. Mascarenas¹, Jeremy S. Edwards^{2,3,4},

¹ Department of Biology, University of New Mexico, Albuquerque, NM 87131

² ³ Cancer Research and Treatment Center, University of New Mexico Health Sciences Center, Albuquerque, NM 87131

³ Molecular Genetics and Microbiology, University of New Mexico Health Sciences Center, Albuquerque, NM 87131

⁴ Department of Chemical and Nuclear Engineering, University of New Mexico, Albuquerque, NM 87131

Correspondence to Michelle N. Costa (mcosta@unm.edu)

Abstract

The membrane skeleton (MSK) is the portion of the cytoskeleton that is closely associated with the cytoplasmic surface of the plasma membrane, varying in general morphology and distribution in different cell types. The MSK is composed of actin filaments and various integral proteins, which provides the cell with structure. It has been hypothesized that MSK rearrangements are associated with an upregulation in proliferative pathways, which is a hallmark of cancer. Based upon these fundamental differences in the cytoskeletal distributions, drug therapies can be tailored to specific tissue types. Although drugs have been designed for maximum potency in the past, emerging evidence suggests that the drug's disassociation rate may be altered to maximize signal as well as reduce desensitization. We hypothesized that optimal drug parameters are dependent on MSK distribution. Using a computational approach, we have tested different dissociation constants in various MSK distributions, with the goal of finding an optimal drug design criteria that is specific for different cell types. Our results suggest that therapeutic techniques should be tailored with tissue specificity in mind.

Keywords: cytoskeleton, picket-fence, drug efficacy, Spatial Monte Carlo

Introduction

The cytoskeleton is the sole structure spanning from the plasma membrane to the nucleus, coordinating all cellular processes in between (Janmey 1998; Janmey, Kas et al. 1998; Kusumi, Nakada et al. 2005). The membrane cytoskeleton (MSK), an interfacial structure of the bulk cytoskeleton located at the plasma membrane, plays an essential role in membrane function as well as cytoskeletal regulation (Fujiwara, Ritchie et al. 2002). Controlling processes such as endocytosis and exocytosis, the MSK provides the plasma membrane with structure yet adaptability (Dai and Sheetz 1999; Gaidarov and Keen 1999; Gaidarov, Santini et al. 1999; Valentijn, Valentijn et al. 2000; Gauthier, Rossier et al. 2009). Experimental evidence has shown that the MSK regulates site specific localization of transmembrane proteins (Bennett 1990; Saxton and Jacobson 1997; Bennett and Baines 2001; Pan, Kao et al. 2006), membrane tension (Togo, Alderton et al. 2000; Doherty and McMahon 2008; Lundmark, Doherty et al. 2008), global cell shape/volume (Linshaw, Fogel et al. 1992; Pedersen, Hoffmann et al. 2001; Sheetz, Sable et al. 2006), and membrane deformation at the leading edge (Svitkina and Borisy 1999; Svitkina and Borisy 1999). Arguably, the most important function of the MSK is the regulation of protein and lipid mobility on the plasma membrane (Fujiwara, Ritchie et al. 2002; Ike, Kosugi et al. 2003; Forgacs, Yook et al. 2004; Kusumi, Nakada et al. 2005). This particular function, an enigma that plagued the fluid mosaic model, is of great consequence for signal transduction.

Fujiwara et al. performed single particle tracking on the 1,2-dioleoyl*sn*-glycero-3-phosphoethanolamine (DOPE), a non-raft lipid, and observed interesting diffusive behavior when the DOPE lipid underwent short-term confinement, followed by long term

“hop” diffusion (Fujiwara, Ritchie et al. 2002). Hypothesizing an explanation to this curious diffusive behavior, cells were treated with latrunculin-A, an actin binding toxin which partially depolarizes filamentous actin, and showed an increase (factor of ~ 1.5) in diffusive compartment sizes. This evidence was indicative of the diffusional restrictions imposed by the MSK and supported the “picket-fence” model, also known as the anchored-protein picket model. The “picket-fence” model (Edidin, Kuo et al. 1991; Kusumi, Nakada et al. 2005) describes the plasma membrane as a landscape of boundaries: the fences are composed of the mesh-work of the MSK, while the pickets are integral proteins that are themselves tethered or confined by the MSK, inhibiting diffusion on the extracellular face.

Recent work using single particle tracking has revealed that the density of the MSK varies with cell type (Murase, Fujiwara et al. 2004). Measurements of time intervals that proteins have spent in confinement yield compartment sizes ranging from 32 nm for CHO-B1 cells to 230 nm for normal rat kidney fibroblast (NRK) cells (Murase, Fujiwara et al. 2004). Similarly, three-dimensional reconstructed images of the plasma membrane in NRK cells and FRSK cells revealed differences in MSK distribution (Morone, Fujiwara et al. 2006). The variation in MSK density appears to be of physiological significance, observed in hippocampal neurons. These specialized cells contain a diffusional barrier highly concentrated with the MSK and transmembrane proteins located between the somatodendritic and axonal domains, limiting the diffusion of even phospholipids (Nakada, Ritchie et al. 2003). Such diffusive barriers have been observed in tight junctions (Dragsten, Blumenthal et al. 1981; Dragsten, Handler et al. 1982; van Meer, Gumbiner et al. 1986; van Meer and Simons 1986), the neck of the bud in budding

yeast (Takizawa, Sil et al. 1997; Takizawa, DeRisi et al. 2000), and between compartments in sperm (Ladha, James et al. 1997). Although few studies have investigated the variation in MSK distribution and its role in cellular function, one could hypothesize that the diversity of MSK densities facilitates cell type specific physiology (Murase, Fujiwara et al. 2004; Morone, Fujiwara et al. 2006).

Variation in the MSK is not only a striking feature observed between cell types, but it is also seen in disease. Morphological changes occur in the MSK during cancer, specifically during metastasis in which cell-cell interactions are often disrupted (Chopra, Fligiel et al. 1990). Studies investigating metastatic characteristics exhibit a decreased interaction of tumor cells with platelets, a trait that is inherently dependent on the lateral mobility of specific receptors (Chopra, Hatfield et al. 1988; Chopra, Timar et al. 1992; Olorundare, Simmons et al. 1992). Antitumorigenic effects were observed when actin-binding proteins were overexpressed, leading to a stabilization of focal adhesion (Nebl, Oh et al. 2000). In addition, oncogenes that alter the actin binding protein expression have been identified (Vandekerckhove, Bauw et al. 1990; Janmey 1998). Another MSK function exploited in cancer, which has not been extensively studied, is the ability to enhance proliferative signal transduction pathways (e.g. clustering) and inhibit regulatory apoptotic pathways (Forgacs, Yook et al. 2004; Doherty and McMahon 2008).

With such variation in the MSK, we propose the concept of tailoring therapeutics with cell specificity in mind, an idea which to our knowledge has not been suggested before. Our hypothesis is that the membrane cytoskeletal distributions have diverse drug binding parameters, k_{off} which correspond to optimal drug inhibition. The motivation for our hypothesis comes from previous work (Woolf and Linderman 2003) done by Woolf et al.

in which performing Monte Carlo (MC) simulations, a minimum rate of G-protein-coupled receptor (GPCR) phosphorylation at an intermediate k_{off} value. Their results were explained in terms of the spatial-temporal dynamics of receptors at different k_{off} parameters.

In this work we have implemented a lattice based spatial kinetic Monte Carlo (SKMC) algorithm which incorporates “picket-fence” boundaries enabling us to test different membrane cytoskeletal distributions. Adding a drug to our system that inhibits epidermal growth factor receptor (ErbB or EGFR) homo and hetero dimerization, we are able to determine the optimal binding parameter k_{off} at a given cytoskeletal distribution. Our results show optimal inhibition at k_{off} values of 100 s^{-1} , 8500 s^{-1} in the case without picket fences and at 100 s^{-1} , and 6000 s^{-1} at a picket fence distribution of 25 corrals/ μm^2 .

Materials and Methods

Spatial Kinetic Monte Carlo (SKMC)

Simulations were performed using a spatial kinetic Monte Carlo (SKMC) algorithm which is a modified null-event lattice based MC, developed by our group (Mayawala, Vlachos et al. 2005; Mayawala, Vlachos et al. 2005). The spatial domain is a two-dimensional square lattice which is implemented with periodic boundary conditions. The algorithm begins by populating receptors onto the lattice. It then computes transition rates for reaction and diffusion rates as shown in Table II. Probabilities are determined based on the normalization transition rate Γ_{\max} for further details see (put Ref). The algorithm proceeds iteratively by selecting an occupied lattice site, choosing a successful (reaction or diffusion) or unsuccessful (null) event based on the probabilities. If a successful event was chosen, the event is executed and species populations as well as

time are updated, if the selected event is a “null-event” the algorithm proceeds until a successful event is chosen. A schematic of our algorithm is shown in Fig 2.

Picket Fences

Spatial boundaries were placed on the lattice to simulate the cytoskeletal distributions interacting with the plasma membrane. Using the term “picket-fences” to describe the cytoskeletal interactions, we quantified the density of boundaries in units of corrals/area, establishing three distributions of 25 corrals/ μm^2 , 100 corrals/ μm^2 , and 400 corrals/ μm^2 as shown in Fig 1. The “picket-fences” reduce the diffusional transition rates of the receptors, as shown from the definition of this rate:

$$\Gamma_{i \rightarrow j}^d = \frac{1}{4} \Gamma^d \sigma_i (1 - \sigma_j) \quad j \in B_i$$

where $\Gamma^d = \frac{4D}{a^2}$ is four times its diffusion coefficient, D , divide by its inner lattice distance, a squared. B_i signifies the set of sites in which diffusion from site σ_i can occur. Diffusion takes place in four directions, each direction with its occupancy function σ_j which is discrete. σ_j equals one if the site is filled, or zero if the site j is empty. This means that Eq. 1 can be either 0 or $\frac{1}{4} \Gamma^d$, depending on the occupancy of the neighboring site. Effective confinement of the receptor occurs when a receptor is neighbored by a “picket-fence”. In this instance, the σ_j term will be equal to one, yielding a diffusion transition rate of zero in that direction.

The membrane cytoskeleton is a very dynamical structure, undergoing rearrangements which enable receptors to escape the confinement of a corral. Single particle tracking shows that receptors hop to a new compartment every 10ms per 230-nm-diameter of

membrane cytoskeletal obstruction (Fujiwara, Ritchie et al. 2002). We simulate the picket-fence dynamics by placing breaks at a time step, Δt_{PF} of 10 ms per $0.04 \mu\text{m}^2$ of picket fences, therefore allowing receptors to diffuse through compartments. After an iteration of the SKMC, breaks are closed and Δt_{PF} is set to zero. This is consistent with the rate kinetics of actin dimerization (Adams, Matov et al. 2004; Vallotton, Gupton et al. 2004; Danuser and Waterman-Storer 2006; Deshpande, McMeeking et al. 2006; Andrews, Lidke et al. 2008) which is on the order of the algorithm's time step of 10 μs .

Drug Binding Kinetics & Parameters

A drug was added to the SKMC model to determine the most effective binding parameter within each picket fence density. This drug was modeled as a monoclonal antibody, inhibiting the formation of homo or heterodimers of the EGFR receptor, also known as ErbB1. A schematic of the primary and secondary inhibition due to the monoclonal antibody are presented in fig. 3. Three receptors belonging to the ErbB family are illustrated in the schematic in fig. 3, however our focus will be on ErbB1 represented as R1u, (receptor 1 unbound) and R1b (receptor 1 bound to ligand). The monoclonal antibody prevents EGF from binding to R1u, ultimately decreasing the probability of dimerization. Arrows highlighted as red indicate direct inhibition and arrows highlighted as blue indicate indirect inhibition.

In every simulation, the drug concentration was varied to maintain an average value of receptor bound drug occupancy of 2.5%, independent of the drug's binding constants (41). The concentration of the drug was determined according to the relationship

$$D = \frac{k_{on}}{k_{off}} \frac{f}{(1-f)}$$

where f is the fraction of receptors bound to ligand at equilibrium ($f = 0.025$). Defining drug occupancy in this way allows the instantaneous drug occupancy to change over the course of the simulation, while maintaining average receptor occupancy constant. This allows us to determine the spatial implications of a particular drug binding parameter.

Each MSK distribution contained the k_{off} values and drug concentrations indicated in Table III. For statistical purposes, ten simulations were performed for each of the ten k_{off} values.

Results

Determining Optimal Drug Parameters

To determine if membrane cytoskeletal distributions have diverse binding parameters, we tested different k_{off} parameters in each MSK density (Fig. 2). The simulations were performed without picket fences, and with picket fence densities of 25 corrals/ μm^2 , 100 corrals/ μm^2 , and 400 corrals/ μm^2 . At a time of two seconds, the total rate of dimerization of EGFR as a function of k_{off} value (Fig. 4) was observed. The most dramatic differences in the rate of dimerization were observed without picket fences (blue line) and at a picket fence distribution of 25 corrals/ μm^2 (pink line). Without picket fences, the optimal drug inhibition values (minimum points, blue line Fig 4) were observed at 100 s^{-1} and 8500 s^{-1} , while the least effective koff parameters were 6000 s^{-1} and 10000 s^{-1} (maximum points, blue line Fig 4). However, the optimal drug binding parameters, k_{off} , occurred at 100 s^{-1} and 6000 s^{-1} (minimum points, pink line Fig_4) for

the 25 corrals/ μm^2 distribution, and the least effective parameter was 4000 s^{-1} and 10000 s^{-1} .

Temporal profiles of optimal and least effective k_{off} Parameters

Next we wanted to determine the effectiveness of both the optimized parameters (minimums) and the least inhibitive parameters (maximums) over time. Fig. 5a shows the temporal profiles of the rate of EGFR dimerization for simulations performed at the selected k_{off} value, without fences. The k_{off} parameters demonstrated in Fig. 5a are the two least effective k_{off} values of 6000 s^{-1} , 10000 s^{-1} and the optimal k_{off} parameters of 100 s^{-1} , and 8500 s^{-1} . Differences are seen in the rate of EGFR dimerization between optimal and least effective parameters, however differences between the optimal values of 100 s^{-1} , and 8500 s^{-1} was marginal. Differences between least effective parameters of 6000 s^{-1} , 10000 s^{-1} were marginal. Similar results were observed in the temporal profiles of simulations with optimized and non-optimized parameters seen in the 25 corrals/ μm^2 distribution. However, the most effective parameters for the 25 corrals/ μm^2 distribution were 6000 s^{-1} , 100 s^{-1} and the least effective k_{off} values were 4000 s^{-1} , 10000 s^{-1} .

Different Predictions between SKMC vs ODE with Drug

We graphed the dimerization profiles using an ODE code without drug (blue line Fig.6A) and with optimal drug (red dots). Results from the SKMC (green line) without drug and SKMC with optimized drug (magenta line) are represented in Fig.6A. Differences in the rate of dimerization between the ODE without drug and ODE with optimized drug were unable to be determined, however, the results from the SKMC showed a decrease in dimerization rate with optimized drug when compared to the same

picket fence density without drug. In Fig.6B, the picket fence density of 100 corrals/ μm^2 yielded a similar result. A clear distinction can be seen between the SKMC with optimal drug (green line Fig 6B) when compared to the instance where there is not drug present (magenta line). In addition, looking at the SKMC model without drug (green line), Fig.6B also reveals a decrease in dimerization rate which is not depicted in the standard ODE model. The difference in dimerization rate can be attributed to the spatial parameters of the SKMC, which is not integrated into the ODE model. The ODE also neglects the heterogeneity of receptors on the membrane, suggesting that the SKMC is a more accurate model for simulating this biological system.

Spatially tracking ErbB1 receptors

Using the SKMC, we wanted to investigate the movement of a ErbB1 receptor in order to determine how different MSK distributions inhibit diffusion rates. In order to do so, we tracked each position on the lattice where the receptor traveled. From this data we determined a MSD plot for each MSK distribution, illustrated in Fig.8. The MSD plot in Fig. 8A shows a movement of the receptor within a domain with out picket fences (Fig. 8A blue line). The displacement of the receptor was dramatically greater than the other three picket fence densities, since the movement of the receptor was not confined by the MSK. Figure 8B shows the MSD plot of the picket fence density of 25 corrals/ μm^2 , which had a displacement that was greater than the picket fence densities of 100 corrals/ μm^2 and 400 corrals/ μm^2 , but still significantly less than the domain with out picket fences (Fig. 8B green line). Due to the confinement of the receptor within a dense MSK distribution, the movement of the receptor simulated in the MSK distributions of 100 corrals/ μm^2 (Fig. 8C red line) and 400 corrals/ μm^2 (Fig. 8D purple line) was limited.

Next we wanted to determine the relationship between the lifetime of the drug bound receptor and the corresponding k_{off} value. By tracking the location of all 40 ErbB1 receptors bound to drug, we were able to determine the life time of the drug as a function of time (Fig.7). As shown in Fig. 8, the relationship between the lifetime of a drug bound receptor and the k_{off} value is exponentially negative. This result indicates that the k_{off} value greatly influences the life time of drug, ultimately influencing the effective area of the drug.

Discussions

Experiments using single particle tracking have shown that different cell types express variations in corral sizes. CHO-B1 cells have corral sizes ranging from 3-4 nm, whereas NSK cells have corral sizes of 320 nm. Based upon the differences in MSK distributions, one would observe variations in the lateral diffusion of membrane bound receptors. Since receptor interactions depend on lateral diffusion, cell types expressing different MSK distributions will have distinct receptor dynamics. These results suggest that drug therapies should be tailored for cytoskeleton distributions.

After simulating various k_{off} values of a drug in different cytoskeleton distributions, the k_{off} parameters that were most effective depended upon the picket fence density. As shown in Fig. 4, the optimal and least effective k_{off} values occurring within the picket fence density of 25 corrals/ μm^2 (pink line) dramatically shifted from the k_{off} parameters that are present in the domain without picket fences. This result implies that the mechanism of the drug may vary depending on the type of cytoskeleton distribution. Different distributions of the picket fences alter the diffusivity of a drug bound receptor,

ultimately affecting the rate of dimerization. Within the picket fence density of 25 corrals/ μm^2 , the diffusion rates of a drug bound receptors are hindered by the MSK meshwork, reducing the dimerization rate and shifting the optimal and least optimal k_{off} values. Since different cell types express a variation in MSK densities, it is important to acknowledge the difference in diffusion rates that occur in different cell types.

However, the domains containing 100 corrals/ μm^2 and 400 corrals/ μm^2 displayed little variation in the rate of dimerization at different k_{off} values. As seen in Fig. 8C and Fig. 8D, the diffusivity of drug bound receptor may be inhibited at these dense picket fence distributions, causing the variations in the k_{off} parameter to have little effect. Previous work has suggested that the receptor concentrations alter the effects of dimerization within a confined corral. We speculate that more variation between k_{off} values would be observed at higher receptor concentration in these picket fence densities.

In addition to addressing tissue specific parameters, drug resistance and desensitization are also important factors that influence the development of a useful drug. Receptor desensitization can be defined as either a loss or reduction in receptor responsiveness after agonist exposure. In the past, drug development has searched for drugs that are potent and have a high affinity for their receptor, viewing desensitization simply as a side effect. However, investigating receptor activation and desensitization as events that can be altered independently can provide better insight into developing new drugs with less side effects and greater applications (Woolf and Linderman 2003).

Experiments performed by Lewis et. al applied a range of full and partial agonists to C-6 glioma cells in order to investigate the role of drug efficacy in agonist-induced desensitization. They observed desensitization in C-6 glioma cells that contained A77636, a compound shown to have a low k_{off} value. Compounds that contain a low k_{off} value result in a long residence time on the receptor, which holds the receptor in an active conformation (Lewis et. al 1998). In the case of a G-protein-coupled receptor (GPCR), holding the receptor in the active state allows phosphorylation to occur, which is known to be the first step in desensitization.

Our results direct the focus of finding a new k_{off} parameter that would reduce the effect of desensitization. Temporal profiles of the simulations without picket fences and the simulations with a picket fence density of 25 corrals/ μm^2 (Fig. 5a and Fig. 5b) demonstrate similar inhibitive behavior between a high k_{off} value and a low k_{off} value. Our data indicates that a compound containing a high k_{off} value may be just as effective as a compound with a low k_{off} value, but reduces the effects of desensitization. The shift in our data between the two picket fence densities also suggests that the efficiency of high k_{off} value may also depend on cell-specific parameters.

Performing simulations using a SKMC model has allowed us to determine the spatial and heterogeneity effects of a biological system that are not addressed in ODE models. As shown in Fig.6, a discrepancy occurred between a comparison between the SKMC and ODE simulations. After adding the optimal drug to the ODE model, the ODE failed to determine a difference in dimerization rate. Although, as shown in the SKMC

model in Fig. 6, there seems to be a discrete difference in dimerization rate that are effected by spatial factors that the SKMC is able to detect.

The work performed in this study would be difficult to test experimentally, due to the viability of cells, cost of supplies and the many variables that would be difficult to control. However, computational biology is a useful tool that can provide predictive results that can be used to guide experimental work. In this work, taking a computational approach has allowed us to explore the complex interactions between membrane receptors and the space around them. We have determined that different MSK distributions contain different k_{off} parameters, as well as identify similar efficiencies between high and low k_{off} values, suggesting new ways of addressing drug desensitization. Our findings suggest that determining specific parameters that are optimal for different cell types and tailoring drugs to meet that criterion may be the latest direction of effective drug design.

Future Work

Our results (Fig. 4) show marginal effects between k_{off} values at distributions of 100 corrals/ μm^2 and 400 corrals/ mm^2 . As shown in preliminary work, increased dimerization in each picket fence density depends on the receptor concentration. A low receptor concentration within a highly dense MSK distribution decreases the rate of dimerization. Alternatively, a high receptor concentration within a highly dense MSK distribution increases dimerization due to the clustering of receptors within a corral. We hypothesize that limited differences in k_{off} values is due to the effects of hindered diffusion; preventing EGFR receptors from interacting with homo and hetero partners. To

test this hypothesis, we will perform the same simulation at a higher receptor concentration of 80 receptors/lattice for both picket fence densities

To understand why some k_{off} parameters demonstrated greater inhibition than others, we will investigate the area covered by a receptor as a function of time. In order to do so, the position on the lattice where ErbB1 receptors are bound to drug will be tracked. In each MSK distribution, we will then determine the MSD for each drug bound receptor. Comparing the MSD of the most effective drug parameters with the least effective drug parameters may explain the mechanism for optimal inhibition. As determined from Fig. 7, the k_{off} value has an impact on the life time of the drug bound receptor, affecting the area traveled by the receptor. We hypothesize that the k_{off} parameters that show the greatest inhibition will transverse a greater portion of the plasma membrane.

Figure Legends

Figure 29 Three picket fence densities

A) 25 corrals/ μm^2 B) 100 corrals/ μm^2 , and C) 400 corrals/ μm^2

Error! Reference source not found.

Demonstrates a sequence of events that can occur within SKMC algorithm

Figure 30 Schematic of Drug Inhibition

Illustrates inhibition mechanism used by a monoclonal antibody. Primary inhibition is highlighted in pink, while secondary inhibition is indicated by blue.

Figure 31 Rate of EGFR Dimerization as a function of k_{off} for each MSK density

MSK distribution without picket fences (blue line) contain optimal k_{off} parameters of 100 s^{-1} , and 8500 s^{-1} and the least optimal k_{off} values of 6000 s^{-1} and 10000 s^{-1} . Optimal drug binding parameters, k_{off} , occurred at 100 s^{-1} and 6000 s^{-1} (minimum points, pink line) for the 25 corrals/ μm^2 distribution, and the least effective parameters were 4000 s^{-1} and 10000 s^{-1} . Total rate of dimerization was defined by the total amount of dimerization events that occurred at two seconds.

Figure 5 Temporal profile of maximum and minimum k_{off} values in MSK

distribution with out picket fences and with a picket fence density of 25 corrals/ μm^2

Without picket fences the k_{off} vales of 6000 s^{-1} (orange line), and 10000 s^{-1} (pink line) represent the least optimal k_{off} parameters. k_{off} parameters of 100 s^{-1} (blue line), and 8500 s^{-1} (green line) represent optimal k_{off} parameters. B) With a picket fence density of 25 corrals/ μm^2 , the most effective parameters for the 25 corrals/ μm^2 distribution were 6000 s^{-1} , 100 s^{-1} and the least effective k_{off} values were 4000 s^{-1} , 10000 s^{-1} .

Figure 6 Comparison between SKMC and ODE simulations

Figure 6A shows the dimerization profile within the 25 corrals/ μm^2 picket fence density. The SKMC illustrates a clear difference between the system without drug (teal line) and with optimized drug (purple line), whereas in the ODE simulation, there is no difference (red dotted line). Similar results are presented in Fig. 6B where the SKMC was

simulated in the 100 corrals/ μm^2 picket fence density. However, in the case without drug, (Fig. 6B teal line) the increased picket fence density revealed a decrease in the rate of dimerization when compared to the picket fence density of 25 corrals/ μm^2 without drug (Fig 6A teal line).

Figure 7 Average time vs. koff value

The average life time of a drug bound receptor as a function of time is demonstrated in this figure.

Figure 8 MSD plots

Fig. 8A shows a MSD plot containing all four picket fence densities. The displacement of the receptor in the domain without picket fences is dramatically greater than all other domains. Fig. 8B demonstrates a scaled down view of the picket fence densities of 25 corrals/ μm^2 , 100 corrals/ μm^2 and 400corrals/ μm^2 . Fig. 8C shows the MSD plot of 100 corrals/ μm^2 and 400corrals/ μm^2 , which are scaled down further. Fig.8D shows the MSD plot for the picket fence density of 400corrals/ μm^2 exclusively.




Figure 1




Figure 2




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

Table I. ErbB Reactions and Reaction Rates

Reactions	Rate Constants (1/s)	
1. $R1u + \text{EGF} \leftrightarrow R1b$	$K_f = 0.003$	$K_b = 0.004$
2. $R3u + \text{NRG} \leftrightarrow R3b$	$K_f = 0.003$	$K_b = 0.004$
3. $R1uR1u + \text{EGF} \leftrightarrow R1bR1u$	$K_f = 0.003$	$K_b = 0.004$
4. $R1bR1u + \text{EGF} \leftrightarrow R1bR1b$	$K_f = 0.003$	$K_b = 0.004$
5. $R1uR2 + \text{EGF} \rightarrow R1bR2$	$K_f = 0.003$	$K_b = 0.004$
6. $R3uR2 + \text{NRG} \leftrightarrow R3bR2$	$K_f = 0.003$	$K_b = 0.00004$
7. $R3uR3u + \text{NRG} \leftrightarrow R3bR3u$	$K_f = 0.003$	$K_b = 0.004$

8. $R3bR3u + NRG \leftarrow \rightarrow R3bR3b$	$K_f = 0.003$	$K_b = 0.004$
9. $R1uR3u + EGF \leftarrow \rightarrow R1bR3u$	$K_f = 0.0015$	$K_b = 0.006$
10. $R1uR3u + NRG \leftarrow \rightarrow R1uR3b$	$K_f = 0.003$	$K_b = 0.004$
11. $R1uR3b + EGF \leftarrow \rightarrow R1bR3b$	$K_f = 0.0015$	$K_b = 0.006$
12. $R1bR3u + NRG \leftarrow \rightarrow R1bR3b$	$K_f = 0.003$	$K_b = 0.004$
13. $R1u + R1u \leftarrow \rightarrow R1uR1u$	$K_f = 0.01$	$K_b = 0.04$
14. $R1b + R1u \leftarrow \rightarrow R1uR1b$	$K_f = 0.01$	$K_b = 0.2$
15. $R1b + R1b \leftarrow \rightarrow R1bR1b$	$K_f = 0.01$	$K_b = 0.04$
16. $R2 + R2 \leftarrow \rightarrow R2R2$	$K_f = 0.01$	$K_b = 1$
17. $R3u + 3Ru \leftarrow \rightarrow R3uR3u$	$K_f = 0.01$	$K_b = 0.4$
18. $R3b + R3u \leftarrow \rightarrow R3bR3u$	$K_f = 0.01$	$K_b = 0.2$
19. $R3b + R3b \leftarrow \rightarrow R3bR3b$	$K_f = 0.01$	$K_b = 0.04$
20. $R1u + R3u \leftarrow \rightarrow R1uR3u$	$K_f = 0.01$	$K_b = 0.04$
21. $R1b + R3u \leftarrow \rightarrow R1bR3u$	$K_f = 0.01$	$K_b = 0.02$
22. $R1u + R3b \leftarrow \rightarrow R1uR3b$	$K_f = 0.01$	$K_b = 0.02$
23. $R1b + R3b \leftarrow \rightarrow R1bR3b$	$K_f = 0.01$	$K_b = 0.04$
24. $R1u + R2 \leftarrow \rightarrow R1uR2$	$K_f = 0.01$	$K_b = 0.8$
25. $R1b + R2 \leftarrow \rightarrow R1bR2$	$K_f = 0.01$	$K_b = 0.4$
26. $R3u + R2 \leftarrow \rightarrow R3uR2$	$K_f = 0.01$	$K_b = 0.8$
27. $R3b + R2 \leftarrow \rightarrow R3bR2$	$K_f = 0.01$	$K_b = 0.4$
28. $R1uR1u \leftarrow \rightarrow p R1uR1u$	$K_f = 0.01$	$K_b = 0.16$
29. $R1uR1u \leftarrow \rightarrow p R1uR1u$	$K_f = 0.03$	$K_b = 0.08$
30. $R1uR1b \leftarrow \rightarrow p R1uR1b$	$K_f = 0.02$	$K_b = 0.08$

31. $R1bR1b \leftarrow \rightarrow pR1bR1b$	$K_f = 0.04$	$K_b = 0.02$
32. $pR1bR1b \leftarrow \rightarrow pR1bpR1b$	$K_f = 0.12$	$K_b = 0.01$
33. $R1uR2 \leftarrow \rightarrow pR1uR2$	$K_f = 0.008$	$K_b = 0.16$
34. $pR1uR2 \leftarrow \rightarrow pR1upR2$	$K_f = 0.024$	$K_b = 0.08$
35. $R1bR2 \leftarrow \rightarrow pR1bR2$	$K_f = 0.024$	$K_b = 0.08$
36. $pR1bR2 \leftarrow \rightarrow pR1bpR2$	$K_f = 0.016$	$K_b = 0.08$
37. $pR2R1b \leftarrow \rightarrow pR2pR1b$	$K_f = 0.048$	$K_b = 0.04$
38. $R3uR2 \leftarrow \rightarrow pR2R3u$	$K_f = 0.008$	$K_b = 0.16$
39. $R3bR2 \leftarrow \rightarrow pR2R3b$	$K_f = 0.016$	$K_b = 0.08$
40. $R1uR3b \leftarrow \rightarrow pR1uR3u$	$K_f = 0.008$	$K_b = 0.16$
41. $R1bR3u \leftarrow \rightarrow pR1bR3u$	$K_f = 0.016$	$K_b = 0.08$
42. $R3bR1u \leftarrow \rightarrow pR1uR3b$	$K_f = 0.016$	$K_b = 0.08$
43. $R1bR3b \leftarrow \rightarrow pR1bR3b$	$K_f = 0.032$	$K_b = 0.02$
44. $R1u + D \rightarrow R1b$	$K_f = 2.6 \cdot 10^3$	$k_{off} = 100$ $k_{off} = 500$ $k_{off} = 1000$ $k_{off} = 1500$ $k_{off} = 2500$ $k_{off} = 3500$ $k_{off} = 4000$ $k_{off} = 6000$ $k_{off} = 8500$ $k_{off} = 10000$

Table II. Membrane Microscopic Events and Transition rates

Microscopic Event

Transition Rate

Diffusion

$$\Gamma_{i \rightarrow j}^D = \frac{1}{4} \Gamma^D \sigma_i (1 - \sigma_j) \quad j \in B_i$$

σ_i is the occupancy(discrete) that is 1, if site i is filled, and 0, if site i is empty (a single index indicating the site is used to simplify notation).

$$\Gamma^D = \frac{D}{a^2}, \text{ where } a \text{ is the microscopic}$$

lattice pixel dimension taken equal to the encounter radius, and D is the diffusivity of a receptor

B_i denotes the set of sites to which diffusion from site

i can occur which includes all 4 first-nearest neighboring sites

Reactions


Ligand Association Reaction

$$\Gamma_i^R = k [S_L] \sigma_i$$

($S_L + M \rightarrow M^*$)

k is the macroscopic reaction rate constant

Ligand Disassociation Reaction



with units as $[s^{-1}]$

$$\Gamma_i^R = k\sigma_i$$

k is the macroscopic reaction rate constant

with units as $[s^{-1}]$

Dimerization Reaction



$$\Gamma_i^R = \frac{k}{2}\sigma_i\sigma_j$$

k is the macroscopic reaction rate constant

with units as $[(\text{receptors/sites})^{-1} s^{-1}]$

Decomposition Reaction




$$\Gamma_i^R = k\sigma_i$$

k is the macroscopic reaction rate constant

with units as $[s^{-1}]$

Phosphorylation/Dephosphorylation
Reaction



$$\Gamma_i^R = k\sigma_i$$

k is the macroscopic reaction rate constant

with units as $[s^{-1}]$

- Adams, M. C., A. Matov, et al. (2004). "Signal analysis of total internal reflection fluorescent speckle microscopy (TIR-FSM) and wide-field epi-fluorescence FSM of the actin cytoskeleton and focal adhesions in living cells." J Microsc 216(Pt 2): 138-52.
- Andrews, N. L., K. A. Lidke, et al. (2008). "Actin restricts FcεRI diffusion and facilitates antigen-induced receptor immobilization." Nat Cell Biol 10(8): 955-63.
- Bennett, V. (1990). "Spectrin-based membrane skeleton: a multipotential adaptor between plasma membrane and cytoplasm." Physiol Rev 70(4): 1029-65.
- Bennett, V. and A. J. Baines (2001). "Spectrin and ankyrin-based pathways: metazoan inventions for integrating cells into tissues." Physiol Rev 81(3): 1353-92.
- Chopra, H., S. E. Fligiel, et al. (1990). "An in vivo study of the role of the tumor cell cytoskeleton in tumor cell-platelet-endothelial cell interactions." Cancer Res 50(23): 7686-96.
- Chopra, H., J. S. Hatfield, et al. (1988). "Role of tumor cytoskeleton and membrane glycoprotein IRGpIIb/IIIa in platelet adhesion to tumor cell membrane and tumor cell-induced platelet aggregation." Cancer Res 48(13): 3787-800.
- Chopra, H., J. Timar, et al. (1992). "Is there a role for the tumor cell integrin alpha IIb beta 3 and cytoskeleton in tumor cell-platelet interaction?" Clin Exp Metastasis 10(2): 125-37.
- Dai, J. and M. P. Sheetz (1999). "Membrane tether formation from blebbing cells." Biophys J 77(6): 3363-70.
- Danuser, G. and C. M. Waterman-Storer (2006). "Quantitative fluorescent speckle microscopy of cytoskeleton dynamics." Annu Rev Biophys Biomol Struct 35: 361-87.
- Deshpande, V. S., R. M. McMeeking, et al. (2006). "A bio-chemo-mechanical model for cell contractility." Proc Natl Acad Sci U S A 103(38): 14015-20.
- Doherty, G. J. and H. T. McMahon (2008). "Mediation, modulation, and consequences of membrane-cytoskeleton interactions." Annu Rev Biophys 37: 65-95.
- Dragsten, P. R., R. Blumenthal, et al. (1981). "Membrane asymmetry in epithelia: is the tight junction a barrier to diffusion in the plasma membrane?" Nature 294(5843): 718-22.
- Dragsten, P. R., J. S. Handler, et al. (1982). "Asymmetry in epithelial cells: is the tight junction a barrier to lateral diffusion in the plasma membrane?" Prog Clin Biol Res 91: 525-36.

- Edidin, M., S. C. Kuo, et al. (1991). "Lateral movements of membrane glycoproteins restricted by dynamic cytoplasmic barriers." Science 254(5036): 1379-82.
- Forgacs, G., S. H. Yook, et al. (2004). "Role of the cytoskeleton in signaling networks." J Cell Sci 117(Pt 13): 2769-75.
- Fujiwara, T., K. Ritchie, et al. (2002). "Phospholipids undergo hop diffusion in compartmentalized cell membrane." J Cell Biol 157(6): 1071-81.
- Gaidarov, I. and J. H. Keen (1999). "Phosphoinositide-AP-2 interactions required for targeting to plasma membrane clathrin-coated pits." J Cell Biol 146(4): 755-64.
- Gaidarov, I., F. Santini, et al. (1999). "Spatial control of coated-pit dynamics in living cells." Nat Cell Biol 1(1): 1-7.
- Gauthier, N. C., O. M. Rossier, et al. (2009). "Plasma Membrane Area Increases with Spread Area by Exocytosis of a GPI Anchored Protein Compartment." Mol Biol Cell.
- Ike, H., A. Kosugi, et al. (2003). "Mechanism of Lck recruitment to the T-cell receptor cluster as studied by single-molecule-fluorescence video imaging." Chemphyschem 4(6): 620-6.
- Janmey, P. A. (1998). "The cytoskeleton and cell signaling: component localization and mechanical coupling." Physiol Rev 78(3): 763-81.
- Janmey, P. A., J. Kas, et al. (1998). "Cytoskeletal networks and filament bundles: regulation by proteins and polycations." Biol Bull 194(3): 334-5; discussion 335-6.
- Kusumi, A., C. Nakada, et al. (2005). "Paradigm shift of the plasma membrane concept from the two-dimensional continuum fluid to the partitioned fluid: high-speed single-molecule tracking of membrane molecules." Annu Rev Biophys Biomol Struct 34: 351-78.
- Ladha, S., P. S. James, et al. (1997). "Lateral mobility of plasma membrane lipids in bull spermatozoa: heterogeneity between surface domains and rigidification following cell death." J Cell Sci 110 (Pt 9): 1041-50.
- Linshaw, M. A., C. A. Fogel, et al. (1992). "Role of cytoskeleton in volume regulation of rabbit proximal tubule in dilute medium." Am J Physiol 262(1 Pt 2): F144-50.
- Lundmark, R., G. J. Doherty, et al. (2008). "Arf family GTP loading is activated by, and generates, positive membrane curvature." Biochem J 414(2): 189-94.
- Mayawala, K., D. G. Vlachos, et al. (2005). "Computational modeling reveals molecular details of epidermal growth factor binding." BMC Cell Biol 6: 41.

- Mayawala, K., D. G. Vlachos, et al. (2005). "Heterogeneities in EGF receptor density at the cell surface can lead to concave up scatchard plot of EGF binding." FEBS Lett 579(14): 3043-7.
- Morone, N., T. Fujiwara, et al. (2006). "Three-dimensional reconstruction of the membrane skeleton at the plasma membrane interface by electron tomography." J Cell Biol 174(6): 851-62.
- Murase, K., T. Fujiwara, et al. (2004). "Ultrafine membrane compartments for molecular diffusion as revealed by single molecule techniques." Biophys J 86(6): 4075-93.
- Nakada, C., K. Ritchie, et al. (2003). "Accumulation of anchored proteins forms membrane diffusion barriers during neuronal polarization." Nat Cell Biol 5(7): 626-32.
- Nebi, T., S. W. Oh, et al. (2000). "Membrane cytoskeleton: PIP(2) pulls the strings." Curr Biol 10(9): R351-4.
- Olorundare, O. E., S. R. Simmons, et al. (1992). "Cytochalasin D and E: effects on fibrinogen receptor movement and cytoskeletal reorganization in fully spread, surface-activated platelets: a correlative light and electron microscopic investigation." Blood 79(1): 99-109.
- Pan, Z., T. Kao, et al. (2006). "A common ankyrin-G-based mechanism retains KCNQ and NaV channels at electrically active domains of the axon." J Neurosci 26(10): 2599-613.
- Pedersen, S. F., E. K. Hoffmann, et al. (2001). "The cytoskeleton and cell volume regulation." Comp Biochem Physiol A Mol Integr Physiol 130(3): 385-99.
- Saxton, M. J. and K. Jacobson (1997). "Single-particle tracking: applications to membrane dynamics." Annu Rev Biophys Biomol Struct 26: 373-99.
- Sheetz, M. P., J. E. Sable, et al. (2006). "Continuous membrane-cytoskeleton adhesion requires continuous accommodation to lipid and cytoskeleton dynamics." Annu Rev Biophys Biomol Struct 35: 417-34.
- Svitkina, T. M. and G. G. Borisy (1999). "Arp2/3 complex and actin depolymerizing factor/cofilin in dendritic organization and treadmilling of actin filament array in lamellipodia." J Cell Biol 145(5): 1009-26.
- Svitkina, T. M. and G. G. Borisy (1999). "Progress in protrusion: the tell-tale scar." Trends Biochem Sci 24(11): 432-6.
- Takizawa, P. A., J. L. DeRisi, et al. (2000). "Plasma membrane compartmentalization in yeast by messenger RNA transport and a septin diffusion barrier." Science 290(5490): 341-4.

- Takizawa, P. A., A. Sil, et al. (1997). "Actin-dependent localization of an RNA encoding a cell-fate determinant in yeast." Nature 389(6646): 90-3.
- Togo, T., J. M. Alderton, et al. (2000). "The mechanism of cell membrane repair." Zygote 8 Suppl 1: S31-2.
- Valentijn, J. A., K. Valentijn, et al. (2000). "Actin coating of secretory granules during regulated exocytosis correlates with the release of rab3D." Proc Natl Acad Sci U S A 97(3): 1091-5.
- Vallotton, P., S. L. Gupton, et al. (2004). "Simultaneous mapping of filamentous actin flow and turnover in migrating cells by quantitative fluorescent speckle microscopy." Proc Natl Acad Sci U S A 101(26): 9660-5.
- van Meer, G., B. Gumbiner, et al. (1986). "The tight junction does not allow lipid molecules to diffuse from one epithelial cell to the next." Nature 322(6080): 639-41.
- van Meer, G. and K. Simons (1986). "The function of tight junctions in maintaining differences in lipid composition between the apical and the basolateral cell surface domains of MDCK cells." EMBO J 5(7): 1455-64.
- Vandekerckhove, J., G. Bauw, et al. (1990). "Comparative two-dimensional gel analysis and microsequencing identifies gelsolin as one of the most prominent downregulated markers of transformed human fibroblast and epithelial cells." J Cell Biol 111(1): 95-102.
- Woolf, P. J. and J. J. Linderman (2003). "Untangling ligand induced activation and desensitization of G-protein-coupled receptors." Biophys J 84(1): 3-13.

Appendix D. Microdomains and the Underlying Cytoskeleton Alter the Efficiency of “Receptor-Sharing”

Michelle Costa¹, Dionisios G. Vlachos³ & Jeremy S. Edwards^{1,2,4,*}

¹ Department of Chemical and Nuclear Engineering, University of New Mexico, Albuquerque, NM 87131

² Cancer Research and Treatment Center, University of New Mexico Health Sciences Center, Albuquerque, NM 87131

³ Department of Chemical Engineering, University of Delaware, Newark, DE 19716

⁴ Molecular Genetics and Microbiology, University of New Mexico Health Sciences Center, Albuquerque, NM 87131

Correspondence to Jeremy S. Edwards (jsedwards@salud.unm.edu)

Abstract

Experimental evidence suggests

Keywords: EGFR, spatial modeling, Clustering. MSK, Picket Fences

Introduction

Signal transduction, the cell's communication network, comes to a crossroads on the highly compartmentalized plasma membrane. Often oversimplified in mathematical models (1-3), the plasma membrane is a facet of heterogeneity that alters receptor organization from transient confinement zones (4, 5) or signal transduction "hot spots" to depletion zones. The spatial heterogeneity of receptors comes as a result of sphingolipid-rich microdomains 'lipid rafts' (6-8), caveolar formation (9, 10), clathrin cages (11-13), and the hindering interactions with the membrane cytoskeleton (14-16). Diffusional limitations are the result herein altering the initiation, and activation of signal transduction pathways (17). Localized concentration gradients become a consequence of the membrane architecture and enhance the efficiency of signal propagation (15, 18).

The notion that the plasma membrane is a highly compartmentalized surface with intrinsic microdomains, came a few years after the fluid mosaic model was proposed by Singer and Nicholson in 1972 (19). Two major hypotheses, the lipid raft (20-22) and the membrane skeleton "picket fence" model (23, 24) were proposed to describe the structure of the plasma membrane. The lipid raft hypothesis was postulated to explain the aggregation of glycosphingolipids in the Golgi apparatus before being sorted to the apical surface of polarized epithelial cells (20). Using cold non-ionic extraction techniques on specialized lipid fractions taken from Golgi membranes, Simons et al. found that the lipid domains were detergent-resistant (21). These domains became known as lipid rafts.

The composition of lipid rafts consists of a highly saturated assembly of cholesterol and sphingolipids which form a lipid order structure ranging from 10-300 nm in size (25, 26). Perhaps one of the more interesting phenomena of lipid rafts is their ability to include or

exclude proteins. The amino acid structure often dictates favorability with hydrophobic modifications increasing affinity for lipid raft domains(18). Signaling complexes formed within the lipid raft microenvironments are therefore, protected by a buffer zone which excludes non-raft enzymes such as membrane phosphatases.

Diffusion within lipid rafts decreases by a factor of three to five as observed on artificial membranes (27-29) and by a factor of two in SPT experiments(5). Crowding effects are thought to be responsible for the decreased diffusivity, which is a mechanism of clustering and a precursor to signaling scaffold formation (10, 30). Experimental techniques have shown lipid rafts intimately involved in signal transduction pathways (18, 31-33). Take for example the inhibition of H-Ras mediated Raf activation, occurring when cells are treated with methyl- β -cyclodextrin, an agent that depletes cholesterol (34). The evidence is suggestive that H-Ras signals only in the presence of lipid rafts (35, 36).

Another insightful theory, describing the structure of the plasma membrane, the membrane skeleton “*picket fence*” model was proposed in an attempt to explain anomalous diffusion (23, 24). Using FRAP techniques to measure diffusion rates in spectrin-deficient mutant mouse erythrocytes, Sheetz et al found that the transmembrane protein band 3 diffuses 10 times faster than in wild type erythrocytes (24). Knowing that the spectrin meshwork, makes up the membrane skeleton, a model describing the hinderance on protein diffusion, the picket fence model was hypothesized (23). The model describes the plasma membrane as a meshwork of “fences”, interactions from the underlying membrane cytoskeleton (MSK), and “pickets” or protruding transmembrane proteins inhibited by the MSK on the cytoplasmic face. This model was further supported by single particle tracking of the DOPE lipid which underwent short term confinement followed by long term “hop” diffusion (14, 37-39).

The MSK is known to vary in structure and density between cell types, with corral sizes ranging from 30 to 230 nm (40, 41). Along with cell specific variation, morphological changes occur in the MSK during disease (42, 43). The density of the MSK has the capacity to inhibit as well as enhance signal transduction pathways (44). Receptor signaling events: oligomerization, crosslinking, and scaffold formation occurring within the constraints of the MSK lead to concentration gradients which facilitate downstream signaling (45, 46). At the same time the MSK can be restrictive hindering diffusion and isolating receptors. Take for example; hippocampal neurons with a diffusional barrier highly concentrated in MSK, phospholipid movement is restricted (37).

Signal transduction and plasma membrane microdomains have typically been studied in isolation. Although growing evidence shows their intimate nature, with coalescing lipids and transient meshes altering signal transduction (47, 48) which comes full circle activating transcription factors that regulate protein expression levels on the cell membrane (34). The motivation for this work is to understand signal transduction in the context of membrane microdomains. Using the adaptively coarse-grained Monte Carlo (ACGMC) we observe signal transduction in terms of our previously proposed “receptor-sharing” mechanism under various microdomain conditions.

Results/Future Work

Effects of Microdomains and MSK on Association Events

To determine the effects of microdomains and the underlying MSK on association events and receptor sharing events, simulations were performed for a lipid raft of size 200nm with diffusivity decreased by 100, for picket fences with densities of 16 corrals/ μm^2 , 100 corrals/ μm^2 , 400 corrals/ μm^2 , and a control (without lipid rafts or an MSK density). Fig 1. shows our results we observe a significant increase between the control and simulations performed with either picket fences or lipid rafts. The most significant increase occurred between the lipid raft and the control. Further investigating these results it was determined that decreasing the diffusivity by a factor 100 is not representative of diffusivity of a receptor through a lipid raft which decreases by a factor of 5. Fig 2. shows the distributions of receptor binding events, from these results we observe an increase in “receptor-sharing” for the lipid raft condition. The picket fence densities seem to inhibit “receptor-sharing”. Simulations with an experimentally determined diffusivity in the lipid raft will be run. This will reduce the number of association events and thus “receptor-sharing”, however I hypothesis that the number of “receptor-sharing” events will increase from the control.

Simulations will also be performed to further elucidate the results shown in Fig 1&2. Performing simulations in which the diffusivity and the size will be varied in lipid rafts will allow us to determine the effects of lipid rafts. Similarly we will perform simulations at different receptor concentrations to determine which picket fence density will increase “receptor-sharing”.

Materials and Methods

Adaptively Coarse-Grained Monte Carlo (ACGMC)

The Adaptively Coarse-Grained Monte Carlo (ACGMC) algorithm extends for three-dimensional spatial modeling building on our previous algorithm, the SKMC, by including nine additional lattices eight of which extend into the cytosol and the ninth on the nucleus. This framework allows us to study the formation of signaling scaffolds and to observe concentration gradients which result from these scaffolds.

The ACGMC, like the SKMC, begins by selecting a spatial location which could be either the cell membrane (1st lattice) or cytosol (2nd-9th lattices) or the nucleus (10th lattice). The selection is made by computing the probabilities for a spatial event (lattices 1-10).

$$P_{Lat\#1} = \frac{\Gamma_{tot, Lat\#1}}{\Gamma_{tot}},$$

$$P_{Lat\#2} = \frac{\Gamma_{tot,Lat\#2}}{\Gamma_{tot}}, \text{and}$$

$$\dots P_{Lat\#10} = \frac{\Gamma_{tot,Lat\#10}}{\Gamma_{tot}} \quad (1),$$

where Γ_{tot} is defined as,


$$\Gamma_{tot} = \sum_{i=1}^{10} \Gamma_{tot,Lat\#i} \quad (2).$$

The probability distribution for the lattices, $\Gamma_{tot,Lat\#i}$, is defined as the sum of all transition rates for all SKMC events; however, the events differ in dimensionality and type as shown in Table ?. For example, on the cell membrane (1st lattice) receptors diffuse in 2D but react with cytosolic species (2nd lattice) on the lattice below such that reaction occurs in 2.5D. In the cytosol (3rd-9th lattice) species are able to diffuse and react in 3D. Cytosolic species in the boundary lattice which borders the cell membrane (2nd lattice) are able to react with neighboring receptors on the cell membrane but are not allowed to diffuse onto the membrane. At the boundary nucleus (10th lattice) species diffuse and react in 2.5D.

In the cytosol the following reaction occurs: $M_1 + M_2 \xrightleftharpoons[kb]{kf} M_{12}$, such that the product M_{12} occupies the single site k. This reaction is valid due to the large site separation distance of 10 nm in comparison to the small molecular masses of cytosolic species Shc, Grb2, and Sos 62 kDa 25 kDa, 11 kDa (49-51).

Once a lattice is chosen an event is selected, either reaction or diffusion for the SKMC algorithm as shown in Fig 3. Events are chosen, as in our previous algorithm(52), with

the exception of dimensionality which changes the way Γ_{\max} and Γ_i^X are computed. The transition probabilities for cytosolic reactions in 3D are



in which reacting species (M_1 and M_2) occupy adjacent sites i and j and k has units of (molecules/site)⁻¹sec⁻¹. Diffusion in 3D is taken into account similarly

$$\Gamma_{i \rightarrow j}^d = \frac{1}{6} \Gamma^d \sigma_i (1 - \sigma_j) \quad j \in B_i \quad (4)$$

where B_i denotes the set of sites to which diffusion from site i can occur. Table I lists all events, both reaction and diffusion, which make up Γ_{\max} . Γ_{\max} is defined as before but multiplying by a factor of 6 for all 3D event, by a factor of 4 for all 2D events, and a factor of 5 for all 2.5D events such that

$$\begin{aligned} \Gamma_{\max} = & 4 \left(\frac{\Gamma^d}{4} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right)_{2D \text{ Events}} + 5 \left(\frac{\Gamma^d}{5} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right)_{2.5D \text{ Events}} \\ & + 6 \left(\frac{\Gamma^d}{6} + \max \left\{ \sum_{\text{all forward reaction events}} \Gamma^r \right\} \right)_{3D \text{ Events}} + \max \left\{ \sum_{\text{all backward reaction events}} \Gamma^r \right\} \quad (5) \end{aligned}$$

The ACGMC operates like a single MC in which time is updated in a “combined”

manner from Γ_{tot} with an average time step as, $\Delta t = \frac{1}{\Gamma_{tot}}$. Execution times vary

depending on a chosen algorithm. The complete algorithm, which is shown in Fig. 3, was implemented in Fortran 90. Due to the stochasticity of the algorithm, 10 simulations with different seeds for the random number generator were used for statistics.

Adaptively Coarse-Graining & Diffusion Transition Rates

In ACGMC, we introduce coarse-graining by grouping microscopic sites into coarse cells, as shown in Fig 4 in which coarse lattice L_C has m_x , m_y , and m_z coarse cells along x, y, z, such that the total number of cells is $m = m_x m_y m_z$. Each coarse cell C_j ($1 \leq j \leq m$) has q_j microscopic sites, in which $q_j = q_j^x q_j^y q_j^z$ is the number of microscopic sites in C_j along x, y and z axis. Diffusion between coarse-grained cells was rigorously derived by (Chatterjee),

$$\bar{C}_m(j \rightarrow i) = \frac{\Gamma_d}{q_j^z (q_j^z + q_i^z)} \eta_j (1 - \bar{\eta}_i) \quad (6)$$

in which, Γ_d is the diffusion transition rate, η_j , is the coarse-grained occupancy function at C_j is defined as

$$\eta_j = \sum_{p \in C_j} \sigma_p \quad (7)$$

While the occupancy of a coarse cell is in terms of the coarse cell coverage,

$$\bar{\eta}_j = \frac{1}{q_j} \sum_{p \in C_j} \sigma_p \quad (8)$$

which for the microscopic case reduces down to either zero for unoccupied or one for occupied.

Validation of Approach

In order to validate our approach we initially placed all molecules within lattices one through nine and allowed them to diffuse a distance of $1 \mu\text{m}$ into lattice ten over time (Fig 5).

Turning off reaction we have reflective boundary conditions at the cytosol membrane boundary (lattice two) and at the cytosol nucleus boundary (lattice 10) and periodic boundary conditions in the plane parallel to the plasma membrane. Using a diffusivity coefficient of $10\mu\text{m}^2 \text{ s}^{-1}$ and a distance $1\mu\text{m}$ and implementing reflective boundary conditions we compared the ACGMC with the PDE solution. Starting with a cytosolic species concentration 105 nM we observe similar predications between the ACGMC and PDE profiles.

Lipid Rafts & Picket Fences.

Lipid rafts are modeled as a square on the lattice with a predefined size. The diffusivity of a receptor through a lipid raft is decreased by an experimentally determined factor. In order to model cytoskeletal interactions with the cell membrane boundaries, “picket fences” were placed on the lattices. The “picket-fences” occupy lattice sites and therefore prevent reaction and diffusion events in the direction of the boundary. Take for example, a receptor neighbored by a “picket fence” on the j th lattice site, it has an occupancy function, $\sigma_j = 1$ of one, yielding a zero transition probability, $\Gamma_{i \rightarrow j}^d = 0$ in the direction of the boundary. Similarly, a receptor separated from its partner by a “picket fence” can not dimerize being that the neighboring lattice site is occupied by the boundary.

Single particle tracking reveals short term receptor confinement followed by long term “hop” diffusion. Cytoskeletal rearrangements and actin depolarization are responsible for “hop” diffusion(15, 40). In order to simulate “hop” diffusion breaks were randomly inserted into the boundaries an average of time step τ_{PF} of 10 milliseconds per $0.04 \mu\text{m}^2$ of picket fences as was observed experimentally (14). Placing breaks in the “picket fence” enables receptors to diffuse out of their corrals. After an iteration of the SKMC

breaks are closed and τ_{PF} is set to zero. This is consistent with the rate kinetics of actin dimerization (16, 53-56) which is on the order of the algorithm's time step, Δt .




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.

1. B. N. Kholodenko, O. V. Demin, G. Moehren, J. B. Hoek, *J Biol Chem* **274**, 30169 (Oct 15, 1999).
2. B. Schoeberl, C. Eichler-Jonsson, E. D. Gilles, G. Muller, *Nat Biotechnol* **20**, 370 (Apr, 2002).
3. S. Sasagawa, Y. Ozaki, K. Fujita, S. Kuroda, *Nat Cell Biol* **7**, 365 (Apr, 2005).
4. I. R. Bates *et al.*, *Biophys J* **91**, 1046 (Aug 1, 2006).
5. C. Dietrich, B. Yang, T. Fujiwara, A. Kusumi, K. Jacobson, *Biophys J* **82**, 274 (Jan, 2002).
6. M. Edidin, *Trends Cell Biol* **11**, 492 (Dec, 2001).
7. M. Edidin, *Sci STKE* **2001**, PE1 (Jan 30, 2001).
8. B. S. Wilson *et al.*, *Mol Biol Cell* **15**, 2580 (Jun, 2004).
9. P. Scheiffele *et al.*, *J Cell Biol* **140**, 795 (Feb 23, 1998).
10. R. G. Anderson, K. Jacobson, *Science* **296**, 1821 (Jun 7, 2002).
11. S. Schein, *J Mol Biol* **387**, 363 (Mar 27, 2009).
12. B. S. Wilson *et al.*, *Methods Mol Biol* **398**, 245 (2007).
13. R. Chadda *et al.*, *Traffic* **8**, 702 (Jun, 2007).
14. T. Fujiwara, K. Ritchie, H. Murakoshi, K. Jacobson, A. Kusumi, *J Cell Biol* **157**, 1071 (Jun 10, 2002).
15. A. Kusumi *et al.*, *Annu Rev Biophys Biomol Struct* **34**, 351 (2005).
16. N. L. Andrews *et al.*, *Nat Cell Biol* **10**, 955 (Aug, 2008).
17. J. Zhang *et al.*, *Micron* **37**, 14 (2006).
18. K. Simons, D. Toomre, *Nat Rev Mol Cell Biol* **1**, 31 (Oct, 2000).
19. S. J. Singer, G. L. Nicolson, *Science* **175**, 720 (Feb 18, 1972).
20. K. Simons, G. van Meer, *Biochemistry* **27**, 6197 (Aug 23, 1988).
21. G. van Meer, K. Simons, *J Cell Biochem* **36**, 51 (Jan, 1988).
22. G. van Meer, E. H. Stelzer, R. W. Wijnaendts-van-Resandt, K. Simons, *J Cell Biol* **105**, 1623 (Oct, 1987).
23. M. P. Sheetz, *Semin Hematol* **20**, 175 (Jul, 1983).
24. M. P. Sheetz, M. Schindler, D. E. Koppel, *Nature* **285**, 510 (Jun 12, 1980).
25. K. Jacobson, C. Dietrich, *Trends Cell Biol* **9**, 87 (Mar, 1999).
26. K. Jacobson, O. G. Mouritsen, R. G. Anderson, *Nat Cell Biol* **9**, 7 (Jan, 2007).
27. A. Filippov, G. Oradd, G. Lindblom, *Biophys J* **86**, 891 (Feb, 2004).
28. A. Filippov, G. Oradd, G. Lindblom, *Biophys J* **90**, 2086 (Mar 15, 2006).
29. G. Oradd, P. W. Westerman, G. Lindblom, *Biophys J* **89**, 315 (Jul, 2005).
30. A. Kusumi, H. Ike, C. Nakada, K. Murase, T. Fujiwara, *Semin Immunol* **17**, 3 (Feb, 2005).
31. J. M. Haugh, *Mol Interv* **2**, 292 (Sep, 2002).
32. Q. Li *et al.*, *J Lipid Res* **46**, 1904 (Sep, 2005).
33. J. R. Muppidi, J. Tschopp, R. M. Siegel, *Immunity* **21**, 461 (Oct, 2004).
34. S. Roy *et al.*, *Nat Cell Biol* **1**, 98 (Jun, 1999).
35. S. J. Plowman, C. Muncke, R. G. Parton, J. F. Hancock, *Proc Natl Acad Sci U S A* **102**, 15500 (Oct 25, 2005).
36. S. J. Plowman, J. F. Hancock, *Biochim Biophys Acta* **1746**, 274 (Dec 30, 2005).
37. C. Nakada *et al.*, *Nat Cell Biol* **5**, 626 (Jul, 2003).
38. K. Ritchie, R. Iino, T. Fujiwara, K. Murase, A. Kusumi, *Mol Membr Biol* **20**, 13 (Jan-Mar, 2003).

39. K. Suzuki, K. Ritchie, E. Kajikawa, T. Fujiwara, A. Kusumi, *Biophys J* **88**, 3659 (May, 2005).
40. K. Murase *et al.*, *Biophys J* **86**, 4075 (Jun, 2004).
41. N. Morone *et al.*, *J Cell Biol* **174**, 851 (Sep 11, 2006).
42. H. Chopra *et al.*, *Cancer Res* **48**, 3787 (Jul 1, 1988).
43. P. A. Janmey, *Physiol Rev* **78**, 763 (Jul, 1998).
44. A. Kusumi, Y. Sako, T. Fujiwara, M. Tomishige, *Methods Cell Biol* **55**, 173 (1998).
45. H. Ike *et al.*, *Chemphyschem* **4**, 620 (Jun 16, 2003).
46. I. Koyama-Honda *et al.*, *Biophys J* **88**, 2126 (Mar, 2005).
47. K. Simons, E. Ikonen, *Science* **290**, 1721 (Dec 1, 2000).
48. K. Simons, E. Ikonen, *Nature* **387**, 569 (Jun 5, 1997).
49. L. Buday, S. E. Egan, P. Rodriguez Viciano, D. A. Cantrell, J. Downward, *The Journal of biological chemistry* **269**, 9019 (Mar 25, 1994).
50. M. Sieh, A. Batzer, J. Schlessinger, A. Weiss, *Molecular and cellular biology* **14**, 4435 (Jul, 1994).
51. F. Yamaguchi, K. Yamaguchi, M. Tokuda, S. Brenner, *FEBS letters* **459**, 105 (Oct 1, 1999).
52. D. G. Kapil Mayawala, Vlachos, Jeremy S. Edwards, *BMC Cell Biology* **6**, (2005).
53. G. Danuser, C. M. Waterman-Storer, *Annu Rev Biophys Biomol Struct* **35**, 361 (2006).
54. M. C. Adams *et al.*, *J Microsc* **216**, 138 (Nov, 2004).
55. P. Vallotton, S. L. Gupton, C. M. Waterman-Storer, G. Danuser, *Proc Natl Acad Sci U S A* **101**, 9660 (Jun 29, 2004).
56. V. S. Deshpande, R. M. McMeeking, A. G. Evans, *Proc Natl Acad Sci U S A* **103**, 14015 (Sep 19, 2006).

Appendix E. Abbreviations Used

ErbB1	Epidermal growth factor receptor
ErbB2	Human epidermal growth factor receptor 2
ErbB3	V-erb-b2 erythroblastic leukemia viral oncogene homolog 3
ErbB4	V-erb-a erythroblastic leukemia viral oncogene homolog 4
EGFR	Epidermal growth factor receptor
ERK	Extracellular regulated kinase
SH2	Src homology 2
Grb2	Growth factor receptor-bound protein 2
SOS	son of the sevenless
Shc	Transforming protein 1 and collagen domain protein
ODE	Ordinary differential equation
MC	Monte Carlo
SKMC	Spatial kinetic Monte Carlo
SSA	Stochastic simulator algorithm
CSNSA	Coupled spatial non-spatial simulation algorithm
PLC γ 1	Phospholipase C γ 1
CPU	Central processing unit
LSSENS	Lewis General Chemical Kinetics and Sensitivity Analysis Code
MSK	Membrane skeleton
FCM	Fuzzy c-means
MSD	Mean squared displacement
SPT	Single particle tracking
FRET	Fluorescence resonance energy transfer

EM Electron microscopy

MAPK Mitogen-activated protein kinase

GAB1 GRB2-associated binding protein 1

Src Tyrosine kinase

FRAP Fluorescence recovery after photobleaching

FRET Forster resonance energy transfer

TEM Transmission electron microscope

STEM Scanning transmission electron microscope

EDX Energy dispersive X-ray

EELS Electron energy loss spectroscopy

ADF Annular dark-field imaging

SEM Scanning electron microscope

GPI Glycosylphosphatidylinositol

DRM Detergent-resistant membrane

DIG Detergent-insoluble glycolipid

LCFAs Long-chain fatty acids

DOPE 1,2-dioleoyl*sn*-glycero-3-phosphoethanolamine

SDE Stochastic differential equations

SPDE stochastic partial differential equations

DMC Dynamic Monte Carlo

PBCs Periodic boundary conditions

CGMC Coarse-graining Monte Carlo

ACGMC Adaptively coarse-grained Monte Carlo

PDE Partial differential equations

GFP green fluorescent protein

PKC ϵ Protein kinase c

INF γ R Interferon- γ receptor

JAK Janus kinase

STAT Signal transducer and activator of transcription pathway

BioXyce Deterministic circuit-based platform

NRK Normal rat kidney fibroblast

GPCR G-protein-coupled receptor

References

- Abad-Rodriguez, J., M. D. Ledesma, et al. (2004). "Neuronal membrane cholesterol loss enhances amyloid peptide generation." *J Cell Biol* 167(5): 953-60.
- Abankwa, D., A. A. Gorfe, et al. (2007). "Ras nanoclusters: molecular structure and assembly." *Semin Cell Dev Biol* 18(5): 599-607.
- Abankwa, D., A. A. Gorfe, et al. (2008). "Mechanisms of Ras membrane organization and signalling: Ras on a rocker." *Cell Cycle* 7(17): 2667-73.
- Abankwa, D., M. Hanzal-Bayer, et al. (2008). "A novel switch region regulates H-ras membrane orientation and signal output." *EMBO J* 27(5): 727-35.
- Abbott, B., R. Abbott, et al. (2005). "Limits on gravitational-wave emission from selected pulsars using LIGO data." *Phys Rev Lett* 94(18): 181103.
- Abbott, B., R. Abbott, et al. (2005). "Upper limits on a stochastic background of gravitational waves." *Phys Rev Lett* 95(22): 221101.
- Abbott, R. G., S. Forrest, et al. (2006). "Simulating the hallmarks of cancer." *Artif Life* 12(4): 617-34.
- Adam, L., S. Ellwood, et al. (1999). "Comparison of *Erysiphe cichoracearum* and *E. cruciferarum* and a survey of 360 *Arabidopsis thaliana* accessions for resistance to these two powdery mildew pathogens." *Mol Plant Microbe Interact* 12(12): 1031-43.
- Adams, J. W., J. Wang, et al. (2008). "Myocardial expression, signaling, and function of GPR22: a protective role for an orphan G protein-coupled receptor." *Am J Physiol Heart Circ Physiol* 295(2): H509-21.
- Adams, M. C., A. Matov, et al. (2004). "Signal analysis of total internal reflection fluorescent speckle microscopy (TIR-FSM) and wide-field epi-fluorescence FSM of the actin cytoskeleton and focal adhesions in living cells." *J Microsc* 216(Pt 2): 138-52.
- Ada-Nguema, A. S., H. Xenias, et al. (2006). "The small GTPase R-Ras regulates organization of actin and drives membrane protrusions through the activity of PLCepsilon." *J Cell Sci* 119(Pt 7): 1307-19.
- Adcox, K., S. S. Adler, et al. (2001). "Centrality dependence of charged particle multiplicity in Au-Au collisions at square root of (s)NN = 130 GeV." *Phys Rev Lett* 86(16): 3500-5.

- Adcox, K., S. S. Adler, et al. (2001). "Measurement of the midrapidity transverse energy distribution from square root of $[(s)NN] = 130$ GeV Au + Au collisions at RHIC." *Phys Rev Lett* 87(5): 052301.
- Adcox, K., S. S. Adler, et al. (2002). "Centrality dependence of $\pi^{+/-}$, $K^{+/-}$, p, and $(-)$ p production from $\sqrt{s(NN)} = 130$ GeV Au + Au collisions at RHIC." *Phys Rev Lett* 88(24): 242301.
- Adcox, K., S. S. Adler, et al. (2002). "Flow measurements via two-particle azimuthal correlations in Au + Au collisions at $\sqrt{s(NN)}=130$ GeV." *Phys Rev Lett* 89(21): 212301.
- Adcox, K., S. S. Adler, et al. (2002). "Measurement of Λ and $\Lambda(\text{macro})$ particles in Au+Au collisions at the square root of $S(NN) = 130$ GeV." *Phys Rev Lett* 89(9): 092302.
- Adcox, K., S. S. Adler, et al. (2002). "Measurement of single electrons and implications for charm production in Au+Au collisions at square root $[\sqrt{s(NN)}] = 130$ GeV." *Phys Rev Lett* 88(19): 192303.
- Adcox, K., S. S. Adler, et al. (2002). "Net charge fluctuations in Au + Au interactions at $\sqrt{s(NN)}=130$ GeV." *Phys Rev Lett* 89(8): 082301.
- Adcox, K., S. S. Adler, et al. (2002). "Suppression of hadrons with large transverse momentum in central Au+Au collisions at root square $[\sqrt{s(NN)}] = 130$ GeV." *Phys Rev Lett* 88(2): 022301.
- Adcox, K., S. S. Adler, et al. (2002). "Transverse-mass dependence of two-pion correlations in Au+Au collisions at square root $[\sqrt{s(NN)}] = 130$ GeV." *Phys Rev Lett* 88(19): 192302.
- Aguado, C., J. Colon, et al. (2008). "Cell type-specific subunit composition of G protein-gated potassium channels in the cerebellum." *J Neurochem* 105(2): 497-511.
- Ahmed, I., A. S. Ponery, et al. (2007). "Morphology, cytoskeletal organization, and myosin dynamics of mouse embryonic fibroblasts cultured on nanofibrillar surfaces." *Mol Cell Biochem* 301(1-2): 241-9.
- Aiello, L. P., A. Clermont, et al. (2006). "Inhibition of PKC beta by oral administration of ruboxistaurin is well tolerated and ameliorates diabetes-induced retinal hemodynamic abnormalities in patients." *Invest Ophthalmol Vis Sci* 47(1): 86-92.
- Aiello, L. P., M. D. Davis, et al. (2006). "Effect of ruboxistaurin on visual loss in patients with diabetic retinopathy." *Ophthalmology* 113(12): 2221-30.

- Ajimura, S., K. Ikeda, et al. (2000). "Asymmetry in the nonmesonic weak decay of polarized (5)(Lambda)He hypernuclei." *Phys Rev Lett* 84(18): 4052-5.
- Aksamitiene, E., J. B. Hoek, et al. (2007). "Multistrip Western blotting to increase quantitative data output." *Electrophoresis* 28(18): 3163-73.
- Albanesi, J. P., H. Fujisaki, et al. (1985). "Monomeric Acanthamoeba myosins I support movement in vitro." *J Biol Chem* 260(15): 8649-52.
- Alhanaty, E. and M. P. Sheetz (1981). "Control of the erythrocyte membrane shape: recovery from the effect of crenating agents." *J Cell Biol* 91(3 Pt 1): 884-8.
- Alhanaty, E. and M. P. Sheetz (1984). "Cell membrane shape control--effects of chloromethyl ketone peptides." *Blood* 63(5): 1203-8.
- Alhanaty, E., M. Snyder, et al. (1984). "Glucose-6-phosphate-dehydrogenase-deficient erythrocytes have an impaired shape recovery mechanism." *Blood* 63(5): 1198-202.
- Allen, J. A., R. A. Halverson-Tamboli, et al. (2007). "Lipid raft microdomains and neurotransmitter signalling." *Nat Rev Neurosci* 8(2): 128-40.
- Allen, J. A., J. Z. Yu, et al. (2005). "Beta-adrenergic receptor stimulation promotes G alpha s internalization through lipid rafts: a study in living cells." *Mol Pharmacol* 67(5): 1493-504.
- Allen, P. G., C. B. Shuster, et al. (1996). "Phalloidin binding and rheological differences among actin isoforms." *Biochemistry* 35(45): 14062-9.
- Allersma, M. W., L. Wang, et al. (2004). "Visualization of regulated exocytosis with a granule-membrane probe using total internal reflection microscopy." *Mol Biol Cell* 15(10): 4658-68.
- Alvarez-Franco, M. and A. S. Paller (1994). "What syndrome is this? Congenital Kasabach-Merritt syndrome." *Pediatr Dermatol* 11(1): 79-81.
- Amanai, T., Y. Nakamura, et al. (2008). "Micro-CT analysis of experimental Candida osteoarthritis in rats." *Mycopathologia* 166(3): 133-41.
- Anderson, R. G. and K. Jacobson (2002). "A role for lipid shells in targeting proteins to caveolae, rafts, and other lipid domains." *Science* 296(5574): 1821-5.
- Andrec, M., B. N. Kholodenko, et al. (2005). "Inference of signaling and gene regulatory networks by steady-state perturbation experiments: structure and accuracy." *J Theor Biol* 232(3): 427-41.

- Andrews, N. L., K. A. Lidke, et al. (2008). "Actin restricts FcεRI diffusion and facilitates antigen-induced receptor immobilization." *Nat Cell Biol* 10(8): 955-63.
- Andrews, R. P., C. L. Kepley, et al. (2001). "Regulation of the very late antigen-4-mediated adhesive activity of normal and nonreleaser basophils: roles for Src, Syk, and phosphatidylinositol 3-kinase." *J Leukoc Biol* 70(5): 776-82.
- Andriotis, A. N., M. Menon, et al. (2003). "Magnetic properties of C60 polymers." *Phys Rev Lett* 90(2): 026801.
- Antal, M., I. Papp, et al. (2004). "Expression of hyperpolarization-activated and cyclic nucleotide-gated cation channel subunit 2 in axon terminals of peptidergic nociceptive primary sensory neurons in the superficial spinal dorsal horn of rats." *Eur J Neurosci* 19(5): 1336-42.
- Aoki, K., E. Kiyokawa, et al. (2008). "Visualization of growth signal transduction cascades in living cells with genetically encoded probes based on Forster resonance energy transfer." *Philos Trans R Soc Lond B Biol Sci* 363(1500): 2143-51.
- Aoki, K., T. Tajima, et al. (2008). "A novel initial codon mutation of the thiazide-sensitive Na-Cl cotransporter gene in a Japanese patient with Gitelman's syndrome." *Endocr J* 55(3): 557-60.
- Apolloni, A., I. A. Prior, et al. (2000). "H-ras but not K-ras traffics to the plasma membrane through the exocytic pathway." *Mol Cell Biol* 20(7): 2475-87.
- Arkin, A. P. (2001). "Synthetic cell biology." *Curr Opin Biotechnol* 12(6): 638-44.
- Arluison, V., S. Hohng, et al. (2007). "Spectroscopic observation of RNA chaperone activities of Hfq in post-transcriptional regulation by a small non-coding RNA." *Nucleic Acids Res* 35(3): 999-1006.
- Arnsmeier, S. L., V. M. Riccardi, et al. (1994). "Familial multiple café au lait spots." *Arch Dermatol* 130(11): 1425-6.
- Arora, P. D., M. W. Chan, et al. (2005). "Separate functions of gelsolin mediate sequential steps of collagen phagocytosis." *Mol Biol Cell* 16(11): 5175-90.
- Arora, P. D., P. A. Janmey, et al. (1999). "A role for gelsolin in stress fiber-dependent cell contraction." *Exp Cell Res* 250(1): 155-67.
- Arsene, I., I. G. Bearden, et al. (2005). "Centrality dependence of charged-particle pseudorapidity distributions from d + Au collisions at $\sqrt{s_{NN}} = 200$ GeV." *Phys Rev Lett* 94(3): 032301.

Artalejo, C. R., M. A. Lemmon, et al. (1997). "Specific role for the PH domain of dynamin-1 in the regulation of rapid endocytosis in adrenal chromaffin cells." *EMBO J* 16(7): 1565-74.

Asher, J., A. Oliver, et al. (2005). "A simple cardiovascular risk score can predict poor outcome in non-heart-beating donor renal transplantation." *Transplant Proc* 37(2): 1044-6.

Axelrod, D. and M. D. Wang (1994). "Reduction-of-dimensionality kinetics at reaction-limited cell surface receptors." *Biophys J* 66(3 Pt 1): 588-600.

Azad, M. B., Y. Chen, et al. (2008). "Hypoxia induces autophagic cell death in apoptosis-competent cells through a mechanism involving BNIP3." *Autophagy* 4(2): 195-204.

Bacallao, R., C. Antony, et al. (1989). "The subcellular organization of Madin-Darby canine kidney cells during the formation of a polarized epithelium." *J Cell Biol* 109(6 Pt 1): 2817-32.

Bakovic, M., M. D. Fullerton, et al. (2007). "Metabolic and molecular aspects of ethanolamine phospholipid biosynthesis: the role of CTP:phosphoethanolamine cytidyltransferase (Pcyt2)." *Biochem Cell Biol* 85(3): 283-300.

Bani-Yaghoub, M. and D. E. Amundsen (2008). "Study and simulation of reaction-diffusion systems affected by interacting signaling pathways." *Acta Biotheor* 56(4): 315-28.

Banizs, B., M. M. Pike, et al. (2005). "Dysfunctional cilia lead to altered ependyma and choroid plexus function, and result in the formation of hydrocephalus." *Development* 132(23): 5329-39.

Baoukina, S., L. Monticelli, et al. (2008). "The molecular mechanism of lipid monolayer collapse." *Proc Natl Acad Sci U S A* 105(31): 10803-8.

Bard, F., R. Barbour, et al. (2003). "Epitope and isotype specificities of antibodies to beta -amyloid peptide for protection against Alzheimer's disease-like neuropathology." *Proc Natl Acad Sci U S A* 100(4): 2023-8.

Bardwell, L., X. Zou, et al. (2007). "Mathematical models of specificity in cell signaling." *Biophys J* 92(10): 3425-41.

Barker, S. A., K. K. Caldwell, et al. (1995). "Wortmannin blocks lipid and protein kinase activities associated with PI 3-kinase and inhibits a subset of responses induced by Fc epsilon R1 cross-linking." *Mol Biol Cell* 6(9): 1145-58.

- Bartynski, W. S., D. S. Whitt, et al. (2007). "Lower cervical nerve root block using CT fluoroscopy in patients with large body habitus: another benefit of the swimmer's position." *AJNR Am J Neuroradiol* 28(4): 706-8.
- Bates, I. R., B. Hebert, et al. (2006). "Membrane lateral diffusion and capture of CFTR within transient confinement zones." *Biophys J* 91(3): 1046-58.
- Bates, I. R., P. W. Wiseman, et al. (2006). "Investigating membrane protein dynamics in living cells." *Biochem Cell Biol* 84(6): 825-31.
- Baudier, J., D. Mochly-Rosen, et al. (1987). "Comparison of S100b protein with calmodulin: interactions with melittin and microtubule-associated tau proteins and inhibition of phosphorylation of tau proteins by protein kinase C." *Biochemistry* 26(10): 2886-93.
- Bauer, D. A., T. L. Graff, et al. (1985). "Differences between proton- and π^- -induced production of the charmonium χ states." *Phys Rev Lett* 54(8): 753-756.
- Bauw, G., H. H. Rasmussen, et al. (1990). "Two-dimensional gel electrophoresis, protein electroblotting and microsequencing: a direct link between proteins and genes." *Electrophoresis* 11(7): 528-36.
- Bearden, I. G., D. Beavis, et al. (2001). "Rapidity dependence of antiproton-to-proton ratios in Au+Au collisions at square root of $(s_{NN}) = 130$ GeV." *Phys Rev Lett* 87(11): 112305.
- Bearden, I. G., D. Beavis, et al. (2002). "Pseudorapidity distributions of charged particles from Au + Au collisions at the maximum RHIC energy, square root $[s(NN)] = 200$ GeV." *Phys Rev Lett* 88(20): 202301.
- Bearden, I. G., D. Beavis, et al. (2005). "Charged meson rapidity distributions in central Au+Au collisions at square root $(s_{NN}) = 200$ GeV." *Phys Rev Lett* 94(16): 162301.
- Beers, B., W. Jaszcz, et al. (1992). "Porokeratosis palmaris et plantaris disseminata. Report of a case with abnormal DNA ploidy in lesional epidermis." *Arch Dermatol* 128(2): 236-9.
- Belanis, L., S. J. Plowman, et al. (2008). "Galectin-1 is a novel structural component and a major regulator of h-ras nanoclusters." *Mol Biol Cell* 19(4): 1404-14.
- Bennett, M. R. (1983). "Development of neuromuscular synapses." *Physiol Rev* 63(3): 915-1048.
- Bennett, V. (1990). "Spectrin-based membrane skeleton: a multipotential adaptor between plasma membrane and cytoplasm." *Physiol Rev* 70(4): 1029-65.

- Bennett, V. and A. J. Baines (2001). "Spectrin and ankyrin-based pathways: metazoan inventions for integrating cells into tissues." *Physiol Rev* 81(3): 1353-92.
- Bennett, V. and L. Chen (2001). "Ankyrins and cellular targeting of diverse membrane proteins to physiological sites." *Curr Opin Cell Biol* 13(1): 61-7.
- Beranger, F., K. Cadwallader, et al. (1994). "Determination of structural requirements for the interaction of Rab6 with RabGDI and Rab geranylgeranyltransferase." *J Biol Chem* 269(18): 13637-43.
- Beranger, F., H. Paterson, et al. (1994). "The effector domain of Rab6, plus a highly hydrophobic C terminus, is required for Golgi apparatus localization." *Mol Cell Biol* 14(1): 744-58.
- Berk, D. A., A. Clark, Jr., et al. (1992). "Analysis of lateral diffusion from a spherical cell surface to a tubular projection." *Biophys J* 61(1): 1-8.
- Berk, D. A. and R. M. Hochmuth (1992). "Lateral mobility of integral proteins in red blood cell tethers." *Biophys J* 61(1): 9-18.
- Betzig, E., G. H. Patterson, et al. (2006). "Imaging intracellular fluorescent proteins at nanometer resolution." *Science* 313(5793): 1642-5.
- Bezdek, J. C. "Pattern recognition with fuzzy objective function algorithms." *Advanced Applications in Pattern Recognition*.
- Biais, N., B. Ladoux, et al. (2008). "Cooperative retraction of bundled type IV pili enables nanonewton force generation." *PLoS Biol* 6(4): e87.
- Bienert, M., B. McCook, et al. (2005). "Sequential FDG PET/CT in 90Y microsphere treatment of unresectable colorectal liver metastases." *Eur J Nucl Med Mol Imaging* 32(6): 723.
- Bienert, M., B. McCook, et al. (2005). "90Y microsphere treatment of unresectable liver metastases: changes in 18F-FDG uptake and tumour size on PET/CT." *Eur J Nucl Med Mol Imaging* 32(7): 778-87.
- Bier, M., B. Teusink, et al. (1996). "Control analysis of glycolytic oscillations." *Biophys Chem* 62(1-3): 15-24.
- Biessen, E. A., A. R. Valentijn, et al. (2000). "Novel hepatotropic prodrugs of the antiviral nucleoside 9-(2-phosphonylmethoxyethyl)adenine with improved pharmacokinetics and antiviral activity." *FASEB J* 14(12): 1784-92.
- Birtwistle, M. R., M. Hatakeyama, et al. (2007). "Ligand-dependent responses of the ErbB signaling network: experimental and modeling analyses." *Mol Syst Biol* 3: 144.

- Bishop, A. T., M. B. Wood, et al. (1995). "Arthrodesis of the ankle with a free vascularized autogenous bone graft. Reconstruction of segmental loss of bone secondary to osteomyelitis, tumor, or trauma." *J Bone Joint Surg Am* 77(12): 1867-75.
- Blackshaw, S., W. P. Kuo, et al. (2003). "MicroSAGE is highly representative and reproducible but reveals major differences in gene expression among samples obtained from similar tissues." *Genome Biol* 4(3): R17.
- Blanchard, P. D., R. A. Angus, et al. (1991). "Pigments and ultrastructures of pigment cells in xanthic sailfin mollies (*Poecilia latipinna*)." *Pigment Cell Res* 4(5-6): 240-6.
- Blinov, M. L., J. R. Faeder, et al. (2004). "BioNetGen: software for rule-based modeling of signal transduction based on the interactions of molecular domains." *Bioinformatics* 20(17): 3289-91.
- Blinov, M. L., J. R. Faeder, et al. (2006). "A network model of early events in epidermal growth factor receptor signaling that accounts for combinatorial complexity." *Biosystems* 83(2-3): 136-51.
- Blinov, M. L., J. R. Faeder, et al. (2005). "'On-the-fly' or 'generate-first' modeling?" *Nat Biotechnol* 23(11): 1344-5; author reply 1345.
- Blinov, M. L., J. Yang, et al. (2006). "Depicting signaling cascades." *Nat Biotechnol* 24(2): 137-8; author reply 138.
- Bluthgen, N., F. J. Bruggeman, et al. (2006). "Effects of sequestration on signal transduction cascades." *FEBS J* 273(5): 895-906.
- Boeneman, K., B. C. Mei, et al. (2009). "Sensing caspase 3 activity with quantum dot-fluorescent protein assemblies." *J Am Chem Soc* 131(11): 3828-9.
- Boldogh, I. R., L. A. Pon, et al. (2007). "Cell-free assays for mitochondria-cytoskeleton interactions." *Methods Cell Biol* 80: 683-706.
- Bollag, G. E., R. A. Roth, et al. (1986). "Protein kinase C directly phosphorylates the insulin receptor in vitro and reduces its protein-tyrosine kinase activity." *Proc Natl Acad Sci U S A* 83(16): 5822-4.
- Borisov, N., E. Aksamitiene, et al. (2009). "Systems-level interactions between insulin-EGF networks amplify mitogenic signaling." *Mol Syst Biol* 5: 256.
- Borisov, N. M., A. S. Chistopolsky, et al. (2008). "Domain-oriented reduction of rule-based network models." *IET Syst Biol* 2(5): 342-51.

- Borisov, N. M., N. I. Markevich, et al. (2005). "Signaling through receptors and scaffolds: independent interactions reduce combinatorial complexity." *Biophys J* 89(2): 951-66.
- Borisov, N. M., N. I. Markevich, et al. (2006). "Trading the micro-world of combinatorial complexity for the macro-world of protein interaction domains." *Biosystems* 83(2-3): 152-66.
- Bortz, A. B., M. H. Kalos, et al. (1975). "New Algorithm for Monte-Carlo Simulation of Ising Spin Systems." *Journal of Computational Physics* 17(1): 10-18.
- Bortz, A. B., M. H. Kalos, et al. (1974). "Time Evolution of a Quenched Binary Alloy - Computer-Simulation of a 2-Dimensional Model System." *Physical Review B* 10(2): 535-541.
- Boyd, N. L., H. Park, et al. (2003). "Chronic shear induces caveolae formation and alters ERK and Akt responses in endothelial cells." *Am J Physiol Heart Circ Physiol* 285(3): H1113-22.
- Bresnick, A. R., P. A. Janmey, et al. (1991). "Evidence that a 27-residue sequence is the actin-binding site of ABP-120." *J Biol Chem* 266(20): 12989-93.
- Bridson, S. J., J. Gandia, et al. (2008). "Plasma membrane diffusion of G protein-coupled receptor oligomers." *Biochim Biophys Acta* 1783(12): 2262-8.
- Brinkerhoff, C. J., J. S. Choi, et al. (2008). "Diffusion-limited reactions in G-protein activation: unexpected consequences of antagonist and agonist competition." *J Theor Biol* 251(4): 561-9.
- Brinkerhoff, C. J. and J. J. Linderman (2005). "Integrin dimerization and ligand organization: key components in integrin clustering for cell adhesion." *Tissue Eng* 11(5-6): 865-76.
- Brinkerhoff, C. J., J. R. Traynor, et al. (2008). "Collision coupling, crosstalk, and compartmentalization in G-protein coupled receptor systems: can a single model explain disparate results?" *J Theor Biol* 255(3): 278-86.
- Brinkerhoff, C. J., P. J. Woolf, et al. (2004). "Monte Carlo simulations of receptor dynamics: insights into cell signaling." *J Mol Histol* 35(7): 667-77.
- Britten, C. D. (2004). "Targeting ErbB receptor signaling: a pan-ErbB approach to cancer." *Mol Cancer Ther* 3(10): 1335-42.
- Broadhead, C. L., G. Betton, et al. (2000). "Prospects for reducing and refining the use of dogs in the regulatory toxicity testing of pharmaceuticals." *Hum Exp Toxicol* 19(8): 440-7.

- Brodey, B. B., C. S. Rosen, et al. (2005). "Reliability and acceptability of automated telephone surveys among Spanish- and English-speaking mental health services recipients." *Ment Health Serv Res* 7(3): 181-4.
- Brodey, B. B., C. S. Rosen, et al. (2004). "Validation of the Addiction Severity Index (ASI) for internet and automated telephone self-report administration." *J Subst Abuse Treat* 26(4): 253-9.
- Brodey, B. B., C. S. Rosen, et al. (2005). "Conversion and validation of the Teen-Addiction Severity Index (T-ASI) for Internet and automated-telephone self-report administration." *Psychol Addict Behav* 19(1): 54-61.
- Brown, C. J., B. D. Hendrich, et al. (1992). "The human XIST gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is highly localized within the nucleus." *Cell* 71(3): 527-42.
- Brown, D. A. and E. London (1998). "Functions of lipid rafts in biological membranes." *Annu Rev Cell Dev Biol* 14: 111-36.
- Brown, D. A. and E. London (1998). "Structure and origin of ordered lipid domains in biological membranes." *J Membr Biol* 164(2): 103-14.
- Brown, D. A. and J. K. Rose (1992). "Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface." *Cell* 68(3): 533-44.
- Brown, G. C., J. B. Hoek, et al. (1997). "Why do protein kinase cascades have more than one level?" *Trends Biochem Sci* 22(8): 288.
- Brown, G. C. and B. N. Kholodenko (1999). "Spatial gradients of cellular phosphoproteins." *FEBS Lett* 457(3): 452-4.
- Brown, G. C., H. V. Westerhoff, et al. (1996). "Molecular control analysis: control within proteins and molecular processes." *J Theor Biol* 182(3): 389-96.
- Brown, M. S. and J. L. Goldstein (1992). "Koch's postulates for cholesterol." *Cell* 71(2): 187-8.
- Brown, W. M., D. L. Christie, et al. (1992). "The rat protein encoded by clone pp63 is a fetuin/alpha 2-HS glycoprotein-like molecule, but is it the tyrosine kinase inhibitor pp63?" *Cell* 68(1): 7-8.
- Brueckl, W. M., L. Herbst, et al. (2006). "Predictive and prognostic factors in small cell lung carcinoma (SCLC)--analysis from routine clinical practice." *Anticancer Res* 26(6C): 4825-32.
- Bruggeman, F. J. and B. N. Kholodenko (2002). "Modular interaction strengths in regulatory networks; an example." *Mol Biol Rep* 29(1-2): 57-61.
- Bruggeman, F. J., H. V. Westerhoff, et al. (2002). "Modular response analysis of cellular regulatory networks." *J Theor Biol* 218(4): 507-20.

Bucki, R. and P. A. Janmey (2006). "Interaction of the gelsolin-derived antibacterial PBP 10 peptide with lipid bilayers and cell membranes." *Antimicrob Agents Chemother* 50(9): 2932-40.

Bucki, R., J. J. Pastore, et al. (2006). "Involvement of the Na⁺/H⁺ exchanger in membrane phosphatidylserine exposure during human platelet activation." *Biochim Biophys Acta* 1761(2): 195-204.

Bucki, R., J. J. Pastore, et al. (2003). "Flavonoid inhibition of platelet procoagulant activity and phosphoinositide synthesis." *J Thromb Haemost* 1(8): 1820-8.

Bui, P., D. Ivan, et al. (2009). "Chondroblastoma of the temporomandibular joint: report of a case and literature review." *J Oral Maxillofac Surg* 67(2): 405-9.

Bulinski, J. C., T. E. McGraw, et al. (1997). "Overexpression of MAP4 inhibits organelle motility and trafficking in vivo." *J Cell Sci* 110 (Pt 24): 3055-64.

Burkhardt, J. K., J. M. McIlvain, Jr., et al. (1993). "Lytic granules from cytotoxic T cells exhibit kinesin-dependent motility on microtubules in vitro." *J Cell Sci* 104 (Pt 1): 151-62.

Burns, A. R., J. M. Oliver, et al. (2008). "Visualizing clathrin-mediated IgE receptor internalization by electron and atomic force microscopy." *Methods Mol Biol* 440: 235-45.

Burroughs, N. J., Z. Lazic, et al. (2006). "Ligand detection and discrimination by spatial relocation: A kinase-phosphatase segregation model of TCR activation." *Biophys J* 91(5): 1619-29.

Burton, T. R., E. S. Henson, et al. (2006). "The pro-cell death Bcl-2 family member, BNIP3, is localized to the nucleus of human glial cells: Implications for glioblastoma multiforme tumor cell survival under hypoxia." *Int J Cancer* 118(7): 1660-9.

Busch, T. M., S. M. Hahn, et al. (2004). "Hypoxia and Photofrin uptake in the intraperitoneal carcinomatosis and sarcomatosis of photodynamic therapy patients." *Clin Cancer Res* 10(14): 4630-8.

Cacace, A. M., N. R. Michaud, et al. (1999). "Identification of constitutive and ras-inducible phosphorylation sites of KSR: implications for 14-3-3 binding, mitogen-activated protein kinase binding, and KSR overexpression." *Mol Cell Biol* 19(1): 229-40.

- Cadwallader, K. A., H. Paterson, et al. (1994). "N-terminally myristoylated Ras proteins require palmitoylation or a polybasic domain for plasma membrane localization." *Mol Cell Biol* 14(7): 4722-30.
- Cai, Y. and M. P. Sheetz (2009). "Force propagation across cells: mechanical coherence of dynamic cytoskeletons." *Curr Opin Cell Biol* 21(1): 47-50.
- Cairo, C. W. (2007). "Signaling by committee: receptor clusters determine pathways of cellular activation." *ACS Chem Biol* 2(10): 652-5.
- Calder, P. C. and P. Yaqoob (2007). "Lipid rafts--composition, characterization, and controversies." *J Nutr* 137(3): 545-7.
- Calderon, E., C. J. Conde, et al. (1987). "Treatment of ordinary and penicillinase-producing strains of *Neisseria gonorrhoeae* in Mexico City." *Diagn Microbiol Infect Dis* 8(1): 13-8.
- Cales, C., J. F. Hancock, et al. (1988). "The cytoplasmic protein GAP is implicated as the target for regulation by the ras gene product." *Nature* 332(6164): 548-51.
- Calloway, N. T., M. Choob, et al. (2007). "Optimized fluorescent trimethoprim derivatives for in vivo protein labeling." *Chembiochem* 8(7): 767-74.
- Cao, G., G. Yang, et al. (2003). "Disruption of the caveolin-1 gene impairs renal calcium reabsorption and leads to hypercalciuria and urolithiasis." *Am J Pathol* 162(4): 1241-8.
- Carey, K. D., A. J. Garton, et al. (2006). "Kinetic analysis of epidermal growth factor receptor somatic mutant proteins shows increased sensitivity to the epidermal growth factor receptor tyrosine kinase inhibitor, erlotinib." *Cancer Res* 66(16): 8163-71.
- Carozzi, A. J., S. Roy, et al. (2002). "Inhibition of lipid raft-dependent signaling by a dystrophy-associated mutant of caveolin-3." *J Biol Chem* 277(20): 17944-9.
- Carriba, P., G. Navarro, et al. (2008). "Detection of heteromerization of more than two proteins by sequential BRET-FRET." *Nat Methods* 5(8): 727-33.
- Carriles, R., K. E. Sheetz, et al. (2008). "Simultaneous multifocal, multiphoton, photon counting microscopy." *Opt Express* 16(14): 10364-71.
- Carver, R. S., M. C. Stevenson, et al. (2002). "Diverse expression of ErbB receptor proteins during rat liver development and regeneration." *Gastroenterology* 123(6): 2017-27.
- Casey, F. P., D. Baird, et al. (2007). "Optimal experimental design in an epidermal growth factor receptor signalling and down-regulation model." *IET Syst Biol* 1(3): 190-202.

Casey, F. P., J. J. Waterfall, et al. (2008). "Variational method for estimating the rate of convergence of Markov-chain Monte Carlo algorithms." *Phys Rev E Stat Nonlin Soft Matter Phys* 78(4 Pt 2): 046704.

Cebers, A., Z. Dogic, et al. (2006). "Counterion-mediated attraction and kinks on loops of semiflexible polyelectrolyte bundles." *Phys Rev Lett* 96(24): 247801.

Celis, J. E., B. Honore, et al. (1990). "Comprehensive computerized 2D gel protein databases offer a global approach to the study of the mammalian cell." *Bioessays* 12(2): 93-7.

Chadda, R., M. T. Howes, et al. (2007). "Cholesterol-sensitive Cdc42 activation regulates actin polymerization for endocytosis via the GEEC pathway." *Traffic* 8(6): 702-17.

Chan, B. L., M. P. Lisanti, et al. (1988). "Insulin-stimulated release of lipoprotein lipase by metabolism of its phosphatidylinositol anchor." *Science* 241(4873): 1670-2.

Chandler, E., E. Hoover, et al. (2009). "High-resolution mosaic imaging with multifocal, multiphoton photon-counting microscopy." *Appl Opt* 48(11): 2067-77.

Chandrasekhar, S., M. Kac, et al. (2000). *Marian Smoluchowski : his life and scientific work*. Warszawa, Polish Scientific Publishers PWN.

Chang, B. L., S. D. Cramer, et al. (2009). "Fine mapping association study and functional analysis implicate a SNP in MSMB at 10q11 as a causal variant for prostate cancer risk." *Hum Mol Genet* 18(7): 1368-75.

Chang, S., T. M. Svitkina, et al. (1999). "Speckle microscopic evaluation of microtubule transport in growing nerve processes." *Nat Cell Biol* 1(7): 399-403.

Chatterjee, A., M. A. Katsoulakis, et al. (2005). "Spatially adaptive grand canonical ensemble Monte Carlo simulations." *Phys Rev E Stat Nonlin Soft Matter Phys* 71(2 Pt 2): 026702.

Chatterjee, A., K. Mayawala, et al. (2005). "Time accelerated Monte Carlo simulations of biological networks using the binomial tau-leap method." *Bioinformatics* 21(9): 2136-7.

Chatterjee, A., M. A. Snyder, et al. (2004). "Mesoscopic modeling of chemical reactivity." *Chemical Engineering Science* 59(22-23): 5559-5567.

Chatterjee, A. and D. Vlachos (2005). "Temporal acceleration of spatially distributed kinetic Monte Carlo simulations." *Journal of Computational Physics* 211: 596-615.

Chatterjee, A. and D. Vlachos (2007). "An overview of spatial microscopic and accelerated kinetic Monte Carlo methods." *J Computer-Aided Mater Des* 14: 253-308.

Chatterjee, A. and D. G. Vlachos (2006). "Multiscale spatial Monte Carlo simulations: multigriding, computational singular perturbation, and hierarchical stochastic closures." *J Chem Phys* 124(6): 64110.

Chatterjee, A. and D. G. Vlachos (2007). "Continuum mesoscopic framework for multiple interacting species and processes on multiple site types and/or crystallographic planes." *J Chem Phys* 127(3): 034705.

Chatterjee, A. and D. G. Vlachos (2007). "Systems tasks in nanotechnology via hierarchical multiscale modeling: Nanopattern formation in heteroepitaxy." *Chemical Engineering Science* 62(18-20): 4852-4863.

Chatterjee, A., D. G. Vlachos, et al. (2004). "Spatially adaptive lattice coarse-grained Monte Carlo simulations for diffusion of interacting molecules." *J Chem Phys* 121(22): 11420-31.

Chatterjee, A., D. G. Vlachos, et al. (2005). "Binomial distribution based tau-leap accelerated stochastic simulation." *J Chem Phys* 122(2): 024112.

Chatterjee, A., D. G. Vlachos, et al. (2005). "Numerical assessment of theoretical error estimates in coarse-grained kinetic Monte Carlo simulations: Application to surface diffusion." *International Journal for Multiscale Computational Engineering* 3(1): 59-70.

Chavrier, P., J. P. Gorvel, et al. (1991). "Hypervariable C-terminal domain of rab proteins acts as a targeting signal." *Nature* 353(6346): 769-72.

Chen, C. and D. Mochly-Rosen (2001). "Opposing effects of delta and xi PKC in ethanol-induced cardioprotection." *J Mol Cell Cardiol* 33(3): 581-5.

Chen, C. H., G. R. Budas, et al. (2008). "Activation of aldehyde dehydrogenase-2 reduces ischemic damage to the heart." *Science* 321(5895): 1493-5.

Chen, D. B., S. Jia, et al. (2006). "Global protein expression profiling underlines reciprocal regulation of caveolin 1 and endothelial nitric oxide synthase expression in ovariectomized sheep uterine artery by estrogen/progesterone replacement therapy." *Biol Reprod* 74(5): 832-8.

Chen, J., J. Dai, et al. (1997). "Loss of cytoskeletal support is not sufficient for anoxic plasma membrane disruption in renal cells." *Am J Physiol* 272(4 Pt 1): C1319-28.

- Chen, J., F. Saeki, et al. (2005). "Gold nanocages: bioconjugation and their potential use as optical imaging contrast agents." *Nano Lett* 5(3): 473-7.
- Chen, J., W. Wang, et al. (2002). "Molecular wires, switches, and memories." *Ann N Y Acad Sci* 960: 69-99.
- Chen, K. F., I. Adachi, et al. (2007). "Search for $B \rightarrow h(*) \nu \nu$ decays at Belle." *Phys Rev Lett* 99(22): 221802.
- Chen, L., H. Hahn, et al. (2001). "Opposing cardioprotective actions and parallel hypertrophic effects of delta PKC and epsilon PKC." *Proc Natl Acad Sci U S A* 98(20): 11114-9.
- Chen, L., L. R. Wright, et al. (2001). "Molecular transporters for peptides: delivery of a cardioprotective epsilonPKC agonist peptide into cells and intact ischemic heart using a transport system, R(7)." *Chem Biol* 8(12): 1123-9.
- Chen, T. Y., P. H. Liu, et al. (2006). "The intracellular domain of amyloid precursor protein interacts with flotillin-1, a lipid raft protein." *Biochem Biophys Res Commun* 342(1): 266-72.
- Chen, W. W., B. Schoeberl, et al. (2009). "Input-output behavior of ErbB signaling pathways as revealed by a mass action model trained against dynamic data." *Mol Syst Biol* 5: 239.
- Cheng, L., M. E. Rivero, et al. (2001). "Evaluation of intraocular pharmacokinetics and toxicity of prinomastat (AG3340) in the rabbit." *J Ocul Pharmacol Ther* 17(3): 295-304.
- Cherry, R. J. (1979). "Rotational and lateral diffusion of membrane proteins." *Biochim Biophys Acta* 559(4): 289-327.
- Cherry, R. J., R. E. Godfrey, et al. (1982). "Mobility of bacteriorhodopsin in lipid vesicles." *Biochem Soc Trans* 10(5): 342-3.
- Chichili, G. R. and W. Rodgers (2007). "Clustering of membrane raft proteins by the actin cytoskeleton." *J Biol Chem* 282(50): 36682-91.
- Chickarmane, V., B. N. Kholodenko, et al. (2007). "Oscillatory dynamics arising from competitive inhibition and multisite phosphorylation." *J Theor Biol* 244(1): 68-76.
- Cho, K. H., S. M. Choo, et al. (2005). "A unified framework for unraveling the functional interaction structure of a biomolecular network based on stimulus-response experimental data." *FEBS Lett* 579(20): 4520-8.

Choi, J., X. Zhang, et al. (2009). "Building alkali-metal-halide layers within a perovskite host by sequential intercalation: $(A(2)Cl)LaNb(2)O(7)$ ($A = Rb, Cs$).
Inorg Chem 48(11): 4811-6.

Chopra, H., S. E. Fligiel, et al. (1990). "An in vivo study of the role of the tumor cell cytoskeleton in tumor cell-platelet-endothelial cell interactions." Cancer Res 50(23): 7686-96.

Chopra, H., J. S. Hatfield, et al. (1988). "Role of tumor cytoskeleton and membrane glycoprotein IRGpIIb/IIIa in platelet adhesion to tumor cell membrane and tumor cell-induced platelet aggregation." Cancer Res 48(13): 3787-800.

Chopra, H., J. Timar, et al. (1991). "The lipoxygenase metabolite 12(S)-HETE induces a cytoskeleton-dependent increase in surface expression of integrin alpha IIb beta 3 on melanoma cells." Int J Cancer 49(5): 774-86.

Chopra, H., J. Timar, et al. (1992). "Is there a role for the tumor cell integrin alpha IIb beta 3 and cytoskeleton in tumor cell-platelet interaction?" Clin Exp Metastasis 10(2): 125-37.

Choquet, D., D. P. Felsenfeld, et al. (1997). "Extracellular matrix rigidity causes strengthening of integrin-cytoskeleton linkages." Cell 88(1): 39-48.

Chung, J. W., K. S. Shin, et al. (1992). "Evidence for the origin of reconstruction of the Mo(001) surface." Phys Rev Lett 69(15): 2228-2231.

Ciruela, F. (2008). "Fluorescence-based methods in the study of protein-protein interactions in living cells." Curr Opin Biotechnol 19(4): 338-43.

Clamp, A. G., S. Ladha, et al. (1994). "Determination of membrane fluidity: a comparison of biophysical methods." Biochem Soc Trans 22(3): 369S.

Clamp, A. G., S. Ladha, et al. (1997). "The influence of dietary lipids on the composition and membrane fluidity of rat hepatocyte plasma membrane." Lipids 32(2): 179-84.

Clapp, A. R., I. L. Medintz, et al. (2005). "Can luminescent quantum dots be efficient energy acceptors with organic dye donors?" J Am Chem Soc 127(4): 1242-50.

Clapp, A. R., I. L. Medintz, et al. (2006). "Forster resonance energy transfer investigations using quantum-dot fluorophores." Chemphyschem 7(1): 47-57.

Clapp, A. R., I. L. Medintz, et al. (2004). "Fluorescence resonance energy transfer between quantum dot donors and dye-labeled protein acceptors." J Am Chem Soc 126(1): 301-10.

- Clapp, A. R., I. L. Medintz, et al. (2005). "Quantum dot-based multiplexed fluorescence resonance energy transfer." *J Am Chem Soc* 127(51): 18212-21.
- Clark, A. G., M. B. Eisen, et al. (2007). "Evolution of genes and genomes on the *Drosophila* phylogeny." *Nature* 450(7167): 203-18.
- Clarkson, B. H., F. F. Feagin, et al. (1991). "Effects of phosphoprotein moieties on the remineralization of human root caries." *Caries Res* 25(3): 166-73.
- Clyde-Smith, J., G. Silins, et al. (2000). "Characterization of RasGRP2, a plasma membrane-targeted, dual specificity Ras/Rap exchange factor." *J Biol Chem* 275(41): 32260-7.
- Cole, N. B., J. Ellenberg, et al. (1998). "Retrograde transport of Golgi-localized proteins to the ER." *J Cell Biol* 140(1): 1-15.
- Cole, N. B. and J. Lippincott-Schwartz (1995). "Organization of organelles and membrane traffic by microtubules." *Curr Opin Cell Biol* 7(1): 55-64.
- Cole, N. B., N. Sciaky, et al. (1996). "Golgi dispersal during microtubule disruption: regeneration of Golgi stacks at peripheral endoplasmic reticulum exit sites." *Mol Biol Cell* 7(4): 631-50.
- Cole, N. B., C. L. Smith, et al. (1996). "Diffusional mobility of Golgi proteins in membranes of living cells." *Science* 273(5276): 797-801.
- Collier, N. C., M. P. Sheetz, et al. (1993). "Concomitant changes in mitochondria and intermediate filaments during heat shock and recovery of chicken embryo fibroblasts." *J Cell Biochem* 52(3): 297-307.
- Collins, J. W., Jr., M. Hoppe, et al. (1991). "A controlled trial of insulin infusion and parenteral nutrition in extremely low birth weight infants with glucose intolerance." *J Pediatr* 118(6): 921-7.
- Collins, S. D., A. Chatterjee, et al. (2008). "Coarse-grained kinetic Monte Carlo models: Complex lattices, multicomponent systems, and homogenization at the stochastic level." *J Chem Phys* 129(18): 184101.
- Conigrave, A. D., K. C. Fernando, et al. (2001). "P2Y(11) receptor expression by human lymphocytes: evidence for two cAMP-linked purinoceptors." *Eur J Pharmacol* 426(3): 157-63.
- Conzelmann, H., J. Saez-Rodriguez, et al. (2006). "A domain-oriented approach to the reduction of combinatorial complexity in signal transduction networks." *BMC Bioinformatics* 7: 34.

Corash, L., S. Spielberg, et al. (1980). "Reduced chronic hemolysis during high-dose vitamin E administration in Mediterranean-type glucose-6-phosphate dehydrogenase deficiency." *N Engl J Med* 303(8): 416-20.

Corash, L. M., M. Sheetz, et al. (1982). "Chronic hemolytic anemia due to glucose-6-phosphate dehydrogenase deficiency or glutathione synthetase deficiency: the role of vitamin E in its treatment." *Ann N Y Acad Sci* 393: 348-60.

Cosgrove, P. and M. P. Sheetz (1982). "Effect of cell shape on extravascular hemolysis." *Blood* 59(2): 421-7.

Coussen, F., D. Choquet, et al. (2002). "Trimers of the fibronectin cell adhesion domain localize to actin filament bundles and undergo rearward translocation." *J Cell Sci* 115(Pt 12): 2581-90.

Crewe, A. V. and J. Wall (1970). "A scanning microscope with 5 Å resolution." *J Mol Biol* 48(3): 375-93.

Crewe, A. V., J. Wall, et al. (1970). "Visibility of Single Atoms." *Science* 168(3937): 1338-1340.

Crewe, N. M. (1969). "Training course: stroke in your family." *Rehabil Rec* 10(1): 32-4.

Crewe, W. and S. M. Crewe (1969). "Possible transmission of bovine cysticercosis by gulls." *Trans R Soc Trop Med Hyg* 63(1): 17.

Cruz, M., J. M. Hernandez, et al. (1999). "Surface redistribution of interferon gamma-receptor and its colocalization with the actin cytoskeleton." *Arch Med Res* 30(2): 97-105.

Curie, M. and M. Smoluchowski (1968). *Studia poświęcone Marii Skłodowskiej-Curie i Marianowi Smoluchowskiemu. Wrocław, Zakład Narodowy im. Ossolińskich.*

Curnow, S. J., D. Scheel-Toellner, et al. (2004). "Inhibition of T cell apoptosis in the aqueous humor of patients with uveitis by IL-6/soluble IL-6 receptor trans-signaling." *J Immunol* 173(8): 5290-7.

Dabora, S. L. and M. P. Sheetz (1988). "Cultured cell extracts support organelle movement on microtubules in vitro." *Cell Motil Cytoskeleton* 10(4): 482-95.

Dabora, S. L. and M. P. Sheetz (1988). "The microtubule-dependent formation of a tubulovesicular network with characteristics of the ER from cultured cell extracts." *Cell* 54(1): 27-35.

- Dai, J. and M. P. Sheetz (1995). "Axon membrane flows from the growth cone to the cell body." *Cell* 83(5): 693-701.
- Dai, J. and M. P. Sheetz (1995). "Mechanical properties of neuronal growth cone membranes studied by tether formation with laser optical tweezers." *Biophys J* 68(3): 988-96.
- Dai, J. and M. P. Sheetz (1995). "Regulation of endocytosis, exocytosis, and shape by membrane tension." *Cold Spring Harb Symp Quant Biol* 60: 567-71.
- Dai, J. and M. P. Sheetz (1998). "Cell membrane mechanics." *Methods Cell Biol* 55: 157-71.
- Dai, J. and M. P. Sheetz (1999). "Membrane tether formation from blebbing cells." *Biophys J* 77(6): 3363-70.
- Dai, J., M. P. Sheetz, et al. (1998). "Membrane tension in swelling and shrinking molluscan neurons." *J Neurosci* 18(17): 6681-92.
- Dai, J., H. P. Ting-Beall, et al. (1999). "Myosin I contributes to the generation of resting cortical tension." *Biophys J* 77(2): 1168-76.
- Dai, J., H. P. Ting-Beall, et al. (1997). "The secretion-coupled endocytosis correlates with membrane tension changes in RBL 2H3 cells." *J Gen Physiol* 110(1): 1-10.
- Damjanovich, S., J. Bot, et al. (1972). "Effect of glycerol on some kinetic parameters of phosphorylase b." *Biochim Biophys Acta* 284(1): 345-8.
- Damjanovich, S., G. Vereb, et al. (1995). "Structural hierarchy in the clustering of HLA class I molecules in the plasma membrane of human lymphoblastoid cells." *Proc Natl Acad Sci U S A* 92(4): 1122-6.
- Danuser, G. and C. M. Waterman-Storer (2003). "Quantitative fluorescent speckle microscopy: where it came from and where it is going." *J Microsc* 211(Pt 3): 191-207.
- Danuser, G. and C. M. Waterman-Storer (2006). "Quantitative fluorescent speckle microscopy of cytoskeleton dynamics." *Annu Rev Biophys Biomol Struct* 35: 361-87.
- Davey, M. L., M. R. Hall, et al. (2000). "Five Crustacean Hyperglycemic Family Hormones of *Penaeus monodon*: Complementary DNA Sequence and Identification in Single Sinus Glands by Electrospray Ionization-Fourier Transform Mass Spectrometry." *Mar Biotechnol (NY)* 2(1): 80-91.
- Davies, P., B. Anderton, et al. (1998). "First one in, last one out: the role of gabaergic transmission in generation and degeneration." *Prog Neurobiol* 55(6): 651-8.

- Davis, C. N., E. Mann, et al. (2006). "MyD88-dependent and -independent signaling by IL-1 in neurons probed by bifunctional Toll/IL-1 receptor domain/BB-loop mimetics." *Proc Natl Acad Sci U S A* 103(8): 2953-8.
- Davis, C. N., I. Tabarean, et al. (2006). "IL-1beta induces a MyD88-dependent and ceramide-mediated activation of Src in anterior hypothalamic neurons." *J Neurochem* 98(5): 1379-89.
- Davis, M. D., M. J. Sheetz, et al. (2009). "Effect of ruboxistaurin on the visual acuity decline associated with long-standing diabetic macular edema." *Invest Ophthalmol Vis Sci* 50(1): 1-4.
- Dawson, J. P., M. B. Berger, et al. (2005). "Epidermal growth factor receptor dimerization and activation require ligand-induced conformational changes in the dimer interface." *Mol Cell Biol* 25(17): 7734-42.
- D'Cruz, I. A., E. Calderon, et al. (1998). "Transthoracic Echocardiographic Visualization of Calcification of the Sinotubular Ridge of the Ascending Aorta." *Echocardiography* 15(5): 425-428.
- D'Cruz, I. A., E. Calderon, et al. (1997). "Left ventricular diastolic compression in acquired immunodeficiency syndrome with large, nonloculated pericardial effusion." *Am Heart J* 133(3): 383-4.
- D'Cruz, I. A., E. Calderon, et al. (1997). "Acquired Ventricular Septal Defect Following Stab Wound Color Flow Doppler Diagnosis." *Echocardiography* 14(4): 409-410.
- De Atauri, P., L. Acerenza, et al. (2001). "Occurrence of paradoxical or sustained control by an enzyme when overexpressed: necessary conditions and experimental evidence with regard to hepatic glucokinase." *Biochem J* 355(Pt 3): 787-93.
- de Figueiredo, P., D. Drecktrah, et al. (2000). "Phospholipase A2 antagonists inhibit constitutive retrograde membrane traffic to the endoplasmic reticulum." *Traffic* 1(6): 504-11.
- de Kort, M., V. Correa, et al. (2000). "Synthesis of potent agonists of the D-myo-inositol 1,4, 5-trisphosphate receptor based on clustered disaccharide polyphosphate analogues of adenophostin A." *J Med Chem* 43(17): 3295-303.
- de Kort, M., A. D. Regenbogen, et al. (2000). "Spirophostins: conformationally restricted analogues of adenophostin A." *Chemistry* 6(15): 2696-704.
- de Rooij, J., A. Kerstens, et al. (2005). "Integrin-dependent actomyosin contraction regulates epithelial cell scattering." *J Cell Biol* 171(1): 153-64.
- De Vos, K. J., V. J. Allan, et al. (2005). "Mitochondrial function and actin regulate dynamin-related protein 1-dependent mitochondrial fission." *Curr Biol* 15(7): 678-83.

De Vos, K. J., J. Sable, et al. (2003). "Expression of phosphatidylinositol (4,5) biphosphate-specific pleckstrin homology domains alters direction but not the level of axonal transport of mitochondria." *Mol Biol Cell* 14(9): 3636-49.

De Vos, K. J. and M. P. Sheetz (2007). "Visualization and quantification of mitochondrial dynamics in living animal cells." *Methods Cell Biol* 80: 627-82.

Deerinck, T. J., B. N. Giepmans, et al. (2007). "Light and electron microscopic localization of multiple proteins using quantum dots." *Methods Mol Biol* 374: 43-53.

del Rio, A., R. Perez-Jimenez, et al. (2009). "Stretching single talin rod molecules activates vinculin binding." *Science* 323(5914): 638-41.

Delehanty, J. B., H. Mattoussi, et al. (2009). "Delivering quantum dots into cells: strategies, progress and remaining issues." *Anal Bioanal Chem* 393(4): 1091-105.

Delehanty, J. B., I. L. Medintz, et al. (2006). "Self-assembled quantum dot-peptide bioconjugates for selective intracellular delivery." *Bioconjug Chem* 17(4): 920-7.

Delmas, D., C. Rebe, et al. (2003). "Resveratrol-induced apoptosis is associated with Fas redistribution in the rafts and the formation of a death-inducing signaling complex in colon cancer cells." *J Biol Chem* 278(42): 41482-90.

Delorme, V., M. Machacek, et al. (2007). "Cofilin activity downstream of Pak1 regulates cell protrusion efficiency by organizing lamellipodium and lamella actin networks." *Dev Cell* 13(5): 646-62.

Demin, O. V., Gorianin, II, et al. (2001). "[Kinetic modeling of energy metabolism and generation of active forms of oxygen in hepatocyte mitochondria]." *Mol Biol (Mosk)* 35(6): 1095-104.

Demin, O. V., H. V. Westerhoff, et al. (1998). "Mathematical modelling of superoxide generation with the bc1 complex of mitochondria." *Biochemistry (Mosc)* 63(6): 634-49.

Deshpande, V. S., R. M. McMeeking, et al. (2006). "A bio-chemo-mechanical model for cell contractility." *Proc Natl Acad Sci U S A* 103(38): 14015-20.

Dey, A., J. Ellenberg, et al. (2000). "A bromodomain protein, MCAP, associates with mitotic chromosomes and affects G(2)-to-M transition." *Mol Cell Biol* 20(17): 6537-49.

Dharmastiti, R., D. C. Barton, et al. (2001). "The wear of oriented UHMWPE under isotropically rough and scratched counterface test conditions." *Biomed Mater Eng* 11(3): 241-56.

- Di Fiore, P. P., O. Segatto, et al. (1990). "EGF receptor and erbB-2 tyrosine kinase domains confer cell specificity for mitogenic signaling." *Science* 248(4951): 79-83.
- Dickson, R. C., R. M. Sheetz, et al. (1981). "Genetic regulation: yeast mutants constitutive for beta-galactosidase activity have an increased level of beta-galactosidase messenger ribonucleic acid." *Mol Cell Biol* 1(11): 1048-56.
- Diebold, Y., L. L. Chen, et al. (2007). "Lymphocytic infiltration and goblet cell marker alteration in the conjunctiva of the MRL/MpJ-Fas(lpr) mouse model of Sjogren's syndrome." *Exp Eye Res* 84(3): 500-12.
- Dietrich, C., L. A. Bagatolli, et al. (2001). "Lipid rafts reconstituted in model membranes." *Biophys J* 80(3): 1417-28.
- Dietrich, C., Z. N. Volovyk, et al. (2001). "Partitioning of Thy-1, GM1, and cross-linked phospholipid analogs into lipid rafts reconstituted in supported model membrane monolayers." *Proc Natl Acad Sci U S A* 98(19): 10642-7.
- Dietrich, C., B. Yang, et al. (2002). "Relationship of lipid rafts to transient confinement zones detected by single particle tracking." *Biophys J* 82(1 Pt 1): 274-84.
- Dimaggio, G. T. and A. H. Sheetz (1983). "The concerns of mothers caring for an infant on an apnea monitor." *MCN Am J Matern Child Nurs* 8(4): 294-7.
- Dmitriev, V. V., N. E. Suzina, et al. (2008). "[Electron microscopic detection and in situ characterization of bacterial nanoforms in extreme biotopes]." *Mikrobiologiya* 77(1): 46-54.
- Dobereiner, H. G., B. Dubin-Thaler, et al. (2004). "Dynamic phase transitions in cell spreading." *Phys Rev Lett* 93(10): 108105.
- Dobereiner, H. G., B. J. Dubin-Thaler, et al. (2005). "Force sensing and generation in cell phases: analyses of complex functions." *J Appl Physiol* 98(4): 1542-6.
- Dobereiner, H. G., B. J. Dubin-Thaler, et al. (2006). "Lateral membrane waves constitute a universal dynamic pattern of motile cells." *Phys Rev Lett* 97(3): 038102.
- Doebbeling, B. N., G. L. Stanley, et al. (1992). "Comparative efficacy of alternative hand-washing agents in reducing nosocomial infections in intensive care units." *N Engl J Med* 327(2): 88-93.
- Doherty, G. J. and H. T. McMahon (2008). "Mediation, modulation, and consequences of membrane-cytoskeleton interactions." *Annu Rev Biophys* 37: 65-95.

- d'Onofrio, A. and I. P. Tomlinson (2007). "A nonlinear mathematical model of cell turnover, differentiation and tumorigenesis in the intestinal crypt." *J Theor Biol* 244(3): 367-74.
- Dorn, G. W., 2nd, M. C. Souroujon, et al. (1999). "Sustained in vivo cardiac protection by a rationally designed peptide that causes epsilon protein kinase C translocation." *Proc Natl Acad Sci U S A* 96(22): 12798-803.
- Douglass, A. D. and R. D. Vale (2005). "Single-molecule microscopy reveals plasma membrane microdomains created by protein-protein networks that exclude or trap signaling molecules in T cells." *Cell* 121(6): 937-50.
- Dovey, H. F., V. John, et al. (2001). "Functional gamma-secretase inhibitors reduce beta-amyloid peptide levels in brain." *J Neurochem* 76(1): 173-81.
- Dragsten, P. R., R. Blumenthal, et al. (1981). "Membrane asymmetry in epithelia: is the tight junction a barrier to diffusion in the plasma membrane?" *Nature* 294(5843): 718-22.
- Dragsten, P. R., J. S. Handler, et al. (1982). "Asymmetry in epithelial cells: is the tight junction a barrier to lateral diffusion in the plasma membrane?" *Prog Clin Biol Res* 91: 525-36.
- Dragsten, P. R., J. S. Handler, et al. (1982). "Fluorescent membrane probes and the mechanism of maintenance of cellular asymmetry in epithelia." *Fed Proc* 41(1): 48-53.
- Dubin-Thaler, B. J., G. Giannone, et al. (2004). "Nanometer analysis of cell spreading on matrix-coated surfaces reveals two distinct cell states and STEPs." *Biophys J* 86(3): 1794-806.
- Dubin-Thaler, B. J., J. M. Hofman, et al. (2008). "Quantification of cell edge velocities and traction forces reveals distinct motility modules during cell spreading." *PLoS ONE* 3(11): e3735.
- Ebert, S., T. Schoeberl, et al. (2008). "Chondroitin sulfate disaccharide stimulates microglia to adopt a novel regulatory phenotype." *J Leukoc Biol* 84(3): 736-40.
- Echigo, A., M. Aoki, et al. (2008). "The excursion of the median nerve during nerve gliding exercise: an observation with high-resolution ultrasonography." *J Hand Ther* 21(3): 221-7; quiz 228.
- Edidin, M. (2001). "Membrane cholesterol, protein phosphorylation, and lipid rafts." *Sci STKE* 2001(67): PE1.
- Edidin, M. (2001). "Near-field scanning optical microscopy, a siren call to biology." *Traffic* 2(11): 797-803.

- Edidin, M. (2001). "Shrinking patches and slippery rafts: scales of domains in the plasma membrane." *Trends Cell Biol* 11(12): 492-6.
- Edidin, M., S. C. Kuo, et al. (1991). "Lateral movements of membrane glycoproteins restricted by dynamic cytoplasmic barriers." *Science* 254(5036): 1379-82.
- Edidin, M., M. C. Zuniga, et al. (1994). "Truncation mutants define and locate cytoplasmic barriers to lateral mobility of membrane glycoproteins." *Proc Natl Acad Sci U S A* 91(8): 3378-82.
- Egerton, R. (2005). *Physical principles of electron microscopy*, Springer.
- Ellis, P. J., R. A. Furlong, et al. (2004). "Modulation of the mouse testis transcriptome during postnatal development and in selected models of male infertility." *Mol Hum Reprod* 10(4): 271-81.
- Emiliusen, L., M. Gough, et al. (2001). "A transcriptional feedback loop for tissue-specific expression of highly cytotoxic genes which incorporates an immunostimulatory component." *Gene Ther* 8(13): 987-98.
- Epstein, E. E., S. L. Soter, et al. (1967). "Mercury: Observations of the 3.4-Millimeter Radio Emission." *Science* 157(3796): 1550-1552.
- Escobar-Alvarez, S., Y. Goldgur, et al. (2009). "Structure and activity of human mitochondrial peptide deformylase, a novel cancer target." *J Mol Biol* 387(5): 1211-28.
- Eswarakumar, V. P., F. Ozcan, et al. (2006). "Attenuation of signaling pathways stimulated by pathologically activated FGF-receptor 2 mutants prevents craniosynostosis." *Proc Natl Acad Sci U S A* 103(49): 18603-8.
- Evans, H. L., J. M. Polski, et al. (2000). "CD5+ true SLL/CLL with plasmacytic differentiation and an unusual 1p36 translocation: case report and review of the literature." *Leuk Lymphoma* 39(5-6): 625-32.
- Faeder, J. R., M. L. Blinov, et al. (2005). "Combinatorial complexity and dynamical restriction of network flows in signal transduction." *Syst Biol (Stevenage)* 2(1): 5-15.
- Faeder, J. R., W. S. Hlavacek, et al. (2003). "Investigation of early events in Fc epsilon RI-mediated signaling using a detailed mathematical model." *J Immunol* 170(7): 3769-81.
- Fahmy, T. M., J. G. Bieler, et al. (2001). "Increased TCR avidity after T cell activation: a mechanism for sensing low-density antigen." *Immunity* 14(2): 135-43.

- Falasca, M., S. K. Logan, et al. (1998). "Activation of phospholipase C gamma by PI 3-kinase-induced PH domain-mediated membrane targeting." *EMBO J* 17(2): 414-22.
- Fang, X., D. Gaudette, et al. (2000). "Lysophospholipid growth factors in the initiation, progression, metastases, and management of ovarian cancer." *Ann N Y Acad Sci* 905: 188-208.
- Felley-Bosco, E., F. C. Bender, et al. (2000). "Caveolin-1 down-regulates inducible nitric oxide synthase via the proteasome pathway in human colon carcinoma cells." *Proc Natl Acad Sci U S A* 97(26): 14334-9.
- Felsenfeld, D. P., D. Choquet, et al. (1996). "Ligand binding regulates the directed movement of beta1 integrins on fibroblasts." *Nature* 383(6599): 438-40.
- Felsenfeld, D. P., P. L. Schwartzberg, et al. (1999). "Selective regulation of integrin--cytoskeleton interactions by the tyrosine kinase Src." *Nat Cell Biol* 1(4): 200-6.
- Fenteany, G., P. A. Janmey, et al. (2000). "Signaling pathways and cell mechanics involved in wound closure by epithelial cell sheets." *Curr Biol* 10(14): 831-8.
- Ferguson, K. M., M. A. Lemmon, et al. (1994). "Crystal structure at 2.2 Å resolution of the pleckstrin homology domain from human dynamin." *Cell* 79(2): 199-209.
- Ferguson, K. M., M. A. Lemmon, et al. (1995). "Structure of the high affinity complex of inositol trisphosphate with a phospholipase C pleckstrin homology domain." *Cell* 83(6): 1037-46.
- Ferguson, K. M., M. A. Lemmon, et al. (1995). "Scratching the surface with the PH domain." *Nat Struct Biol* 2(9): 715-8.
- Ferre, S., F. Ciruela, et al. (2008). "Adenosine A1-A2A receptor heteromers: new targets for caffeine in the brain." *Front Biosci* 13: 2391-9.
- Ferre, S., C. Quiroz, et al. (2008). "An update on adenosine A2A-dopamine D2 receptor interactions: implications for the function of G protein-coupled receptors." *Curr Pharm Des* 14(15): 1468-74.
- Feynman, R. P., N. Metropolis, et al. (1949). "Equations of State of Elements Based on the Generalized Fermi-Thomas Theory." *Phys Rev E Stat Nonlin Soft Matter Phys* 75(10): 1561-1573.
- Filippov, A., G. Oradd, et al. (2004). "Lipid lateral diffusion in ordered and disordered phases in raft mixtures." *Biophys J* 86(2): 891-6.
- Filippov, A., G. Oradd, et al. (2006). "Sphingomyelin structure influences the lateral diffusion and raft formation in lipid bilayers." *Biophys J* 90(6): 2086-92.

- Fink, A. L., L. J. Calciano, et al. (1994). "Classification of acid denaturation of proteins: intermediates and unfolded states." *Biochemistry* 33(41): 12504-11.
- Firestone, A. R., F. F. Feagin, et al. (1993). "In vitro demineralization by strains of *Actinomyces viscosus* and *Streptococcus sobrinus* of sound and demineralized root surfaces." *J Dent Res* 72(8): 1180-3.
- Fitzgerald, J. B., B. Schoeberl, et al. (2006). "Systems biology and combination therapy in the quest for clinical efficacy." *Nat Chem Biol* 2(9): 458-66.
- Flicker, P. F., G. Peltz, et al. (1985). "Site-specific inhibition of myosin-mediated motility in vitro by monoclonal antibodies." *J Cell Biol* 100(4): 1024-30.
- Forgacs, G., S. H. Yook, et al. (2004). "Role of the cytoskeleton in signaling networks." *J Cell Sci* 117(Pt 13): 2769-75.
- Forrester, K., S. Ambs, et al. (1996). "Nitric oxide-induced p53 accumulation and regulation of inducible nitric oxide synthase expression by wild-type p53." *Proc Natl Acad Sci U S A* 93(6): 2442-7.
- Franco, R., V. Casado, et al. (2008). "G-protein-coupled receptor heteromers: function and ligand pharmacology." *Br J Pharmacol* 153 Suppl 1: S90-8.
- Frankel, D. J., J. R. Pfeiffer, et al. (2006). "Revealing the topography of cellular membrane domains by combined atomic force microscopy/fluorescence imaging." *Biophys J* 90(7): 2404-13.
- Freeman, P. and A. E. Gear (1971). "Probabilistic Objective Function for R and D Portfolio Selection." *Operational Research Quarterly* 22(3): 253-&.
- Friedman, H. M., S. Sheetz, et al. (1986). "Combination chemotherapy and radiation therapy. The medical management of epidural spinal cord compression from testicular cancer." *Arch Intern Med* 146(3): 509-12.
- Frings, L., K. Wagner, et al. (2008). "Early detection of behavioral side effects of antiepileptic treatment using handheld computers." *Epilepsy Behav* 13(2): 402-6.
- Frost, F. J., R. L. Calderon, et al. (1998). "A two-year follow-up survey of antibody to *Cryptosporidium* in Jackson County, Oregon following an outbreak of waterborne disease." *Epidemiol Infect* 121(1): 213-7.
- Frost, F. J., A. A. de la Cruz, et al. (1998). "Comparisons of ELISA and Western blot assays for detection of *Cryptosporidium* antibody." *Epidemiol Infect* 121(1): 205-11.
- Fu, H., L. Bjorkman, et al. (2004). "The two neutrophil members of the formylpeptide receptor family activate the NADPH-oxidase through signals that differ in sensitivity to a gelsolin derived phosphoinositide-binding peptide." *BMC Cell Biol* 5(1): 50.

Fuchs, E., Y. M. Chan, et al. (1994). "Cracks in the foundation: keratin filaments and genetic disease." *Trends Cell Biol* 4(9): 321-6.

Fujita, H., P. G. Allen, et al. (1999). "Induction of apoptosis by gelsolin truncates." *Ann N Y Acad Sci* 886: 217-20.

Fujiwara, H., Y. Tanaka, et al. (1997). "Anthocyanin 5-aromatic acyltransferase from *Gentiana triflora*. Purification, characterization and its role in anthocyanin biosynthesis." *Eur J Biochem* 249(1): 45-51.

Fujiwara, H., Y. Tanaka, et al. (1998). "cDNA cloning, gene expression and subcellular localization of anthocyanin 5-aromatic acyltransferase from *Gentiana triflora*." *Plant J* 16(4): 421-31.

Fujiwara, S., S. Kusumi, et al. (2000). "Prevalence of anti-hepatitis C virus antibody and chronic liver disease among atomic bomb survivors." *Radiat Res* 154(1): 12-9.

Fujiwara, S., G. B. Sharp, et al. (2003). "Prevalence of hepatitis B virus infection among atomic bomb survivors." *Radiat Res* 159(6): 780-6.

Fujiwara, T., K. Ritchie, et al. (2002). "Phospholipids undergo hop diffusion in compartmentalized cell membrane." *J Cell Biol* 157(6): 1071-81.

Fulton, A. M., J. Mellor, et al. (1985). "Variants within the yeast Ty sequence family encode a class of structurally conserved proteins." *Nucleic Acids Res* 13(11): 4097-112.

Funaki, M., L. DiFransico, et al. (2006). "PI 4,5-P2 stimulates glucose transport activity of GLUT4 in the plasma membrane of 3T3-L1 adipocytes." *Biochim Biophys Acta* 1763(8): 889-99.

Fushman, D., S. Cahill, et al. (1995). "Solution structure of pleckstrin homology domain of dynamin by heteronuclear NMR spectroscopy." *Proc Natl Acad Sci U S A* 92(3): 816-20.

Gaidarov, G. M. and N. Alekseeva (2007). "[The medical economic approaches to the enhancement of the effectiveness of the functioning of the multi-field hospital in the conditions of the mandatory medical insurance]." *Probl Sotsialnoi Gig Zdravookhranenniai Istor Med*(5): 29-30.

Gaidarov, G. M. and S. V. Makarov (2008). "[The application of the social economic zoning in the study of malignant neoplasm mortality]." *Probl Sotsialnoi Gig Zdravookhranenniai Istor Med*(4): 5-9.

- Gaidarov, G. M., T. P. Maslauskene, et al. (2008). "[The historical aspects of the antituberculous service activities in the Irkutsk oblast]." *Probl Sotsialnoi Gig Zdravookhranennii i Istor Med*(3): 57-9.
- Gaidarov, I. and J. H. Keen (1999). "Phosphoinositide-AP-2 interactions required for targeting to plasma membrane clathrin-coated pits." *J Cell Biol* 146(4): 755-64.
- Gaidarov, I., J. G. Krupnick, et al. (1999). "Arrestin function in G protein-coupled receptor endocytosis requires phosphoinositide binding." *EMBO J* 18(4): 871-81.
- Gaidarov, I., F. Santini, et al. (1999). "Spatial control of coated-pit dynamics in living cells." *Nat Cell Biol* 1(1): 1-7.
- Gaietta, G. M., B. N. Giepmans, et al. (2006). "Golgi twins in late mitosis revealed by genetically encoded tags for live cell imaging and correlated electron microscopy." *Proc Natl Acad Sci U S A* 103(47): 17777-82.
- Galbraith, C. G. and M. P. Sheetz (1997). "A micromachined device provides a new bend on fibroblast traction forces." *Proc Natl Acad Sci U S A* 94(17): 9114-8.
- Galbraith, C. G. and M. P. Sheetz (1998). "Forces on adhesive contacts affect cell function." *Curr Opin Cell Biol* 10(5): 566-71.
- Galbraith, C. G. and M. P. Sheetz (1999). "Keratocytes pull with similar forces on their dorsal and ventral surfaces." *J Cell Biol* 147(6): 1313-24.
- Galbraith, C. G. and M. P. Sheetz (2001). "Cell traction." *Curr Protoc Cell Biol* Chapter 12: Unit 12 3.
- Galbraith, C. G., K. M. Yamada, et al. (2002). "The relationship between force and focal complex development." *J Cell Biol* 159(4): 695-705.
- Gallegos, A. M., S. M. Storey, et al. (2006). "Structure and cholesterol dynamics of caveolae/raft and nonraft plasma membrane domains." *Biochemistry* 45(39): 12100-16.
- Gandia, J., J. Galino, et al. (2008). "Detection of higher-order G protein-coupled receptor oligomers by a combined BRET-BiFC technique." *FEBS Lett* 582(20): 2979-84.
- Gandia, J., C. Lluís, et al. (2008). "Light resonance energy transfer-based methods in the study of G protein-coupled receptor oligomerization." *Bioessays* 30(1): 82-9.
- Ganguly, S., T. J. Pucadyil, et al. (2008). "Actin cytoskeleton-dependent dynamics of the human serotonin1A receptor correlates with receptor signaling." *Biophys J* 95(1): 451-63.

- Garcia Rubira, J. C., D. Romero Chacon, et al. (1992). "[Prevalence of coronary disease in patients with aortic stenosis]." *Rev Esp Cardiol* 45(7): 427-31.
- Garcia-Blanco, M. A., D. D. Miller, et al. (1995). "Nuclear spreads: I. Visualization of bipartite ribosomal RNA domains." *J Cell Biol* 128(1-2): 15-27.
- Gaspar, R., Jr., G. Panyi, et al. (1994). "Effects of bretylium tosylate on voltage-gated potassium channels in human T lymphocytes." *Mol Pharmacol* 46(4): 762-6.
- Gauthier, N. C., O. M. Rossier, et al. (2009). "Plasma Membrane Area Increases with Spread Area by Exocytosis of a GPI Anchored Protein Compartment." *Mol Biol Cell*.
- Gear, A. E., A. G. Lockett, et al. (1971). "Analysis of Some Portfolio Selection Models for R+D." *Ieee Transactions on Engineering Management* Em18(2): 66-&.
- Gear, A. R. L. (1971). "Changes in Particle Volume and Numbers during Mitochondrial Biogenesis in Regenerating Liver." *Federation Proceedings* 30(3): 1225-&.
- Gear, C. W. (1969). "A Simple Set of Test Matrices for Eigenvalue Programs." *Mathematics of Computation* 23(105): 119-&.
- Gear, C. W. (1971). "Automatic Integration of Ordinary Differential Equations." *Communications of the Acm* 14(3): 176-&.
- Gear, C. W. (1971). "Difsub for Solution of Ordinary Differential Equations [D2]." *Communications of the Acm* 14(3): 185-&.
- Gear, C. W. (1971). "Simultaneous Numerical Solution of Differential-Algebraic Equations." *Ieee Transactions on Circuit Theory* Ct18(1): 89-&.
- Gear, M. and Whitehea.R (1970). "Chronic Gastritis and Gastric Ulcer." *Journal of Pathology* 101(4): P8-&.
- Gear, M. W. L., S. C. Truelove, et al. (1971). "Gastric Ulcer and Gastritis." *Gut* 12(8): 639-&.
- Gear, R. B., D. R. Heppner, et al. (1971). "Eei-Manufacturers 500/550 Kv Cable Research Project - Pothead No-1 - High Pressure Oil Paper Pipe Type." *Ieee Transactions on Power Apparatus and Systems* Pa90(1): 199-&.
- Gear, R. W. (1971). "Missing Point - Reply." *Journal of the American Dental Association* 83(5): 971-&.
- Gelles, J., B. J. Schnapp, et al. (1988). "Tracking kinesin-driven movements with nanometre-scale precision." *Nature* 331(6155): 450-3.

- Genadiev, T., D. Gaidarov, et al. (2007). "[Laparoscopic pelvic lymph node dissection-- method for prostate cancer staging]." *Khirurgiia (Sofia)*(6): 13-7.
- Georges, P. C., W. J. Miller, et al. (2006). "Matrices with compliance comparable to that of brain tissue select neuronal over glial growth in mixed cortical cultures." *Biophys J* 90(8): 3012-8.
- Giannone, G., B. J. Dubin-Thaler, et al. (2004). "Periodic lamellipodial contractions correlate with rearward actin waves." *Cell* 116(3): 431-43.
- Giannone, G., B. J. Dubin-Thaler, et al. (2007). "Lamellipodial actin mechanically links myosin activity with adhesion-site formation." *Cell* 128(3): 561-75.
- Giannone, G., G. Jiang, et al. (2003). "Talin1 is critical for force-dependent reinforcement of initial integrin-cytoskeleton bonds but not tyrosine kinase activation." *J Cell Biol* 163(2): 409-19.
- Giannone, G. and M. P. Sheetz (2006). "Substrate rigidity and force define form through tyrosine phosphatase and kinase pathways." *Trends Cell Biol* 16(4): 213-23.
- Gibson, E. M., E. S. Henson, et al. (2002). "Epidermal growth factor protects epithelial-derived cells from tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis by inhibiting cytochrome c release." *Cancer Res* 62(2): 488-96.
- Gibson, E. M., E. S. Henson, et al. (2002). "MEK kinase 1 induces mitochondrial permeability transition leading to apoptosis independent of cytochrome c release." *J Biol Chem* 277(12): 10573-80.
- Giddings, V. L., S. M. Kurtz, et al. (2001). "A small punch test technique for characterizing the elastic modulus and fracture behavior of PMMA bone cement used in total joint replacement." *Biomaterials* 22(13): 1875-81.
- Giepmans, B. N., S. R. Adams, et al. (2006). "The fluorescent toolbox for assessing protein location and function." *Science* 312(5771): 217-24.
- Giepmans, B. N., T. J. Deerinck, et al. (2005). "Correlated light and electron microscopic imaging of multiple endogenous proteins using Quantum dots." *Nat Methods* 2(10): 743-9.
- Giffard, C. J., S. Ladha, et al. (1996). "Interaction of nisin with planar lipid bilayers monitored by fluorescence recovery after photobleaching." *J Membr Biol* 151(3): 293-300.

- Gill, S. R., T. A. Schroer, et al. (1991). "Dynactin, a conserved, ubiquitously expressed component of an activator of vesicle motility mediated by cytoplasmic dynein." *J Cell Biol* 115(6): 1639-50.
- Gillespie, D. (1977). "Exact Stochastic Simulation of Coupled Chemical Reactions." *The Journal of Physical Chemistry* 18(25): 2340-2361.
- Gillespie, D. T. (1976). "General Method for Numerically Simulating Stochastic Time Evolution of Coupled Chemical-Reactions." *Journal of Computational Physics* 22(4): 403-434.
- Gillespie, D. T. (1977). "Concerning Validity of Stochastic Approach to Chemical-Kinetics." *Journal of Statistical Physics* 16(3): 311-318.
- Gillespie, D. T. (2007). "Stochastic simulation of chemical kinetics." *Annu Rev Phys Chem* 58: 35-55.
- Gillespie, D. T., S. Lampoudi, et al. (2007). "Effect of reactant size on discrete stochastic chemical kinetics." *J Chem Phys* 126(3): 034302.
- Gilmartin, L., C. A. Tarleton, et al. (2008). "A comparison of inflammatory mediators released by basophils of asthmatic and control subjects in response to high-affinity IgE receptor aggregation." *Int Arch Allergy Immunol* 145(3): 182-92.
- Gilmer, T. M., L. Cable, et al. (2008). "Impact of common epidermal growth factor receptor and HER2 variants on receptor activity and inhibition by lapatinib." *Cancer Res* 68(2): 571-9.
- Girach, A., L. P. Aiello, et al. (2009). "Sustained moderate visual loss as a predictive end point for visual loss in non-proliferative diabetic retinopathy." *Eye* 23(1): 209-14.
- Glogauer, M., P. Arora, et al. (1998). "The role of actin-binding protein 280 in integrin-dependent mechanoprotection." *J Biol Chem* 273(3): 1689-98.
- Goebert, D. A., M. Y. Ng, et al. (1991). "Traumatic spinal cord injury in Hawaii." *Hawaii Med J* 50(2): 44, 47-8, 50.
- Goldman, E. R., A. R. Clapp, et al. (2004). "Multiplexed toxin analysis using four colors of quantum dot fluororeagents." *Anal Chem* 76(3): 684-8.
- Goldman, E. R., H. Mattoussi, et al. (2005). "Fluoroimmunoassays using antibody-conjugated quantum dots." *Methods Mol Biol* 303: 19-34.
- Goldman, E. R., I. L. Medintz, et al. (2006). "Luminescent quantum dots in immunoassays." *Anal Bioanal Chem* 384(3): 560-3.

- Goldman, E. R., I. L. Medintz, et al. (2005). "A hybrid quantum dot-antibody fragment fluorescence resonance energy transfer-based TNT sensor." *J Am Chem Soc* 127(18): 6744-51.
- Goldschmidt-Clermont, P. J. and P. A. Janmey (1991). "Profilin, a weak CAP for actin and RAS." *Cell* 66(3): 419-21.
- Goldstein, B., D. Coombs, et al. (2008). "Kinetic proofreading model." *Adv Exp Med Biol* 640: 82-94.
- Goldstein, B. and M. Dembo (1995). "Approximating the effects of diffusion on reversible reactions at the cell surface: ligand-receptor kinetics." *Biophys J* 68(4): 1222-30.
- Goldstein, B., J. R. Faeder, et al. (2004). "Mathematical and computational models of immune-receptor signalling." *Nat Rev Immunol* 4(6): 445-56.
- Goldstein, B., J. R. Faeder, et al. (2002). "Modeling the early signaling events mediated by FcepsilonRI." *Mol Immunol* 38(16-18): 1213-9.
- Golgher, D., F. Korangy, et al. (2001). "An immunodominant MHC class II-restricted tumor antigen is conformation dependent and binds to the endoplasmic reticulum chaperone, calreticulin." *J Immunol* 167(1): 147-55.
- Gonzalez, E., A. Nagiel, et al. (2004). "Small interfering RNA-mediated down-regulation of caveolin-1 differentially modulates signaling pathways in endothelial cells." *J Biol Chem* 279(39): 40659-69.
- Goraya, T. A., N. Masada, et al. (2008). "Kinetic properties of Ca²⁺/calmodulin-dependent phosphodiesterase isoforms dictate intracellular cAMP dynamics in response to elevation of cytosolic Ca²⁺." *Cell Signal* 20(2): 359-74.
- Gorfe, A. A., M. Hanzal-Bayer, et al. (2007). "Structure and dynamics of the full-length lipid-modified H-Ras protein in a 1,2-dimyristoylglycero-3-phosphocholine bilayer." *J Med Chem* 50(4): 674-84.
- Gou, J. P., T. Gotow, et al. (1998). "Regulation of neurofilament interactions in vitro by natural and synthetic polypeptides sharing Lys-Ser-Pro sequences with the heavy neurofilament subunit NF-H: neurofilament crossbridging by antiparallel sidearm overlapping." *Med Biol Eng Comput* 36(3): 371-87.
- Gough, M. J., A. A. Melcher, et al. (2001). "Macrophages orchestrate the immune response to tumor cell death." *Cancer Res* 61(19): 7240-7.

Grady, W. M., L. L. Myeroff, et al. (1999). "Mutational inactivation of transforming growth factor beta receptor type II in microsatellite stable colon cancers." *Cancer Res* 59(2): 320-4.

Graham, T. E., J. R. Pfeiffer, et al. (1998). "MEK and ERK activation in ras-disabled RBL-2H3 mast cells and novel roles for geranylgeranylated and farnesylated proteins in Fc epsilonRI-mediated signaling." *J Immunol* 161(12): 6733-44.

Granoff, D. M., A. Chacko, et al. (1989). "Immunogenicity of Haemophilus influenzae type b polysaccharide-outer membrane protein conjugate vaccine in patients who acquired Haemophilus disease despite previous vaccination with type b polysaccharide vaccine." *J Pediatr* 114(6): 925-33.

Granoff, D. M., K. Sheetz, et al. (1989). "Host and bacterial factors associated with Haemophilus influenzae type b disease in Minnesota children vaccinated with type b polysaccharide vaccine." *J Infect Dis* 159(5): 908-16.

Granoff, D. M., K. E. Sheetz, et al. (1988). "Further immunologic evaluation of children who develop haemophilus disease despite previous vaccination with type b polysaccharide vaccine." *Monogr Allergy* 23: 256-68.

Grau, S., I. Krauss, et al. (2008). "Hip abductor weakness is not the cause for iliotibial band syndrome." *Int J Sports Med* 29(7): 579-83.

Grau, S., C. Maiwald, et al. (2008). "The influence of matching populations on kinematic and kinetic variables in runners with iliotibial band syndrome." *Res Q Exerc Sport* 79(4): 450-7.

Grau, S., C. Maiwald, et al. (2008). "What are causes and treatment strategies for patellar-tendinopathy in female runners?" *J Biomech* 41(9): 2042-6.

Green, C. R., K. O. Anderson, et al. (2003). "The unequal burden of pain: confronting racial and ethnic disparities in pain." *Pain Med* 4(3): 277-94.

Greene, E. L. and M. S. Paller (1994). "Calcium and free radicals in hypoxia/reoxygenation injury of renal epithelial cells." *Am J Physiol* 266(1 Pt 2): F13-20.

Gu, B. J., W. Zhang, et al. (2001). "A Glu-496 to Ala polymorphism leads to loss of function of the human P2X7 receptor." *J Biol Chem* 276(14): 11135-42.

Gu, B. J., W. Y. Zhang, et al. (2000). "Expression of P2X(7) purinoceptors on human lymphocytes and monocytes: evidence for nonfunctional P2X(7) receptors." *Am J Physiol Cell Physiol* 279(4): C1189-97.

Guess, H. A. and J. H. Gillespie (1977). "Diffusion Approximations to Linear Stochastic Difference Equations with Stationary Coefficients." *Journal of Applied Probability* 14(1): 58-74.

Guo, X. N., A. Rajput, et al. (2007). "Mutant PIK3CA-bearing colon cancer cells display increased metastasis in an orthotopic model." *Cancer Res* 67(12): 5851-8.

Gupton, S. L., K. L. Anderson, et al. (2005). "Cell migration without a lamellipodium: translation of actin dynamics into cell movement mediated by tropomyosin." *J Cell Biol* 168(4): 619-31.

Gutierrez, L., A. I. Magee, et al. (1989). "Post-translational processing of p21ras is two-step and involves carboxyl-methylation and carboxy-terminal proteolysis." *EMBO J* 8(4): 1093-8.

Haaland, D. M., H. D. Jones, et al. (2009). "Hyperspectral confocal fluorescence imaging: exploring alternative multivariate curve resolution approaches." *Appl Spectrosc* 63(3): 271-9.

Haggbloom, M. M., L. J. Nohynek, et al. (1994). "Transfer of polychlorophenol-degrading *Rhodococcus chlorophenolicus* (Apajalahti et al. 1986) to the genus *Mycobacterium* as *Mycobacterium chlorophenolicum* comb. nov." *Int J Syst Bacteriol* 44(3): 485-93.

Hahn, F., C. Maiwald, et al. (2008). "Changes in plantar pressure distribution after Achilles tendon augmentation with flexor hallucis longus transfer." *Clin Biomech (Bristol, Avon)* 23(1): 109-16.

Hahn, F., P. Meyer, et al. (2008). "Treatment of chronic achilles tendinopathy and ruptures with flexor hallucis tendon transfer: clinical outcome and MRI findings." *Foot Ankle Int* 29(8): 794-802.

Hahn, J., B. Maier, et al. (2005). "Transformation proteins and DNA uptake localize to the cell poles in *Bacillus subtilis*." *Cell* 122(1): 59-71.

Haig, A. J., M. E. Geisser, et al. (2003). "The effect of order of testing in functional performance in persons with and without chronic back pain." *J Occup Rehabil* 13(2): 115-23.

Halasz, J., A. Holczbauer, et al. (2006). "Claudin-1 and claudin-2 differentiate fetal and embryonal components in human hepatoblastoma." *Hum Pathol* 37(5): 555-61.

Hall, A., C. Cales, et al. (1988). "Analysis of mammalian ras effector function." *Cold Spring Harb Symp Quant Biol* 53 Pt 2: 855-62.

Hall, A. L., B. S. Wilson, et al. (1997). "Relationship of ligand-receptor dynamics to actin polymerization in RBL-2H3 cells transfected with the human formyl peptide receptor." *J Leukoc Biol* 62(4): 535-46.

Hamm-Alvarez, S. F., B. E. Alayof, et al. (1994). "Coordinate depression of bradykinin receptor recycling and microtubule-dependent transport by taxol." *Proc Natl Acad Sci U S A* 91(16): 7812-6.

Hamm-Alvarez, S. F., P. Y. Kim, et al. (1993). "Regulation of vesicle transport in CV-1 cells and extracts." *J Cell Sci* 106 (Pt 3): 955-66.

Hamm-Alvarez, S. F. and M. P. Sheetz (1998). "Microtubule-dependent vesicle transport: modulation of channel and transporter activity in liver and kidney." *Physiol Rev* 78(4): 1109-29.

Han, G. W., S. Sri Krishna, et al. (2006). "Crystal structure of the ApbE protein (TM1553) from *Thermotoga maritima* at 1.58 Å resolution." *Proteins* 64(4): 1083-90.

Hanahan, D. and R. A. Weinberg (2000). "The hallmarks of cancer." *Cell* 100(1): 57-70.

Hancock, J. F. (1993). "Anti-Ras drugs come of age." *Curr Biol* 3(11): 770-2.

Hancock, J. F. (1995). "Prenylation and palmitoylation analysis." *Methods Enzymol* 255: 237-45.

Hancock, J. F. (1995). "Reticulocyte lysate assay for in vitro translation and posttranslational modification of Ras proteins." *Methods Enzymol* 255: 60-5.

Hancock, J. F. (2003). "Ras proteins: different signals from different locations." *Nat Rev Mol Cell Biol* 4(5): 373-84.

Hancock, J. F. (2004). "GPI-anchor synthesis: Ras takes charge." *Dev Cell* 6(6): 743-5.

Hancock, J. F. (2007). "PA promoted to manager." *Nat Cell Biol* 9(6): 615-7.

Hancock, J. F., K. Cadwallader, et al. (1991). "Methylation and proteolysis are essential for efficient membrane binding of prenylated p21K-ras(B)." *EMBO J* 10(3): 641-6.

Hancock, J. F., K. Cadwallader, et al. (1991). "A CAAX or a CAAL motif and a second signal are sufficient for plasma membrane targeting of ras proteins." *EMBO J* 10(13): 4033-9.

Hancock, J. F., A. I. Magee, et al. (1989). "All ras proteins are polyisoprenylated but only some are palmitoylated." *Cell* 57(7): 1167-77.

- Hancock, J. F., C. J. Marshall, et al. (1988). "Mutant but not normal p21 ras elevates inositol phospholipid breakdown in two different cell systems." *Oncogene* 3(2): 187-93.
- Hancock, J. F. and R. G. Parton (2005). "Ras plasma membrane signalling platforms." *Biochem J* 389(Pt 1): 1-11.
- Hancock, J. F., H. Paterson, et al. (1990). "A polybasic domain or palmitoylation is required in addition to the CAAX motif to localize p21ras to the plasma membrane." *Cell* 63(1): 133-9.
- Hancock, J. F. and I. A. Prior (2005). "Electron microscopic imaging of Ras signaling domains." *Methods* 37(2): 165-72.
- Hancock, R. L. (1985). "Theoretical mechanisms for synthesis of carcinogen-induced embryonic proteins: XIV. Mutational and non-mutational mechanisms as subsets of a more general mechanism. Part C. A defined cancer mutation." *Med Hypotheses* 18(3): 199-206.
- Hannan, L. A., M. P. Lisanti, et al. (1993). "Correctly sorted molecules of a GPI-anchored protein are clustered and immobile when they arrive at the apical surface of MDCK cells." *J Cell Biol* 120(2): 353-8.
- Hannocks, M. J., L. Oliver, et al. (1992). "Regulation of proteolytic activity in human bone marrow stromal cells by basic fibroblast growth factor, interleukin-1, and transforming growth factor beta." *Blood* 79(5): 1178-84.
- Hansen, M., I. A. Prior, et al. (2003). "C-terminal sequences in R-Ras are involved in integrin regulation and in plasma membrane microdomain distribution." *Biochem Biophys Res Commun* 311(4): 829-38.
- Hansson, G. C., K. Simons, et al. (1986). "Two strains of the Madin-Darby canine kidney (MDCK) cell line have distinct glycosphingolipid compositions." *EMBO J* 5(3): 483-9.
- Harding, A., N. Giles, et al. (2003). "Mechanism of mitosis-specific activation of MEK1." *J Biol Chem* 278(19): 16747-54.
- Harding, A. and J. F. Hancock (2008). "Ras nanoclusters: combining digital and analog signaling." *Cell Cycle* 7(2): 127-34.
- Harding, A., V. Hsu, et al. (2003). "Identification of residues and domains of Raf important for function in vivo and in vitro." *J Biol Chem* 278(46): 45519-27.
- Harding, A., T. Tian, et al. (2005). "Subcellular localization determines MAP kinase signal output." *Curr Biol* 15(9): 869-73.

- Harley, B. A., H. D. Kim, et al. (2008). "Microarchitecture of three-dimensional scaffolds influences cell migration behavior via junction interactions." *Biophys J* 95(8): 4013-24.
- Hartley, A. H., T. A. Kahn, et al. (1994). "Proceedings of the Society for Pediatric Dermatology meeting. Hilton Head, South Carolina, June 22-25, 1994." *Pediatr Dermatol* 11(4): 365-70.
- Hartsough, M. T., D. K. Morrison, et al. (2002). "Nm23-H1 metastasis suppressor phosphorylation of kinase suppressor of Ras via a histidine protein kinase pathway." *J Biol Chem* 277(35): 32389-99.
- Hathcock, J. J., E. Rusinova, et al. (2006). "Phospholipid surfaces regulate the delivery of substrate to tissue factor:VIIa and the removal of product." *Blood Cells Mol Dis* 36(2): 194-8.
- Haugh, J. M. (2002). "Localization of receptor-mediated signal transduction pathways: the inside story." *Mol Interv* 2(5): 292-307.
- Haugh, J. M. (2002). "A unified model for signal transduction reactions in cellular membranes." *Biophys J* 82(2): 591-604.
- Haugh, J. M., A. C. Huang, et al. (1999). "Internalized epidermal growth factor receptors participate in the activation of p21(ras) in fibroblasts." *J Biol Chem* 274(48): 34350-60.
- Haugh, J. M. and D. A. Lauffenburger (1997). "Physical modulation of intracellular signaling processes by locational regulation." *Biophys J* 72(5): 2014-31.
- Haugh, J. M. and D. A. Lauffenburger (1998). "Analysis of receptor internalization as a mechanism for modulating signal transduction." *J Theor Biol* 195(2): 187-218.
- Haugh, J. M., K. Schooler, et al. (1999). "Effect of epidermal growth factor receptor internalization on regulation of the phospholipase C-gamma1 signaling pathway." *J Biol Chem* 274(13): 8958-65.
- Haugh, J. M., A. Wells, et al. (2000). "Mathematical modeling of epidermal growth factor receptor signaling through the phospholipase C pathway: mechanistic insights and predictions for molecular interventions." *Biotechnol Bioeng* 70(2): 225-38.
- Haylock, D. N., L. B. To, et al. (1992). "Ex vivo expansion and maturation of peripheral blood CD34+ cells into the myeloid lineage." *Blood* 80(6): 1405-12.
- Hazzalin, C. A. and L. C. Mahadevan (2002). "MAPK-regulated transcription: a continuously variable gene switch?" *Nat Rev Mol Cell Biol* 3(1): 30-40.

Hegener, O., L. Prenner, et al. (2004). "Dynamics of beta2-adrenergic receptor-ligand complexes on living cells." *Biochemistry* 43(20): 6190-9.

Heinrich, R., B. G. Neel, et al. (2002). "Mathematical models of protein kinase signal transduction." *Mol Cell* 9(5): 957-70.

Hendriks, B. S., L. K. Opresko, et al. (2003). "Quantitative analysis of HER2-mediated effects on HER2 and epidermal growth factor receptor endocytosis: distribution of homo- and heterodimers depends on relative HER2 levels." *J Biol Chem* 278(26): 23343-51.

Heneberg, P. and P. Draber (2005). "Regulation of cys-based protein tyrosine phosphatases via reactive oxygen and nitrogen species in mast cells and basophils." *Curr Med Chem* 12(16): 1859-71.

Heneberg, P., P. Lebduska, et al. (2006). "Topography of plasma membrane microdomains and its consequences for mast cell signaling." *Eur J Immunol* 36(10): 2795-806.

Henis, Y. I., J. F. Hancock, et al. (2009). "Ras acylation, compartmentalization and signaling nanoclusters (Review)." *Mol Membr Biol* 26(1): 80-92.

Henson, E. S., E. M. Gibson, et al. (2003). "Increased expression of Mcl-1 is responsible for the blockage of TRAIL-induced apoptosis mediated by EGF/ErbB1 signaling pathway." *J Cell Biochem* 89(6): 1177-92.

Henson, E. S. and S. B. Gibson (2006). "Surviving cell death through epidermal growth factor (EGF) signal transduction pathways: implications for cancer therapy." *Cell Signal* 18(12): 2089-97.

Henson, E. S., X. Hu, et al. (2006). "Herceptin sensitizes ErbB2-overexpressing cells to apoptosis by reducing antiapoptotic Mcl-1 expression." *Clin Cancer Res* 12(3 Pt 1): 845-53.

Henson, E. S., J. B. Johnston, et al. (2008). "The role of TRAIL death receptors in the treatment of hematological malignancies." *Leuk Lymphoma* 49(1): 27-35.

Henson, J. H., T. M. Svitkina, et al. (1999). "Two components of actin-based retrograde flow in sea urchin coelomocytes." *Mol Biol Cell* 10(12): 4075-90.

Hernandez-Hansen, V., J. D. Bard, et al. (2005). "Increased expression of genes linked to FcepsilonRI Signaling and to cytokine and chemokine production in Lyn-deficient mast cells." *J Immunol* 175(12): 7880-8.

Hernandez-Hansen, V., G. A. Mackay, et al. (2004). "The Src kinase Lyn is a negative regulator of mast cell proliferation." *J Leukoc Biol* 75(1): 143-51.

Hernandez-Hansen, V., A. J. Smith, et al. (2004). "Dysregulated FcepsilonRI signaling and altered Fyn and SHIP activities in Lyn-deficient mast cells." *J Immunol* 173(1): 100-12.

Hernandez-Sanchez, B. A., T. J. Boyle, et al. (2006). "Synthesizing biofunctionalized nanoparticles to image cell signaling pathways." *IEEE Trans Nanobioscience* 5(4): 222-30.

Hildebrand, D. C., N. J. Palleroni, et al. (1994). "*Pseudomonas flavescens* sp. nov., isolated from walnut blight cankers." *Int J Syst Bacteriol* 44(3): 410-5.

Hindmarsh, A. (1980). "LSODE and LSODI, Two New Initial Value Ordinary Differential Equation Solvers." *ACM SIGNUM Newsletter* 15: 10-11.

Hinow, P., S. E. Wang, et al. (2007). "relocating job wise? A mathematical model separates quantitatively the cytostatic and cytotoxic effects of a HER2 tyrosine kinase inhibitor." *Theor Biol Med Model* 4: 14.

Hiroyama, T., K. Mihařada, et al. (2008). "Establishment of mouse embryonic stem cell-derived erythroid progenitor cell lines able to produce functional red blood cells." *PLoS ONE* 3(2): e1544.

Hiroyama, T., K. Sudo, et al. (2008). "Human umbilical cord-derived cells can often serve as feeder cells to maintain primate embryonic stem cells in a state capable of producing hematopoietic cells." *Cell Biol Int* 32(1): 1-7.

Hlavacek, W. S., J. R. Faeder, et al. (2003). "The complexity of complexes in signal transduction." *Biotechnol Bioeng* 84(7): 783-94.

Hlavacek, W. S., A. Redondo, et al. (2002). "Kinetic proofreading in receptor-mediated transduction of cellular signals: receptor aggregation, partially activated receptors, and cytosolic messengers." *Bull Math Biol* 64(5): 887-911.

Hochmuth, F. M., J. Y. Shao, et al. (1996). "Deformation and flow of membrane into tethers extracted from neuronal growth cones." *Biophys J* 70(1): 358-69.

Hochmuth, R. M. and D. A. Berk (1984). "Analytical solutions for shear deformation and flow of red cell membrane." *J Biomech Eng* 106(1): 2-9.

Hochmuth, R. M., D. A. Berk, et al. (1983). "Viscous flow of cytoplasm and red cell membrane: membrane recovery and tether contraction." *Ann N Y Acad Sci* 416: 207-24.

Hoek, J. B. and B. N. Kholodenko (1998). "The intracellular signaling network as a target for ethanol." *Alcohol Clin Exp Res* 22(5 Suppl): 224S-230S.

- Hofman, E. G., M. O. Ruonala, et al. (2008). "EGF induces coalescence of different lipid rafts." *J Cell Sci* 121(Pt 15): 2519-28.
- Hogue-Angeletti, R. A. and P. B. Sheetz (1978). "A soluble lipid-protein complex from bovine adrenal medulla chromaffin granules." *J Biol Chem* 253(16): 5613-6.
- Holbro, T., G. Civenni, et al. (2003). "The ErbB receptors and their role in cancer progression." *Exp Cell Res* 284(1): 99-110.
- Holowka, D., D. Sil, et al. (2007). "Insights into immunoglobulin E receptor signaling from structurally defined ligands." *Immunol Rev* 217: 269-79.
- Hong, W. K., G. M. O'Donoghue, et al. (1985). "Sequential response patterns to chemotherapy and radiotherapy in head and neck cancer: potential impact of treatment in advanced laryngeal cancer." *Prog Clin Biol Res* 201: 191-7.
- Honn, K. V., I. M. Grossi, et al. (1989). "Lipoxygenase regulation of membrane expression of tumor cell glycoproteins and subsequent metastasis." *Adv Prostaglandin Thromboxane Leukot Res* 19: 439-43.
- Hornberg, J. J., B. Binder, et al. (2005). "Control of MAPK signalling: from complexity to what really matters." *Oncogene* 24(36): 5533-42.
- Horvat, R. D., S. Nelson, et al. (1999). "Intrinsically fluorescent luteinizing hormone receptor demonstrates hormone-driven aggregation." *Biochem Biophys Res Commun* 255(2): 382-5.
- Hsia, A. Y., E. Masliah, et al. (1999). "Plaque-independent disruption of neural circuits in Alzheimer's disease mouse models." *Proc Natl Acad Sci U S A* 96(6): 3228-33.
- Hsieh, M., S. Yang, et al. (2008). "Stochastic Simulations of ErbB Receptor Dimerization: Predicted contributions of membrane topography and receptor conformation." Manuscript submitted for publication.
- Hsieh, M. Y., S. Yang, et al. (2008). "Stochastic simulations of ErbB homo and heterodimerisation: potential impacts of receptor conformational state and spatial segregation." *IET Syst Biol* 2(5): 256-72.
- Hsu, F. C., J. Sun, et al. (2009). "A novel prostate cancer susceptibility locus at 19q13." *Cancer Res* 69(7): 2720-3.
- Hu, K., L. Ji, et al. (2007). "Differential transmission of actin motion within focal adhesions." *Science* 315(5808): 111-5.

- Hu, K., D. Mochly-Rosen, et al. (2000). "Evidence for functional role of epsilonPKC isozyme in the regulation of cardiac Ca(2+) channels." *Am J Physiol Heart Circ Physiol* 279(6): H2658-64.
- Huang, C. Y. and J. E. Ferrell, Jr. (1996). "Ultrasensitivity in the mitogen-activated protein kinase cascade." *Proc Natl Acad Sci U S A* 93(19): 10078-83.
- Huang, D. C., C. J. Marshall, et al. (1993). "Plasma membrane-targeted ras GTPase-activating protein is a potent suppressor of p21ras function." *Mol Cell Biol* 13(4): 2420-31.
- Huang, H., J. Sylvan, et al. (2005). "Cell stiffness and receptors: evidence for cytoskeletal subnetworks." *Am J Physiol Cell Physiol* 288(1): C72-80.
- Hubbell, H. R., P. C. Sheetz, et al. (1991). "Heterogeneous nuclear RNA from hairy cell leukemia patients activates 2',5'-oligoadenylate synthetase." *Anticancer Res* 11(5): 1927-32.
- Hubner, S., A. D. Couvillon, et al. (1998). "Enhancement of phosphoinositide 3-kinase (PI 3-kinase) activity by membrane curvature and inositol-phospholipid-binding peptides." *Eur J Biochem* 258(2): 846-53.
- Huitfeldt, H. S., A. Heyden, et al. (1994). "Gene activation studied by immunological methods." *Environ Health Perspect* 102 Suppl 6: 205-7.
- Huitfeldt, H. S., E. Skarpen, et al. (1996). "Differential distribution of Met and epidermal growth factor receptor in normal and carcinogen-treated rat liver." *J Histochem Cytochem* 44(3): 227-33.
- Hynes, N. E. and H. A. Lane (2005). "ERBB receptors and cancer: the complexity of targeted inhibitors." *Nat Rev Cancer* 5(5): 341-54.
- Iarovaia, O. V., A. Bystritskiy, et al. (2004). "Visualization of individual DNA loops and a map of loop domains in the human dystrophin gene." *Nucleic Acids Res* 32(7): 2079-86.
- Ichinose, J., M. Morimatsu, et al. (2006). "Covalent immobilization of epidermal growth factor molecules for single-molecule imaging analysis of intracellular signaling." *Biomaterials* 27(18): 3343-50.
- Iijima, Y., Y. Nakamura, et al. (2008). "Metabolite annotations based on the integration of mass spectral information." *Plant J* 54(5): 949-62.
- Ike, H., A. Kosugi, et al. (2003). "Mechanism of Lck recruitment to the T-cell receptor cluster as studied by single-molecule-fluorescence video imaging." *Chemphyschem* 4(6): 620-6.

Ikeda, S., M. Ushio-Fukai, et al. (2005). "Novel role of ARF6 in vascular endothelial growth factor-induced signaling and angiogenesis." *Circ Res* 96(4): 467-75.

Ikonen, E., J. B. de Almeida, et al. (1997). "Myosin II is associated with Golgi membranes: identification of p200 as nonmuscle myosin II on Golgi-derived vesicles." *J Cell Sci* 110 (Pt 18): 2155-64.

Ikonen, E., K. Fiedler, et al. (1995). "Prohibitin, an antiproliferative protein, is localized to mitochondria." *FEBS Lett* 358(3): 273-7.

Ikonen, E., R. G. Parton, et al. (1993). "Transcytosis of the polymeric immunoglobulin receptor in cultured hippocampal neurons." *Curr Biol* 3(10): 635-44.

Ikonen, E., R. G. Parton, et al. (1996). "Analysis of the role of p200-containing vesicles in post-Golgi traffic." *Mol Biol Cell* 7(6): 961-74.

Ikonen, E. and K. Simons (1998). "Protein and lipid sorting from the trans-Golgi network to the plasma membrane in polarized cells." *Semin Cell Dev Biol* 9(5): 503-9.

Ikonen, E., M. Tagaya, et al. (1995). "Different requirements for NSF, SNAP, and Rab proteins in apical and basolateral transport in MDCK cells." *Cell* 81(4): 571-80.

Ilowite, J. S., M. I. Baskin, et al. (1991). "Delivered dose and regional distribution of aerosolized pentamidine using different delivery systems." *Chest* 99(5): 1139-44.

Ilowite, J. S., W. D. Bennett, et al. (1989). "Permeability of the bronchial mucosa to ^{99m}Tc-DTPA in asthma." *Am Rev Respir Dis* 139(5): 1139-43.

Im, M. Y., S. H. Lee, et al. (2008). "Scaling behavior of the first arrival time of a random-walking magnetic domain." *Phys Rev Lett* 100(16): 167204.

Inder, K., A. Harding, et al. (2008). "Activation of the MAPK module from different spatial locations generates distinct system outputs." *Mol Biol Cell* 19(11): 4776-84.

Irie, K., T. Tadauchi, et al. (2002). "The Khd1 protein, which has three KH RNA-binding motifs, is required for proper localization of ASH1 mRNA in yeast." *EMBO J* 21(5): 1158-67.

Ito-Kuwa, S., K. Nakamura, et al. (2008). "Diversity of laccase among *Cryptococcus neoformans* serotypes." *Microbiol Immunol* 52(10): 492-8.

Ivanova, P. T., S. B. Milne, et al. (2004). "LIPID arrays: new tools in the understanding of membrane dynamics and lipid signaling." *Mol Interv* 4(2): 86-96.
Jacobson, K. and C. Dietrich (1999). "Looking at lipid rafts?" *Trends Cell Biol* 9(3): 87-91.

- Jacobson, K., O. G. Mouritsen, et al. (2007). "Lipid rafts: at a crossroad between cell biology and physics." *Nat Cell Biol* 9(1): 7-14.
- Jain, K. A., U. M. Hamper, et al. (1988). "Comparison of transvaginal and transabdominal sonography in the detection of early pregnancy and its complications." *AJR Am J Roentgenol* 151(6): 1139-43.
- Jambhekar, A., K. McDermott, et al. (2005). "Unbiased selection of localization elements reveals cis-acting determinants of mRNA bud localization in *Saccharomyces cerevisiae*." *Proc Natl Acad Sci U S A* 102(50): 18005-10.
- Janes, K. A., J. G. Albeck, et al. (2005). "A systems model of signaling identifies a molecular basis set for cytokine-induced apoptosis." *Science* 310(5754): 1646-53.
- Janes, K. A., J. G. Albeck, et al. (2003). "A high-throughput quantitative multiplex kinase assay for monitoring information flow in signaling networks: application to sepsis-apoptosis." *Mol Cell Proteomics* 2(7): 463-73.
- Janmey, P. A. (1991). "Mechanical properties of cytoskeletal polymers." *Curr Opin Cell Biol* 3(1): 4-11.
- Janmey, P. A. (1995). "Protein regulation by phosphatidylinositol lipids." *Chem Biol* 2(2): 61-5.
- Janmey, P. A. (1998). "The cytoskeleton and cell signaling: component localization and mechanical coupling." *Physiol Rev* 78(3): 763-81.
- Janmey, P. A., J. Kas, et al. (1998). "Cytoskeletal networks and filament bundles: regulation by proteins and polycations." *Biol Bull* 194(3): 334-5; discussion 335-6.
- Janmey, P. A. and P. K. Kinnunen (2006). "Biophysical properties of lipids and dynamic membranes." *Trends Cell Biol* 16(10): 538-46.
- Janmey, P. A. and U. Lindberg (2004). "Cytoskeletal regulation: rich in lipids." *Nat Rev Mol Cell Biol* 5(8): 658-66.
- Janmey, P. A., M. E. McCormick, et al. (2007). "Negative normal stress in semiflexible biopolymer gels." *Nat Mater* 6(1): 48-51.
- Janmey, P. A., J. V. Shah, et al. (1998). "Viscoelasticity of intermediate filament networks." *Subcell Biochem* 31: 381-97.
- Janmey, P. A., J. V. Shah, et al. (2001). "Actin filament networks." *Results Probl Cell Differ* 32: 181-99.

- Janmey, P. A., T. P. Stossel, et al. (1998). "Deconstructing gelsolin: identifying sites that mimic or alter binding to actin and phosphoinositides." *Chem Biol* 5(4): R81-5.
- Janmey, P. A., W. Xian, et al. (1999). "Controlling cytoskeleton structure by phosphoinositide-protein interactions: phosphoinositide binding protein domains and effects of lipid packing." *Chem Phys Lipids* 101(1): 93-107.
- Janssen, R. A., P. N. Kim, et al. (2003). "Overexpression of kinase suppressor of Ras upregulates the high-molecular-weight tropomyosin isoforms in ras-transformed NIH 3T3 fibroblasts." *Mol Cell Biol* 23(5): 1786-97.
- Jaumot, M. and J. F. Hancock (2001). "Protein phosphatases 1 and 2A promote Raf-1 activation by regulating 14-3-3 interactions." *Oncogene* 20(30): 3949-58.
- Jaumot, M., J. Yan, et al. (2002). "The linker domain of the Ha-Ras hypervariable region regulates interactions with exchange factors, Raf-1 and phosphoinositide 3-kinase." *J Biol Chem* 277(1): 272-8.
- Jayasuriya, A. C., S. Ghosh, et al. (2003). "A study of piezoelectric and mechanical anisotropies of the human cornea." *Biosens Bioelectron* 18(4): 381-7.
- Jenniskens, P., M. H. Shaddad, et al. (2009). "The impact and recovery of asteroid 2008 TC(3)." *Nature* 458(7237): 485-8.
- Jensen, P. R., J. L. Snoep, et al. (1995). "Molecular biology for flux control." *Biochem Soc Trans* 23(2): 367-70.
- Ji, S., C. Song, et al. (2003). "Interference of magnetic and anisotropic tensor susceptibility reflections in resonant X-ray scattering of GdB4." *Phys Rev Lett* 91(25): 257205.
- Jiang, G., G. Giannone, et al. (2003). "Two-piconewton slip bond between fibronectin and the cytoskeleton depends on talin." *Nature* 424(6946): 334-7.
- Jiang, G., A. H. Huang, et al. (2006). "Rigidity sensing at the leading edge through alphavbeta3 integrins and RPTPalph." *Biophys J* 90(5): 1804-9.
- Jiang, M. Y. and M. P. Sheetz (1994). "Mechanics of myosin motor: force and step size." *Bioessays* 16(8): 531-2.
- Jiang, M. Y. and M. P. Sheetz (1995). "Cargo-activated ATPase activity of kinesin." *Biophys J* 68(4 Suppl): 283S-284S; discussion 285S.
- Jiang, X. and A. Sorkin (2002). "Coordinated traffic of Grb2 and Ras during epidermal growth factor receptor endocytosis visualized in living cells." *Mol Biol Cell* 13(5): 1522-35.

Jin, X., P. H. Mathers, et al. (2001). "Vertical bias in dendritic trees of non-pyramidal neocortical neurons expressing GAD67-GFP in vitro." *Cereb Cortex* 11(7): 666-78.

Jirakulsomchok, D., J. H. Yu, et al. (1983). "Effects of adriamycin on calcium concentration and morphology of mouse salivary glands." *J Oral Pathol* 12(6): 491-501.

Johnson, C. M., G. R. Chichili, et al. (2008). "Compartmentalization of phosphatidylinositol 4,5-bisphosphate signaling evidenced using targeted phosphatases." *J Biol Chem* 283(44): 29920-8.

Johnson, K. A., B. E. Powers, et al. (1989). "Splénomegaly in dogs. Predictors of neoplasia and survival after splenectomy." *J Vet Intern Med* 3(3): 160-6.

Johnson, M. B., E. W. Heineke, et al. (1993). "MDL 29311. Antioxidant with marked lipid- and glucose-lowering activity in diabetic rats and mice." *Diabetes* 42(8): 1179-86.

Joo, C., H. Balci, et al. (2008). "Advances in single-molecule fluorescence methods for molecular biology." *Annu Rev Biochem* 77: 51-76.

Joslin, E. J., L. K. Opresko, et al. (2007). "EGF-receptor-mediated mammary epithelial cell migration is driven by sustained ERK signaling from autocrine stimulation." *J Cell Sci* 120(Pt 20): 3688-99.

Jury, E. C., P. S. Kabouridis, et al. (2004). "Altered lipid raft-associated signaling and ganglioside expression in T lymphocytes from patients with systemic lupus erythematosus." *J Clin Invest* 113(8): 1176-87.

Kabouridis, P. S. and E. C. Jury (2008). "Lipid rafts and T-lymphocyte function: implications for autoimmunity." *FEBS Lett* 582(27): 3711-8.

Kaise, H. and N. Kohno (2006). "[Bisphosphonates for bone metastasis of malignant tumor]." *Clin Calcium* 16(4): 655- 64.

Kalaidzidis, Y. (2007). "Intracellular objects tracking." *Eur J Cell Biol* 86(9): 569-78.

Kalaidzidis, Y. (2009). "Multiple objects tracking in fluorescence microscopy." *J Math Biol* 58(1-2): 57-80.

Kanaani, J., G. Patterson, et al. (2008). "A palmitoylation cycle dynamically regulates partitioning of the GABA-synthesizing enzyme GAD65 between ER-Golgi and post-Golgi membranes." *J Cell Sci* 121(Pt 4): 437-49.

- Kang, J. S., G. Kim, et al. (2005). "Spatial chemical inhomogeneity and local electronic structure of Mn-doped Ge ferromagnetic semiconductors." *Phys Rev Lett* 94(14): 147202.
- Kania, D. R., H. Kornblum, et al. (1992). "Characterization of an x-ray-flux source for the production of high-energy-density plasmas." *Phys Rev A* 46(12): 7853-7868.
- Karashchuk, G. N., D. L. Kakuev, et al. (1999). "[Biochemical characteristics of bovine retina nucleoside diphosphate kinase]." *Bioorg Khim* 25(7): 513-9.
- Karni-Schmidt, O., A. Friedler, et al. (2007). "Energy-dependent nucleolar localization of p53 in vitro requires two discrete regions within the p53 carboxyl terminus." *Oncogene* 26(26): 3878-91.
- Karolyi, G., J. Skapinyecz, et al. (1972). "Effect of ^{60}Co -gamma-rays on the infrared spectra of DNA." *Acta Biochim Biophys Acad Sci Hung* 7(2): 179-85.
- Katsoulakis, M. A., A. J. Majda, et al. (2005). "Multiscale Couplings in Prototype Hybrid Deterministic/Stochastic Systems: Part I, Deterministic Closures." *Comm. Math. Sci* 2(2): 255-294.
- Keller, P., D. Toomre, et al. (2001). "Multicolour imaging of post-Golgi sorting and trafficking in live cells." *Nat Cell Biol* 3(2): 140-9.
- Kepley, C. L., J. C. Cambier, et al. (2000). "Negative regulation of FcepsilonRI signaling by FcgammaRII costimulation in human blood basophils." *J Allergy Clin Immunol* 106(2): 337-48.
- Kepley, C. L., J. R. Pfeiffer, et al. (1998). "The identification and characterization of umbilical cord blood-derived human basophils." *J Leukoc Biol* 64(4): 474-83.
- Kepley, C. L., B. S. Wilson, et al. (1998). "Identification of the Fc epsilonRI-activated tyrosine kinases Lyn, Syk, and Zap-70 in human basophils." *J Allergy Clin Immunol* 102(2): 304-15.
- Kepley, C. L., L. Youssef, et al. (1999). "Syk deficiency in nonreleaser basophils." *J Allergy Clin Immunol* 104(2 Pt 1): 279-84.
- Kepley, C. L., L. Youssef, et al. (2000). "Multiple defects in Fc epsilon RI signaling in Syk-deficient nonreleaser basophils and IL-3-induced recovery of Syk expression and secretion." *J Immunol* 165(10): 5913-20.
- Khaidukov, S. V., I. V. Kholodenko, et al. (2003). "[Phenotype characteristic of the ionomycin-resistant CD4⁺ subset of the peripheral blood T-lymphocytes]." *Tsitologiya* 45(3): 249-54.

Khan, S. and M. P. Sheetz (1997). "Force effects on biochemical kinetics." *Annu Rev Biochem* 66: 785-805.

Khankan, A. A., T. Murakami, et al. (2008). "Hepatocellular carcinoma treated with radio frequency ablation: an early evaluation with magnetic resonance imaging." *J Magn Reson Imaging* 27(3): 546-51.

Kholodenko, B., O. Demin, et al. (1999). "Quantification of Short Term Signaling by the Epidermal Growth Factor Receptor." *The Journal of Biological Chemistry* 274(42): 30169-30181.

Kholodenko, B. N. (2000). "Negative feedback and ultrasensitivity can bring about oscillations in the mitogen-activated protein kinase cascades." *Eur J Biochem* 267(6): 1583-8.

Kholodenko, B. N. (2002). "MAP kinase cascade signaling and endocytic trafficking: a marriage of convenience?" *Trends Cell Biol* 12(4): 173-7.

Kholodenko, B. N. (2003). "Four-dimensional organization of protein kinase signaling cascades: the roles of diffusion, endocytosis and molecular motors." *J Exp Biol* 206(Pt 12): 2073-82.

Kholodenko, B. N. (2006). "Cell-signalling dynamics in time and space." *Nat Rev Mol Cell Biol* 7(3): 165-76.

Kholodenko, B. N. (2007). "Untangling the signalling wires." *Nat Cell Biol* 9(3): 247-9.

Kholodenko, B. N., G. C. Brown, et al. (2000). "Diffusion control of protein phosphorylation in signal transduction pathways." *Biochem J* 350 Pt 3: 901-7.

Kholodenko, B. N., M. Cascante, et al. (1998). "Metabolic design: how to engineer a living cell to desired metabolite concentrations and fluxes." *Biotechnol Bioeng* 59(2): 239-47.

Kholodenko, B. N., M. Cascante, et al. (1993). "Dramatic changes in control properties that accompany channelling and metabolite sequestration." *FEBS Lett* 336(3): 381-4.

Kholodenko, B. N., M. Cascante, et al. (1995). "Control theory of metabolic channelling." *Mol Cell Biochem* 143(2): 151-68.

Kholodenko, B. N., O. V. Demin, et al. (1999). "Quantification of short term signaling by the epidermal growth factor receptor." *J Biol Chem* 274(42): 30169-81.

Kholodenko, B. N., O. V. Demin, et al. (1993). "'Channelled' pathways can be more sensitive to specific regulatory signals." *FEBS Lett* 320(1): 75-8.

- Kholodenko, B. N., J. B. Hoek, et al. (2000). "Why cytoplasmic signalling proteins should be recruited to cell membranes." *Trends Cell Biol* 10(5): 173-8.
- Kholodenko, B. N., J. B. Hoek, et al. (1997). "Quantification of information transfer via cellular signal transduction pathways." *FEBS Lett* 414(2): 430-4.
- Kholodenko, B. N., A. Kiyatkin, et al. (2002). "Untangling the wires: a strategy to trace functional interactions in signaling and gene networks." *Proc Natl Acad Sci U S A* 99(20): 12841-6.
- Kholodenko, B. N. and W. Kolch (2008). "Giving space to cell signaling." *Cell* 133(4): 566-7.
- Kholodenko, B. N., D. Molenaar, et al. (1995). "Defining control coefficients in non-ideal metabolic pathways." *Biophys Chem* 56(3): 215-26.
- Kholodenko, B. N., J. M. Rohwer, et al. (1998). "Subtleties in control by metabolic channelling and enzyme organization." *Mol Cell Biochem* 184(1-2): 311-20.
- Kholodenko, B. N., N. Sakamoto, et al. (1996). "Strong control on the transit time in metabolic channelling." *FEBS Lett* 389(2): 123-5.
- Kholodenko, B. N. and H. M. Sauro (2008). "Spatio-temporal dynamics of protein modification cascades." *SEB Exp Biol Ser* 61: 141-59.
- Kholodenko, B. N., H. M. Sauro, et al. (1994). "Control by enzymes, coenzymes and conserved moieties. A generalisation of the connectivity theorem of metabolic control analysis." *Eur J Biochem* 225(1): 179-86.
- Kholodenko, B. N., H. M. Sauro, et al. (1995). "Coenzyme cycles and metabolic control analysis: the determination of the elasticity coefficients from the generalised connectivity theorem." *Biochem Mol Biol Int* 35(3): 615-25.
- Kholodenko, B. N., S. Schuster, et al. (1998). "Control analysis of metabolic systems involving quasi-equilibrium reactions." *Biochim Biophys Acta* 1379(3): 337-52.
- Kholodenko, B. N., S. Schuster, et al. (1995). "Composite control of cell function: metabolic pathways behaving as single control units." *FEBS Lett* 368(1): 1-4.
- Kholodenko, B. N. and H. V. Westerhoff (1993). "Metabolic channelling and control of the flux." *FEBS Lett* 320(1): 71-4.
- Kholodenko, B. N. and H. V. Westerhoff (1994). "Control theory of one enzyme." *Biochim Biophys Acta* 1208(2): 294-305.

- Kholodenko, B. N. and H. V. Westerhoff (1995). "The macroworld versus the microworld of biochemical regulation and control." *Trends Biochem Sci* 20(2): 52-4.
- Kholodenko, B. N., H. V. Westerhoff, et al. (1994). "Rate limitation within a single enzyme is directly related to enzyme intermediate levels." *FEBS Lett* 349(1): 131-4.
- Kholodenko, B. N., H. V. Westerhoff, et al. (1996). "Effect of channelling on the concentration of bulk-phase intermediates as cytosolic proteins become more concentrated." *Biochem J* 313 (Pt 3): 921-6.
- Kholodenko, B. N., H. V. Westerhoff, et al. (1995). "Control in channelled pathways. A matrix method calculating the enzyme control coefficients." *Biophys Chem* 53(3): 247-58.
- Kholodenko, B. N., H. V. Westerhoff, et al. (2000). "Engineering a living cell to desired metabolite concentrations and fluxes: pathways with multifunctional enzymes." *Metab Eng* 2(1): 1-13.
- Kholodenko, I. V., A. A. Buzdin, et al. (2006). "Mouse retinal progenitor cell (RPC) cocultivation with retinal pigment epithelial cell culture affects features of RPC differentiation." *Biochemistry (Mosc)* 71(7): 767-74.
- Kholodenko, I. V., R. V. Kholodenko, et al. (2008). "Ganglioside GM1-binding sites in interleukin-4: a photoaffinity labeling study." *Dokl Biochem Biophys* 418: 31-5.
- Kholodenko, R., I. Kholodenko, et al. (2007). "Anti-apoptotic effect of retinoic acid on retinal progenitor cells mediated by a protein kinase A-dependent mechanism." *Cell Res* 17(2): 151-62.
- Kholodenko, V. and E. Freire (1999). "A simple method to measure the absolute heat capacity of proteins." *Anal Biochem* 270(2): 336-8.
- Kholodenko, Y., E. A. Gooding, et al. (1999). "Heme protein dynamics revealed by geminate nitric oxide recombination in mutants of iron and cobalt myoglobin." *Biochemistry* 38(18): 5918-24.
- Kim, D. H., S. B. Choe, et al. (2003). "Direct observation of Barkhausen avalanche in Co thin films." *Phys Rev Lett* 90(8): 087203.
- Kim, H. and K. J. Shin (1999). "Exact Solution of the Reversible Diffusion-Influenced Reaction for an Isolated Pair in Three Dimensions." *Physical Review Letters* 82(7).
- Kim, H. D., T. W. Guo, et al. (2008). "Epidermal growth factor-induced enhancement of glioblastoma cell migration in 3D arises from an intrinsic increase in speed but an

extrinsic matrix- and proteolysis-dependent increase in persistence." *Mol Biol Cell* 19(10): 4249-59.

Kimura, T., J. Zhang, et al. (1997). "Syk-independent tyrosine phosphorylation and association of the protein tyrosine phosphatases SHP-1 and SHP-2 with the high affinity IgE receptor." *J Immunol* 159(9): 4426-34.

Kinoshita, E., E. Kinoshita-Kikuta, et al. (2008). "Separation of phosphoprotein isotypes having the same number of phosphate groups using phosphate-affinity SDS-PAGE." *Proteomics* 8(15): 2994-3003.

Kinzer-Ursem, T. L. and J. J. Linderman (2007). "Both ligand- and cell-specific parameters control ligand agonism in a kinetic model of G protein-coupled receptor signaling." *PLoS Comput Biol* 3(1): e6.

Kinzer-Ursem, T. L., K. L. Sutton, et al. (2006). "Multiple receptor states are required to describe both kinetic binding and activation of neutrophils via N-formyl peptide receptor ligands." *Cell Signal* 18(10): 1732-47.

Kiselar, J. G., P. A. Janmey, et al. (2003). "Structural analysis of gelsolin using synchrotron protein footprinting." *Mol Cell Proteomics* 2(10): 1120-32.

Kiss, Z., Y. Luo, et al. (1986). "Catalytic unit-independent phosphorylation and dephosphorylation of type II regulatory subunit of cyclic AMP-dependent protein kinase in rat liver plasma membranes." *Biochem J* 234(1): 163-8.

Kiyatkin, A., E. Aksamitiene, et al. (2006). "Scaffolding protein Grb2-associated binder 1 sustains epidermal growth factor-induced mitogenic and survival signaling by multiple positive feedback loops." *J Biol Chem* 281(29): 19925-38.

Klein, P., D. Mattoon, et al. (2004). "A structure-based model for ligand binding and dimerization of EGF receptors." *Proc Natl Acad Sci U S A* 101(4): 929-34.

Knecht, V., H. J. Risselada, et al. (2008). "Electrophoretic mobility does not always reflect the charge on an oil droplet." *J Colloid Interface Sci* 318(2): 477-86.

Ko, K. W., A. Paul, et al. (2005). "Endothelial lipase modulates HDL but has no effect on atherosclerosis development in apoE^{-/-} and LDLR^{-/-} mice." *J Lipid Res* 46(12): 2586-94.

Kodama, K., N. Ushida, et al. (1991). "Measurement of the relative branching fraction $\Gamma(D0 \rightarrow K \mu \nu) / \Gamma(D0 \rightarrow \mu X)$." *Phys Rev Lett* 66(14): 1819-1822.

Kolch, W., M. Calder, et al. (2005). "When kinases meet mathematics: the systems biology of MAPK signalling." *FEBS Lett* 579(8): 1891-5.

- Komarova, N. L., X. Zou, et al. (2005). "A theoretical framework for specificity in cell signaling." *Mol Syst Biol* 1: 2005 0023.
- Komori, S., Y. Ito, et al. (2008). "A long-term user of cosmetic cream containing estrogen developed breast cancer and endometrial hyperplasia." *Menopause* 15(6): 1191-2.
- Koppel, D. E. and M. P. Sheetz (1981). "Fluorescence photobleaching does not alter the lateral mobility of erythrocyte membrane glycoproteins." *Nature* 293(5828): 159-61.
- Koppel, D. E. and M. P. Sheetz (1983). "A localized pattern photobleaching method for the concurrent analysis of rapid and slow diffusion processes." *Biophys J* 43(2): 175-81.
- Koppel, D. E., M. P. Sheetz, et al. (1980). "Lateral diffusion in biological membranes. A normal-mode analysis of diffusion on a spherical surface." *Biophys J* 30(1): 187-92.
- Koppel, D. E., M. P. Sheetz, et al. (1981). "Matrix control of protein diffusion in biological membranes." *Proc Natl Acad Sci U S A* 78(6): 3576-80.
- Kostic, A., J. Sap, et al. (2007). "RPTPalpha is required for rigidity-dependent inhibition of extension and differentiation of hippocampal neurons." *J Cell Sci* 120(Pt 21): 3895-904.
- Kostic, A. and M. P. Sheetz (2006). "Fibronectin rigidity response through Fyn and p130Cas recruitment to the leading edge." *Mol Biol Cell* 17(6): 2684-95.
- Kovbasnjuk, O., M. Edidin, et al. (2001). "Role of lipid rafts in Shiga toxin 1 interaction with the apical surface of Caco-2 cells." *J Cell Sci* 114(Pt 22): 4025-31.
- Koyama-Honda, I., K. Ritchie, et al. (2005). "Fluorescence imaging for monitoring the colocalization of two single molecules in living cells." *Biophys J* 88(3): 2126-36.
- Kozminski, K. G., L. Beven, et al. (2003). "Interaction between a Ras and a Rho GTPase couples selection of a growth site to the development of cell polarity in yeast." *Mol Biol Cell* 14(12): 4958-70.
- Krauss, I., S. Grau, et al. (2008). "Sex-related differences in foot shape." *Ergonomics* 51(11): 1693-709.
- Krauss, S. W., D. Mochly-Rosen, et al. (1987). "Exposure of HeLa DNA polymerase alpha to protein kinase C affects its catalytic properties." *J Biol Chem* 262(8): 3432-5.
- Kucik, D. F., E. L. Elson, et al. (1989). "Forward transport of glycoproteins on leading lamellipodia in locomoting cells." *Nature* 340(6231): 315-7.

- Kucik, D. F., E. L. Elson, et al. (1990). "Cell migration does not produce membrane flow." *J Cell Biol* 111(4): 1617-22.
- Kucik, D. F., E. L. Elson, et al. (1999). "Weak dependence of mobility of membrane protein aggregates on aggregate size supports a viscous model of retardation of diffusion." *Biophys J* 76(1 Pt 1): 314-22.
- Kucik, D. F., S. C. Kuo, et al. (1991). "Preferential attachment of membrane glycoproteins to the cytoskeleton at the leading edge of lamella." *J Cell Biol* 114(5): 1029-36.
- Kumar, A., S. Dietrich, et al. (1997). "Genetic relatedness of *Burkholderia* (*Pseudomonas*) *cepacia* isolates from five cystic fibrosis centers in Michigan." *Respir Med* 91(8): 485-92.
- Kumar, J., H. P. Erickson, et al. (1998). "Ultrastructural and biochemical properties of the 120-kDa form of chick kinectin." *J Biol Chem* 273(48): 31738-43.
- Kumar, J., I. Toyoshima, et al. (1998). "Isolation and characterization of kinectin." *Methods Enzymol* 298: 185-97.
- Kumar, J., H. Yu, et al. (1995). "Kinectin, an essential anchor for kinesin-driven vesicle motility." *Science* 267(5205): 1834-7.
- Kumar, N., R. Afeyan, et al. (2008). "Multipathway model enables prediction of kinase inhibitor cross-talk effects on migration of Her2-overexpressing mammary epithelial cells." *Mol Pharmacol* 73(6): 1668-78.
- Kumar, N., M. H. Zaman, et al. (2006). "A high-throughput migration assay reveals HER2-mediated cell migration arising from increased directional persistence." *Biophys J* 91(4): L32-4.
- Kummer, U., B. Krajnc, et al. (2005). "Transition from stochastic to deterministic behavior in calcium oscillations." *Biophys J* 89(3): 1603-11.
- Kundu, A. K., M. Nagaoka, et al. (2003). "IGF-1 induces growth, survival and morphological change of primary hepatocytes on a galactose-bared polymer through both MAPK and beta-catenin pathways." *Cell Struct Funct* 28(4): 255-63.
- Kuo, S. C., J. Gelles, et al. (1991). "A model for kinesin movement from nanometer-level movements of kinesin and cytoplasmic dynein and force measurements." *J Cell Sci Suppl* 14: 135-8.
- Kuo, S. C. and M. P. Sheetz (1992). "Optical tweezers in cell biology." *Trends Cell Biol* 2(4): 116-8.

- Kuo, S. C. and M. P. Sheetz (1993). "Force of single kinesin molecules measured with optical tweezers." *Science* 260(5105): 232-4.
- Kurtz, S. M., C. L. Muhlstein, et al. (2000). "Surface morphology and wear mechanisms of four clinically relevant biomaterials after hip simulator testing." *J Biomed Mater Res* 52(3): 447-59.
- Kusumi, A., H. Ike, et al. (2005). "Single-molecule tracking of membrane molecules: plasma membrane compartmentalization and dynamic assembly of raft-philic signaling molecules." *Semin Immunol* 17(1): 3-21.
- Kusumi, A., C. Nakada, et al. (2005). "Paradigm Shift of the Plasma Membrane Concept from the Two-Dimensional Continuum Fluid to the Partitioned Fluid: High-Speed Single-Molecule Tracking of Membrane Molecules." *Annu. Rev. Biophys. Biomol. Struct* 34: 351-378.
- Kusumi, A., Y. Sako, et al. (1998). "Application of laser tweezers to studies of the fences and tethers of the membrane skeleton that regulate the movements of plasma membrane proteins." *Methods Cell Biol* 55: 173-94.
- Kusumi, A., Y. Sako, et al. (1993). "Confined lateral diffusion of membrane receptors as studied by single particle tracking (nanovid microscopy). Effects of calcium-induced differentiation in cultured epithelial cells." *Biophys J* 65(5): 2021-40.
- Kusumi, E., M. Kami, et al. (2006). "Hepatic injury following reduced intensity unrelated cord blood transplantation for adult patients with hematological diseases." *Biol Blood Marrow Transplant* 12(12): 1302-9.
- Kwik, J., S. Boyle, et al. (2003). "Membrane cholesterol, lateral mobility, and the phosphatidylinositol 4,5-bisphosphate-dependent organization of cell actin." *Proc Natl Acad Sci U S A* 100(24): 13964-9.
- Ladbury, J. E., M. A. Lemmon, et al. (1995). "Measurement of the binding of tyrosyl phosphopeptides to SH2 domains: a reappraisal." *Proc Natl Acad Sci U S A* 92(8): 3199-203.
- Ladha, S., P. S. James, et al. (1997). "Lateral mobility of plasma membrane lipids in bull spermatozoa: heterogeneity between surface domains and rigidification following cell death." *J Cell Sci* 110 (Pt 9): 1041-50.
- Ladha, S., A. R. Mackie, et al. (1994). "Cheek cell membrane fluidity measured by fluorescence recovery after photobleaching and steady-state fluorescence anisotropy." *J Membr Biol* 142(2): 223-8.
- Ladha, S., A. R. Mackie, et al. (1996). "Lateral diffusion in planar lipid bilayers: a fluorescence recovery after photobleaching investigation of its modulation by lipid

composition, cholesterol, or alamethicin content and divalent cations." *Biophys J* 71(3): 1364-73.

Lafont, F., K. Simons, et al. (1995). "Dissecting the molecular mechanisms of polarized membrane traffic: reconstitution of three transport steps in epithelial cells using streptolysin-O permeabilization." *Cold Spring Harb Symp Quant Biol* 60: 753-62.

Lag, M., E. Skarpen, et al. (2000). "Cell-specific expression of CCAAT/enhancer-binding protein delta (C/EBP delta) in epithelial lung cells." *Exp Lung Res* 26(5): 383-99.

Laidmae, I., M. E. McCormick, et al. (2006). "Stability, sterility, coagulation, and immunologic studies of salmon coagulation proteins with potential use for mammalian wound healing and cell engineering." *Biomaterials* 27(34): 5771-9.

Laing, S. J., S. J. Oliver, et al. (2008). "Neutrophil-degranulation and lymphocyte-subset response after 48 hr of fluid and/or energy restriction." *Int J Sport Nutr Exerc Metab* 18(5): 443-56.

Lajoie, P., E. A. Partridge, et al. (2007). "Plasma membrane domain organization regulates EGFR signaling in tumor cells." *J Cell Biol* 179(2): 341-56.

Lam, V., J. Kalesnikoff, et al. (2003). "IgE alone stimulates mast cell adhesion to fibronectin via pathways similar to those used by IgE + antigen but distinct from those used by Steel factor." *Blood* 102(4): 1405-13.

Lambert, A., R. Sadir, et al. (2000). "In vivo subcellular target compartments of interferon-gamma and interferon-gamma receptor (alpha- and beta-chains) in mouse liver." *Cytokine* 12(6): 715-9.

Lambert, T. N., N. L. Andrews, et al. (2007). "Water-soluble germanium(0) nanocrystals: cell recognition and near-infrared photothermal conversion properties." *Small* 3(4): 691-9.

Lane, T. M., W. Ansell, et al. (2004). "Long-term outcomes in patients with prostate cancer managed with intermittent androgen suppression." *Urol Int* 73(2): 117-22.

Lara, M., E. Ortega, et al. (2001). "Overcoming the signaling defect of Lyn-sequestering, signal-curtailing FcepsilonRI dimers: aggregated dimers can dissociate from Lyn and form signaling complexes with Syk." *J Immunol* 167(8): 4329-37.

Lasserre, R., X. J. Guo, et al. (2008). "Raft nanodomains contribute to Akt/PKB plasma membrane recruitment and activation." *Nat Chem Biol* 4(9): 538-47.

- Lauffenburger, D. A., E. M. Fallon, et al. (1998). "Scratching the (cell) surface: cytokine engineering for improved ligand/receptor trafficking dynamics." *Chem Biol* 5(10): R257-63.
- Lauffenburger, D. A., J. Linderman, et al. (1987). "Analysis of mammalian cell growth factor receptor dynamics." *Ann N Y Acad Sci* 506: 147-62.
- Lauffenburger, D. A. and J. J. Linderman (1993). *Receptors Models for Binding, Trafficking, and Signaling*. New York, Oxford University Press.
- LaVoo, E. J. and A. S. Paller (1994). "Common skin problems during the first year of life." *Pediatr Clin North Am* 41(5): 1105-19.
- Lebedeva, T., N. Anikeeva, et al. (2004). "Major histocompatibility complex class I-intercellular adhesion molecule-1 association on the surface of target cells: implications for antigen presentation to cytotoxic T lymphocytes." *Immunology* 113(4): 460-71.
- Lee, D. R., K. Shin, et al. (2003). "X-ray scattering from freestanding polymer films with geometrically curved surfaces." *Phys Rev Lett* 90(18): 185503.
- Lee, G., S. Hong, et al. (2001). "Structure of the Ba-induced Si(111)-(3 x 2) reconstruction." *Phys Rev Lett* 87(5): 056104.
- Lee, J., S. Eggert, et al. (2004). "Real space imaging of one-dimensional standing waves: direct evidence for a Luttinger liquid." *Phys Rev Lett* 93(16): 166403.
- Lee, J. T., P. Brafford, et al. (2008). "Unraveling the mysteries of IGF-1 signaling in melanoma." *J Invest Dermatol* 128(6): 1358-60.
- Lee, K. I., S. J. Joo, et al. (2007). "Kondo effect in magnetic tunnel junctions." *Phys Rev Lett* 98(10): 107202.
- Lee, R. J., D. E. Lujan, et al. (1997). "Cooperation between the Fc epsilonR1 and formyl peptide receptor signaling pathways in RBL(FPR) cells: the contribution of receptor-specific Ca²⁺ mobilization responses." *Biochem Biophys Res Commun* 235(3): 812-9.
- Leem, K., S. Y. Park, et al. (2003). "Effects of Jaoga-Yukmiwon(R), a Korean herbal medicine, on chondrocyte proliferation and longitudinal bone growth in adolescent male rats." *Phytother Res* 17(9): 1113-6.
- Legewie, S., B. Schoeberl, et al. (2007). "Competing docking interactions can bring about bistability in the MAPK cascade." *Biophys J* 93(7): 2279-88.

- Lei, M. G., X. Tan, et al. (2005). "Regulation of cellular caveolin-1 protein expression in murine macrophages by microbial products." *Infect Immun* 73(12): 8136-43.
- Leigh, J. P. and R. M. Sheetz (1989). "Prevalence of back pain among fulltime United States workers." *Br J Ind Med* 46(9): 651-7.
- Lemmon, M. A., Z. Bu, et al. (1997). "Two EGF molecules contribute additively to stabilization of the EGFR dimer." *EMBO J* 16(2): 281-94.
- Lemmon, M. A., K. M. Ferguson, et al. (1995). "Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin homology domain." *Proc Natl Acad Sci U S A* 92(23): 10472-6.
- Lemmon, M. A., K. M. Ferguson, et al. (1996). "PH domains: diverse sequences with a common fold recruit signaling molecules to the cell surface." *Cell* 85(5): 621-4.
- Lemmon, M. A., J. E. Ladbury, et al. (1994). "Independent binding of peptide ligands to the SH2 and SH3 domains of Grb2." *J Biol Chem* 269(50): 31653-8.
- Lemmon, M. A., D. Pinchasi, et al. (1997). "Kit receptor dimerization is driven by bivalent binding of stem cell factor." *J Biol Chem* 272(10): 6311-7.
- Lemmon, M. A. and J. Schlessinger (1994). "Regulation of signal transduction and signal diversity by receptor oligomerization." *Trends Biochem Sci* 19(11): 459-63.
- Lemmon, M. A. and J. Schlessinger (1998). "Transmembrane signaling by receptor oligomerization." *Methods Mol Biol* 84: 49-71.
- Lenne, P. F., L. Wawrezinieck, et al. (2006). "Dynamic molecular confinement in the plasma membrane by microdomains and the cytoskeleton meshwork." *EMBO J* 25(14): 3245-56.
- Lewis, J. (2008). "From signals to patterns: space, time, and mathematics in developmental biology." *Science* 322(5900): 399-403.
- Li, Q., M. Wang, et al. (2005). "Docosahexaenoic acid changes lipid composition and interleukin-2 receptor signaling in membrane rafts." *J Lipid Res* 46(9): 1904-13.
- Liepina, I., P. Janmey, et al. (2004). "Towards gelsolin amyloid formation." *Biopolymers* 76(6): 543-8.
- Lifshitz, J., P. A. Janmey, et al. (2006). "Photon correlation spectroscopy of brain mitochondrial populations: application to traumatic brain injury." *Exp Neurol* 197(2): 318-29.

- Lillemeier, B. F., J. R. Pfeiffer, et al. (2006). "Plasma membrane-associated proteins are clustered into islands attached to the cytoskeleton." *Proc Natl Acad Sci U S A* 103(50): 18992-7.
- Lin, M. F., J. W. Carlson, et al. (2007). "Revisiting the protein-coding gene catalog of *Drosophila melanogaster* using 12 fly genomes." *Genome Res* 17(12): 1823-36.
- Linderman, J. J. and D. A. Lauffenburger (1986). "Analysis of intracellular receptor/ligand sorting. Calculation of mean surface and bulk diffusion times within a sphere." *Biophys J* 50(2): 295-305.
- Lindesmith, L., J. M. McIlvain, Jr., et al. (1997). "Phosphotransferases associated with the regulation of kinesin motor activity." *J Biol Chem* 272(36): 22929-33.
- Lindesmith, L. C., J. Kumar, et al. (2001). "Identification of kinesin-associated proteins." *Methods Mol Biol* 164: 205-12.
- Linggi, B. and G. Carpenter (2006). "ErbB receptors: new insights on mechanisms and biology." *Trends Cell Biol* 16(12): 649-56.
- Link, T. M., L. S. Steinbach, et al. (2003). "Osteoarthritis: MR imaging findings in different stages of disease and correlation with clinical findings." *Radiology* 226(2): 373-81.
- Linshaw, M. A., C. A. Fogel, et al. (1992). "Role of cytoskeleton in volume regulation of rabbit proximal tubule in dilute medium." *Am J Physiol* 262(1 Pt 2): F144-50.
- Lippincott-Schwartz, J., N. Altan-Bonnet, et al. (2003). "Photobleaching and photoactivation: following protein dynamics in living cells." *Nat Cell Biol Suppl*: S7-14.
- Lippincott-Schwartz, J., N. Cole, et al. (1998). "Unravelling Golgi membrane traffic with green fluorescent protein chimeras." *Trends Cell Biol* 8(1): 16-20.
- Lippincott-Schwartz, J. and N. B. Cole (1995). "Roles for microtubules and kinesin in membrane traffic between the endoplasmic reticulum and the Golgi complex." *Biochem Soc Trans* 23(3): 544-8.
- Lippincott-Schwartz, J., N. B. Cole, et al. (1998). "Building a secretory apparatus: role of ARF1/COPI in Golgi biogenesis and maintenance." *Histochem Cell Biol* 109(5-6): 449-62.
- Lippincott-Schwartz, J., N. B. Cole, et al. (1995). "Kinesin is the motor for microtubule-mediated Golgi-to-ER membrane traffic." *J Cell Biol* 128(3): 293-306.

Lippincott-Schwartz, J., J. Glickman, et al. (1991). "Forskolin inhibits and reverses the effects of brefeldin A on Golgi morphology by a cAMP-independent mechanism." *J Cell Biol* 112(4): 567-77.

Lippincott-Schwartz, J. and G. H. Patterson (2003). "Development and use of fluorescent protein markers in living cells." *Science* 300(5616): 87-91.

Lippincott-Schwartz, J. and G. H. Patterson (2008). "Fluorescent proteins for photoactivation experiments." *Methods Cell Biol* 85: 45-61.

Lisanti, M. and E. Rodriguez-Boulan (1989). "Sorting of apical surface proteins and lipids by epithelial cells: is there a common mechanism?" *Soc Gen Physiol Ser* 44: 167-73.

Lisanti, M. P., I. W. Caras, et al. (1989). "A glycosphospholipid membrane anchor acts as an apical targeting signal in polarized epithelial cells." *J Cell Biol* 109(5): 2145-56.

Lisanti, M. P., I. W. Caras, et al. (1990). "Vectorial apical delivery and slow endocytosis of a glycolipid-anchored fusion protein in transfected MDCK cells." *Proc Natl Acad Sci U S A* 87(19): 7419-23.

Lisanti, M. P., I. W. Caras, et al. (1991). "Fusion proteins containing a minimal GPI-attachment signal are apically expressed in transfected MDCK cells." *J Cell Sci* 99 (Pt 3): 637-40.

Lisanti, M. P., J. C. Darnell, et al. (1989). "The distribution of glycosyl-phosphatidylinositol anchored proteins is differentially regulated by serum and insulin." *Biochem Biophys Res Commun* 164(2): 824-32.

Lisanti, M. P., M. C. Field, et al. (1991). "Mannosamine, a novel inhibitor of glycosylphosphatidylinositol incorporation into proteins." *EMBO J* 10(8): 1969-77.

Lisanti, M. P., A. Le Bivic, et al. (1990). "Preferred apical distribution of glycosyl-phosphatidylinositol (GPI) anchored proteins: a highly conserved feature of the polarized epithelial cell phenotype." *J Membr Biol* 113(2): 155-67.

Lisanti, M. P., A. Le Bivic, et al. (1989). "Steady-state distribution and biogenesis of endogenous Madin-Darby canine kidney glycoproteins: evidence for intracellular sorting and polarized cell surface delivery." *J Cell Biol* 109(5): 2117-27.

Lisanti, M. P. and E. Rodriguez-Boulan (1990). "Glycosphospholipid membrane anchoring provides clues to the mechanism of protein sorting in polarized epithelial cells." *Trends Biochem Sci* 15(3): 113-8.

Lisanti, M. P. and E. Rodriguez-Boulan (1991). "Polarized sorting of GPI-linked proteins in epithelia and membrane microdomains." *Cell Biol Int Rep* 15(11): 1023-49.

Lisanti, M. P., E. Rodriguez-Boulton, et al. (1990). "Emerging functional roles for the glycosyl-phosphatidylinositol membrane protein anchor." *J Membr Biol* 117(1): 1-10.

Lisanti, M. P., M. Sargiacomo, et al. (1988). "Polarized apical distribution of glycosyl-phosphatidylinositol-anchored proteins in a renal epithelial cell line." *Proc Natl Acad Sci U S A* 85(24): 9557-61.

Liu, J., X. B. Wang, et al. (2002). "Caveolin-1 expression enhances endothelial capillary tubule formation." *J Biol Chem* 277(12): 10661-8.

Liu, W., J. Sun, et al. (2009). "Association of a germ-line copy number variation at 2p24.3 and risk for aggressive prostate cancer." *Cancer Res* 69(6): 2176-9.

Liu, Y., Y. Zhu, et al. (2004). "Laminar flow activates peroxisome proliferator-activated receptor-gamma in vascular endothelial cells." *Circulation* 110(9): 1128-33.

Long, S. B., P. J. Hancock, et al. (2001). "The crystal structure of human protein farnesyltransferase reveals the basis for inhibition by CaaX tetrapeptides and their mimetics." *Proc Natl Acad Sci U S A* 98(23): 12948-53.

Lopez, L. A. and M. P. Sheetz (1993). "Steric inhibition of cytoplasmic dynein and kinesin motility by MAP2." *Cell Motil Cytoskeleton* 24(1): 1-16.

Lopez, L. A. and M. P. Sheetz (1995). "A microtubule-associated protein (MAP2) kinase restores microtubule motility in embryonic brain." *J Biol Chem* 270(21): 12511-7.

Louboutin, J. P., J. M. Navenot, et al. (1998). "X-linked vacuolated myopathy: membrane attack complex deposition on the surface membrane of injured muscle fibers is not accompanied by S-protein." *Muscle Nerve* 21(7): 932-5.

Lu, Y. B., K. Franze, et al. (2006). "Viscoelastic properties of individual glial cells and neurons in the CNS." *Proc Natl Acad Sci U S A* 103(47): 17759-64.

Lu, Z., S. Ghosh, et al. (2003). "Downregulation of caveolin-1 function by EGF leads to the loss of E-cadherin, increased transcriptional activity of beta-catenin, and enhanced tumor cell invasion." *Cancer Cell* 4(6): 499-515.

Luna, E. J., K. N. Prestonjamas, et al. (2001). "Motile Membrane Skeletons: What Neutrophils and Muscle Have in Common." *Cell Mol Biol Lett* 6(2): 219.

Lund, E. K., L. J. Harvey, et al. (1999). "Effects of dietary fish oil supplementation on the phospholipid composition and fluidity of cell membranes from human volunteers." *Ann Nutr Metab* 43(5): 290-300.

- Lundmark, R., G. J. Doherty, et al. (2008). "The GTPase-activating protein GRAF1 regulates the CLIC/GEEC endocytic pathway." *Curr Biol* 18(22): 1802-8.
- Lundmark, R., G. J. Doherty, et al. (2008). "Arf family GTP loading is activated by, and generates, positive membrane curvature." *Biochem J* 414(2): 189-94.
- Lusa, S., T. S. Blom, et al. (2001). "Depletion of rafts in late endocytic membranes is controlled by NPC1-dependent recycling of cholesterol to the plasma membrane." *J Cell Sci* 114(Pt 10): 1893-900.
- Ma, L., L. C. Cantley, et al. (1998). "Corequirement of specific phosphoinositides and small GTP-binding protein Cdc42 in inducing actin assembly in Xenopus egg extracts." *J Cell Biol* 140(5): 1125-36.
- Mac Gabhann, F. and A. S. Popel (2005). "Differential binding of VEGF isoforms to VEGF receptor 2 in the presence of neuropilin-1: a computational model." *Am J Physiol Heart Circ Physiol* 288(6): H2851-60.
- Mac Neil, S., M. Wagner, et al. (1992). "Signal transduction in murine B16 melanoma cells." *Melanoma Res* 2(3): 197-206.
- Maciejewski-Lenoir, D., J. G. Richman, et al. (2006). "Langerhans cells release prostaglandin D2 in response to nicotinic acid." *J Invest Dermatol* 126(12): 2637-46.
- Mackay, K. and D. Mochly-Rosen (2001). "Arachidonic acid protects neonatal rat cardiac myocytes from ischaemic injury through epsilon protein kinase C." *Cardiovasc Res* 50(1): 65-74.
- Mackay, K. and D. Mochly-Rosen (2001). "Localization, anchoring, and functions of protein kinase C isozymes in the heart." *J Mol Cell Cardiol* 33(7): 1301-7.
- Mackinnon, W. B., L. Huschtscha, et al. (1994). "Correlation of cellular differentiation in human colorectal carcinoma and adenoma cell lines with metabolite profiles determined by ¹H magnetic resonance spectroscopy." *Int J Cancer* 59(2): 248-61.
- Maeyama, R., T. Fukuyama, et al. (2007). "[A long-term survivor case with good quality of life due to biliary bypass and gemcitabine for advanced pancreatic cancer]." *Gan To Kagaku Ryoho* 34(11): 1873-5.
- Maffucci, T., A. Brancaccio, et al. (2003). "Insulin induces phosphatidylinositol-3-phosphate formation through TC10 activation." *EMBO J* 22(16): 4178-89.
- Magee, A. I., L. Gutierrez, et al. (1989). "Targeting of oncoproteins to membranes by fatty acylation." *J Cell Sci Suppl* 11: 149-60.

- Magee, A. I., C. M. Newman, et al. (1992). "Lipid modifications and function of the ras superfamily of proteins." *Biochem Soc Trans* 20(2): 497-9.
- Maguy, A., T. E. Hebert, et al. (2006). "Involvement of lipid rafts and caveolae in cardiac ion channel function." *Cardiovasc Res* 69(4): 798-807.
- Mahama, P. A. and J. J. Linderman (1994). "A Monte Carlo study of the dynamics of G-protein activation." *Biophys J* 67(3): 1345-57.
- Mahoney, N. M., P. A. Janmey, et al. (1997). "Structure of the profilin-poly-L-proline complex involved in morphogenesis and cytoskeletal regulation." *Nat Struct Biol* 4(11): 953-60.
- Maier, B., I. Chen, et al. (2004). "DNA transport into *Bacillus subtilis* requires proton motive force to generate large molecular forces." *Nat Struct Mol Biol* 11(7): 643-9.
- Maier, B., M. Koomey, et al. (2004). "A force-dependent switch reverses type IV pilus retraction." *Proc Natl Acad Sci U S A* 101(30): 10961-6.
- Maier, B., L. Potter, et al. (2002). "Single pilus motor forces exceed 100 pN." *Proc Natl Acad Sci U S A* 99(25): 16012-7.
- Maiwald, C., S. Grau, et al. (2008). "Reproducibility of plantar pressure distribution data in barefoot running." *J Appl Biomech* 24(1): 14-23.
- Maiwald, M. (2008). "Alcohol-based hand hygiene and nosocomial infection rates." *Infect Control Hosp Epidemiol* 29(6): 579-80; author reply 580-2.
- Maiwald, T. and J. Timmer (2008). "Dynamical modeling and multi-experiment fitting with PottersWheel." *Bioinformatics* 24(18): 2037-43.
- Malcolm, M. J. and J. R. Gear (1971). "Biosynthesis of Chelidonic Acid." *Canadian Journal of Biochemistry* 49(4): 412-&.
- Mandiyani, V., R. O'Brien, et al. (1996). "Thermodynamic studies of SHC phosphotyrosine interaction domain recognition of the NPXpY motif." *J Biol Chem* 271(9): 4770-5.
- Manley, S., J. M. Gillette, et al. (2008). "High-density mapping of single-molecule trajectories with photoactivated localization microscopy." *Nat Methods* 5(2): 155-7.
- Mao, S. Y., T. Yamashita, et al. (1995). "Chemical cross-linking of IgE-receptor complexes in RBL-2H3 cells." *Biochemistry* 34(6): 1968-77.
- Marani, M., D. Hancock, et al. (2004). "Role of Bim in the survival pathway induced by Raf in epithelial cells." *Oncogene* 23(14): 2431-41.

- Marguet, D., P. F. Lenne, et al. (2006). "Dynamics in the plasma membrane: how to combine fluidity and order." *EMBO J* 25(15): 3446-57.
- Marhl, M., M. Gosak, et al. (2008). "Spatio-temporal modelling explains the effect of reduced plasma membrane Ca^{2+} efflux on intracellular Ca^{2+} oscillations in hepatocytes." *J Theor Biol* 252(3): 419-26.
- Markevich, N. I., J. B. Hoek, et al. (2004). "Signaling switches and bistability arising from multisite phosphorylation in protein kinase cascades." *J Cell Biol* 164(3): 353-9.
- Markevich, N. I., G. Moehren, et al. (2004). "Signal processing at the Ras circuit: what shapes Ras activation patterns?" *Syst Biol (Stevenage)* 1(1): 104-13.
- Markevich, N. I., M. A. Tsyganov, et al. (2006). "Long-range signaling by phosphoprotein waves arising from bistability in protein kinase cascades." *Mol Syst Biol* 2: 61.
- Marro, J., A. B. Bortz, et al. (1975). "Time Evolution of a Quenched Binary Alloy .2. Computer-Simulation of a 3-Dimensional Model System." *Physical Review B* 12(6): 2000-2011.
- Martenson, C., K. Stone, et al. (1993). "Fast axonal transport is required for growth cone advance." *Nature* 366(6450): 66-9.
- Martenson, C. H., A. Odom, et al. (1995). "The effect of acrylamide and other sulfhydryl alkylators on the ability of dynein and kinesin to translocate microtubules in vitro." *Toxicol Appl Pharmacol* 133(1): 73-81.
- Martenson, C. H., M. P. Sheetz, et al. (1995). "In vitro acrylamide exposure alters growth cone morphology." *Toxicol Appl Pharmacol* 131(1): 119-29.
- Martinez-Pinna, J., I. S. Gurung, et al. (2005). "Direct voltage control of signaling via P2Y1 and other G α q-coupled receptors." *J Biol Chem* 280(2): 1490-8.
- Masada, N., A. Ciruela, et al. (2009). "Distinct mechanisms of regulation by Ca^{2+} /calmodulin of type 1 and 8 adenylyl cyclases support their different physiological roles." *J Biol Chem* 284(7): 4451-63.
- Masuda, A., K. Nakamura, et al. (2008). "Serum autotaxin measurement in haematological malignancies: a promising marker for follicular lymphoma." *Br J Haematol* 143(1): 60-70.
- Matko, J., A. Bodnar, et al. (2002). "GPI-microdomains (membrane rafts) and signaling of the multi-chain interleukin-2 receptor in human lymphoma/leukemia T cell lines." *Eur J Biochem* 269(4): 1199-208.

- Matsubara, H., Y. Yamada, et al. (2008). "Potential for HER-2/neu molecular targeted therapy for invasive bladder carcinoma: comparative study of immunohistochemistry and fluorescent in situ hybridization." *Oncol Rep* 19(1): 57-63.
- Matsumoto, T., A. Andoh, et al. (2008). "Multivariate analysis for factors predicting rapid response of leukocytapheresis in patients with steroid-resistant ulcerative colitis: a multicenter prospective open-label study." *Ther Apher Dial* 12(6): 484-90.
- Mattoon, D., P. Klein, et al. (2004). "The tethered configuration of the EGF receptor extracellular domain exerts only a limited control of receptor function." *Proc Natl Acad Sci U S A* 101(4): 923-8.
- Matyja, A., M. Rozanska, et al. (2007). "Observation of $B(0) \rightarrow D^{(*)} \tau^{+} \nu(\tau)$ decay at Belle." *Phys Rev Lett* 99(19): 191807.
- Mauch, M., S. Grau, et al. (2008). "Foot morphology of normal, underweight and overweight children." *Int J Obes (Lond)* 32(7): 1068-75.
- May, R. M. (2004). "Uses and abuses of mathematics in biology." *Science* 303(5659): 790-3.
- Mayawala, K., D. G. Vlachos, et al. (2005). "Computational modeling reveals molecular details of epidermal growth factor binding." *BMC Cell Biol* 6: 41.
- Mayawala, K., D. G. Vlachos, et al. (2005). "Heterogeneities in EGF receptor density at the cell surface can lead to concave up scatchard plot of EGF binding." *FEBS Lett* 579(14): 3043-7.
- Mayawala, K., D. G. Vlachos, et al. (2006). "Spatial modeling of dimerization reaction dynamics in the plasma membrane: Monte Carlo vs. continuum differential equations." *Biophys Chem* 121(3): 194-208.
- Mazighi, M., A. Pelle, et al. (2004). "IL-10 inhibits vascular smooth muscle cell activation in vitro and in vivo." *Am J Physiol Heart Circ Physiol* 287(2): H866-71.
- McClatchey, A. I., P. Van den Bergh, et al. (1992). "Temperature-sensitive mutations in the III-IV cytoplasmic loop region of the skeletal muscle sodium channel gene in paramyotonia congenita." *Cell* 68(4): 769-74.
- McDiarmid, M. A., S. Engelhardt, et al. (2004). "Health effects of depleted uranium on exposed Gulf War veterans: a 10-year follow-up." *J Toxicol Environ Health A* 67(4): 277-96.
- McDiarmid, M. A., S. M. Engelhardt, et al. (2009). "Surveillance results of depleted uranium-exposed Gulf War I veterans: sixteen years of follow-up." *J Toxicol Environ Health A* 72(1): 14-29.

- McDiarmid, M. A., S. M. Engelhardt, et al. (2007). "Health surveillance of Gulf War I veterans exposed to depleted uranium: updating the cohort." *Health Phys* 93(1): 60-73.
- McDiarmid, M. A., S. M. Engelhardt, et al. (2006). "Biological monitoring and surveillance results of Gulf War I veterans exposed to depleted uranium." *Int Arch Occup Environ Health* 79(1): 11-21.
- McDiarmid, M. A., K. Squibb, et al. (2001). "Surveillance of depleted uranium exposed Gulf War veterans: health effects observed in an enlarged "friendly fire" cohort." *J Occup Environ Med* 43(12): 991-1000.
- McGoldrick, C. and M. Sheetz (1998). "Organelle motility and membrane network formation from cultured mammalian cells." *Methods Enzymol* 298: 353-60.
- McIlvain, J. M., Jr., J. K. Burkhardt, et al. (1994). "Regulation of kinesin activity by phosphorylation of kinesin-associated proteins." *J Biol Chem* 269(29): 19176-82.
- McIlvain, J. M., Jr., C. Lamb, et al. (1993). "Microtubule motor-dependent formation of tubulovesicular networks from endoplasmic reticulum and Golgi membranes." *Methods Cell Biol* 39: 227-36.
- McIlvain, J. M., Jr. and M. P. Sheetz (1992). "Formation of endoplasmic reticulum networks in vitro." *Methods Enzymol* 219: 72-80.
- McIntosh, A. L., A. M. Gallegos, et al. (2003). "Fluorescence and multiphoton imaging resolve unique structural forms of sterol in membranes of living cells." *J Biol Chem* 278(8): 6384-403.
- McIntyre, C. L., A. H. Sheetz, et al. (2005). "Administration of epinephrine for life-threatening allergic reactions in school settings." *Pediatrics* 116(5): 1134-40.
- McKay, M. M. and D. K. Morrison (2007). "Caspase-dependent cleavage disrupts the ERK cascade scaffolding function of KSR1." *J Biol Chem* 282(36): 26225-34.
- McLemore, M. R., C. Miaskowski, et al. (2008). "Rules of tumor cell development and their application to biomarkers for ovarian cancer." *Oncol Nurs Forum* 35(3): 403-9.
- McPherson, R. A., A. Harding, et al. (1999). "Interactions of c-Raf-1 with phosphatidylserine and 14-3-3." *Oncogene* 18(26): 3862-9.
- Mecham, R. P., L. Whitehouse, et al. (1991). "Ligand affinity of the 67-kD elastin/laminin binding protein is modulated by the protein's lectin domain: visualization of elastin/laminin-receptor complexes with gold-tagged ligands." *J Cell Biol* 113(1): 187-94.

- Medintz, I. L., L. Berti, et al. (2007). "A reactive peptidic linker for self-assembling hybrid quantum dot-DNA bioconjugates." *Nano Lett* 7(6): 1741-8.
- Medintz, I. L., A. R. Clapp, et al. (2006). "Proteolytic activity monitored by fluorescence resonance energy transfer through quantum-dot-peptide conjugates." *Nat Mater* 5(7): 581-9.
- Medintz, I. L., A. R. Clapp, et al. (2003). "Self-assembled nanoscale biosensors based on quantum dot FRET donors." *Nat Mater* 2(9): 630-8.
- Medintz, I. L., J. H. Konnert, et al. (2004). "A fluorescence resonance energy transfer-derived structure of a quantum dot-protein bioconjugate nanoassembly." *Proc Natl Acad Sci U S A* 101(26): 9612-7.
- Medintz, I. L. and H. Mattoussi (2009). "Quantum dot-based resonance energy transfer and its growing application in biology." *Phys Chem Chem Phys* 11(1): 17-45.
- Medintz, I. L., H. Mattoussi, et al. (2008). "Potential clinical applications of quantum dots." *Int J Nanomedicine* 3(2): 151-67.
- Medintz, I. L., T. Pons, et al. (2008). "Intracellular delivery of quantum dot-protein cargos mediated by cell penetrating peptides." *Bioconjug Chem* 19(9): 1785-95.
- Medintz, I. L., T. Pons, et al. (2008). "Interactions between Redox Complexes and Semiconductor Quantum Dots Coupled via a Peptide Bridge." *J Am Chem Soc* 130(49): 16745-16756.
- Medintz, I. L., K. E. Sapsford, et al. (2006). "Designer variable repeat length polypeptides as scaffolds for surface immobilization of quantum dots." *J Phys Chem B* 110(22): 10683-90.
- Medintz, I. L., K. E. Sapsford, et al. (2005). "Decoration of discretely immobilized cowpea mosaic virus with luminescent quantum dots." *Langmuir* 21(12): 5501-10.
- Medintz, I. L., S. A. Trammell, et al. (2004). "Reversible modulation of quantum dot photoluminescence using a protein-bound photochromic fluorescence resonance energy transfer acceptor." *J Am Chem Soc* 126(1): 30-1.
- Medintz, I. L., H. T. Uyeda, et al. (2005). "Quantum dot bioconjugates for imaging, labelling and sensing." *Nat Mater* 4(6): 435-46.
- Mehta, N. N., M. Sheetz, et al. (2009). "Selective PKC beta inhibition with ruboxistaurin and endothelial function in type-2 diabetes mellitus." *Cardiovasc Drugs Ther* 23(1): 17-24.

- Mei, B. C., K. Susumu, et al. (2009). "Polyethylene glycol-based bidentate ligands to enhance quantum dot and gold nanoparticle stability in biological media." *Nat Protoc* 4(3): 412-23.
- Meilhac, N., L. Le Guyader, et al. (2006). "Detection of confinement and jumps in single-molecule membrane trajectories." *Phys Rev E Stat Nonlin Soft Matter Phys* 73(1 Pt 1): 011915.
- Mendoza, F. J., E. S. Henson, et al. (2005). "MEKK1-induced apoptosis is mediated by Smac/Diablo release from the mitochondria." *Biochem Biophys Res Commun* 331(4): 1089-98.
- Merz, A. J., M. So, et al. (2000). "Pilus retraction powers bacterial twitching motility." *Nature* 407(6800): 98-102.
- Mesarovic, M. D., S. N. Sreenath, et al. (2004). "Search for organising principles: understanding in systems biology." *Syst Biol (Stevenage)* 1(1): 19-27.
- Meshel, A. S., Q. Wei, et al. (2005). "Basic mechanism of three-dimensional collagen fibre transport by fibroblasts." *Nat Cell Biol* 7(2): 157-64.
- Metoki, R., T. Horibe, et al. (2008). "[A case of splenic inflammatory pseudotumor]." *Nippon Shokakibyo Gakkai Zasshi* 105(2): 257-64.
- Metropolis, N., A. W. Rosenbluth, et al. (1953). "Equation of State Calculations by Fast Computing Machines." *Journal of Chemical Physics* 21(6): 1087-1092.
- Metsikko, K. and K. Simons (1986). "The budding mechanism of spikeless vesicular stomatitis virus particles." *EMBO J* 5(8): 1913-20.
- Metsikko, K., G. van Meer, et al. (1986). "Reconstitution of the fusogenic activity of vesicular stomatitis virus." *EMBO J* 5(13): 3429-35.
- Metzger, H., B. Goldstein, et al. (1994). "Quantitative aspects of receptor aggregation." *Adv Exp Med Biol* 365: 175-83.
- Mhadeshwar, A. B., J. Ludwig, et al. (2004). "Surface reactivity as a many body multiscale problem." *Abstracts of Papers of the American Chemical Society* 227: U1087-U1087.
- Michaud, N. R., M. Therrien, et al. (1997). "KSR stimulates Raf-1 activity in a kinase-independent manner." *Proc Natl Acad Sci U S A* 94(24): 12792-6.
- Michel, C. M., M. M. Murray, et al. (2004). "EEG source imaging." *Clin Neurophysiol* 115(10): 2195-222.

- Michel, V. and M. Bakovic (2007). "Lipid rafts in health and disease." *Biol Cell* 99(3): 129-40.
- Michel, V. and M. Bakovic (2009). "The solute carrier 44A1 is a mitochondrial protein and mediates choline transport." *FASEB J*.
- Michel, V., Z. Yuan, et al. (2006). "Choline transport for phospholipid synthesis." *Exp Biol Med (Maywood)* 231(5): 490-504.
- Micheloud, D., M. Calderon, et al. (2007). "Intravenous immunoglobulin therapy in severe lupus myocarditis: good outcome in three patients." *Ann Rheum Dis* 66(7): 986-7.
- Mildaziene, V., R. Baniene, et al. (1995). "Calcium indirectly increases the control exerted by the adenine nucleotide translocator over 2-oxoglutarate oxidation in rat heart mitochondria." *Arch Biochem Biophys* 324(1): 130-4.
- Miller, J. H., F. Zheng, et al. (2005). "A model of cytokine shedding induced by low doses of gamma radiation." *Radiat Res* 163(3): 337-42.
- Miller, K. E. and M. P. Sheetz (2000). "Characterization of myosin V binding to brain vesicles." *J Biol Chem* 275(4): 2598-606.
- Miller, K. E. and M. P. Sheetz (2004). "Axonal mitochondrial transport and potential are correlated." *J Cell Sci* 117(Pt 13): 2791-804.
- Miller, K. E. and M. P. Sheetz (2006). "Direct evidence for coherent low velocity axonal transport of mitochondria." *J Cell Biol* 173(3): 373-81.
- Miller, L. W., Y. Cai, et al. (2005). "In vivo protein labeling with trimethoprim conjugates: a flexible chemical tag." *Nat Methods* 2(4): 255-7.
- Miller, L. W., J. Sable, et al. (2004). "Methotrexate conjugates: a molecular in vivo protein tag." *Angew Chem Int Ed Engl* 43(13): 1672-5.
- Miller, M. M., C. R. Sweeney, et al. (1999). "Effects of blood contamination of cerebrospinal fluid on western blot analysis for detection of antibodies against *Sarcocystis neurona* and on albumin quotient and immunoglobulin G index in horses." *J Am Vet Med Assoc* 215(1): 67-71.
- Minemoto, Y., J. Gannon, et al. (2001). "Characterization of adriamycin-induced G2 arrest and its abrogation by caffeine in FL-amnion cells with or without p53." *Exp Cell Res* 262(1): 37-48.
- Miyaji, M., Z. X. Jin, et al. (2005). "Role of membrane sphingomyelin and ceramide in platform formation for Fas-mediated apoptosis." *J Exp Med* 202(2): 249-59.

Mocanu, M. M., Z. Fazekas, et al. (2005). "Associations of ErbB2, beta1-integrin and lipid rafts on Herceptin (Trastuzumab) resistant and sensitive tumor cell lines." *Cancer Lett* 227(2): 201-12.

Mochly-Rosen, D. (1995). "Localization of protein kinases by anchoring proteins: a theme in signal transduction." *Science* 268(5208): 247-51.

Mochly-Rosen, D., A. I. Basbaum, et al. (1987). "Distinct cellular and regional localization of immunoreactive protein kinase C in rat brain." *Proc Natl Acad Sci U S A* 84(13): 4660-4.

Mochly-Rosen, D., J. A. Fagin, et al. (2001). "Spontaneous occurrence of an inhibitor of protein kinase C localization in a thyroid cancer cell line: role in thyroid tumorigenesis." *Adv Enzyme Regul* 41: 87-97.

Mochly-Rosen, D. and D. E. Koshland, Jr. (1987). "Domain structure and phosphorylation of protein kinase C." *J Biol Chem* 262(5): 2291-7.
Mochly-Rosen, D. and D. E. Koshland, Jr. (1988). "A general procedure for screening inhibitory antibodies: application for identifying anti-protein kinase C antibodies." *Anal Biochem* 170(1): 31-7.

Moehren, G., N. Markevich, et al. (2002). "Temperature dependence of the epidermal growth factor receptor signaling network can be accounted for by a kinetic model." *Biochemistry* 41(1): 306-20.

Molotkovskaya, I. M., R. V. Kholodenko, et al. (2002). "Influence of gangliosides on the IL-2- and IL-4-dependent cell proliferation." *Neurochem Res* 27(7-8): 761-70.

Monine, M. I., A. M. Berezhkovskii, et al. (2005). "Ligand accumulation in autocrine cell cultures." *Biophys J* 88(4): 2384-90.

Mooseker, M. S., K. A. Conzelman, et al. (1989). "Characterization of intestinal microvillar membrane disks: detergent-resistant membrane sheets enriched in associated brush border myosin I (110K-calmodulin)." *J Cell Biol* 109(3): 1153-61.

Mora, R., V. L. Bonilha, et al. (1999). "Caveolin-2 localizes to the golgi complex but redistributes to plasma membrane, caveolae, and rafts when co-expressed with caveolin-1." *J Biol Chem* 274(36): 25708-17.

Morales, C. P., R. F. Souza, et al. (2002). "Hallmarks of cancer progression in Barrett's oesophagus." *Lancet* 360(9345): 1587-9.

Morimatsu, M., H. Takagi, et al. (2007). "Multiple-state reactions between the epidermal growth factor receptor and Grb2 as observed by using single-molecule analysis." *Proc Natl Acad Sci U S A* 104(46): 18013-8.

- Morone, N., T. Fujiwara, et al. (2006). "Three-dimensional reconstruction of the membrane skeleton at the plasma membrane interface by electron tomography." *J Cell Biol* 174(6): 851-62.
- Morone, N., C. Nakada, et al. (2008). "Three-dimensional molecular architecture of the plasma-membrane-associated cytoskeleton as reconstructed by freeze-etch electron tomography." *Methods Cell Biol* 88: 207-36.
- Morrison, D. K. (2001). "KSR: a MAPK scaffold of the Ras pathway?" *J Cell Sci* 114(Pt 9): 1609-12.
- Morrison, D. K. and R. J. Davis (2003). "Regulation of MAP kinase signaling modules by scaffold proteins in mammals." *Annu Rev Cell Dev Biol* 19: 91-118.
- Morton, C. O., A. Hayes, et al. (2007). "Global phenotype screening and transcript analysis outlines the inhibitory mode(s) of action of two amphibian-derived, alpha-helical, cationic peptides on *Saccharomyces cerevisiae*." *Antimicrob Agents Chemother* 51(11): 3948-59.
- Moses-Kolko, E. L., C. C. Meltzer, et al. (2005). "No interruption of lactation is needed after (11)C-WAY 100635 or (11)C-raclopride PET." *J Nucl Med* 46(10): 1765.
- Muller, J., A. M. Cacace, et al. (2000). "Identification of B-KSR1, a novel brain-specific isoform of KSR1 that functions in neuronal signaling." *Mol Cell Biol* 20(15): 5529-39.
- Muller, J., S. Ory, et al. (2001). "C-TAK1 regulates Ras signaling by phosphorylating the MAPK scaffold, KSR1." *Mol Cell* 8(5): 983-93.
- Muller, J., D. A. Ritt, et al. (2003). "Functional analysis of C-TAK1 substrate binding and identification of PKP2 as a new C-TAK1 substrate." *EMBO J* 22(17): 4431-42.
- Munoz-Garcia, J., Z. Neufeld, et al. (2009). "Positional information generated by spatially distributed signaling cascades." *PLoS Comput Biol* 5(3): e1000330.
- Munro, E. (2007). "The microtubules dance and the spindle poles swing." *Cell* 129(3): 457-8.
- Munro, S. (2003). "Lipid rafts: elusive or illusive?" *Cell* 115(4): 377-88.

- Munro, S., B. J. Bast, et al. (1992). "The B lymphocyte surface antigen CD75 is not an alpha-2,6-sialyltransferase but is a carbohydrate antigen, the production of which requires the enzyme." *Cell* 68(6): 1003.
- Munro, S. and H. R. Pelham (1986). "An Hsp70-like protein in the ER: identity with the 78 kd glucose-regulated protein and immunoglobulin heavy chain binding protein." *Cell* 46(2): 291-300.
- Munro, S. and H. R. Pelham (1987). "A C-terminal signal prevents secretion of luminal ER proteins." *Cell* 48(5): 899-907.
- Muppidi, J. R., J. Tschopp, et al. (2004). "Life and death decisions: secondary complexes and lipid rafts in TNF receptor family signal transduction." *Immunity* 21(4): 461-5.
- Murakoshi, H., R. Iino, et al. (2004). "Single-molecule imaging analysis of Ras activation in living cells." *Proc Natl Acad Sci U S A* 101(19): 7317-22.
- Murase, K., T. Fujiwara, et al. (2004). "Ultrafine membrane compartments for molecular diffusion as revealed by single molecule techniques." *Biophys J* 86(6): 4075-93.
- Nagasaka, A., H. Matsue, et al. (2008). "Osteopontin is produced by mast cells and affects IgE-mediated degranulation and migration of mast cells." *Eur J Immunol* 38(2): 489-99.
- Nagy, P., G. Vereb, et al. (2002). "Lipid rafts and the local density of ErbB proteins influence the biological role of homo- and heteroassociations of ErbB2." *J Cell Sci* 115(Pt 22): 4251-62.
- Nakada, C., N. Morone, et al. (2006). "[Membrane skeleton: interaction of the plasma membrane with the cytoskeleton]." *Tanpakushitsu Kakusan Koso* 51(6 Suppl): 672-82.
- Nakada, C., K. Ritchie, et al. (2003). "Accumulation of anchored proteins forms membrane diffusion barriers during neuronal polarization." *Nat Cell Biol* 5(7): 626-32.
- Nakamori, M., M. Iwahashi, et al. (2008). "Laparoscopic resection for gastrointestinal stromal tumors of the stomach." *Am J Surg* 196(3): 425-9.
- Nakamura, K., H. Aoki, et al. (2008). "Compartment syndrome with thrombosis of common iliac artery after gynecologic surgery." *Obstet Gynecol* 112(2 Pt 2): 486-8.
- Nakamura, K., K. Igarashi, et al. (2008). "Validation of an autotaxin enzyme immunoassay in human serum samples and its application to hypoalbuminemia differentiation." *Clin Chim Acta* 388(1-2): 51-8.

Nakamura, K., M. Nangaku, et al. (2008). "Analysis of serum and urinary lysophospholipase D/autotaxin in nephrotic syndrome." *Clin Chem Lab Med* 46(1): 150-1.

Nakamura, S., Y. Aoki, et al. (2008). "Sox9b/sox9a2-EGFP transgenic medaka reveals the morphological reorganization of the gonads and a common precursor of both the female and male supporting cells." *Mol Reprod Dev* 75(3): 472-6.

Nakamura, S., M. Nambu, et al. (2008). "Effect of controlled release of fibroblast growth factor-2 from chitosan/fucoidan micro complex-hydrogel on in vitro and in vivo vascularization." *J Biomed Mater Res A* 85(3): 619-27.

Nakamura, T., K. Aoki, et al. (2008). "FRET imaging and in silico simulation: analysis of the signaling network of nerve growth factor-induced neuritogenesis." *Brain Cell Biol* 36(1-4): 19-30.

National Research Council (U.S.). Committee on Solids. and R. Smoluchowski (1951). *Phase transformations in solids*. New York,, Wiley.

Navarro, G., P. Carriba, et al. (2008). "Detection of heteromers formed by cannabinoid CB1, dopamine D2, and adenosine A2A G-protein-coupled receptors by combining bimolecular fluorescence complementation and bioluminescence energy transfer." *ScientificWorldJournal* 8: 1088-97.

Nebi, T., S. W. Oh, et al. (2000). "Membrane cytoskeleton: PIP(2) pulls the strings." *Curr Biol* 10(9): R351-4.

Nebi, T., K. N. Pestonjamasp, et al. (2002). "Proteomic analysis of a detergent-resistant membrane skeleton from neutrophil plasma membranes." *J Biol Chem* 277(45): 43399-409.

Nehls, S., E. L. Snapp, et al. (2000). "Dynamics and retention of misfolded proteins in native ER membranes." *Nat Cell Biol* 2(5): 288-95.

Nelson, S., R. D. Horvat, et al. (1999). "Characterization of an intrinsically fluorescent gonadotropin-releasing hormone receptor and effects of ligand binding on receptor lateral diffusion." *Endocrinology* 140(2): 950-7.

Newman, C. M., T. Giannakouros, et al. (1992). "Post-translational processing of *Schizosaccharomyces pombe* YPT proteins." *J Biol Chem* 267(16): 11329-36.

Nezu, U., A. Nakamura, et al. (2008). "Comparative Study of Effectiveness of Multiple-Daily Injections of Insulin Versus Twice-Daily Injections of Biphasic Insulin in Patients with Type 2 Diabetes." *Endocr J*.

- Nguyen, A., W. R. Burack, et al. (2002). "Kinase suppressor of Ras (KSR) is a scaffold which facilitates mitogen-activated protein kinase activation in vivo." *Mol Cell Biol* 22(9): 3035-45.
- Nguyen, H. L., D. Gruber, et al. (1998). "Stabilization and functional modulation of microtubules by microtubule-associated protein 4." *Biol Bull* 194(3): 354-7.
- Nicke, B., J. Bastien, et al. (2005). "Involvement of MINK, a Ste20 family kinase, in Ras oncogene-induced growth arrest in human ovarian surface epithelial cells." *Mol Cell* 20(5): 673-85.
- Niebuhr, K., S. Giuriato, et al. (2002). "Conversion of PtdIns(4,5)P(2) into PtdIns(5)P by the *S.flexneri* effector IpgD reorganizes host cell morphology." *EMBO J* 21(19): 5069-78.
- Niehaus, A. M., D. G. Vlachos, et al. (2008). "Microscopic simulation of membrane molecule diffusion on corrallled membrane surfaces." *Biophys J* 94(5): 1551-64.
- Nielsen, U. B. and B. Schoeberl (2005). "Using computational modeling to drive the development of targeted therapeutics." *IDrugs* 8(10): 822-6.
- Nigg, E., M. Kessler, et al. (1979). "Labeling of human erythrocyte membranes with eosin probes used for protein diffusion measurements: inhibition of anion transport and photo-oxidative inactivation of acetylcholinesterase." *Biochim Biophys Acta* 550(2): 328-40.
- Nikonov, S. S., R. Kholodenko, et al. (2006). "Physiological features of the S- and M-cone photoreceptors of wild-type mice from single-cell recordings." *J Gen Physiol* 127(4): 359-74.
- Nishizaka, T., Q. Shi, et al. (2000). "Position-dependent linkages of fibronectin-integrin-cytoskeleton." *Proc Natl Acad Sci U S A* 97(2): 692-7.
- Nouso, K., H. Tanaka, et al. (2008). "Cost-effectiveness of the surveillance program of hepatocellular carcinoma depends on the medical circumstances." *J Gastroenterol Hepatol* 23(3): 437-44.
- Oback, B., A. T. Wiersema, et al. (2003). "Cloned cattle derived from a novel zona-free embryo reconstruction system." *Cloning Stem Cells* 5(1): 3-12.
- Oguma, K., H. Oshima, et al. (2008). "Activated macrophages promote Wnt signalling through tumour necrosis factor-alpha in gastric tumour cells." *EMBO J* 27(12): 1671-81.
- Oh, S. W., R. K. Pope, et al. (2003). "Archvillin, a muscle-specific isoform of supervillin, is an early expressed component of the costameric membrane skeleton." *J Cell Sci* 116(Pt 11): 2261-75.

- Ohama, K., I. Kusumi, et al. (1986). "Chromosome abnormalities in 1355 induced abortuses." *Hiroshima J Med Sci* 35(2): 135-41.
- Ohara, M., Y. Shimizu, et al. (2008). "Safety and usefulness of emergency maternal transport using helicopter." *J Obstet Gynaecol Res* 34(2): 189-94.
- Oida, T., Y. Sako, et al. (1993). "Fluorescence lifetime imaging microscopy (flimscopy). Methodology development and application to studies of endosome fusion in single cells." *Biophys J* 64(3): 676-85.
- Okigaki, M., C. Davis, et al. (2003). "Pyk2 regulates multiple signaling events crucial for macrophage morphology and migration." *Proc Natl Acad Sci U S A* 100(19): 10740-5.
- Oliver, A. D., C. B. Wilson, et al. (1973). "Transient postprandial paresis associated with arteriovenous malformations of the spinal cord. Report of two cases." *J Neurosurg* 39(5): 652-5.
- Oliver, J. E., T. J. Aitman, et al. (1989). "Insulin-like growth factor I gene expression in the rat ovary is confined to the granulosa cells of developing follicles." *Endocrinology* 124(6): 2671-9.
- Oliver, J. M., D. L. Burg, et al. (1994). "Inhibition of mast cell Fc epsilon R1-mediated signaling and effector function by the Syk-selective inhibitor, piceatannol." *J Biol Chem* 269(47): 29697-703.
- Oliver, J. M., C. L. Kepley, et al. (2000). "Immunologically mediated signaling in basophils and mast cells: finding therapeutic targets for allergic diseases in the human Fc epsilon R1 signaling pathway." *Immunopharmacology* 48(3): 269-81.
- Oliver, J. M., J. R. Pfeiffer, et al. (2004). "Membrane receptor mapping: the membrane topography of Fc(epsilon)RI signaling." *Subcell Biochem* 37: 3-34.
- Oliver, L. J., M. Keeton, et al. (1989). "Regulation and secretion of plasminogen activators and their inhibitors in a human leukemic cell line (K562)." *Blood* 74(4): 1321-7.
- Oliver, L. J., D. B. Rifkin, et al. (1990). "Long-term culture of human bone marrow stromal cells in the presence of basic fibroblast growth factor." *Growth Factors* 3(3): 231-6.
- Oliver, M. J., A. R. Wilson, et al. (1990). "Acute pancreatitis and gastric volvulus occurring in a congenital diaphragmatic hernia." *J Pediatr Surg* 25(12): 1240-1.

- Oliver, S. J., S. J. Laing, et al. (2007). "Endurance running performance after 48 h of restricted fluid and/or energy intake." *Med Sci Sports Exerc* 39(2): 316-22.
- Oliver, S. J., S. J. Laing, et al. (2008). "Saliva indices track hypohydration during 48h of fluid restriction or combined fluid and energy restriction." *Arch Oral Biol* 53(10): 975-80.
- Oliver, S. J., S. J. Laing, et al. (2007). "Salivary immunoglobulin A response at rest and after exercise following a 48 h period of fluid and/or energy restriction." *Br J Nutr* 97(6): 1109-16.
- Oliver, W. J., B. D. Graham, et al. (1958). "Lack of scientific validity of body surface as basis for parenteral fluid dosage." *J Am Med Assoc* 167(10): 1211-8.
- Olorundare, O. E., S. R. Simmons, et al. (1992). "Cytochalasin D and E: effects on fibrinogen receptor movement and cytoskeletal reorganization in fully spread, surface-activated platelets: a correlative light and electron microscopic investigation." *Blood* 79(1): 99-109.
- Ongchin, M., E. Sharratt, et al. (2009). "The Effects of Epidermal Growth Factor Receptor Activation and Attenuation of the TGFbeta Pathway in an Orthotopic Model of Colon Cancer." *J Surg Res*.
- Oradd, G., P. W. Westerman, et al. (2005). "Lateral diffusion coefficients of separate lipid species in a ternary raft-forming bilayer: a Pfg-NMR multinuclear study." *Biophys J* 89(1): 315-20.
- Orr, G., D. Hu, et al. (2005). "Cholesterol dictates the freedom of EGF receptors and HER2 in the plane of the membrane." *Biophys J* 89(2): 1362-73.
- Ortega, E., M. Lara, et al. (1999). "Lyn dissociation from phosphorylated Fc epsilon RI subunits: a new regulatory step in the Fc epsilon RI signaling cascade revealed by studies of Fc epsilon RI dimer signaling activity." *J Immunol* 162(1): 176-85.
- Ortega, F., J. L. Garces, et al. (2006). "Bistability from double phosphorylation in signal transduction. Kinetic and structural requirements." *FEBS J* 273(17): 3915-26.
- Orton, R., O. Sturm, et al. (2005). "Computational modelling of the receptor-tyrosine-kinase-activated MAPK pathway." *Biochem. J.* 392: 249-261.
- Ory, S. and D. K. Morrison (2004). "Signal transduction: implications for Ras-dependent ERK signaling." *Curr Biol* 14(7): R277-8.

- Ory, S., M. Zhou, et al. (2003). "Protein phosphatase 2A positively regulates Ras signaling by dephosphorylating KSR1 and Raf-1 on critical 14-3-3 binding sites." *Curr Biol* 13(16): 1356-64.
- Ota, I., D. Zoukhri, et al. (2003). "Alpha 1-adrenergic and cholinergic agonists activate MAPK by separate mechanisms to inhibit secretion in lacrimal gland." *Am J Physiol Cell Physiol* 284(1): C168-78.
- Ota, Y., T. Aoki, et al. (2008). "Determination of deep surgical margin based on anatomical architecture for local control of squamous cell carcinoma of the buccal mucosa." *Oral Oncol*.
- Ozaki, Y., S. Sasagawa, et al. (2005). "Dynamic characteristics of transient responses." *J Biochem* 137(6): 659-63.
- Ozcan, F., P. Klein, et al. (2006). "On the nature of low- and high-affinity EGF receptors on living cells." *Proc Natl Acad Sci U S A* 103(15): 5735-40.
- Paar, J. M., N. T. Harris, et al. (2002). "Bivalent ligands with rigid double-stranded DNA spacers reveal structural constraints on signaling by Fc epsilon RI." *J Immunol* 169(2): 856-64.
- Paller, A. S. (1994). "Histology of lipoid proteinosis." *JAMA* 272(7): 564-5.
- Paller, A. S. (1994). "Laboratory tests for ichthyosis." *Dermatol Clin* 12(1): 99-107.
- Paller, A. S., V. Nanda, et al. (1994). "Leukocyte adhesion deficiency: recurrent childhood skin infections." *J Am Acad Dermatol* 31(2 Pt 2): 316-9.
- Paller, A. S., A. J. Syder, et al. (1994). "Genetic and clinical mosaicism in a type of epidermal nevus." *N Engl J Med* 331(21): 1408-15.
- Paller, K. A. and A. R. Mayes (1994). "New-association priming of word identification in normal and amnesic subjects." *Cortex* 30(1): 53-73.
- Paller, M. S. (1994). "Bone marrow transplantation nephropathy." *J Lab Clin Med* 124(3): 315-7.
- Paller, M. S. (1994). "The cell biology of reperfusion injury in the kidney." *J Investig Med* 42(4): 632-9.
- Paller, M. S. (1994). "Lateral mobility of Na,K-ATPase and membrane lipids in renal cells. Importance of cytoskeletal integrity." *J Membr Biol* 142(1): 127-35.
- Paller, M. S. and E. L. Greene (1994). "Role of calcium in reperfusion injury of the kidney." *Ann N Y Acad Sci* 723: 59-70.

- Paller, M. S. and B. E. Hedlund (1994). "Extracellular iron chelators protect kidney cells from hypoxia/reoxygenation." *Free Radic Biol Med* 17(6): 597-603.
- Paller, M. S. and H. S. Jacob (1994). "Cytochrome P-450 mediates tissue-damaging hydroxyl radical formation during reoxygenation of the kidney." *Proc Natl Acad Sci U S A* 91(15): 7002-6.
- Pallera, A. M., J. B. Schweitzer, et al. (1994). "192 IgG-saporin causes a major loss of synaptic content in rat olfactory bulb." *Exp Neurol* 127(2): 265-77.
- Palleros, D. R., L. Shi, et al. (1994). "hsp70-protein complexes. Complex stability and conformation of bound substrate protein." *J Biol Chem* 269(18): 13107-14.
- Palmieri, S. J., T. Nebl, et al. (2000). "Mutant Rac1B expression in Dictyostelium: effects on morphology, growth, endocytosis, development, and the actin cytoskeleton." *Cell Motil Cytoskeleton* 46(4): 285-304.
- Pan, Z., T. Kao, et al. (2006). "A common ankyrin-G-based mechanism retains KCNQ and NaV channels at electrically active domains of the axon." *J Neurosci* 26(10): 2599-613.
- Park, D. S., H. Lee, et al. (2001). "Prolactin negatively regulates caveolin-1 gene expression in the mammary gland during lactation, via a Ras-dependent mechanism." *J Biol Chem* 276(51): 48389-97.
- Park, D. S., B. Razani, et al. (2001). "Evidence that Myc isoforms transcriptionally repress caveolin-1 gene expression via an INR-dependent mechanism." *Biochemistry* 40(11): 3354-62.
- Park, I. H., E. M. Hwang, et al. (2003). "Lovastatin enhances Abeta production and senile plaque deposition in female Tg2576 mice." *Neurobiol Aging* 24(5): 637-43.
- Park, S., K. J. Shin, et al. (2005). "Diffusion-influenced excited-state reversible transfer reactions, $A^* + \text{Bright harpoon} \rightarrow \text{left harpoon} + C^* + D$, with two different lifetimes: theories and simulations." *J Chem Phys* 123(3): 34507.
- Parolini, I., M. Sargiacomo, et al. (1999). "Expression of caveolin-1 is required for the transport of caveolin-2 to the plasma membrane. Retention of caveolin-2 at the level of the golgi complex." *J Biol Chem* 274(36): 25718-25.
- Parton, R. G. and J. F. Hancock (2001). "Caveolin and Ras function." *Methods Enzymol* 333: 172-83.
- Parton, R. G. and J. F. Hancock (2004). "Lipid rafts and plasma membrane microorganization: insights from Ras." *Trends Cell Biol* 14(3): 141-7.

- Pastore, J. J., M. Funaki, et al. (2005). "Flavonoid-mediated inhibition of actin polymerization in cold-activated platelets." *Platelets* 16(6): 362-7.
- Pathak, A., V. S. Deshpande, et al. (2008). "The simulation of stress fibre and focal adhesion development in cells on patterned substrates." *J R Soc Interface* 5(22): 507-24.
- Patterson, G. H., K. Hirschberg, et al. (2008). "Transport through the Golgi apparatus by rapid partitioning within a two-phase membrane system." *Cell* 133(6): 1055-67.
- Patterson, G. H. and J. Lippincott-Schwartz (2002). "A photoactivatable GFP for selective photolabeling of proteins and cells." *Science* 297(5588): 1873-7.
- Patterson, G. H. and J. Lippincott-Schwartz (2004). "Selective photolabeling of proteins using photoactivatable GFP." *Methods* 32(4): 445-50.
- Paul, J. T., E. S. Henson, et al. (2005). "Cyclin D expression in chronic lymphocytic leukemia." *Leuk Lymphoma* 46(9): 1275-85.
- Payne, E., M. R. Bowles, et al. (2001). "Human papillomavirus type 6b virus-like particles are able to activate the Ras-MAP kinase pathway and induce cell proliferation." *J Virol* 75(9): 4150-7.
- Pbert, L., S. K. Osganian, et al. (2006). "A school nurse-delivered adolescent smoking cessation intervention: a randomized controlled trial." *Prev Med* 43(4): 312-20.
- Pedersen, S. F., E. K. Hoffmann, et al. (2001). "The cytoskeleton and cell volume regulation." *Comp Biochem Physiol A Mol Integr Physiol* 130(3): 385-99.
- Pedraza-Chaverri, J., P. Calderon, et al. (1993). "Electrophoretic analysis of serum and urinary proteins in rats with aminonucleoside-induced nephrotic syndrome." *Ren Fail* 15(2): 149-55.
- Peletier, M. A., H. V. Westerhoff, et al. (2003). "Control of spatially heterogeneous and time-varying cellular reaction networks: a new summation law." *J Theor Biol* 225(4): 477-87.
- Pelham, H. R. and S. Munro (1993). "Sorting of membrane proteins in the secretory pathway." *Cell* 75(4): 603-5.
- Pentcheva, T. and M. Edidin (2001). "Clustering of peptide-loaded MHC class I molecules for endoplasmic reticulum export imaged by fluorescence resonance energy transfer." *J Immunol* 166(11): 6625-32.

- Perc, M., A. K. Green, et al. (2008). "Establishing the stochastic nature of intracellular calcium oscillations from experimental data." *Biophys Chem* 132(1): 33-8.
- Perez, T. D., M. Tamada, et al. (2008). "Immediate-early signaling induced by E-cadherin engagement and adhesion." *J Biol Chem* 283(8): 5014-22.
- Peters, R. and R. J. Cherry (1982). "Lateral and rotational diffusion of bacteriorhodopsin in lipid bilayers: experimental test of the Saffman-Delbruck equations." *Proc Natl Acad Sci U S A* 79(14): 4317-21.
- Petko, M., G. Veress, et al. (2004). "Commissural propriospinal connections between the lateral aspects of laminae III-IV in the lumbar spinal cord of rats." *J Comp Neurol* 480(4): 364-77.
- Petrasek, J., J. Mravec, et al. (2006). "PIN proteins perform a rate-limiting function in cellular auxin efflux." *Science* 312(5775): 914-8.
- Philipp, U., E. T. Roussos, et al. (2008). "A Mena invasion isoform potentiates EGF-induced carcinoma cell invasion and metastasis." *Dev Cell* 15(6): 813-28.
- Piessens, P. W., M. C. King, et al. (1995). "A statewide institute to deliver professional development programs to school health personnel in Massachusetts." *J Sch Health* 65(5): 176-80.
- Pimenta, W. P., I. M. Calderon, et al. (2004). "Subclinical abnormalities of glucose metabolism in Brazilian women with a history of gestational diabetes mellitus." *Acta Obstet Gynecol Scand* 83(12): 1152-8.
- Pimplikar, S. W., E. Ikonen, et al. (1994). "Basolateral protein transport in streptolysin O-permeabilized MDCK cells." *J Cell Biol* 125(5): 1025-35.
- Pincet, F. (2007). "Membrane recruitment of scaffold proteins drives specific signaling." *PLoS One* 2(10): e977.
- Planck, M., P. J. W. Debye, et al. (1914). *Vorträge über die kinetische theorie der materie und der elektrizität, gehalten in Göttingen auf einladung der Kommission der Wolfskehlstiftung. Leipzig, Berlin, B.G. Teubner.*
- Plowman, S. J., N. Ariotti, et al. (2008). "Electrostatic interactions positively regulate K-Ras nanocluster formation and function." *Mol Cell Biol* 28(13): 4377-85.
- Plowman, S. J. and J. F. Hancock (2005). "Ras signaling from plasma membrane and endomembrane microdomains." *Biochim Biophys Acta* 1746(3): 274-83.

- Plowman, S. J., C. Muncke, et al. (2005). "H-ras, K-ras, and inner plasma membrane raft proteins operate in nanoclusters with differential dependence on the actin cytoskeleton." *Proc Natl Acad Sci U S A* 102(43): 15500-5.
- Pohl, J., K. Pollmann, et al. (2004). "Antibiotic prophylaxis after variceal hemorrhage reduces incidence of early rebleeding." *Hepatogastroenterology* 51(56): 541-6.
- Pohl, J., A. Ring, et al. (2004). "New concepts of cellular fatty acid uptake: role of fatty acid transport proteins and of caveolae." *Proc Nutr Soc* 63(2): 259-62.
- Pohl, J., A. Ring, et al. (2004). "Long-chain fatty acid uptake into adipocytes depends on lipid raft function." *Biochemistry* 43(14): 4179-87.
- Pohl, J., A. Ring, et al. (2004). "Role of FATP in parenchymal cell fatty acid uptake." *Biochim Biophys Acta* 1686(1-2): 1-6.
- Pohl, J., A. Ring, et al. (2005). "FAT/CD36-mediated long-chain fatty acid uptake in adipocytes requires plasma membrane rafts." *Mol Biol Cell* 16(1): 24-31.
- Pons, T., I. L. Medintz, et al. (2007). "On the quenching of semiconductor quantum dot photoluminescence by proximal gold nanoparticles." *Nano Lett* 7(10): 3157-64.
- Pons, T., I. L. Medintz, et al. (2006). "Solution-phase single quantum dot fluorescence resonance energy transfer." *J Am Chem Soc* 128(47): 15324-31.
- Pons, T., H. T. Uyeda, et al. (2006). "Hydrodynamic dimensions, electrophoretic mobility, and stability of hydrophilic quantum dots." *J Phys Chem B* 110(41): 20308-16.
- Ponti, A., M. Machacek, et al. (2004). "Two distinct actin networks drive the protrusion of migrating cells." *Science* 305(5691): 1782-6.
- Ponti, A., A. Matov, et al. (2005). "Periodic patterns of actin turnover in lamellipodia and lamellae of migrating epithelial cells analyzed by quantitative Fluorescent Speckle Microscopy." *Biophys J* 89(5): 3456-69.
- Ponti, A., P. Vallotton, et al. (2003). "Computational analysis of F-actin turnover in cortical actin meshworks using fluorescent speckle microscopy." *Biophys J* 84(5): 3336-52.
- Poorthuis, B. J., T. P. van der Krift, et al. (1980). "Phospholipid transfer activities in Morris hepatomas and the specific contribution of the phosphatidylcholine exchange protein." *Biochim Biophys Acta* 600(2): 376-86.
- Popov, A. V., N. Agmon, et al. (2004). "Influence of diffusion on the kinetics of excited-state association--dissociation reactions: comparison of theory and simulation." *J Chem Phys* 120(13): 6111-6.

- Porfiri, E., T. Evans, et al. (1995). "Purification of baculovirus-expressed recombinant Ras and Rap proteins." *Methods Enzymol* 255: 13-21.
- Porfiri, E., T. Evans, et al. (1994). "Prenylation of Ras proteins is required for efficient hSOS1-promoted guanine nucleotide exchange." *J Biol Chem* 269(36): 22672-7.
- Porfiri, E. and J. F. Hancock (1995). "Stimulation of nucleotide exchange on Ras- and Rho-related proteins by small GTP-binding protein GDP dissociation stimulator." *Methods Enzymol* 256: 85-90.
- Porter, M. E., J. M. Scholey, et al. (1987). "Characterization of the microtubule movement produced by sea urchin egg kinesin." *J Biol Chem* 262(6): 2794-802.
- Powell, S. K., M. P. Lisanti, et al. (1991). "Thy-1 expresses two signals for apical localization in epithelial cells." *Am J Physiol* 260(4 Pt 1): C715-20.
- Presley, J. F., N. B. Cole, et al. (1997). "ER-to-Golgi transport visualized in living cells." *Nature* 389(6646): 81-5.
- Presley, J. F., C. Smith, et al. (1998). "Golgi membrane dynamics." *Mol Biol Cell* 9(7): 1617-26.
- Preston, R. J. (2002). "Quantitation of molecular endpoints for the dose-response component of cancer risk assessment." *Toxicol Pathol* 30(1): 112-6.
- Preston, R. J. (2005). "Extrapolations are the Achilles heel of risk assessment." *Mutat Res* 589(3): 153-7.
- Prior, I. A. and J. F. Hancock (2001). "Compartmentalization of Ras proteins." *J Cell Sci* 114(Pt 9): 1603-8.
- Prior, I. A., A. Harding, et al. (2001). "GTP-dependent segregation of H-ras from lipid rafts is required for biological activity." *Nat Cell Biol* 3(4): 368-75.
- Prior, I. A., C. Muncke, et al. (2003). "Direct visualization of Ras proteins in spatially distinct cell surface microdomains." *J Cell Biol* 160(2): 165-70.
- Puklin-Faucher, E. and M. P. Sheetz (2009). "The mechanical integrin cycle." *J Cell Sci* 122(Pt 2): 179-86.
- Pulcini, J., A. Sheetz, et al. (2008). "Establishing a practice-based research network: lessons from the Massachusetts experience." *J Sch Health* 78(3): 172-4.

- Puthanveettil, S. V., F. J. Monje, et al. (2008). "A new component in synaptic plasticity: upregulation of kinesin in the neurons of the gill-withdrawal reflex." *Cell* 135(5): 960-73.
- Qian, H. and J. A. Cooper (2008). "Temporal cooperativity and sensitivity amplification in biological signal transduction." *Biochemistry* 47(7): 2211-20.
- Qian, H., M. P. Sheetz, et al. (1991). "Single particle tracking. Analysis of diffusion and flow in two-dimensional systems." *Biophys J* 60(4): 910-21.
- Quilter, C. R., S. C. Blott, et al. (2007). "Porcine maternal infanticide as a model for puerperal psychosis." *Am J Med Genet B Neuropsychiatr Genet* 144B(7): 862-8.
- Quilter, C. R., C. L. Gilbert, et al. (2008). "Gene expression profiling in porcine maternal infanticide: a model for puerperal psychosis." *Am J Med Genet B Neuropsychiatr Genet* 147B(7): 1126-37.
- Radhakrishnan, K. (1991). "Combustion Kinetics and Sensitivity Analysis Computations," in *Numerical Approaches to Combustion Modeling*, ES.Oran and JP Boris." *AIAA J*: 83-128.
- Radhakrishnan, K. (1999). "LSENS: Multipurpose Kinetics and Sensitivity Analysis Code." *AIAA J* 41: 848-855.
- Radhakrishnan, K. and A. Hindmarsh (1993). "Description and Use of LSODE, the Livermore Solver for Ordinary Differential Equations." *NASA RP-1327*.
- Rainey, P. B., I. P. Thompson, et al. (1994). "Genome and fatty acid analysis of *Pseudomonas stutzeri*." *Int J Syst Bacteriol* 44(1): 54-61.
- Rajput, A., I. Dominguez San Martin, et al. (2008). "Characterization of HCT116 human colon cancer cells in an orthotopic model." *J Surg Res* 147(2): 276-81.
- Rajput, A., A. P. Koterba, et al. (2007). "A novel mechanism of resistance to epidermal growth factor receptor antagonism in vivo." *Cancer Res* 67(2): 665-73.
- Rao, P. N., H. Yu, et al. (1997). "Assignment of the human kinectin gene (KTN1), encoding a kinesin-binding protein, to chromosome 14 band q22.1 by in situ hybridization." *Cytogenet Cell Genet* 79(3-4): 196-7.
- Raucher, D. and M. P. Sheetz (1999). "Characteristics of a membrane reservoir buffering membrane tension." *Biophys J* 77(4): 1992-2002.
- Raucher, D. and M. P. Sheetz (1999). "Membrane expansion increases endocytosis rate during mitosis." *J Cell Biol* 144(3): 497-506.

- Raucher, D. and M. P. Sheetz (2000). "Cell spreading and lamellipodial extension rate is regulated by membrane tension." *J Cell Biol* 148(1): 127-36.
- Raucher, D. and M. P. Sheetz (2001). "Phospholipase C activation by anesthetics decreases membrane-cytoskeleton adhesion." *J Cell Sci* 114(Pt 20): 3759-66.
- Raucher, D., T. Stauffer, et al. (2000). "Phosphatidylinositol 4,5-bisphosphate functions as a second messenger that regulates cytoskeleton-plasma membrane adhesion." *Cell* 100(2): 221-8.
- Razani, B., T. P. Combs, et al. (2002). "Caveolin-1-deficient mice are lean, resistant to diet-induced obesity, and show hypertriglyceridemia with adipocyte abnormalities." *J Biol Chem* 277(10): 8635-47.
- Read, M., P. Small, et al. (2009). "Evaluating parent satisfaction of school nursing services." *J Sch Nurs* 25(3): 205-13.
- Reagan, D. R., B. N. Doebbeling, et al. (1991). "Elimination of coincident *Staphylococcus aureus* nasal and hand carriage with intranasal application of mupirocin calcium ointment." *Ann Intern Med* 114(2): 101-6.
- Refsnes, M., E. V. Thrane, et al. (2001). "Mechanisms in fluoride-induced interleukin-8 synthesis in human lung epithelial cells." *Toxicology* 167(2): 145-58.
- Reijenga, K. A., H. V. Westerhoff, et al. (2002). "Control analysis for autonomously oscillating biochemical networks." *Biophys J* 82(1 Pt 1): 99-108.
- Reinker, S., R. M. Altman, et al. (2006). "Parameter estimation in stochastic biochemical reactions." *Syst Biol (Stevenage)* 153(4): 168-78.
- Resat, H., J. A. Ewald, et al. (2003). "An integrated model of epidermal growth factor receptor trafficking and signal transduction." *Biophys J* 85(2): 730-43.
- Rheault, T. R., T. R. Caferro, et al. (2009). "Thienopyrimidine-based dual EGFR/ErbB-2 inhibitors." *Bioorg Med Chem Lett* 19(3): 817-20.
- Richman, J. G., M. Kanemitsu-Parks, et al. (2007). "Nicotinic acid receptor agonists differentially activate downstream effectors." *J Biol Chem* 282(25): 18028-36.
- Rios, J. D., D. Ferdman, et al. (2002). "Role of Ca²⁺ and protein kinase C in cholinergic, and alpha1-adrenergic agonists and EGF stimulated mitogen-activated protein kinase activity in lacrimal gland." *Adv Exp Med Biol* 506(Pt A): 185-90.
- Risselada, H. J., A. E. Mark, et al. (2008). "Application of mean field boundary potentials in simulations of lipid vesicles." *J Phys Chem B* 112(25): 7438-47.

- Risselada, H. J. and S. J. Marrink (2008). "The molecular face of lipid rafts in model membranes." *Proc Natl Acad Sci U S A* 105(45): 17367-72.
- Ritchie, K., R. Iino, et al. (2003). "The fence and picket structure of the plasma membrane of live cells as revealed by single molecule techniques (Review)." *Mol Membr Biol* 20(1): 13-8.
- Ritchie, K. and A. Kusumi (2003). "Single-particle tracking image microscopy." *Methods Enzymol* 360: 618-34.
- Ritchie, K. and A. Kusumi (2004). "Role of the membrane skeleton in creation of microdomains." *Subcell Biochem* 37: 233-45.
- Ritchie, K., X. Y. Shan, et al. (2005). "Detection of non-Brownian diffusion in the cell membrane in single molecule tracking." *Biophys J* 88(3): 2266-77.
- Ritt, D. A., I. O. Daar, et al. (2006). "KSR regulation of the Raf-MEK-ERK cascade." *Methods Enzymol* 407: 224-37.
- Ritt, D. A., M. Zhou, et al. (2007). "CK2 Is a component of the KSR1 scaffold complex that contributes to Raf kinase activation." *Curr Biol* 17(2): 179-84.
- Robillard, M. S., A. R. Valentijn, et al. (2000). "The First Solid-Phase Synthesis of a Peptide-Tethered Platinum(II) Complex This research was supported by the Council for Chemical Sciences of The Netherlands Organization for Scientific Research (CW-NWO) and by The Netherlands Foundation for Technical Sciences (STW). Support and sponsorship by COST Action D8/00097 (biocoordination chemistry) is kindly acknowledged. The authors thank Johnson & Matthey (Reading, UK) for their generous loan of K(2)PtCl(4)." *Angew Chem Int Ed Engl* 39(17): 3096-3099.
- Robles, N. R., J. Gonzalez-Reymundo, et al. (1996). "[Use of averaged concentration of urea and whole body urea clearance in prescribing the length of a dialysis session]." *An Med Interna* 13(1): 21-4.
- Rodriguez-Boulan, E., P. J. Salas, et al. (1989). "Methods to estimate the polarized distribution of surface antigens in cultured epithelial cells." *Methods Cell Biol* 32: 37-56.
- Roess, D. A., R. D. Horvat, et al. (2000). "Luteinizing hormone receptors are self-associated in the plasma membrane." *Endocrinology* 141(12): 4518-23.
- Rohwer, J. M., P. W. Postma, et al. (1998). "Implications of macromolecular crowding for signal transduction and metabolite channeling." *Proc Natl Acad Sci U S A* 95(18): 10547-52.

- Ron, D., C. H. Chen, et al. (1994). "Cloning of an intracellular receptor for protein kinase C: a homolog of the beta subunit of G proteins." *Proc Natl Acad Sci U S A* 91(3): 839-43.
- Rosenthal, P. B., X. Zhang, et al. (1998). "Structure of the haemagglutinin-esterase-fusion glycoprotein of influenza C virus." *Nature* 396(6706): 92-6.
- Rossman, J. S., N. G. Stoicheva, et al. (2006). "POLKADOTS are foci of functional interactions in T-Cell receptor-mediated signaling to NF-kappaB." *Mol Biol Cell* 17(5): 2166-76.
- Rotblat, B., I. A. Prior, et al. (2004). "Three separable domains regulate GTP-dependent association of H-ras with the plasma membrane." *Mol Cell Biol* 24(15): 6799-810.
- Rouger, K., J. P. Louboutin, et al. (2001). "X-linked vacuolated myopathy : TNF-alpha and IFN-gamma expression in muscle fibers with MHC class I on sarcolemma." *Am J Pathol* 158(2): 355-9.
- Rowlinson, S. W., H. Yoshizato, et al. (2008). "An agonist-induced conformational change in the growth hormone receptor determines the choice of signalling pathway." *Nat Cell Biol* 10(6): 740-7.
- Roy, R., S. Hohng, et al. (2008). "A practical guide to single-molecule FRET." *Nat Methods* 5(6): 507-16.
- Roy, S., A. Lane, et al. (1997). "Activity of plasma membrane-recruited Raf-1 is regulated by Ras via the Raf zinc finger." *J Biol Chem* 272(32): 20139-45.
- Roy, S., R. Luetterforst, et al. (1999). "Dominant-negative caveolin inhibits H-Ras function by disrupting cholesterol-rich plasma membrane domains." *Nat Cell Biol* 1(2): 98-105.
- Roy, S., R. A. McPherson, et al. (1998). "14-3-3 facilitates Ras-dependent Raf-1 activation in vitro and in vivo." *Mol Cell Biol* 18(7): 3947-55.
- Roy, S., S. Plowman, et al. (2005). "Individual palmitoyl residues serve distinct roles in H-ras trafficking, microlocalization, and signaling." *Mol Cell Biol* 25(15): 6722-33.
- Roy, S., B. Wyse, et al. (2002). "H-Ras signaling and K-Ras signaling are differentially dependent on endocytosis." *Mol Cell Biol* 22(14): 5128-40.
- Russell, T., J. M. Oliver, et al. (2008). "Differential expression of Ikaros isoforms in monozygotic twins with MLL-rearranged precursor-B acute lymphoblastic leukemia." *J Pediatr Hematol Oncol* 30(12): 941-4.

Sachs, K., S. Itani, et al. (2009). "Learning cyclic signaling pathway structures while minimizing data requirements." *Pac Symp Biocomput*: 63-74.

Sadir, R., A. Lambert, et al. (2001). "Caveolae and clathrin-coated vesicles: two possible internalization pathways for IFN-gamma and IFN-gamma receptor." *Cytokine* 14(1): 19-26.

Sadir, R., H. Lortat-Jacob, et al. (2000). "Internalization and nuclear translocation of IFN-gamma and IFN-gammaR: an ultrastructural approach." *Cytokine* 12(6): 711-4.

Saez-Rodriguez, J., A. Goldsipe, et al. (2008). "Flexible informatics for linking experimental data to mathematical models via DataRail." *Bioinformatics* 24(6): 840-7.

Saffman, P. G. and M. Delbruck (1975). "Brownian motion in biological membranes." *Proc Natl Acad Sci U S A* 72(8): 3111-3.

Sagawa, K., T. Kimura, et al. (1997). "The protein-tyrosine phosphatase SHP-2 associates with tyrosine-phosphorylated adhesion molecule PECAM-1 (CD31)." *J Biol Chem* 272(49): 31086-91.

Sagawa, K., W. Swaim, et al. (1997). "Aggregation of the high affinity IgE receptor results in the tyrosine phosphorylation of the surface adhesion protein PECAM-1 (CD31)." *J Biol Chem* 272(20): 13412-8.

Saito, N., Y. Usui, et al. (2008). "Carbon nanotubes for biomaterials in contact with bone." *Curr Med Chem* 15(5): 523-7.

Sakai, M., S. Aoki, et al. (2008). "Silent white matter lesion in linear scleroderma en coup de sabre." *J Comput Assist Tomogr* 32(5): 822-4.

Sakaki, Y., B. N. Kholodenko, et al. (2005). "The International Consortium on Systems Biology of Receptor Tyrosine Kinase Regulatory Networks." *Syst Biol (Stevenage)* 152(2): 53-4.

Sakamoto, H., K. Yoshimura, et al. (2008). "Genetic variation in PSCA is associated with susceptibility to diffuse-type gastric cancer." *Nat Genet* 40(6): 730-40.

Sako, Y., J. Ichinose, et al. (2003). "Optical bioimaging: from living tissue to a single molecule: single-molecule visualization of cell signaling processes of epidermal growth factor receptor." *J Pharmacol Sci* 93(3): 253-8.

Salani, B., L. Briatore, et al. (2008). "Caveolin-1 down-regulation inhibits insulin-like growth factor-I receptor signal transduction in H9C2 rat cardiomyoblasts." *Endocrinology* 149(2): 461-5.

Saltiel, A. R. and C. R. Kahn (2001). "Insulin signalling and the regulation of glucose and lipid metabolism." *Nature* 414(6865): 799-806.

Samuelson, J. T., P. E. Schwarze, et al. (2007). "Regulation of rat alveolar type 2 cell proliferation in vitro involves type II cAMP-dependent protein kinase." *Am J Physiol Lung Cell Mol Physiol* 292(1): L232-9.

Sanchez, F. A., D. D. Kim, et al. (2008). "Internalization of eNOS via caveolae regulates PAF-induced inflammatory hyperpermeability to macromolecules." *Am J Physiol Heart Circ Physiol* 295(4): H1642-8.

Santos, S. D., P. J. Verveer, et al. (2007). "Growth factor-induced MAPK network topology shapes Erk response determining PC-12 cell fate." *Nat Cell Biol* 9(3): 324-30.

Sapsford, K. E., I. L. Medintz, et al. (2004). "Surface-immobilized self-assembled protein-based quantum dot nanoassemblies." *Langmuir* 20(18): 7720-8.

Sargiacomo, M., M. Lisanti, et al. (1989). "Integral and peripheral protein composition of the apical and basolateral membrane domains in MDCK cells." *J Membr Biol* 107(3): 277-86.

Sasagawa, S., Y. Ozaki, et al. (2005). "Prediction and validation of the distinct dynamics of transient and sustained ERK activation." *Nat Cell Biol* 7(4): 365-73.

Sasagawa, S., T. Takabatake, et al. (2002). "Axes establishment during eye morphogenesis in *Xenopus* by coordinate and antagonistic actions of BMP4, Shh, and RA." *Genesis* 33(2): 86-96.

Sasagawa, T., M. Okita, et al. (1999). "Abnormal serum lysophospholipids in multiple myeloma patients." *Lipids* 34(1): 17-21.

Sasai, K., T. Akagi, et al. (2007). "O6-methylguanine-DNA methyltransferase is downregulated in transformed astrocyte cells: implications for anti-glioma therapies." *Mol Cancer* 6: 36.

Sasai, K., K. Kakumoto, et al. (2007). "The Ras-MAPK pathway downregulates Caveolin-1 in rodent fibroblast but not in human fibroblasts: implications in the resistance to oncogene-mediated transformation." *Oncogene* 26(3): 449-55.

Saunders, M. P., R. Wilson, et al. (2009). "Vandetanib with FOLFIRI in patients with advanced colorectal adenocarcinoma: results from an open-label, multicentre Phase I study." *Cancer Chemother Pharmacol*.

Sauro, H. M. and B. N. Kholodenko (2004). "Quantitative analysis of signaling networks." *Prog Biophys Mol Biol* 86(1): 5-43.

- Sawada, Y., K. Nakamura, et al. (2001). "Rap1 is involved in cell stretching modulation of p38 but not ERK or JNK MAP kinase." *J Cell Sci* 114(Pt 6): 1221-7.
- Sawada, Y. and M. P. Sheetz (2002). "Force transduction by Triton cytoskeletons." *J Cell Biol* 156(4): 609-15.
- Sawada, Y., M. Tamada, et al. (2006). "Force sensing by mechanical extension of the Src family kinase substrate p130Cas." *Cell* 127(5): 1015-26.
- Saxton, M. J. and K. Jacobson (1997). "Single-particle tracking: applications to membrane dynamics." *Annu Rev Biophys Biomol Struct* 26: 373-99.
- Schad, A., K. Schindler, et al. (2008). "Application of a multivariate seizure detection and prediction method to non-invasive and intracranial long-term EEG recordings." *Clin Neurophysiol* 119(1): 197-211.
- Schafer, D. A., J. Gelles, et al. (1991). "Transcription by single molecules of RNA polymerase observed by light microscopy." *Nature* 352(6334): 444-8.
- Schaff, J., C. C. Fink, et al. (1997). "A general computational framework for modeling cellular structure and function." *Biophys J* 73(3): 1135-46.
- Schaid, D. J., S. K. McDonnell, et al. (2006). "Pooled genome linkage scan of aggressive prostate cancer: results from the International Consortium for Prostate Cancer Genetics." *Hum Genet* 120(4): 471-85.
- Schechtman, D. and D. Mochly-Rosen (2001). "Adaptor proteins in protein kinase C-mediated signal transduction." *Oncogene* 20(44): 6339-47.
- Schechtman, D. and D. Mochly-Rosen (2002). "Isozyme-specific inhibitors and activators of protein kinase C." *Methods Enzymol* 345: 470-89.
- Scheel-Toellner, D., K. Wang, et al. (2004). "Clustering of death receptors in lipid rafts initiates neutrophil spontaneous apoptosis." *Biochem Soc Trans* 32(Pt 5): 679-81.
- Scheel-Toellner, D., K. Wang, et al. (2004). "Reactive oxygen species limit neutrophil life span by activating death receptor signaling." *Blood* 104(8): 2557-64.
- Scheel-Toellner, D., K. Q. Wang, et al. (2004). "Early events in spontaneous neutrophil apoptosis." *Biochem Soc Trans* 32(Pt3): 461-4.
- Scheiffele, P., P. Verkade, et al. (1998). "Caveolin-1 and -2 in the exocytic pathway of MDCK cells." *J Cell Biol* 140(4): 795-806.

- Schein, S. (2009). "Architecture of clathrin fullerene cages reflects a geometric constraint--the head-to-tail exclusion rule--and a preference for asymmetry." *J Mol Biol* 387(2): 363-75.
- Schelhaas, M., H. Ewers, et al. (2008). "Human papillomavirus type 16 entry: retrograde cell surface transport along actin-rich protrusions." *PLoS Pathog* 4(9): e1000148.
- Schenk, D., R. Barbour, et al. (1999). "Immunization with amyloid-beta attenuates Alzheimer-disease-like pathology in the PDAPP mouse." *Nature* 400(6740): 173-7.
- Scheving, L. A., R. Buchanan, et al. (2007). "Dexamethasone modulates ErbB tyrosine kinase expression and signaling through multiple and redundant mechanisms in cultured rat hepatocytes." *Am J Physiol Gastrointest Liver Physiol* 293(3): G552-9.
- Scheving, L. A., M. C. Stevenson, et al. (2002). "Integral role of the EGF receptor in HGF-mediated hepatocyte proliferation." *Biochem Biophys Res Commun* 290(1): 197-203.
- Scheving, L. A., L. Zhang, et al. (2006). "The emergence of ErbB2 expression in cultured rat hepatocytes correlates with enhanced and diversified EGF-mediated signaling." *Am J Physiol Gastrointest Liver Physiol* 291(1): G16-25.
- Schindler, M., D. E. Koppel, et al. (1980). "Modulation of membrane protein lateral mobility by polyphosphates and polyamines." *Proc Natl Acad Sci U S A* 77(3): 1457-61.
- Schlessinger, J., I. Lax, et al. (1995). "Regulation of growth factor activation by proteoglycans: what is the role of the low affinity receptors?" *Cell* 83(3): 357-60.
- Schlessinger, J. and M. A. Lemmon (2003). "SH2 and PTB domains in tyrosine kinase signaling." *Sci STKE* 2003(191): RE12.
- Schlessinger, J. and M. A. Lemmon (2006). "Nuclear signaling by receptor tyrosine kinases: the first robin of spring." *Cell* 127(1): 45-8.
- Schmidt, C. E., A. F. Horwitz, et al. (1993). "Integrin-cytoskeletal interactions in migrating fibroblasts are dynamic, asymmetric, and regulated." *J Cell Biol* 123(4): 977-91.
- Schnapp, B. J., B. Crise, et al. (1990). "Delayed start-up of kinesin-driven microtubule gliding following inhibition by adenosine 5'-[beta,gamma-imido]triphosphate." *Proc Natl Acad Sci U S A* 87(24): 10053-7.
- Schnapp, B. J., J. Gelles, et al. (1988). "Nanometer-scale measurements using video light microscopy." *Cell Motil Cytoskeleton* 10(1-2): 47-53.

- Schnapp, B. J., R. D. Vale, et al. (1985). "Single microtubules from squid axoplasm support bidirectional movement of organelles." *Cell* 40(2): 455-62.
- Schnapp, B. J., R. D. Vale, et al. (1986). "Microtubules and the mechanism of directed organelle movement." *Ann N Y Acad Sci* 466: 909-18.
- Schoeberl, B., C. Eichler-Jonsson, et al. (2002). "Computational modeling of the dynamics of the MAP kinase cascade activated by surface and internalized EGF receptors." *Nat Biotechnol* 20(4): 370-5.
- Schoeberl, B., E. Pace, et al. (2006). "A data-driven computational model of the ErbB receptor signaling network." *Conf Proc IEEE Eng Med Biol Soc* 1: 53-4.
- Schroder, K., P. J. Hertzog, et al. (2004). "Interferon-gamma: an overview of signals, mechanisms and functions." *J Leukoc Biol* 75(2): 163-89.
- Schroder, W. A., M. Buck, et al. (2007). "Human Sin1 contains Ras-binding and pleckstrin homology domains and suppresses Ras signalling." *Cell Signal* 19(6): 1279-89.
- Schroeder, F., A. M. Gallegos, et al. (2001). "Recent advances in membrane microdomains: rafts, caveolae, and intracellular cholesterol trafficking." *Exp Biol Med (Maywood)* 226(10): 873-90.
- Schroeder, R. J., S. N. Ahmed, et al. (1998). "Cholesterol and sphingolipid enhance the Triton X-100 insolubility of glycosylphosphatidylinositol-anchored proteins by promoting the formation of detergent-insoluble ordered membrane domains." *J Biol Chem* 273(2): 1150-7.
- Schroer, T. A., B. J. Schnapp, et al. (1988). "The role of kinesin and other soluble factors in organelle movement along microtubules." *J Cell Biol* 107(5): 1785-92.
- Schroer, T. A. and M. P. Sheetz (1991). "Functions of microtubule-based motors." *Annu Rev Physiol* 53: 629-52.
- Schroer, T. A. and M. P. Sheetz (1991). "Two activators of microtubule-based vesicle transport." *J Cell Biol* 115(5): 1309-18.
- Schroer, T. A., E. R. Steuer, et al. (1989). "Cytoplasmic dynein is a minus end-directed motor for membranous organelles." *Cell* 56(6): 937-46.
- Schubert, W., P. G. Frank, et al. (2001). "Caveolae-deficient endothelial cells show defects in the uptake and transport of albumin in vivo." *J Biol Chem* 276(52): 48619-22.

- Schulman, J. D., L. Corash, et al. (1979). "Reduced chronic hemolysis in Mediterranean glucose-6-phosphate dehydrogenase deficiency after vitamin E therapy." *Prog Clin Biol Res* 34: 381-7.
- Schuster, S., B. N. Kholodenko, et al. (2000). "Cellular information transfer regarded from a stoichiometry and control analysis perspective." *Biosystems* 55(1-3): 73-81.
- Schwarze, P. E., N. M. Johnsen, et al. (1996). "The use of isolated lung cells in in vitro pulmonary toxicology: studies of DNA damage, apoptosis and alteration of gene expression." *Cent Eur J Public Health* 4 Suppl: 6-10.
- Schwarze, P. E., M. Lag, et al. (2000). "Role of signal transduction pathways in lung inflammatory responses." *Toxicol Lett* 112-113: 165-70.
- Sciaky, N., J. Presley, et al. (1997). "Golgi tubule traffic and the effects of brefeldin A visualized in living cells." *J Cell Biol* 139(5): 1137-55.
- Scott, J. D. and T. Pawson (2000). "Cell communication: the inside story." *Sci Am* 282(6): 72-9.
- Scott, J. P., S. F. Oliver, et al. (2006). "Practical asymmetric synthesis of a gamma-secretase inhibitor exploiting substrate-controlled intramolecular nitrile oxide-olefin cycloaddition." *J Org Chem* 71(8): 3086-92.
- Seabra, M. C., M. S. Brown, et al. (1992). "Purification of component A of Rab geranylgeranyl transferase: possible identity with the choroideremia gene product." *Cell* 70(6): 1049-57.
- Sekar, R. B. and A. Periasamy (2003). "Fluorescence resonance energy transfer (FRET) microscopy imaging of live cell protein localizations." *J Cell Biol* 160(5): 629-33.
- Sellers, J. R., J. A. Spudich, et al. (1985). "Light chain phosphorylation regulates the movement of smooth muscle myosin on actin filaments." *J Cell Biol* 101(5 Pt 1): 1897-902.
- Sengupta, K., H. Aranda-Espinoza, et al. (2006). "Spreading of neutrophils: from activation to migration." *Biophys J* 91(12): 4638-48.
- Seubert, P., R. Barbour, et al. (2008). "Antibody capture of soluble Abeta does not reduce cortical Abeta amyloidosis in the PDAPP mouse." *Neurodegener Dis* 5(2): 65-71.
- Shah, J. V., L. A. Flanagan, et al. (2000). "Bidirectional translocation of neurofilaments along microtubules mediated in part by dynein/dynactin." *Mol Biol Cell* 11(10): 3495-508.

- Shah, J. V., L. Z. Wang, et al. (1998). "Interaction of vimentin with actin and phospholipids." *Biol Bull* 194(3): 402-5.
- Shalin, S. C., C. M. Hernandez, et al. (2006). "Kinase suppressor of Ras1 compartmentalizes hippocampal signal transduction and subserves synaptic plasticity and memory formation." *Neuron* 50(5): 765-79.
- Shalom-Feuerstein, R., S. J. Plowman, et al. (2008). "K-ras nanoclustering is subverted by overexpression of the scaffold protein galectin-3." *Cancer Res* 68(16): 6608-16.
- Shamash, J., G. Dancey, et al. (2005). "Chlorambucil and lomustine (CL56) in absolute hormone refractory prostate cancer: re-induction of endocrine sensitivity an unexpected finding." *Br J Cancer* 92(1): 36-40.
- Shamash, J., A. Davies, et al. (2008). "A phase II study investigating the re-induction of endocrine sensitivity following chemotherapy in androgen-independent prostate cancer." *Br J Cancer* 98(1): 22-4.
- Shamash, J., T. Powles, et al. (2007). "GAMEC--a new intensive protocol for untreated poor prognosis and relapsed or refractory germ cell tumours." *Br J Cancer* 97(3): 308-14.
- Shamash, J., T. Powles, et al. (2007). "A phase II study using a topoisomerase I-based approach in patients with multiply relapsed germ-cell tumours." *Ann Oncol* 18(5): 925-30.
- Shamash, J., T. Powles, et al. (2005). "Prognostic factors in cytokine-refractory renal cell carcinoma treated with irinotecan, Cisplatin, and mitomycin chemotherapy." *J Clin Oncol* 23(6): 1323-5.
- Shamash, J., J. P. Steele, et al. (2003). "IPM chemotherapy in cytokine refractory renal cell cancer." *Br J Cancer* 88(10): 1516-21.
- Shankaran, H., H. Resat, et al. (2007). "Cell surface receptors for signal transduction and ligand transport: a design principles study." *PLoS Comput Biol* 3(6): e101.
- Shankaran, H., H. S. Wiley, et al. (2006). "Modeling the effects of HER/ErbB1-3 coexpression on receptor dimerization and biological response." *Biophys J* 90(11): 3993-4009.
- Shankaran, H., H. S. Wiley, et al. (2007). "Receptor downregulation and desensitization enhance the information processing ability of signalling receptors." *BMC Syst Biol* 1: 48.
- Shatos, M. A., J. D. Rios, et al. (2001). "Isolation, characterization, and propagation of rat conjunctival goblet cells in vitro." *Invest Ophthalmol Vis Sci* 42(7): 1455-64.

Shaw, G. L., P. Wilson, et al. (2007). "International study into the use of intermittent hormone therapy in the treatment of carcinoma of the prostate: a meta-analysis of 1446 patients." *BJU Int* 99(5): 1056-65.

Sheetz, A. H. (2003). "Developing school health services in Massachusetts: a public health model." *J Sch Nurs* 19(4): 204-11.

Sheetz, A. H. and M. S. Blum (1998). "Medication administration in schools: the Massachusetts experience." *J Sch Health* 68(3): 94-8.

Sheetz, A. H., P. G. Goldman, et al. (2004). "Guidelines for managing life-threatening food allergies in Massachusetts schools." *J Sch Health* 74(5): 155-60.

Sheetz, A. H. and C. L. McIntyre (2005). "Anaphylaxis experienced by school children offers opportunities for ED nurse, school nurse collaboration." *J Emerg Nurs* 31(1): 102-4; quiz 122.

Sheetz, D. A. (1997). "Caffeine and chronic back pain." *Arch Phys Med Rehabil* 78(7): 786.

Sheetz, J. H. and L. Menaker (1979). "Morphological and functional study of the effect of isoproterenol on salivary gland cells." *Cell Tissue Res* 203(2): 321-9.

Sheetz, J. H., A. H. Morgan, et al. (1983). "Morphological and biochemical changes in the rat parotid gland after compensatory and isoproterenol-induced enlargement." *Arch Oral Biol* 28(5): 441-5.

Sheetz, K. and P. J. Lynch (1991). "Ichthyosis and dermatophyte fungal infection." *J Am Acad Dermatol* 24(2 Pt 1): 321.

Sheetz, K. E., E. E. Hoover, et al. (2008). "Advancing multifocal nonlinear microscopy: development and application of a novel multibeam Yb:KGd(WO₄)₂ oscillator." *Opt Express* 16(22): 17574-84.

Sheetz, K. K., A. T. Bishop, et al. (1995). "The arterial blood supply of the distal radius and ulna and its potential use in vascularized pedicled bone grafts." *J Hand Surg [Am]* 20(6): 902-14.

Sheetz, M., E. Elson, et al. (1990). "To flow or not to flow?" *Nature* 345(6270): 28.

Sheetz, M., D. Swanson, et al. (2009). "Physical presence during gamma stereotactic radiosurgery." *Health Phys* 96(2 Suppl): S11-5.

Sheetz, M. J. (1995). "MDL 29311, a phenolic antioxidant, interferes with the interaction of apoC with VLDL: a possible explanation for its triglyceride-lowering effect." *J Lipid Res* 36(12): 2609-21.

- Sheetz, M. J., R. L. Barnhart, et al. (1994). "MDL 29311, an analog of probucol, decreases triglycerides in rats by increasing hepatic clearance of very-low-density lipoprotein." *Metabolism* 43(2): 233-40.
- Sheetz, M. J. and M. A. Bowman (2008). "Pediatric palliative care: an assessment of physicians' confidence in skills, desire for training, and willingness to refer for end-of-life care." *Am J Hosp Palliat Care* 25(2): 100-5.
- Sheetz, M. J. and G. L. King (2002). "Molecular understanding of hyperglycemia's adverse effects for diabetic complications." *JAMA* 288(20): 2579-88.
- Sheetz, M. J. and H. S. Tager (1988). "Characterization of a glucagon receptor-linked protease from canine hepatic plasma membranes. Partial purification, kinetic analysis, and determination of sites for hormone processing." *J Biol Chem* 263(35): 19210-7.
- Sheetz, M. J. and H. S. Tager (1988). "Receptor-linked proteolysis of membrane-bound glucagon yields a membrane-associated hormone fragment." *J Biol Chem* 263(17): 8509-14.
- Sheetz, M. P. (1979). "DNase-I-dependent dissociation of erythrocyte cytoskeletons." *J Cell Biol* 81(1): 266-70.
- Sheetz, M. P. (1979). "Integral membrane protein interaction with Triton cytoskeletons of erythrocytes." *Biochim Biophys Acta* 557(1): 122-34.
- Sheetz, M. P. (1983). "Membrane skeletal dynamics: role in modulation of red cell deformability, mobility of transmembrane proteins, and shape." *Semin Hematol* 20(3): 175-88.
- Sheetz, M. P. (1987). "What are the functions of kinesin?" *Bioessays* 7(4): 165-8.
- Sheetz, M. P. (1988). "Muscle-bound bacteria and weak worms." *Nature* 331(6153): 212-3.
- Sheetz, M. P. (1992). "Can indirect cost funding be improved?" *FASEB J* 6(14): 3234-5.
- Sheetz, M. P. (1993). "Glycoprotein motility and dynamic domains in fluid plasma membranes." *Annu Rev Biophys Biomol Struct* 22: 417-31.
- Sheetz, M. P. (1994). "Cell migration by graded attachment to substrates and contraction." *Semin Cell Biol* 5(3): 149-55.
- Sheetz, M. P. (1995). "Cellular plasma membrane domains." *Mol Membr Biol* 12(1): 89-91.

- Sheetz, M. P. (1996). "Microtubule motor complexes moving membranous organelles." *Cell Struct Funct* 21(5): 369-73.
- Sheetz, M. P. (1998). "Laser tweezers in cell biology. Introduction." *Methods Cell Biol* 55: xi-xii.
- Sheetz, M. P. (1999). "Motor and cargo interactions." *Eur J Biochem* 262(1): 19-25.
- Sheetz, M. P. (2001). "Cell control by membrane-cytoskeleton adhesion." *Nat Rev Mol Cell Biol* 2(5): 392-6.
- Sheetz, M. P. and E. Alhanaty (1983). "Bilayer sensor model of erythrocyte shape control." *Ann N Y Acad Sci* 416: 58-65.
- Sheetz, M. P., N. L. Baumrind, et al. (1990). "Concentration of membrane antigens by forward transport and trapping in neuronal growth cones." *Cell* 61(2): 231-41.
- Sheetz, M. P., S. M. Block, et al. (1986). "Myosin movement in vitro: a quantitative assay using oriented actin cables from *Nitella*." *Methods Enzymol* 134: 531-44.
- Sheetz, M. P. and J. Casaly (1980). "2,3-Diphosphoglycerate and ATP dissociate erythrocyte membrane skeletons." *J Biol Chem* 255(20): 9955-60.
- Sheetz, M. P. and J. Casaly (1981). "Phosphate metabolite regulation of spectrin interactions." *Scand J Clin Lab Invest Suppl* 156: 117-22.
- Sheetz, M. P., R. Chasan, et al. (1984). "ATP-dependent movement of myosin in vitro: characterization of a quantitative assay." *J Cell Biol* 99(5): 1867-71.
- Sheetz, M. P. and J. Dai (1996). "Modulation of membrane dynamics and cell motility by membrane tension." *Trends Cell Biol* 6(3): 85-9.
- Sheetz, M. P., P. Febroriello, et al. (1982). "Triphosphoinositide increases glycoprotein lateral mobility in erythrocyte membranes." *Nature* 296(5852): 91-3.
- Sheetz, M. P., D. Felsenfeld, et al. (1999). "Cell migration as a five-step cycle." *Biochem Soc Symp* 65: 233-43.
- Sheetz, M. P., D. P. Felsenfeld, et al. (1998). "Cell migration: regulation of force on extracellular-matrix-integrin complexes." *Trends Cell Biol* 8(2): 51-4.
- Sheetz, M. P. and D. E. Koppel (1979). "Membrane damage caused by irradiation of fluorescent concanavalin A." *Proc Natl Acad Sci U S A* 76(7): 3314-7.
- Sheetz, M. P. and S. C. Kuo (1993). "Tracking nanometer movements of single motor molecules." *Methods Cell Biol* 39: 129-36.

- Sheetz, M. P. and C. H. Martenson (1991). "Axonal transport: beyond kinesin and cytoplasmic dynein." *Curr Opin Neurobiol* 1(3): 393-8.
- Sheetz, M. P., K. K. Pfister, et al. (1998). "Mechanisms of trafficking in axons and dendrites: implications for development and neurodegeneration." *Prog Neurobiol* 55(6): 577-94.
- Sheetz, M. P., J. E. Sable, et al. (2006). "Continuous membrane-cytoskeleton adhesion requires continuous accommodation to lipid and cytoskeleton dynamics." *Annu Rev Biophys Biomol Struct* 35: 417-34.
- Sheetz, M. P. and D. Sawyer (1978). "Triton shells of intact erythrocytes." *J Supramol Struct* 8(4): 399-412.
- Sheetz, M. P., D. Sawyer, et al. (1978). "The ATP-dependent red cell membrane shape change: a molecular explanation." *Prog Clin Biol Res* 21: 431-52.
- Sheetz, M. P., M. Schindler, et al. (1980). "Lateral mobility of integral membrane proteins is increased in spherocytic erythrocytes." *Nature* 285(5765): 510-1.
- Sheetz, M. P. and S. J. Singer (1977). "On the mechanism of ATP-induced shape changes in human erythrocyte membranes. I. The role of the spectrin complex." *J Cell Biol* 73(3): 638-46.
- Sheetz, M. P. and J. A. Spudich (1983). "Movement of myosin-coated fluorescent beads on actin cables in vitro." *Nature* 303(5912): 31-5.
- Sheetz, M. P. and J. A. Spudich (1983). "Movement of myosin-coated structures on actin cables." *Cell Motil* 3(5-6): 485-9.
- Sheetz, M. P., E. R. Steuer, et al. (1989). "The mechanism and regulation of fast axonal transport." *Trends Neurosci* 12(11): 474-8.
- Sheetz, M. P., S. Turney, et al. (1989). "Nanometre-level analysis demonstrates that lipid flow does not drive membrane glycoprotein movements." *Nature* 340(6231): 284-8.
- Sheetz, M. P., R. Vale, et al. (1986). "Vesicle movements and microtubule-based motors." *J Cell Sci Suppl* 5: 181-8.
- Sheetz, M. P., R. Vale, et al. (1987). "Movements of vesicles on microtubules." *Ann N Y Acad Sci* 493: 409-16.
- Sheetz, M. P., W. P. Wang, et al. (1984). "Polyphosphoinositides as regulators of membrane skeletal stability." *Kroc Found Ser* 16: 87-94.

- Sheetz, M. P., D. B. Wayne, et al. (1992). "Extension of filopodia by motor-dependent actin assembly." *Cell Motil Cytoskeleton* 22(3): 160-9.
- Sheetz, R. M. and R. C. Dickson (1980). "Mutations affecting synthesis of beta-galactosidase activity in the yeast *Kluyveromyces lactis*." *Genetics* 95(4): 877-90.
- Sheetz, R. M. and R. C. Dickson (1981). "Lac4 is the structural gene for beta-galactosidase in *Kluyveromyces lactis*." *Genetics* 98(4): 729-45.
- Shenderov, A. D. and M. P. Sheetz (1997). "Inversely correlated cycles in speed and turning in an ameba: an oscillatory model of cell locomotion." *Biophys J* 72(5): 2382-9.
- Shepard, K. A., A. P. Gerber, et al. (2003). "Widespread cytoplasmic mRNA transport in yeast: identification of 22 bud-localized transcripts using DNA microarray analysis." *Proc Natl Acad Sci U S A* 100(20): 11429-34.
- Shetty, S., J. B. Gladden, et al. (2002). "Tumor necrosis factor-related apoptosis inducing ligand (TRAIL) up-regulates death receptor 5 (DR5) mediated by NFkappaB activation in epithelial derived cell lines." *Apoptosis* 7(5): 413-20.
- Shi, L., D. R. Palleros, et al. (1994). "Protein conformational changes induced by 1,1'-bis(4-anilino-5-naphthalenesulfonic acid): preferential binding to the molten globule of DnaK." *Biochemistry* 33(24): 7536-46.
- Shim, J., T. F. Bersano-Begey, et al. (2003). "Micro- and nanotechnologies for studying cellular function." *Curr Top Med Chem* 3(6): 687-703.
- Shinar, J., R. Shinar, et al. (1989). "Nature of long-range atomic H motion in a-Si:H." *Phys Rev Lett* 62(17): 2001-2004.
- Shrank, W. H., S. M. Asch, et al. (2006). "Physicians' perceived knowledge of and responsibility for managing patients' out-of-pocket costs for prescription drugs." *Ann Pharmacother* 40(9): 1534-40.
- Sil, D., J. B. Lee, et al. (2007). "Trivalent ligands with rigid DNA spacers reveal structural requirements for IgE receptor signaling in RBL mast cells." *ACS Chem Biol* 2(10): 674-84.
- Simmons, P. J., B. Masinovsky, et al. (1992). "Vascular cell adhesion molecule-1 expressed by bone marrow stromal cells mediates the binding of hematopoietic progenitor cells." *Blood* 80(2): 388-95.
- Simon, V. R., S. L. Karmon, et al. (1997). "Mitochondrial inheritance: cell cycle and actin cable dependence of polarized mitochondrial movements in *Saccharomyces cerevisiae*." *Cell Motil Cytoskeleton* 37(3): 199-210.

- Simon, V. R. and L. A. Pon (1996). "Actin-based organelle movement." *Experientia* 52(12): 1117-22.
- Simon, V. R., T. C. Swayne, et al. (1995). "Actin-dependent mitochondrial motility in mitotic yeast and cell-free systems: identification of a motor activity on the mitochondrial surface." *J Cell Biol* 130(2): 345-54.
- Simons, K. and E. Ikonen (1997). "Functional rafts in cell membranes." *Nature* 387(6633): 569-72.
- Simons, K. and E. Ikonen (2000). "How cells handle cholesterol." *Science* 290(5497): 1721-6.
- Simons, K. and D. Toomre (2000). "Lipid rafts and signal transduction." *Nat Rev Mol Cell Biol* 1(1): 31-9.
- Simons, M., E. Ikonen, et al. (1995). "Intracellular routing of human amyloid protein precursor: axonal delivery followed by transport to the dendrites." *J Neurosci Res* 41(1): 121-8.
- Simonsz, H. J., R. A. Crone, et al. (1985). "Bielschowsky head-tilt test--I. Ocular counterrolling and Bielschowsky head-tilt test in 23 cases of superior oblique palsy." *Vision Res* 25(12): 1977-82.
- Sims, T. N., T. J. Soos, et al. (2007). "Opposing effects of PKC θ and WASp on symmetry breaking and relocation of the immunological synapse." *Cell* 129(4): 773-85.
- Singer, S. J. (1972). "A fluid lipid-globular protein mosaic model of membrane structure." *Ann N Y Acad Sci* 195: 16-23.
- Singer, S. J. and G. L. Nicolson (1972). "The fluid mosaic model of the structure of cell membranes." *Science* 175(23): 720-31.
- Skelly, J. V., D. A. Suter, et al. (1990). "Conformational effects of nucleotide exchange in ras p21 proteins as studied by fluorescence spectroscopy." *FEBS Lett* 262(1): 127-30.
- Smith, A. M., G. Ruan, et al. (2006). "Engineering luminescent quantum dots for in vivo molecular and cellular imaging." *Ann Biomed Eng* 34(1): 3-14.
- Smith, M. G., V. R. Simon, et al. (1995). "Organelle-cytoskeletal interactions: actin mutations inhibit meiosis-dependent mitochondrial rearrangement in the budding yeast *Saccharomyces cerevisiae*." *Mol Biol Cell* 6(10): 1381-96.

- Smith, R. A., V. Cokkinides, et al. (2008). "Cancer Screening in the United States, 2008: A Review of Current American Cancer Society Guidelines and Cancer Screening Issues." *CA Cancer J Clin*.
- Smoluchowski (1917). "Versuch einer mathematischen Theorie der Koagulationskinetik kolloider Lösungen." *Zeitschrift für Physikalische Chemie* 92: 129-168.
- Smoluchowski, A. (1949). "[History of medicine.]." *Zdrow Publiczne* 65(3-4): 86-9.
- Smoluchowski, A. (1949). "[Institute of scientific medical publications.]." *Zdrav Rev Vestn Minist Zdrav* 24(6-7): 140.
- Smoluchowski, A. (1949). "[Institute of scientific medical publications.]." *Zdrav Rev Vestn Minist Zdrav* 24(6-7): 140.
- Smoluchowski, A. (1949). "[Medical Institute of Scientific Publications.]." *Zdrow Publiczne* 65(3-4): 103.
- Smoluchowski, A. (1951). "[Principles of tissue therapy.]." *Med Weter* 7(1): 31-3.
- Smoluchowski, I. (1992). *Brussels Griffon champions, 1952-1990*. Kings Beach, CA, Camino Book Co.
- Smoluchowski, L., L. Tolstoy, et al. (1988). *Lev and Sonya : the story of the Tolstoy marriage*. New York, Paragon.
- Smoluchowski, M. (1924). *Pisma. Z polecenia Polskiej Akademji Umiej*etno*sci zgromadzone i wydane przez W*adys*awa Natansona i Jana Stocka*. W Krakowie,, Druk. Uniwersytetu Jagiello*nskigo.
- Smoluchowski, M. and R. Fürth (1923). *Abhandlungen über die Brownsche bewegung und verwandte erscheinungen*. Leipzig,, Akademische verlagsgesellschaft m.b.h.
- Smoluchowski, M. and W. a. a. Krajewski (1956). *Wybór pism filozoficznych*. Warszawa, Pa*ństwowe Wydawn. Naukowe.
- Smoluchowski, M. V. (1917). "Versuch einer mathematischen Theorie der Koagulationskinetik kolloider Lösungen." *Zeitschrift für Physikalische Chemie* 92: 129-168.
- Smoluchowski, R. (1935). *Fine structure of X-ray absorption edges of alloys*. Groningen [etc.], J. B. Wolters' uitgevers-maatschappij n. v.
- Smoluchowski, R. (1965). "Radiation Sintering of Lunar Dust." *Science* 150(3699): 1025-1026.

- Smoluchowski, R. (1968). "Mars: Retention of Ice." *Science* 159(3821): 1348-1350.
- Smoluchowski, R. (1970). "Jupiter's Convection and Its Red Spot." *Science* 168(3937): 1340-1342.
- Smoluchowski, R. (1978). "Amorphous Ice on Saturnian Rings and on Icy Satellites: Its Formation, Stability, and Observability." *Science* 201(4358): 809-811.
- Smoluchowski, R. (1983). *The solar system : the sun, planets, and life*. New York, Scientific American Library : Distributed by W.H. Freeman.
- Smoluchowski, R. (1983). "Solar System Ice: Amorphous or Crystalline?" *Science* 222(4620): 161-163.
- Smoluchowski, R., J. N. Bahcall, et al. (1986). *The Galaxy and the solar system*. Tucson, University of Arizona Press.
- Smoluchowski, R. and R. W. Turner (1949). "Absorption corrections in X-ray studies of preferred orientation." *Rev Sci Instrum* 20(3): 173.
- Snedecor, G. W. and W. G. Cochran (1989). *Statistical methods*. Ames, Iowa State University Press.
- Snoep, J. L., F. Bruggeman, et al. (2006). "Towards building the silicon cell: a modular approach." *Biosystems* 83(2-3): 207-16.
- Snoep, J. L., P. R. Jensen, et al. (1994). "How to determine control of growth rate in a chemostat. Using metabolic control analysis to resolve the paradox." *Biochem Mol Biol Int* 33(5): 1023-32.
- Snyder, M. A., A. Chatterjee, et al. (2005). "Net-event kinetic Monte Carlo for overcoming stiffness in spatially homogeneous and distributed systems." *Computers & Chemical Engineering* 29(4): 701-712.
- Sodergren, E., G. M. Weinstock, et al. (2006). "The genome of the sea urchin *Strongylocentrotus purpuratus*." *Science* 314(5801): 941-52.
- Sofia, M. J., N. Allanson, et al. (1999). "Discovery of novel disaccharide antibacterial agents using a combinatorial library approach." *J Med Chem* 42(17): 3193-8.
- Solomon, P. S., R. C. Lee, et al. (2004). "Pathogenicity of *Stagonospora nodorum* requires malate synthase." *Mol Microbiol* 53(4): 1065-73.
- Song, Y., Y. Zhang, et al. (2004). "Continuum diffusion reaction rate calculations of wild-type and mutant mouse acetylcholinesterase: adaptive finite element analysis." *Biophys J* 87(3): 1558-66.

- Song, Y., Y. Zhang, et al. (2004). "Finite element solution of the steady-state Smoluchowski equation for rate constant calculations." *Biophys J* 86(4): 2017-29.
- Sontag, E., A. Kiyatkin, et al. (2004). "Inferring dynamic architecture of cellular networks using time series of gene expression, protein and metabolite data." *Bioinformatics* 20(12): 1877-86.
- Sorokin, A., M. A. Lemmon, et al. (1994). "Stabilization of an active dimeric form of the epidermal growth factor receptor by introduction of an inter-receptor disulfide bond." *J Biol Chem* 269(13): 9752-9.
- Sosinsky, G. E., B. N. Giepmans, et al. (2007). "Markers for correlated light and electron microscopy." *Methods Cell Biol* 79: 575-91.
- Sotoudeh, M., Y. S. Li, et al. (2002). "Induction of apoptosis in vascular smooth muscle cells by mechanical stretch." *Am J Physiol Heart Circ Physiol* 282(5): H1709-16.
- Southwick, S. M., S. R. Axelrod, et al. (2003). "Twenty-four-hour urine cortisol in combat veterans with PTSD and comorbid borderline personality disorder." *J Nerv Ment Dis* 191(4): 261-2.
- Spack, E., Jr. and M. Edidin (1986). "The class I MHC antigens of erythrocytes: a serologic and biochemical study." *J Immunol* 136(8): 2943-52.
- Spack, E. G., Jr., B. Packard, et al. (1986). "Hydrophobic adsorption chromatography to reduce nonspecific staining by rhodamine-labeled antibodies." *Anal Biochem* 158(1): 233-7.
- Spencer, S. L., R. A. Gerety, et al. (2006). "Modeling somatic evolution in tumorigenesis." *PLoS Comput Biol* 2(8): e108.
- Spiliotis, E. T., H. Manley, et al. (2000). "Selective export of MHC class I molecules from the ER after their dissociation from TAP." *Immunity* 13(6): 841-51.
- Spivak-Kroizman, T., M. A. Lemmon, et al. (1994). "Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation." *Cell* 79(6): 1015-24.
- Spudich, J. A., S. J. Kron, et al. (1985). "Movement of myosin-coated beads on oriented filaments reconstituted from purified actin." *Nature* 315(6020): 584-6.
- Stebbins, E. G. and D. Mochly-Rosen (2001). "Binding specificity for RACK1 resides in the V5 region of beta II protein kinase C." *J Biol Chem* 276(32): 29644-50.

- Steck, T. L. and J. Yu (1973). "Selective solubilization of proteins from red blood cell membranes by protein perturbants." *J Supramol Struct* 1(3): 220-32.
- Steeber, D. A., P. Engel, et al. (1997). "Ligation of L-selectin through conserved regions within the lectin domain activates signal transduction pathways and integrin function in human, mouse, and rat leukocytes." *J Immunol* 159(2): 952-63.
- Stelling, J. and B. N. Kholodenko (2009). "Signaling cascades as cellular devices for spatial computations." *J Math Biol* 58(1-2): 35-55.
- Sterba, R. E. and M. P. Sheetz (1998). "Basic laser tweezers." *Methods Cell Biol* 55: 29-41.
- Steuer, E. R., L. Wordeman, et al. (1990). "Localization of cytoplasmic dynein to mitotic spindles and kinetochores." *Nature* 345(6272): 266-8.
- Stites, E. C., P. C. Trampont, et al. (2007). "Network analysis of oncogenic Ras activation in cancer." *Science* 318(5849): 463-7.
- Stokes, G. M., A. D. Milner, et al. (1986). "Ventilatory response to increased dead spaces in the first week of life." *Pediatr Pulmonol* 2(2): 89-93.
- Stokoe, D., S. G. Macdonald, et al. (1994). "Activation of Raf as a result of recruitment to the plasma membrane." *Science* 264(5164): 1463-7.
- Storm, C., J. J. Pastore, et al. (2005). "Nonlinear elasticity in biological gels." *Nature* 435(7039): 191-4.
- Straus, A. H., V. B. Valero, et al. (1997). "Glycosphingolipid antigens from *Leishmania (L.) amazonensis* amastigotes. Binding of anti-glycosphingolipid monoclonal antibodies in vitro and in vivo." *Braz J Med Biol Res* 30(3): 395-9.
- Stroud, A. N., L. M. Welter, et al. (1969). "Scanning electron microscopy of cells." *Science* 164(881): 830-2.
- Subach, F. V., G. H. Patterson, et al. (2009). "Photoactivatable mCherry for high-resolution two-color fluorescence microscopy." *Nat Methods* 6(2): 153-9.
- Subczynski, W. K., J. Widomska, et al. (2007). "Saturation-recovery electron paramagnetic resonance discrimination by oxygen transport (DOT) method for characterizing membrane domains." *Methods Mol Biol* 398: 143-57.
- Subtil, A., I. Gaidarov, et al. (1999). "Acute cholesterol depletion inhibits clathrin-coated pit budding." *Proc Natl Acad Sci U S A* 96(12): 6775-80.

- Suenaga, A., M. Hatakeyama, et al. (2009). "Molecular dynamics simulations reveal that Tyr-317 phosphorylation reduces Shc binding affinity for phosphotyrosyl residues of epidermal growth factor receptor." *Biophys J* 96(6): 2278-88.
- Suenaga, A., A. B. Kiyatkin, et al. (2004). "Tyr-317 phosphorylation increases Shc structural rigidity and reduces coupling of domain motions remote from the phosphorylation site as revealed by molecular dynamics simulations." *J Biol Chem* 279(6): 4657-62.
- Sugano, K., S. Nakamura, et al. (2008). "Cross-sectional analysis of germline BRCA1 and BRCA2 mutations in Japanese patients suspected to have hereditary breast/ovarian cancer." *Cancer Sci* 99(10): 1967-76.
- Sun, J., S. L. Zheng, et al. (2009). "Sequence variants at 22q13 are associated with prostate cancer risk." *Cancer Res* 69(1): 10-5.
- Sung, J., K. J. Shin, et al. (1997). "Many-particle effects on the relaxation kinetics of fast reversible reactions of the type $A + B \rightleftharpoons C$." *J. Chem. Phys.* 107(22).
- Susumu, K., H. T. Uyeda, et al. (2007). "Design of biotin-functionalized luminescent quantum dots." *J Biomed Biotechnol* 2007: 90651.
- Susumu, K., H. T. Uyeda, et al. (2007). "Enhancing the stability and biological functionalities of quantum dots via compact multifunctional ligands." *J Am Chem Soc* 129(45): 13987-96.
- Suzuki, H., R. Sasaki, et al. (2008). "Metabolic profiling of flavonoids in *Lotus japonicus* using liquid chromatography Fourier transform ion cyclotron resonance mass spectrometry." *Phytochemistry* 69(1): 99-111.
- Suzuki, K., K. Ritchie, et al. (2005). "Rapid hop diffusion of a G-protein-coupled receptor in the plasma membrane as revealed by single-molecule techniques." *Biophys J* 88(5): 3659-80.
- Suzuki, K. and M. P. Sheetz (2001). "Binding of cross-linked glycosylphosphatidylinositol-anchored proteins to discrete actin-associated sites and cholesterol-dependent domains." *Biophys J* 81(4): 2181-9.
- Suzuki, K., R. E. Sterba, et al. (2000). "Outer membrane monolayer domains from two-dimensional surface scanning resistance measurements." *Biophys J* 79(1): 448-59.
- Suzuki, K. G., T. K. Fujiwara, et al. (2007). "Dynamic recruitment of phospholipase C gamma at transiently immobilized GPI-anchored receptor clusters induces IP3-Ca²⁺ signaling: single-molecule tracking study 2." *J Cell Biol* 177(4): 731-42.

- Suzuki, K. G., T. K. Fujiwara, et al. (2007). "GPI-anchored receptor clusters transiently recruit Lyn and G alpha for temporary cluster immobilization and Lyn activation: single-molecule tracking study 1." *J Cell Biol* 177(4): 717-30.
- Svitkina, T. M. and G. G. Borisy (1999). "Arp2/3 complex and actin depolymerizing factor/cofilin in dendritic organization and treadmilling of actin filament array in lamellipodia." *J Cell Biol* 145(5): 1009-26.
- Svitkina, T. M. and G. G. Borisy (1999). "Progress in protrusion: the tell-tale scar." *Trends Biochem Sci* 24(11): 432-6.
- Syder, A. J., Q. C. Yu, et al. (1994). "Genetic mutations in the K1 and K10 genes of patients with epidermolytic hyperkeratosis. Correlation between location and disease severity." *J Clin Invest* 93(4): 1533-42.
- Szabo, A., R. Zwanzig, et al. (1988). "Diffusion-controlled reactions with mobile traps." *Phys Rev Lett* 61(21): 2496-2499.
- Tai, K., S. D. Bond, et al. (2003). "Finite element simulations of acetylcholine diffusion in neuromuscular junctions." *Biophys J* 84(4): 2234-41.
- Takabatake, Y., T. Takabatake, et al. (2002). "Conserved expression control and shared activity between cognate T-box genes Tbx2 and Tbx3 in connection with Sonic hedgehog signaling during Xenopus eye development." *Dev Growth Differ* 44(4): 257-71.
- Takaoka, A., Y. Mitani, et al. (2000). "Cross talk between interferon-gamma and -alpha/beta signaling components in caveolar membrane domains." *Science* 288(5475): 2357-60.
- Takizawa, N., T. C. Smith, et al. (2006). "Supervillin modulation of focal adhesions involving TRIP6/ZRP-1." *J Cell Biol* 174(3): 447-58.
- Takizawa, P. A., J. L. DeRisi, et al. (2000). "Plasma membrane compartmentalization in yeast by messenger RNA transport and a septin diffusion barrier." *Science* 290(5490): 341-4.
- Takizawa, P. A., A. Sil, et al. (1997). "Actin-dependent localization of an RNA encoding a cell-fate determinant in yeast." *Nature* 389(6646): 90-3.
- Takizawa, P. A. and R. D. Vale (2000). "The myosin motor, Myo4p, binds Ash1 mRNA via the adapter protein, She3p." *Proc Natl Acad Sci U S A* 97(10): 5273-8.
- Tamada, M., T. D. Perez, et al. (2007). "Two distinct modes of myosin assembly and dynamics during epithelial wound closure." *J Cell Biol* 176(1): 27-33.

- Tamada, M., M. P. Sheetz, et al. (2004). "Activation of a signaling cascade by cytoskeleton stretch." *Dev Cell* 7(5): 709-18.
- Tanaka, K., H. Toyoda, et al. (2008). "Duodenal metastasis of malignant melanoma observed by magnification endoscopy." *Endoscopy* 40 Suppl 2: E6-7.
- Tanaka, N., K. Aoki, et al. (2008). "Molecular basis for peroxisomal localization of tetrameric carbonyl reductase." *Structure* 16(3): 388-97.
- Tanaka, Y., K. Yonekura, et al. (1996). "Molecular and biochemical characterization of three anthocyanin synthetic enzymes from *Gentiana triflora*." *Plant Cell Physiol* 37(5): 711-6.
- Tanase, M., N. Biais, et al. (2007). "Magnetic tweezers in cell biology." *Methods Cell Biol* 83: 473-93.
- Tandon, R., I. Levental, et al. (2007). "HIV infection changes glomerular podocyte cytoskeletal composition and results in distinct cellular mechanical properties." *Am J Physiol Renal Physiol* 292(2): F701-10.
- Tang, J. X. and P. A. Janmey (1998). "Two distinct mechanisms of actin bundle formation." *Biol Bull* 194(3): 406-8.
- Tang, Q. and M. Edidin (2001). "Vesicle trafficking and cell surface membrane patchiness." *Biophys J* 81(1): 196-203.
- Tank, D. W., W. J. Fredericks, et al. (1985). "Electric field-induced redistribution and postfield relaxation of low density lipoprotein receptors on cultured human fibroblasts." *J Cell Biol* 101(1): 148-57.
- Tank, D. W., R. L. Haganir, et al. (1983). "Patch-recorded single-channel currents of the purified and reconstituted Torpedo acetylcholine receptor." *Proc Natl Acad Sci U S A* 80(16): 5129-33.
- Tank, D. W., C. Miller, et al. (1982). "Isolated-patch recording from liposomes containing functionally reconstituted chloride channels from Torpedo electroplax." *Proc Natl Acad Sci U S A* 79(24): 7749-53.
- Tank, D. W., E. S. Wu, et al. (1982). "Lateral diffusion of gramicidin C in phospholipid multibilayers. Effects of cholesterol and high gramicidin concentration." *Biophys J* 40(2): 129-35.
- Tank, D. W., E. S. Wu, et al. (1982). "Enhanced molecular diffusibility in muscle membrane blebs: release of lateral constraints." *J Cell Biol* 92(1): 207-12.

- Tappia, P. S., S. Ladha, et al. (1997). "The influence of membrane fluidity, TNF receptor binding, cAMP production and GTPase activity on macrophage cytokine production in rats fed a variety of fat diets." *Mol Cell Biochem* 166(1-2): 135-43.
- Tarca, A. L., V. J. Carey, et al. (2007). "Machine learning and its applications to biology." *PLoS Comput Biol* 3(6): e116.
- Taylor, S. W., E. Fahy, et al. (2003). "Characterization of the human heart mitochondrial proteome." *Nat Biotechnol* 21(3): 281-6.
- Teesalu, K., O. Uibo, et al. (2001). "Increased levels of IgA antibodies against desmin in children with coeliac disease." *Int Arch Allergy Immunol* 126(2): 157-66.
- Teis, D., W. Wunderlich, et al. (2002). "Localization of the MP1-MAPK scaffold complex to endosomes is mediated by p14 and required for signal transduction." *Dev Cell* 3(6): 803-14.
- Temple, G. S., S. Oliver, et al. (1991). "Detection of micro-organisms using a two bottle blood culture system: a twelve month study." *Med Lab Sci* 48(3): 178-82.
- Tepavcevic, V., R. R. Hodges, et al. (2003). "Signal transduction pathways used by EGF to stimulate protein secretion in rat lacrimal gland." *Invest Ophthalmol Vis Sci* 44(3): 1075-81.
- Teske, A. and M. Smoluchowski (1955). Marian Smoluchowski; *zycie i twórczo*s*c. W Krakowie], Pa*ństwowo Wydaw. Naukowe.
- Teske, A. and M. Smoluchowski (1977). Marian Smoluchowski : Leben und Werk. Wroc*aw, Zak*ad Narodowy Imienia Ossoli*nskich Wydawn. Polskiej Akademii Nauk.
- Therrien, M., N. R. Michaud, et al. (1996). "KSR modulates signal propagation within the MAPK cascade." *Genes Dev* 10(21): 2684-95.
- Therrien, M., D. K. Morrison, et al. (2000). "A genetic screen for modifiers of a kinase suppressor of Ras-dependent rough eye phenotype in *Drosophila*." *Genetics* 156(3): 1231-42.
- Thomas, R., C. Gonzalez, et al. (2004). "A novel assay to determine the sequence preference and affinity of DNA minor groove binding compounds." *Nucleic Acids Res* 32(1): e8.
- Thrane, E. V., R. Becher, et al. (1997). "Differential distribution and increased levels of ras proteins during lung development." *Exp Lung Res* 23(1): 35-49.

- Thrane, E. V., M. Refsnes, et al. (2001). "Fluoride-induced apoptosis in epithelial lung cells involves activation of MAP kinases p38 and possibly JNK." *Toxicol Sci* 61(1): 83-91.
- Thrane, E. V., P. E. Schwarze, et al. (2001). "Persistent versus transient map kinase (ERK) activation in the proliferation of lung epithelial type 2 cells." *Exp Lung Res* 27(4): 387-400.
- Thuduppathy, G. R., J. W. Craig, et al. (2006). "Evidence that membrane insertion of the cytosolic domain of Bcl-xL is governed by an electrostatic mechanism." *J Mol Biol* 359(4): 1045-58.
- Tian, T., A. Harding, et al. (2007). "Plasma membrane nanoswitches generate high-fidelity Ras signal transduction." *Nat Cell Biol* 9(8): 905-14.
- Tien, T. Q., M. Maiwald, et al. (2008). "Microexternal cavity tapered lasers at 670 nm with 5 W peak power and nearly diffraction-limited beam quality." *Opt Lett* 33(22): 2692-4.
- Tienari, P. J., B. De Strooper, et al. (1996). "Neuronal sorting and processing of amyloid precursor protein: implications for Alzheimer's disease." *Cold Spring Harb Symp Quant Biol* 61: 575-85.
- Tienari, P. J., B. De Strooper, et al. (1996). "The beta-amyloid domain is essential for axonal sorting of amyloid precursor protein." *EMBO J* 15(19): 5218-29.
- Tienari, P. J., N. Ida, et al. (1997). "Intracellular and secreted Alzheimer beta-amyloid species are generated by distinct mechanisms in cultured hippocampal neurons." *Proc Natl Acad Sci U S A* 94(8): 4125-30.
- Timar, J., H. Chopra, et al. (1992). "Calcium channel blocker treatment of tumor cells induces alterations in the cytoskeleton, mobility of the integrin alpha IIb beta 3 and tumor-cell-induced platelet aggregation." *J Cancer Res Clin Oncol* 118(6): 425-34.
- Todryk, S., A. Melcher, et al. (2001). "Cell death associated with genetic prodrug activation therapy of colorectal cancer." *Cancer Lett* 174(1): 25-33.
- Togo, T., J. M. Alderton, et al. (2000). "The mechanism of cell membrane repair." *Zygote* 8 Suppl 1: S31-2.
- Togo, T., T. B. Krasieva, et al. (2000). "A decrease in membrane tension precedes successful cell-membrane repair." *Mol Biol Cell* 11(12): 4339-46.
- Tolhurst, G., C. Vial, et al. (2005). "Interplay between P2Y(1), P2Y(12), and P2X(1) receptors in the activation of megakaryocyte cation influx currents by ADP: evidence that the primary megakaryocyte represents a fully functional model of platelet P2 receptor signaling." *Blood* 106(5): 1644-51.

- Tomlinson, I., P. Sasieni, et al. (2002). "How many mutations in a cancer?" *Am J Pathol* 160(3): 755-8.
- Tomlinson, I. P., M. R. Novelli, et al. (1996). "The mutation rate and cancer." *Proc Natl Acad Sci U S A* 93(25): 14800-3.
- Toomre, D., P. Keller, et al. (1999). "Dual-color visualization of trans-Golgi network to plasma membrane traffic along microtubules in living cells." *J Cell Sci* 112 (Pt 1): 21-33.
- Toomre, D., J. A. Steyer, et al. (2000). "Fusion of constitutive membrane traffic with the cell surface observed by evanescent wave microscopy." *J Cell Biol* 149(1): 33-40.
- Toyoshima, I. and M. P. Sheetz (1996). "Kinectin distribution in chicken nervous system." *Neurosci Lett* 211(3): 171-4.
- Toyoshima, I., M. Sugawara, et al. (1998). "Kinesin and cytoplasmic dynein in spinal spheroids with motor neuron disease." *J Neurol Sci* 159(1): 38-44.
- Toyoshima, I., H. Yu, et al. (1992). "Kinectin, a major kinesin-binding protein on ER." *J Cell Biol* 118(5): 1121-31.
- Tran, A. D., T. P. Marmo, et al. (2007). "HDAC6 deacetylation of tubulin modulates dynamics of cellular adhesions." *J Cell Sci* 120(Pt 8): 1469-79.
- Trosko, J. E., C. C. Chang, et al. (2004). "Ignored hallmarks of carcinogenesis: stem cells and cell-cell communication." *Ann N Y Acad Sci* 1028: 192-201.
- Tsuchimoto, M., M. Aoki, et al. (2004). "The changes of gene expression in honeybee (*Apis mellifera*) brains associated with ages." *Zoolog Sci* 21(1): 23-8.
- Tsukada, Y., K. Aoki, et al. (2008). "Quantification of local morphodynamics and local GTPase activity by edge evolution tracking." *PLoS Comput Biol* 4(11): e1000223.
- Tubbs, R. S., J. W. Custis, et al. (2005). "Landmarks for the greater petrosal nerve." *Clin Anat* 18(3): 210-4.
- Tubbs, R. S., E. G. Salter, et al. (2005). "Novel surgical approach to the carpal tunnel: cadaveric feasibility study." *Clin Anat* 18(5): 350-6.
- Tubbs, R. S., E. G. Salter, et al. (2006). "An unusual case of a mediastinal mass in a cadaver." *Clin Anat* 19(2): 151-3.
- Tubbs, R. S., J. Sheetz, et al. (2003). "Accessory obturator nerves with bilateral pseudoganglia in man." *Ann Anat* 185(6): 571-2.

Tubbs, R. S., E. C. Tyler-Kabara, et al. (2006). "Additional vascular compression of the brachial plexus in a cadaver with a cervical rib: case illustration." *Surg Radiol Anat* 28(1): 112-3.

Turner, P. R., M. P. Sheetz, et al. (1984). "Fertilization increases the polyphosphoinositide content of sea urchin eggs." *Nature* 310(5976): 414-5.

Ueno, K., S. Nakamura, et al. (2008). "Electric-field-induced superconductivity in an insulator." *Nat Mater* 7(11): 855-8.

Umemura, Y. M., M. Vrljic, et al. (2008). "Both MHC class II and its GPI-anchored form undergo hop diffusion as observed by single-molecule tracking." *Biophys J* 95(1): 435-50.

Upadhyaya, A. and M. P. Sheetz (2004). "Tension in tubulovesicular networks of Golgi and endoplasmic reticulum membranes." *Biophys J* 86(5): 2923-8.

Ushio-Fukai, M., L. Zuo, et al. (2005). "cAbl tyrosine kinase mediates reactive oxygen species- and caveolin-dependent AT1 receptor signaling in vascular smooth muscle: role in vascular hypertrophy." *Circ Res* 97(8): 829-36.

Usui, Y., K. Aoki, et al. (2008). "Carbon nanotubes with high bone-tissue compatibility and bone-formation acceleration effects." *Small* 4(2): 240-6.

Usuki, F., E. Fujita, et al. (2008). "Methylmercury activates ASK1/JNK signaling pathways, leading to apoptosis due to both mitochondria- and endoplasmic reticulum (ER)-generated processes in myogenic cell lines." *Neurotoxicology* 29(1): 22-30.

Usuki, F., S. Ishiura, et al. (1997). "Expanded CTG repeats in myotonin protein kinase suppresses myogenic differentiation." *Neuroreport* 8(17): 3749-53.

Usuki, F., N. Takahashi, et al. (2000). "Differential signaling pathways following oxidative stress in mutant myotonin protein kinase cDNA-transfected C2C12 cell lines." *Biochem Biophys Res Commun* 267(3): 739-43.

Uyeda, H. T., I. L. Medintz, et al. (2005). "Synthesis of compact multidentate ligands to prepare stable hydrophilic quantum dot fluorophores." *J Am Chem Soc* 127(11): 3870-8.

Vale, R. D., T. S. Reese, et al. (1985). "Identification of a novel force-generating protein, kinesin, involved in microtubule-based motility." *Cell* 42(1): 39-50.

Vale, R. D., B. J. Schnapp, et al. (1985). "Different axoplasmic proteins generate movement in opposite directions along microtubules in vitro." *Cell* 43(3 Pt 2): 623-32.

- Vale, R. D., B. J. Schnapp, et al. (1985). "Movement of organelles along filaments dissociated from the axoplasm of the squid giant axon." *Cell* 40(2): 449-54.
- Vale, R. D., B. J. Schnapp, et al. (1985). "Organelle, bead, and microtubule translocations promoted by soluble factors from the squid giant axon." *Cell* 40(3): 559-69.
- Vale, R. D., A. G. Szent-Gyorgyi, et al. (1984). "Movement of scallop myosin on Nitella actin filaments: regulation by calcium." *Proc Natl Acad Sci U S A* 81(21): 6775-8.
- Valentijn, J. A., L. T. Gien, et al. (2000). "An evaluation of the expression, subcellular localization, and function of rab4 in the exocrine pancreas." *Biochem Biophys Res Commun* 268(3): 847-52.
- Valentijn, J. A., K. Valentijn, et al. (2000). "Actin coating of secretory granules during regulated exocytosis correlates with the release of rab3D." *Proc Natl Acad Sci U S A* 97(3): 1091-5.
- Vallee, R. B. and M. P. Sheetz (1996). "Targeting of motor proteins." *Science* 271(5255): 1539-44.
- Vallotton, P., S. L. Gupton, et al. (2004). "Simultaneous mapping of filamentous actin flow and turnover in migrating cells by quantitative fluorescent speckle microscopy." *Proc Natl Acad Sci U S A* 101(26): 9660-5.
- Vallotton, P., A. Ponti, et al. (2003). "Recovery, visualization, and analysis of actin and tubulin polymer flow in live cells: a fluorescent speckle microscopy study." *Biophys J* 85(2): 1289-306.
- Vamosi, G., A. Bodnar, et al. (2004). "IL-2 and IL-15 receptor alpha-subunits are coexpressed in a supramolecular receptor cluster in lipid rafts of T cells." *Proc Natl Acad Sci U S A* 101(30): 11082-7.
- van Dam, K., J. van der Vlag, et al. (1993). "The sum of the control coefficients of all enzymes on the flux through a group-transfer pathway can be as high as two." *Eur J Biochem* 212(3): 791-9.
- van der Meersch, J. J. and R. D. Simons (1966). "[Testing of percutaneous steroid resorption by application of the Locacorten ointment under plastic occlusive dressings]." *Dermatologica* 132(6): 460-75.
- van Golde, L. M., V. Oldenburg, et al. (1980). "Phospholipid transfer proteins in rat lung. Identification of a protein specific for phosphatidylglycerol." *J Biol Chem* 255(13): 6011-3.

- van Heeswijk, W. C., B. M. Bakker, et al. (1999). "Live control of the living cell." *Biochem Soc Trans* 27(2): 261-4.
- Van Komen, J. S., S. Mishra, et al. (2007). "Early and dynamic polarization of T cell membrane rafts and constituents prior to TCR stop signals." *J Immunol* 179(10): 6845-55.
- van Meer, G., J. Davoust, et al. (1985). "Parameters affecting low-pH-mediated fusion of liposomes with the plasma membrane of cells infected with influenza virus." *Biochemistry* 24(14): 3593-602.
- van Meer, G., B. Gumbiner, et al. (1986). "The tight junction does not allow lipid molecules to diffuse from one epithelial cell to the next." *Nature* 322(6080): 639-41.
- van Meer, G., B. J. Poorthuis, et al. (1980). "Transbilayer distribution and mobility of phosphatidylcholine in intact erythrocyte membranes. A study with phosphatidylcholine exchange protein." *Eur J Biochem* 103(2): 283-8.
- van Meer, G. and K. Simons (1982). "Viruses budding from either the apical or the basolateral plasma membrane domain of MDCK cells have unique phospholipid compositions." *EMBO J* 1(7): 847-52.
- van Meer, G. and K. Simons (1983). "An efficient method for introducing defined lipids into the plasma membrane of mammalian cells." *J Cell Biol* 97(5 Pt 1): 1365-74.
- van Meer, G. and K. Simons (1986). "The function of tight junctions in maintaining differences in lipid composition between the apical and the basolateral cell surface domains of MDCK cells." *EMBO J* 5(7): 1455-64.
- van Meer, G. and K. Simons (1988). "Lipid polarity and sorting in epithelial cells." *J Cell Biochem* 36(1): 51-8.
- van Meer, G., K. Simons, et al. (1981). "Phospholipid asymmetry in Semliki Forest virus grown on baby hamster kidney (BHK-21) cells." *Biochemistry* 20(7): 1974-81.
- van Meer, G., E. H. Stelzer, et al. (1987). "Sorting of sphingolipids in epithelial (Madin-Darby canine kidney) cells." *J Cell Biol* 105(4): 1623-35.
- Vandekerckhove, J., G. Bauw, et al. (1990). "Comparative two-dimensional gel analysis and microsequencing identifies gelsolin as one of the most prominent downregulated markers of transformed human fibroblast and epithelial cells." *J Cell Biol* 111(1): 95-102.

- Varadhachary, A. S., M. Edidin, et al. (2001). "Phosphatidylinositol 3'-kinase blocks CD95 aggregation and caspase-8 cleavage at the death-inducing signaling complex by modulating lateral diffusion of CD95." *J Immunol* 166(11): 6564-9.
- Vatsurina, A. V., T. Z. Esikova, et al. (2005). "[Corrosion of pipe steel samples and conjugated conversion of sulfur compounds by thiobacteria *Halothiobacillus neapolitanus* DSM 15147]." *Prikl Biokhim Mikrobiol* 41(5): 564-7.
- Vegners, R., I. Shestakova, et al. (1995). "Use of a gel-forming dipeptide derivative as a carrier for antigen presentation." *J Pept Sci* 1(6): 371-8.
- Velazquez, C., I. Valette, et al. (1995). "Identification of immunogenic epitopes of the 170-kDa subunit adhesin of *Entamoeba histolytica* in patients with invasive amebiasis." *J Eukaryot Microbiol* 42(5): 636-41.
- Vereb, G., J. Matko, et al. (2004). "Cytometry of fluorescence resonance energy transfer." *Methods Cell Biol* 75: 105-52.
- Vereb, G., Jr., G. Panyi, et al. (1990). "Effect of cyclosporin A on the membrane potential and Ca²⁺ level of human lymphoid cell lines and mouse thymocytes." *Biochim Biophys Acta* 1019(2): 159-65.
- Vereb, G., J. Szollosi, et al. (2003). "Dynamic, yet structured: The cell membrane three decades after the Singer-Nicolson model." *Proc Natl Acad Sci U S A* 100(14): 8053-8.
- Verkhovsky, A. B., T. M. Svitkina, et al. (1999). "Network contraction model for cell translocation and retrograde flow." *Biochem Soc Symp* 65: 207-22.
- Verkhovsky, A. B., T. M. Svitkina, et al. (1999). "Self-polarization and directional motility of cytoplasm." *Curr Biol* 9(1): 11-20.
- Vial, C. and R. J. Evans (2005). "Disruption of lipid rafts inhibits P2X₁ receptor-mediated currents and arterial vasoconstriction." *J Biol Chem* 280(35): 30705-11.
- Vickery, B. H. and J. P. Bennett (1968). "The cervix and its secretion in mammals." *Physiol Rev* 48(1): 135-54.
- Vijayan, K., D. E. Discher, et al. (2005). "Interactions of membrane-active peptides with thick, neutral, nonzwitterionic bilayers." *J Phys Chem B* 109(30): 14356-64.
- Vogel, V. and M. Sheetz (2006). "Local force and geometry sensing regulate cell functions." *Nat Rev Mol Cell Biol* 7(4): 265-75.
- Vogel, V. and M. P. Sheetz (2009). "Cell fate regulation by coupling mechanical cycles to biochemical signaling pathways." *Curr Opin Cell Biol* 21(1): 38-46.

- Volmer, J., D. Abbott, et al. (2001). "Measurement of the charged pion electromagnetic form factor." *Phys Rev Lett* 86(9): 1713-6.
- Von Bonsdorff, C. H., S. D. Fuller, et al. (1985). "Apical and basolateral endocytosis in Madin-Darby canine kidney (MDCK) cells grown on nitrocellulose filters." *EMBO J* 4(11): 2781-92.
- von Wichert, G., B. Haimovich, et al. (2003). "Force-dependent integrin-cytoskeleton linkage formation requires downregulation of focal complex dynamics by Shp2." *EMBO J* 22(19): 5023-35.
- von Wichert, G., G. Jiang, et al. (2003). "RPTP- α acts as a transducer of mechanical force on α 5 β 3-integrin-cytoskeleton linkages." *J Cell Biol* 161(1): 143-53.
- von Wichert, G., D. Krndija, et al. (2008). "Focal adhesion kinase mediates defects in the force-dependent reinforcement of initial integrin-cytoskeleton linkages in metastatic colon cancer cell lines." *Eur J Cell Biol* 87(1): 1-16.
- von Wichert, G. and M. P. Sheetz (2005). "Mechanisms of disease: the biophysical interpretation of the ECM affects physiological and pathophysiological cellular behavior." *Z Gastroenterol* 43(12): 1329-36.
- Wada, H., K. Masuda, et al. (2008). "Adult T-cell progenitors retain myeloid potential." *Nature* 452(7188): 768-72.
- Walker, R. A., E. T. O'Brien, et al. (1997). "n-ethylmaleimide and ethacrynic acid inhibit kinesin binding to microtubules in a motility assay." *Cell Motil Cytoskeleton* 37(4): 289-99.
- Walker, R. A. and M. P. Sheetz (1993). "Cytoplasmic microtubule-associated motors." *Annu Rev Biochem* 62: 429-51.
- Waller, A., K. L. Sutton, et al. (2004). "Receptor binding kinetics and cellular responses of six N-formyl peptide agonists in human neutrophils." *Biochemistry* 43(25): 8204-16.
- Wang, C., J. E. Mahaffey, et al. (1979). "Hyperfunctioning supernumerary parathyroid glands." *Surg Gynecol Obstet* 148(5): 711-4.
- Wang, D., S. Y. Gou, et al. (1992). "Reaction rate enhancement by surface diffusion of adsorbates." *Biophys Chem* 43(2): 117-37.
- Wang, D. L., B. S. Wung, et al. (1995). "Mechanical strain induces monocyte chemotactic protein-1 gene expression in endothelial cells. Effects of mechanical strain on monocyte adhesion to endothelial cells." *Circ Res* 77(2): 294-302.

- Wang, J., R. Bright, et al. (2004). "Cell-specific role for epsilon- and betaI-protein kinase C isozymes in protecting cortical neurons and astrocytes from ischemia-like injury." *Neuropharmacology* 47(1): 136-45.
- Wang, J., A. Rajput, et al. (2009). "Knockdown of ron kinase inhibits mutant phosphatidylinositol 3-kinase and reduces metastasis in human colon carcinoma." *J Biol Chem* 284(16): 10912-22.
- Wang, L., M. A. Bittner, et al. (2008). "The structural and functional implications of linked SNARE motifs in SNAP25." *Mol Biol Cell* 19(9): 3944-55.
- Wang, L. Z., J. Gorlin, et al. (2000). "Purification of salmon clotting factors and their use as tissue sealants." *Thromb Res* 100(6): 537-48.
- Wang, M. D. and D. Axelrod (1994). "Time-lapse total internal reflection fluorescence video of acetylcholine receptor cluster formation on myotubes." *Dev Dyn* 201(1): 29-40.
- Wang, Y., J. Chang, et al. (2004). "Shear stress and VEGF activate IKK via the Flk-1/Cbl/Akt signaling pathway." *Am J Physiol Heart Circ Physiol* 286(2): H685-92.
- Wang, Y., H. Miao, et al. (2002). "Interplay between integrins and FLK-1 in shear stress-induced signaling." *Am J Physiol Cell Physiol* 283(5): C1540-7.
- Wang, Y., J. Y. Shyy, et al. (2008). "Fluorescence proteins, live-cell imaging, and mechanobiology: seeing is believing." *Annu Rev Biomed Eng* 10: 1-38.
- Wang, Z., S. Khan, et al. (1995). "Single cytoplasmic dynein molecule movements: characterization and comparison with kinesin." *Biophys J* 69(5): 2011-23.
- Wang, Z. and M. P. Sheetz (1999). "One-dimensional diffusion on microtubules of particles coated with cytoplasmic dynein and immunoglobulins." *Cell Struct Funct* 24(5): 373-83.
- Wang, Z. and M. P. Sheetz (2000). "The C-terminus of tubulin increases cytoplasmic dynein and kinesin processivity." *Biophys J* 78(4): 1955-64.
- Waterman-Storer, C. M. and G. Danuser (2002). "New directions for fluorescent speckle microscopy." *Curr Biol* 12(18): R633-40.
- Waterston, R. H., K. Lindblad-Toh, et al. (2002). "Initial sequencing and comparative analysis of the mouse genome." *Nature* 420(6915): 520-62.
- Ways, D. K. and M. J. Sheetz (2000). "The role of protein kinase C in the development of the complications of diabetes." *Vitam Horm* 60: 149-93.

- Weaver, F. E., H. Polster, et al. (1990). "Normal band 3-cytoskeletal interactions are maintained on tanktreading erythrocytes." *Biophys J* 58(6): 1427-36.
- Webb, W. W., L. S. Barak, et al. (1981). "Molecular mobility on the cell surface." *Biochem Soc Symp*(46): 191-205.
- Wehr, H., D. Gorska, et al. (1977). "[Cholesterol esterification catalysed by LCAT and erythrocyte lipid pattern in Wilson's disease]." *Pol Tyg Lek* 32(15): 549-51.
- Wei, Z., V. S. Deshpande, et al. (2008). "Analysis and interpretation of stress fiber organization in cells subject to cyclic stretch." *J Biomech Eng* 130(3): 031009.
- Werner, I., L. A. Deanovic, et al. (2002). "Toxicity of stormwater runoff after dormant spray application of diazinon and esfenvalerate (Asana) in a French prune orchard, Glenn county, California, USA." *Bull Environ Contam Toxicol* 68(1): 29-36.
- Westerhoff, H. V., W. M. Getz, et al. (2002). "Bioinformatics, cellular flows, and calculation." *Ernst Schering Res Found Workshop*(38): 221-43.
- Westerhoff, H. V., J. H. Hofmeyr, et al. (1994). "Getting to the inside of cells using metabolic control analysis." *Biophys Chem* 50(3): 273-83.
- Westerhoff, H. V., B. N. Kholodenko, et al. (1995). "Elusive control." *J Bioenerg Biomembr* 27(5): 491-7.
- Westerhoff, H. V., A. Kolodkin, et al. (2009). "Systems biology towards life in silico: mathematics of the control of living cells." *J Math Biol* 58(1-2): 7-34.
- Westerhoff, H. V. and B. O. Palsson (2004). "The evolution of molecular biology into systems biology." *Nat Biotechnol* 22(10): 1249-52.
- Widschwendter, M. and P. A. Jones (2002). "DNA methylation and breast carcinogenesis." *Oncogene* 21(35): 5462-82.
- Wiley, A., D. Katsaros, et al. (2006). "Aberrant promoter methylation of multiple genes in malignant ovarian tumors and in ovarian tumors with low malignant potential." *Cancer* 107(2): 299-308.
- Wiley, J. S., C. E. Gargett, et al. (1998). "Partial agonists and antagonists reveal a second permeability state of human lymphocyte P2Z/P2X7 channel." *Am J Physiol* 275(5 Pt 1): C1224-31.
- Wilkins, S. J., S. Yoong, et al. (2008). "Mtx2 directs zebrafish morphogenetic movements during epiboly by regulating microfilament formation." *Dev Biol* 314(1): 12-22.

- Wilson, A. D. and D. W. Oliver (2009). "Local anaesthetic calculation--reply to letter." *J Plast Reconstr Aesthet Surg* 62(2): 266.
- Wilson, B. S., G. G. Deanin, et al. (1991). "Regulation of IgE receptor-mediated secretion from RBL-2H3 mast cells by GTP binding-proteins and calcium." *Biochem Biophys Res Commun* 174(3): 1064-9.
- Wilson, B. S., G. G. Deanin, et al. (1989). "Depletion of guanine nucleotides with mycophenolic acid suppresses IgE receptor-mediated degranulation in rat basophilic leukemia cells." *J Immunol* 143(1): 259-65.
- Wilson, B. S., N. Kapp, et al. (1995). "Distinct functions of the Fc epsilon R1 gamma and beta subunits in the control of Fc epsilon R1-mediated tyrosine kinase activation and signaling responses in RBL-2H3 mast cells." *J Biol Chem* 270(8): 4013-22.
- Wilson, B. S. and J. M. Oliver (2002). "Effector roles of IgE antibodies: targeting allergen to the high-affinity IgE receptor for Fc epsilon RI-dependent signaling and antigen presentation." *Clin Allergy Immunol* 16: 197-232.
- Wilson, B. S., J. R. Pfeiffer, et al. (2000). "Observing FcepsilonRI signaling from the inside of the mast cell membrane." *J Cell Biol* 149(5): 1131-42.
- Wilson, B. S., J. R. Pfeiffer, et al. (2002). "FcepsilonRI signaling observed from the inside of the mast cell membrane." *Mol Immunol* 38(16-18): 1259-68.
- Wilson, B. S., J. R. Pfeiffer, et al. (2007). "Exploring membrane domains using native membrane sheets and transmission electron microscopy." *Methods Mol Biol* 398: 245-61.
- Wilson, B. S., J. R. Pfeiffer, et al. (1998). "Calcium-dependent clustering of inositol 1,4,5-trisphosphate receptors." *Mol Biol Cell* 9(6): 1465-78.
- Wilson, B. S., J. R. Pfeiffer, et al. (2001). "High resolution mapping of mast cell membranes reveals primary and secondary domains of Fc(epsilon)RI and LAT." *J Cell Biol* 154(3): 645-58.
- Wilson, B. S., J. Seagrave, et al. (1991). "Impaired secretion and increased insolubilization of IgE-receptor complexes in mycophenolic acid-treated (guanine nucleotide-depleted) RBL-2H3 mast cells." *J Cell Physiol* 149(3): 403-7.
- Wilson, B. S., S. L. Steinberg, et al. (2004). "Markers for detergent-resistant lipid rafts occupy distinct and dynamic domains in native membranes." *Mol Biol Cell* 15(6): 2580-92.

- Wilson, C., J. Yang, et al. (2005). "Overexpression of genes on 16q associated with cisplatin resistance of testicular germ cell tumor cell lines." *Genes Chromosomes Cancer* 43(2): 211-6.
- Wilson, D. J., B. B. Norman, et al. (1991). "Evaluation of multiple reticulorumen selenium pellets as a health risk in growing Hereford steers." *Am J Vet Res* 52(11): 1866-70.
- Wilson, E. L., P. Jacobs, et al. (1985). "Plasminogen activator as a prognostic factor in hematological malignancies." *Haematol Blood Transfus* 29: 197-9.
- Wilson, G., C. G. Sheps, et al. (1982). "Effects of hospital revenue bonds on hospital planning and operations." *N Engl J Med* 307(23): 1426-30.
- Wilson, G. F., F. C. Richardson, et al. (1996). "Identification and characterization of a Ca^{2+} -sensitive nonspecific cation channel underlying prolonged repetitive firing in Aplysia neurons." *J Neurosci* 16(11): 3661-71.
- Wilson, G. N. and W. J. Oliver (1988). "Further delineation of the G syndrome: a manageable genetic cause of infantile dysphagia." *J Med Genet* 25(3): 157-63.
- Wilson, R., A. J. Allen, et al. (1995). "The translocation, folding, assembly and redox-dependent degradation of secretory and membrane proteins in semi-permeabilized mammalian cells." *Biochem J* 307 (Pt 3): 679-87.
- Wilson, T. J., K. K. Thomsen, et al. (2003). "Detection of 3-hydroxykynurenine in a plant pathogenic fungus." *Biochem J* 371(Pt 3): 783-8.
- Winter, C. G., B. Wang, et al. (2001). "Drosophila Rho-associated kinase (Drok) links Frizzled-mediated planar cell polarity signaling to the actin cytoskeleton." *Cell* 105(1): 81-91.
- Wirkner, K., D. Stanchev, et al. (2005). "Regulation of human recombinant P2X3 receptors by ecto-protein kinase C." *J Neurosci* 25(34): 7734-42.
- Wisniewska, A., J. Draus, et al. (2003). "Is a fluid-mosaic model of biological membranes fully relevant? Studies on lipid organization in model and biological membranes." *Cell Mol Biol Lett* 8(1): 147-59.
- Wisniewska, A. and W. K. Subczynski (2006). "Accumulation of macular xanthophylls in unsaturated membrane domains." *Free Radic Biol Med* 40(10): 1820-6.
- Wisniewska, A. and W. K. Subczynski (2006). "Distribution of macular xanthophylls between domains in a model of photoreceptor outer segment membranes." *Free Radic Biol Med* 41(8): 1257-65.

- Wisniewska, A. and W. K. Subczynski (2008). "The liquid-ordered phase in sphingomyelincholesterol membranes as detected by the discrimination by oxygen transport (DOT) method." *Cell Mol Biol Lett* 13(3): 430-51.
- Wolf-Yadlin, A., N. Kumar, et al. (2006). "Effects of HER2 overexpression on cell signaling networks governing proliferation and migration." *Mol Syst Biol* 2: 54.
- Wong, G. C., J. X. Tang, et al. (2000). "Hierarchical self-assembly of F-actin and cationic lipid complexes: stacked three-layer tubule networks." *Science* 288(5473): 2035-9.
- Wood, E. R., L. M. Shewchuk, et al. (2008). "6-Ethynylthieno[3,2-d]- and 6-ethynylthieno[2,3-d]pyrimidin-4-anilines as tunable covalent modifiers of ErbB kinases." *Proc Natl Acad Sci U S A* 105(8): 2773-8.
- Wood, E. R., A. T. Truesdale, et al. (2004). "A unique structure for epidermal growth factor receptor bound to GW572016 (Lapatinib): relationships among protein conformation, inhibitor off-rate, and receptor activity in tumor cells." *Cancer Res* 64(18): 6652-9.
- Wood, E. R., A. T. Truesdale, et al. (2004). "A unique structure for epidermal growth factor receptor bound to GW572016 (Lapatinib): relationships among protein conformation, inhibitor off-rate, and receptor activity in tumor cells." *Cancer Res* 64(18): 6652-9.
- Woodyard, L. W. and J. E. Sheetz (1993). "Critical pathway patient outcomes: the missing standard." *J Nurs Care Qual* 8(1): 51-7.
- Woolf, P. J., T. P. Kenakin, et al. (2001). "Uncovering biases in high throughput screens of G-protein coupled receptors." *J Theor Biol* 208(4): 403-18.
- Woolf, P. J. and J. J. Linderman (2003). "Self organization of membrane proteins via dimerization." *Biophys Chem* 104(1): 217-27.
- Woolf, P. J. and J. J. Linderman (2003). "Untangling ligand induced activation and desensitization of G-protein-coupled receptors." *Biophys J* 84(1): 3-13.
- Woolf, P. J. and J. J. Linderman (2004). "An algebra of dimerization and its implications for G-protein coupled receptor signaling." *J Theor Biol* 229(2): 157-68.
- Wordeman, L., E. R. Steuer, et al. (1991). "Chemical subdomains within the kinetochore domain of isolated CHO mitotic chromosomes." *J Cell Biol* 114(2): 285-94.

- Wriggers, W., J. X. Tang, et al. (1998). "Cofilin and gelsolin segment-1: molecular dynamics simulation and biochemical analysis predict a similar actin binding mode." *J Mol Biol* 282(5): 921-32.
- Wu, D., T. L. Foreman, et al. (2002). "Protein kinase cepsilon has the potential to advance the recurrence of human prostate cancer." *Cancer Res* 62(8): 2423-9.
- Wu, D. and D. M. Terrian (2002). "Regulation of caveolin-1 expression and secretion by a protein kinase cepsilon signaling pathway in human prostate cancer cells." *J Biol Chem* 277(43): 40449-55.
- Wu, E. S., D. W. Tank, et al. (1982). "Unconstrained lateral diffusion of concanavalin A receptors on bulbous lymphocytes." *Proc Natl Acad Sci U S A* 79(16): 4962-6.
- Wu, Z. L. and M. S. Paller (1994). "Iron loading enhances susceptibility to renal ischemia in rats." *Ren Fail* 16(4): 471-80.
- Xiao, G. Q., Y. Qu, et al. (2001). "Evidence for functional role of epsilonPKC isozyme in the regulation of cardiac Na(+) channels." *Am J Physiol Cell Physiol* 281(5): C1477-86.
- Xing, Y., A. M. Smith, et al. (2006). "Molecular profiling of single cancer cells and clinical tissue specimens with semiconductor quantum dots." *Int J Nanomedicine* 1(4): 473-81.
- Xu, J., L. Dimitrov, et al. (2005). "A combined genomewide linkage scan of 1,233 families for prostate cancer-susceptibility genes conducted by the international consortium for prostate cancer genetics." *Am J Hum Genet* 77(2): 219-29.
- Xu, J., W. H. Schwarz, et al. (1998). "Mechanical properties of actin filament networks depend on preparation, polymerization conditions, and storage of actin monomers." *Biophys J* 74(5): 2731-40.
- Xue, M., G. Hsieh, et al. (2007). "Activated N-formyl peptide receptor and high-affinity IgE receptor occupy common domains for signaling and internalization." *Mol Biol Cell* 18(4): 1410-20.
- Yalamanchili, N., D. E. Zak, et al. (2006). "Quantifying gene network connectivity in silico: scalability and accuracy of a modular approach." *Syst Biol (Stevenage)* 153(4): 236-46.
- Yamada, M., S. Momoshima, et al. (2008). "Diffusion-tensor neuronal fiber tractography and manganese-enhanced MR imaging of primate visual pathway in the common marmoset: preliminary results." *Radiology* 249(3): 855-64.

- Yamada, S., S. Shiono, et al. (2003). "Control mechanism of JAK/STAT signal transduction pathway." *FEBS Lett* 534(1-3): 190-6.
- Yamamoto, T., T. Kozasa, et al. (2008). "Genomic analyses of bovine viral diarrhea viruses isolated from cattle imported into Japan between 1991 and 2005." *Vet Microbiol* 127(3-4): 386-91.
- Yamashita, T., S. Y. Mao, et al. (1994). "Aggregation of the high-affinity IgE receptor and enhanced activity of p53/56lyn protein-tyrosine kinase." *Proc Natl Acad Sci U S A* 91(23): 11251-5.
- Yan, J., S. Roy, et al. (1998). "Ras isoforms vary in their ability to activate Raf-1 and phosphoinositide 3-kinase." *J Biol Chem* 273(37): 24052-6.
- Yang, H. C., V. Simon, et al. (2001). "Visualization of mitochondrial movement in yeast." *Methods Cell Biol* 65: 333-51.
- Yang, J., M. I. Monine, et al. (2008). "Kinetic Monte Carlo method for rule-based modeling of biochemical networks." *Phys Rev E Stat Nonlin Soft Matter Phys* 78(3 Pt 1): 031910.
- Yang, S., M. A. Raymond-Stintz, et al. (2007). "Mapping ErbB receptors on breast cancer cell membranes during signal transduction." *J Cell Sci* 120(Pt 16): 2763-73.
- Yaqoob, P. and P. C. Calder (2007). "Fatty acids and immune function: new insights into mechanisms." *Br J Nutr* 98 Suppl 1: S41-5.
- Yarden, Y. and M. X. Sliwkowski (2001). "Untangling the ErbB signalling network." *Nat Rev Mol Cell Biol* 2(2): 127-37.
- Yauch, R. L., D. P. Felsenfeld, et al. (1997). "Mutational evidence for control of cell adhesion through integrin diffusion/clustering, independent of ligand binding." *J Exp Med* 186(8): 1347-55.
- Yedovitzky, M., D. Mochly-Rosen, et al. (1997). "Translocation inhibitors define specificity of protein kinase C isoenzymes in pancreatic beta-cells." *J Biol Chem* 272(3): 1417-20.
- Yeung, T., P. C. Georges, et al. (2005). "Effects of substrate stiffness on cell morphology, cytoskeletal structure, and adhesion." *Cell Motil Cytoskeleton* 60(1): 24-34.
- Yip, S., K. S. Aboody, et al. (2003). "Neural stem cell biology may be well suited for improving brain tumor therapies." *Cancer J* 9(3): 189-204.

- Yonekura-Sakakibara, K., Y. Tanaka, et al. (2000). "Molecular and biochemical characterization of a novel hydroxycinnamoyl-CoA: anthocyanin 3-O-glucoside-6"-O-acyltransferase from *Perilla frutescens*." *Plant Cell Physiol* 41(4): 495-502.
- Yoo, G. H., J. Washington, et al. (2002). "The effects of exogenous p53 overexpression on HPV-immortalized and carcinogen transformed oral keratinocytes." *Cancer* 94(1): 159-66.
- Youssef, L. A., M. Schuyler, et al. (2007). "Histamine release from the basophils of control and asthmatic subjects and a comparison of gene expression between "releaser" and "nonreleaser" basophils." *J Immunol* 178(7): 4584-94.
- Youssef, L. A., B. S. Wilson, et al. (2002). "Proteasome-dependent regulation of Syk tyrosine kinase levels in human basophils." *J Allergy Clin Immunol* 110(3): 366-73.
- Yu, H., C. V. Nicchitta, et al. (1995). "Characterization of kinectin, a kinesin-binding protein: primary sequence and N-terminal topogenic signal analysis." *Mol Biol Cell* 6(2): 171-83.
- Yu, H., I. Toyoshima, et al. (1992). "Kinesin and cytoplasmic dynein binding to brain microsomes." *J Biol Chem* 267(28): 20457-64.
- Yu, J., D. A. Fischman, et al. (1973). "Selective solubilization of proteins and phospholipids from red blood cell membranes by nonionic detergents." *J Supramol Struct* 1(3): 233-48.
- Yu, J. Z., R. H. Dave, et al. (2009). "Cytosolic Galpha s acts as an intracellular messenger to increase microtubule dynamics and promote neurite outgrowth." *J Biol Chem*.
- Zaal, K. J., C. L. Smith, et al. (1999). "Golgi membranes are absorbed into and reemerge from the ER during mitosis." *Cell* 99(6): 589-601.
- Zalom, F. G., I. Werner, et al. (2004). "Organophosphate dormant spray pest control efficacy, pesticide concentration and toxicity in storm runoff." *Bull Environ Contam Toxicol* 73(2): 299-305.
- Zennami, K., Y. Yamada, et al. (2008). "Solitary brain metastasis from pT1, G3 bladder cancer." *Int J Urol* 15(1): 96-8.
- Zhang, D., J. Suen, et al. (2005). "Tetrameric mouse acetylcholinesterase: continuum diffusion rate calculations by solving the steady-state Smoluchowski equation using finite element methods." *Biophys J* 88(3): 1659-65.

- Zhang, J., K. Leiderman, et al. (2006). "Characterizing the topography of membrane receptors and signaling molecules from spatial patterns obtained using nanometer-scale electron-dense probes and electron microscopy." *Micron* 37(1): 14-34.
- Zhang, J., S. L. Steinberg, et al. (2008). "Markov random field modeling of the spatial distribution of proteins on cell membranes." *Bull Math Biol* 70(1): 297-321.
- Zhang, L. M., Z. P. Zhang, et al. (2007). "Measurement of D0-D0 mixing parameters in $D^0 \rightarrow K_S \pi^+ \pi^-$ decays." *Phys Rev Lett* 99(13): 131803.
- Zhang, R., J. Wiley, et al. (1989). "Rare clonal karyotypic variants in primary cultures of human breast carcinoma cells." *Cancer Res* 49(2): 444-9.
- Zhang, X., G. Jiang, et al. (2008). "Talin depletion reveals independence of initial cell spreading from integrin activation and traction." *Nat Cell Biol* 10(9): 1062-8.
- Zhang, X., P. B. Rosenthal, et al. (1999). "X-ray crystallographic determination of the structure of the influenza C virus haemagglutinin-esterase-fusion glycoprotein." *Acta Crystallogr D Biol Crystallogr* 55(Pt 5): 945-61.
- Zhang, Z. H., J. A. Johnson, et al. (1997). "C2 region-derived peptides of beta-protein kinase C regulate cardiac Ca^{2+} channels." *Circ Res* 80(5): 720-9.
- Zhang, Z. J., T. G. Berbos, et al. (1996). "Loss of nucleus basalis magnocellularis, but not septal, cholinergic neurons correlates with passive avoidance impairment in rats treated with 192-saporin." *Neurosci Lett* 203(3): 214-8.
- Zhang, Z. J., D. A. Lappi, et al. (1998). "Selective lesion of the cholinergic basal forebrain causes a loss of cortical neuropeptide Y and somatostatin neurons." *Brain Res* 800(2): 198-206.
- Zhao, Y., I. Gaidarov, et al. (2007). "Phosphoinositide 3-kinase C2alpha links clathrin to microtubule-dependent movement." *J Biol Chem* 282(2): 1249-56.
- Zheng, J., S. M. Cahill, et al. (1996). "Identification of the binding site for acidic phospholipids on the pH domain of dynamin: implications for stimulation of GTPase activity." *J Mol Biol* 255(1): 14-21.
- Zheng, S. L., V. L. Stevens, et al. (2009). "Two independent prostate cancer risk-associated Loci at 11q13." *Cancer Epidemiol Biomarkers Prev* 18(6): 1815-20.
- Zhong, H., S. M. Wade, et al. (2003). "A spatial focusing model for G protein signals. Regulator of G protein signaling (RGS) protein-mediated kinetic scaffolding." *J Biol Chem* 278(9): 7278-84.

- Zhou, M., D. A. Horita, et al. (2002). "Solution structure and functional analysis of the cysteine-rich C1 domain of kinase suppressor of Ras (KSR)." *J Mol Biol* 315(3): 435-46.
- Zhu, D., W. C. Xiong, et al. (2006). "Lipid rafts serve as a signaling platform for nicotinic acetylcholine receptor clustering." *J Neurosci* 26(18): 4841-51.
- Zhu, D. M., M. L. Dustin, et al. (2007). "Analysis of two-dimensional dissociation constant of laterally mobile cell adhesion molecules." *Biophys J* 92(3): 1022-34.
- Zhu, Y., H. Liao, et al. (2001). "LDL-activated p38 in endothelial cells is mediated by Ras." *Arterioscler Thromb Vasc Biol* 21(7): 1159-64.
- Zou, X., M. Liu, et al. (2008). "Robustness analysis of EGFR signaling network with a multi-objective evolutionary algorithm." *Biosystems* 91(1): 245-61.
- Zundel, W., C. Schindler, et al. (2000). "Loss of PTEN facilitates HIF-1-mediated gene expression." *Genes Dev* 14(4): 391-6.
- Zundel, W., L. M. Swiersz, et al. (2000). "Caveolin 1-mediated regulation of receptor tyrosine kinase-associated phosphatidylinositol 3-kinase activity by ceramide." *Mol Cell Biol* 20(5): 1507-14.
- Zuo, L., M. Ushio-Fukai, et al. (2005). "Caveolin-1 is essential for activation of Rac1 and NAD(P)H oxidase after angiotensin II type 1 receptor stimulation in vascular smooth muscle cells: role in redox signaling and vascular hypertrophy." *Arterioscler Thromb Vasc Biol* 25(9): 1824-30.
- Zurzolo, C., M. P. Lisanti, et al. (1993). "Glycosylphosphatidylinositol-anchored proteins are preferentially targeted to the basolateral surface in Fischer rat thyroid epithelial cells." *J Cell Biol* 121(5): 1031-9.