

10-1-2014

# Cocaine Intoxication and Thyroid Storm: Similarity in Presentation and Implications for Treatment

Mary E. Lacy

Kristina M. Utzschneider

Follow this and additional works at: [https://digitalrepository.unm.edu/hostpitalmed\\_pubs](https://digitalrepository.unm.edu/hostpitalmed_pubs)

---

## Recommended Citation

Lacy, Mary E., Utzschneider, Kristina M. Journal of Investigative Medicine High Impact Case Reports. October-December 2014: 1-5.  
doi:10.1177/23247096145548

This Article is brought to you for free and open access by the Internal Medicine at UNM Digital Repository. It has been accepted for inclusion in Hospital Medicine by an authorized administrator of UNM Digital Repository. For more information, please contact [disc@unm.edu](mailto:disc@unm.edu).

# Cocaine Intoxication and Thyroid Storm: Similarity in Presentation and Implications for Treatment

Journal of Investigative Medicine High  
Impact Case Reports  
October-December 2014: 1–5  
© 2014 American Federation for  
Medical Research  
DOI: 10.1177/2324709614554836  
hic.sagepub.com



Mary E. Lacy, MD<sup>1</sup>, and Kristina M. Utzschneider, MD<sup>1,2</sup>

## Abstract

**Introduction.** Cocaine, a widely used sympathomimetic drug, causes thermoregulatory and cardiac manifestations that can mimic a life-threatening thyroid storm. **Case.** A man presented to the emergency department requesting only cocaine detoxification. He reported symptoms over the last few years including weight loss and diarrhea, which he attributed to ongoing cocaine use. On presentation he had an elevated temperature of 39.4°C and a heart rate up to 130 beats per minute. Examination revealed the presence of an enlarged, nontender goiter with bilateral continuous bruits. He was found to have thyrotoxicosis by labs and was treated for thyroid storm and cocaine intoxication concurrently. The patient was ultimately diagnosed with Graves' disease and treated with iodine-131 therapy. **Conclusion.** Cocaine use should be considered a possible trigger for thyroid storm. Recognition of thyroid storm is critical because of the necessity for targeted therapy and the significant mortality associated with the condition if left untreated.

## Keywords

thyrotoxicosis, Graves' disease, sympathetic nervous system, clinical decision making

Thyroid storm is a diagnosis that typically involves an underlying thyrotoxicosis along with a precipitant, such as infection or surgery, with resultant hyperpyrexia and cardiovascular and central nervous system dysfunction.<sup>1</sup> Cocaine induces a vasoconstrictive and sympathomimetic response<sup>2</sup> that can clinically mimic thyroid storm. We report the case of a man who presented to the emergency department requesting cocaine detoxification and was found to have concurrent thyroid storm.

## Case Presentation

Mr S presented to a local emergency department requesting cocaine detoxification. He reported a cocaine binge that ended the night prior to admission and regular cocaine use over the past year. He noted a constellation of symptoms, all of which he attributed to cocaine use, including weight loss, diarrhea, mental "fogginess," diaphoresis, tremor, and palpitations. Past medical history was significant for type 2 diabetes treated with metformin and hypertension treated with lisinopril. His family history was notable for type 1 diabetes in his mother and alcoholism in his father. He had no family history of thyroid abnormalities.

On evaluation, his vital signs were the following: height 170 cm (67 in.), weight 56 kg (124 pounds), rectal temperature of 39.4°C (102.9°F), respiratory rate of 22 breaths per minute, blood pressure of 146/51 mm Hg, and heart rates up

to 130 beats per minute. His exam was notable for cachexia, normally sized and reactive pupils, absence of exophthalmos, a diffusely enlarged and nontender thyroid with bilateral continuous bruits, tachycardia with a systolic flow murmur, clear lungs, an intention tremor, and global hyperreflexia. He had significant inattention and distractibility requiring frequent redirection, though he was fully oriented. Laboratory results revealed a thyroid-stimulating hormone (TSH) of <0.01  $\mu$ U/mL (normal range = 0.27–4.20), a free T4 of 7.77 ng/dL (0.93–1.70), and a total T3 of 651 ng/dL (85–184). A urine toxicology screen was positive for cocaine and an infectious workup was negative. Chart review confirmed a 9-kg (20-pound) weight loss over the previous 10 months.

The patient was treated concurrently for cocaine intoxication with benzodiazepines and for thyroid storm with hydrocortisone and propylthiouracil followed by saturated solution of potassium iodine. By the following day, his vital signs had normalized and he was notably less distractible. The patient

<sup>1</sup>University of Washington, Seattle, WA, USA

<sup>2</sup>VA Puget Sound Health Care System, Seattle, WA, USA


## Corresponding Author:

Kristina M. Utzschneider, VA Puget Sound Health Care System (151),  
1660 S. Columbian Way, Seattle, WA 98108, USA.  
Email: kutzschn@u.washington.edu

**Table 1.** Composite Vital Signs and Laboratory Values.

	Day 1	Day 2	Day 38
TSH (normal = 0.27-4.20 $\mu$ IU/mL)	<0.01		<0.01
Free T4 (normal = 0.93-1.70 ng/dL)	7.77	6.89	1.67
Total T3 (normal = 85-184 ng/dL)	651		251
Heart rate (beats per minute)	130	97	81
Temperature ( $^{\circ}$ C)	39.4	37.9	36.7
Weight (kg)	56	57	64

Abbreviations: TSH, thyroid-stimulating hormone; T3, triiodothyronine; T4, thyroxine.

**Figure 1.** Thyroid uptake and scan.

Thyroid scan reflecting uptake of  $^{99m}\text{Tc}$ -pertechnetate seen from anterior, anterior with markers, left anterior oblique, and right anterior oblique views (clockwise from top left). The upper pole of the right lobe contains an area of photopenia consistent with a cold nodule. He had increased uptake with 4-hour and 24-hour values of 45% (normal = 4% to 20%) and 60% (normal = 10% to 30%), respectively.

was discharged on methimazole and propranolol. He was referred for outpatient substance abuse treatment.

On follow-up in endocrine clinic, he had remained cocaine-free. He underwent a cool-off period on continued methimazole and propranolol. Laboratory studies revealed hyperthyroidism but the patient was no longer severely symptomatic and was gaining weight (full results are shown in Table 1). A thyroid uptake and scan 3 months after initial presentation revealed diffusely increased uptake (Figure 1) along with a focus of photopenia within the upper pole of the right lobe consistent with a cold nodule. He was treated with iodine-131 for Graves' disease immediately following the

uptake and scan. He missed his endocrinology appointments postablation and is currently lost to follow-up.

## Discussion

The patient presented with classic signs and symptoms of thyroid storm that could have been attributed solely to his recent cocaine use. A detailed physical exam revealed a symmetrically enlarged thyroid and bruit pathognomonic for Graves' disease.

This case raises several interesting topics including the similar signs and symptoms of acute cocaine intoxication and thyroid storm, the possible interactions between cocaine and the thyroid, and the optimal treatment of thyroid storm in the setting of acute cocaine ingestion.

## Diagnosis of Thyroid Storm

The diagnosis of thyroid storm is clinical, a combination of patient presentation and laboratory values confirming hyperthyroidism. The degree to which the thyroid hormone levels are elevated does not determine the presence or absence of thyroid storm. Brooks et al showed that serum triiodothyronine (T3) and thyroxine (T4) were similar in patients with uncomplicated thyrotoxicosis and thyroid storm.<sup>3,4</sup> While patients with thyroid storm did have higher free T4 levels, this laboratory value alone does not establish the presence of life-threatening thyroid storm.<sup>4</sup> Burch and Wartofsky defined diagnostic criteria for diagnosis of thyroid storm in 1993 to aid clinicians in making the diagnosis and decisions on treatment (see Table 2).<sup>1</sup>

Cocaine intoxication and thyroid storm lead to clinical pictures with overlapping signs, including hyperpyrexia, tachycardia, and central nervous system disturbances. Because even with early diagnosis and targeted treatment the mortality rate of thyroid storm ranges from 10% to 30%, clinicians should consider assessing for coexisting thyrotoxic state in patients presenting with presumed severe cocaine intoxication.<sup>5</sup> In the case of Mr S, his score based on diagnostic criteria for thyroid storm (Table 2) was high at 70, and a detailed physical exam revealed a pathologically enlarged thyroid suggesting the need for further laboratory testing, ultimately leading to accurate diagnosis and appropriate treatment.

## Physiology of Thyroid Storm

The mechanisms for the hyperadrenergic state seen in thyroid storm are complex and vary by organ system.<sup>6,7</sup> Thyrotoxic states have been associated with normal or diminished plasma and urine norepinephrine levels, suggesting the mechanism is not explained by primary adrenergic activation.<sup>6</sup> Studies have also shown changes in quantity and affinity of  $\beta$ -adrenergic receptors and, perhaps more important, an augmentation in postreceptor signaling cascades ultimately leading to a state of adrenergic hyperresponsiveness.<sup>6,7</sup>

**Table 2.** Diagnostic Criteria for Thyroid Storm<sup>a</sup>.

Thermoregulatory dysfunction (pyrexia); temperature in degrees Celsius		Cardiovascular dysfunction (tachycardia); heart rate in beats per minute	
37.2-37.7	5	90-109	5
37.8-38.2	10	110-119	10
38.3-38.8	15	120-129	15
38.9-39.3	20	130-139	20
39.4-39.9	25	≥140	25
≥40.0	30	Cardiovascular dysfunction (heart failure)	
Central nervous system effects		Absent	0
Absent	0	Mild (pedal edema)	5
Mild (agitation)	10	Moderate (bibasilar rales)	10
Moderate (delirium, psychosis, extreme lethargy)	20	Severe (pulmonary edema)	15
Severe (seizure, coma)	30	Cardiovascular dysfunction (atrial fibrillation)	
Gastrointestinal-hepatic dysfunction		Absent	0
Absent	0	Present	10
Moderate (diarrhea, nausea/vomiting, abdominal pain)	10	Precipitant history	
Severe (unexplained jaundice)	20	Negative	0
		Positive	10

<sup>a</sup>Adapted from Burch and Wartofsky.<sup>1</sup> Points from each category are added together. Points are calculated to attribute symptoms to thyrotoxicosis when it is not possible to determine true cause of symptoms. A score less than 25 is unlikely to indicate thyroid storm, a score of greater 45 is highly indicative, and a score of 25 to 44 is intermediate.

Common precipitants of thyroid storm include infection, surgery, trauma, and radioactive iodine. There are several theories as to how triggers could precipitate thyroid storm. These include sudden decreased affinity of thyroid hormone binding proteins leading to higher levels of active hormone, increased thyroid hormone uptake into cells, and transient loss of central sympathetic nervous system downregulation in the setting of adrenergic hyperresponsiveness.<sup>5,6,8</sup>

## Cocaine and the Thyroid Axis

Cocaine inhibits presynaptic reuptake of biogenic amines.<sup>2</sup> It is the inhibition of dopamine reuptake that is thought to explain the euphoric and addictive properties of cocaine.<sup>2</sup> The additional inhibition of norepinephrine reuptake induced by cocaine explains the sympathetic overdrive that is observed in cocaine intoxication. It is possible that this stimulant effect of cocaine intoxication in a patient with the adrenergic hyperresponsiveness associated with thyrotoxicosis could trigger thyroid storm, as exemplified by our patient.

The interplay of cocaine use and thyroid function is complex and poorly understood. Cocaine- and amphetamine-regulated transcript (CART) peptides were first discovered by looking for brain transcripts that were increased following acute cocaine or methamphetamine exposure.<sup>9</sup> CART peptides have since been shown in rats to innervate hypothalamic neurons and exhibit a stimulatory effect on transcription of thyrotropin-releasing hormone (TRH).<sup>10</sup> Furthermore, several studies have documented a diminished thyroid-stimulating

hormone (TSH) response to TRH in people with concurrent cocaine abuse.<sup>11-13</sup> After 30 days of abstinence, however, TRH produced an increased TSH response.<sup>13</sup> A hypothesis for this interaction is that chronic cocaine use could result in a dopamine and norepinephrine-mediated increase in TRH levels that leads to downregulation of TRH receptors and decreased TSH.<sup>11</sup>

Despite these connections between thyroid hormone regulation and cocaine use, acute as well as repeated cocaine stimulation in rats was not shown to produce a change in serum T3 and T4 levels.<sup>14</sup> Thyroid function tests in 22 heavy cocaine users studied prior to admission to a treatment center showed no statistically significant difference when compared to a control population.<sup>15</sup> Conversely, in a retrospective study that looked specifically at patients with diagnosed hyperthyroidism (90% subclinical), there was an increased incidence of cocaine use.<sup>16</sup> It was theorized that an enhanced responsiveness to cocaine's stimulatory effects could contribute to increased use of the drug.<sup>16</sup>

## Treatment

Benzodiazepines are the mainstay of treatment for cocaine intoxication and have beneficial effects on cocaine-induced chest pain as well as central nervous system disturbances.<sup>2</sup> The treatment of thyroid storm requires a multistep approach with identification and treatment of any precipitant, as well as treatment aimed at decreasing thyroid hormone production, thyroid hormone release, and the peripheral effects of existing thyroid hormone.<sup>17</sup>

Perhaps the most effective immediate treatment for thyroid storm is blockage of the adrenergic effects created by excess thyroid hormone. This is most often accomplished with high-dose propranolol, a nonselective  $\beta$ -blocker, which also functions to block some peripheral conversion of T4 to T3.<sup>5</sup> While the frequently cited contraindication of  $\beta$ -blockade in a patient with cocaine use has been reasonably called into question,<sup>18,19</sup> it was propranolol that was originally shown to increase cocaine-associated coronary vasoconstriction.<sup>20</sup> Interestingly, there are cases of increased  $\alpha$ -adrenergic effects following labetalol administration in cases of pheochromocytoma, raising concerns even about the use of  $\beta$ -blockers with  $\alpha$ -blocking action in extreme hyperadrenergic states.<sup>21</sup> To the best of our knowledge, there exists no data on the safety of the use of  $\beta$ -blockade in patients presenting with a mixed picture of thyroid storm and cocaine intoxication to guide clinical decision making regarding use of selective or nonselective  $\beta$ -blockers.

Thionamides are the preferred agents to block new thyroid hormone production and act via inhibition of thyroid peroxidase. While propylthiouracil (PTU) has less convenient dosing requirements and a higher toxicity profile compared to methimazole, PTU is often preferred in the acute management of thyroid storm because it additionally decreases peripheral conversion of T4 to T3 by as much as 20% to 30% in the acute setting.<sup>22</sup> Iodine administration acutely acts to block organification of iodide and synthesis and release of thyroid hormone via the Wolff Chaikoff effect. Iodine should typically be administered 30 to 60 minutes after thionamide therapy has been started to prevent stimulation of new hormone synthesis. Hydrocortisone administration is also recommended by current guidelines and treats possible relative adrenal insufficiency while also decreasing T4 to T3 conversion.<sup>23</sup>

## Conclusion

The case of Mr S highlights the importance of developing a broad differential diagnosis to prevent premature closure. Our patient presented with a reported cocaine binge and displayed many signs and symptoms that could be directly explained by cocaine intoxication alone. Other diagnoses to consider in a patient with substance abuse, tachycardia, and fever include infectious etiologies, such as pneumonia or endocarditis, or pulmonary embolus. A thorough history and physical examination prompted consideration of thyrotoxicosis as an alternative explanation for his historical weight loss and diarrhea as well as his current fever, tachycardia, and mental status changes. The high morbidity and mortality associated with thyroid storm necessitates that clinicians consider this diagnosis in the differential. Mr S was promptly diagnosed and treated for thyroid storm as well as cocaine intoxication and ultimately received more definitive treatment for his Graves' disease.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Department of Veteran Affairs.

## References

1. Burch HB, Wartofsky L. Life-threatening thyrotoxicosis. Thyroid storm. *Endocrinol Metab Clin North Am*. 1993;22:263-277.
2. Zimmerman JL. Cocaine intoxication. *Crit Care Clin*. 2012;28:517-526.
3. Brooks MH, Waldstein SS, Bronsky D, Sterling K. Serum triiodothyronine concentration in thyroid storm. *J Clin Endocrinol Metab*. 1975;40:339-341.
4. Brooks MH, Waldstein SS. Free thyroxine concentrations in thyroid storm. *Ann Intern Med*. 1980;93:694-697.
5. Nayak B, Burman K. Thyrotoxicosis and thyroid storm. *Endocrinol Metab Clin North Am*. 2006;35:663-686.
6. Silva JE, Bianco SD. Thyroid-adrenergic interactions: physiological and clinical implications. *Thyroid*. 2008;18:157-165.
7. Bilezikian JP, Loeb JN. The influence of hyperthyroidism and hypothyroidism on alpha- and beta-adrenergic receptor systems and adrenergic responsiveness. *Endocr Rev*. 1983;4:378-388.
8. Tietgens ST, Leinung MC. Thyroid storm. *Med Clin North Am*. 1995;79:169-184.
9. Douglass J, McKinzie AA, Couceyro P. PCR differential display identifies a rat brain mRNA that is transcriptionally regulated by cocaine and amphetamine. *J Neurosci*. 1995;15(3 pt 2):2471-2481.
10. Fekete C, Mihály E, Luo LG, et al. Association of cocaine- and amphetamine-regulated transcript-immunoreactive elements with thyrotropin-releasing hormone-synthesizing neurons in the hypothalamic paraventricular nucleus and its role in the regulation of the hypothalamic-pituitary-thyroid axis during fasting. *J Neurosci*. 2000;20:9224-9234.
11. Dackis CA, Gold MS. New concepts in cocaine addiction: the dopamine depletion hypothesis. *Neurosci Biobehav Rev*. 1985;9:469-477.
12. Giannini AJ, Malone DA, Loiselle RH, Price WA. Blunting of TSH response to TRH in chronic cocaine and phencyclidine abusers. *J Clin Psychiatry*. 1987;48:25-26.
13. Vescovi PP, Pezzarossa A. Thyrotropin-releasing hormone-induced GH release after cocaine withdrawal in cocaine addicts. *Neuropeptides*. 1999;33:522-525.
14. Budziszewska B, Jaworska-Feil L, Lasoń W. The effect of repeated amphetamine and cocaine administration on adrenal, gonadal and thyroid hormone levels in the rat plasma. *Exp Clin Endocrinol Diabetes*. 1996;104:334-338.
15. Dhopes VP, Burke WM, Maany I, Ravi NV. Effect of cocaine on thyroid functions. *Am J Drug Alcohol Abuse*. 1991;17:423-427.
16. Vasudevan VN, Black RD, Bowman B, Solomon SS. Does subclinical hyperthyroidism predispose to cocaine addiction? *Endocr Pract*. 2012;18:478-482.

17. Chiha M, Samarasinghe S, Kabaker AS. Thyroid storm: an updated review [published online August 5, 2013]. *J Intensive Care Med*. doi:10.1177/0885066613498053.
18. Dattilo PB, Hailpern SM, Fearon K, Sohal D, Nordin C. Beta-blockers are associated with reduced risk of myocardial infarction after cocaine use. *Ann Emerg Med*. 2008;51:117-125.
19. Rangel C, Shu RG, Lazar LD, Vittinghoff E, Hsue PY, Marcus GM. Beta-blockers for chest pain associated with recent cocaine use. *Arch Intern Med*. 2010;170:874-879.
20. Lange RA, Cigarroa RG, Flores ED, et al. Potentiation of cocaine-induced coronary vasoconstriction by beta-adrenergic blockade. *Ann Intern Med*. 1990;112:897-903.
21. Briggs RS, Birtwell AJ, Pohl JE. Hypertensive response to labetalol in phaeochromocytoma. *Lancet*. 1978;1:1045-1046.
22. Cooper DS, Saxe VC, Meskell M, Maloof F, Ridgway EC. Acute effects of propylthiouracil (PTU) on thyroidal iodide organification and peripheral iodothyronine deiodination: correlation with serum PTU levels measured by radio-immunoassay. *J Clin Endocrinol Metab*. 1982;54:101-107.
23. Bahn Chair RS, Burch HB, Cooper DS, et al. Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists. *Thyroid*. 2011;21:593-646.